

Poster presentation

Open Access

## Preliminary validation of the Juvenile Arthritis Multidimensional Assessment Report (JAMAR) in 403 clinic patients

N Solari\*<sup>1</sup>, G Filocamo<sup>1</sup>, B Schiappapietra<sup>1</sup>, A Consolaro<sup>1</sup>, S Magni-Manzoni<sup>2</sup>, S Viola<sup>1</sup>, N Ruperto<sup>1</sup>, C Saad-Magalhaes<sup>1</sup>, D Tani<sup>1</sup>, S Serpico<sup>1</sup>, A Martini<sup>1</sup> and A Ravelli<sup>1</sup>

Address: <sup>1</sup>IRCCS G Gaslini, Genova, Italy and <sup>2</sup>IRCCS Policlinico S Matteo, Pavia, Italy

\* Corresponding author

from 15<sup>th</sup> Paediatric Rheumatology European Society (PreS) Congress  
London, UK. 14–17 September 2008

Published: 15 September 2008

*Pediatric Rheumatology* 2008, **6**(Suppl 1):P106 doi:10.1186/1546-0096-6-S1-P106

This abstract is available from: <http://www.ped-rheum.com/content/6/S1/P106>

© 2008 Solari et al; licensee BioMed Central Ltd.

### Objective

To provide preliminary evidence of validity of the JAMAR in a large sample of JIA patients.

### Methods

The JAMAR includes: Juvenile Arthritis Functionality Scale (JAFS) (0 = normal; 30 = worst), Pediatric Rheumatology Quality of Life (PRQL) questionnaire (0 = best; 30 = worst); visual analogue scales (0 = best; 10 = worst) for parent/patient rating of well-being (WB), pain (P), disease activity (DA); parent/patient assessment of morning stiffness, disease status, satisfaction about disease outcome.

### Results

The JAMAR was completed by 403 parents in 696 visits between March 2007 and February 2008. All completers reported that JAMAR was easy to understand and fill. Completion time was < 10 minutes. Results of quantita-

tive JAMAR assessments are shown in the table 1 together with assessments made by attending physician.

### Conclusion

The JAMAR proved to be feasible and to have good face and content validity. Parents and physicians revealed fair concordance in assessing overall disease activity. Around half and 2/3 of patients, respectively, were judged to be in remission or in satisfactory state by parents.

**Table 1: Percentage of parents reporting morning stiffness, remission, continued activity, flare, and satisfaction about outcome was 34.3, 47.9, 29.5, 22.6, and 72.1, respectively.**

	Parent WB	Parent DA	Parent P	JAFS	PRQL	MD global	No. swollen joints	No. tender joints	No. restricted joints	No. active joints
Mean	2.3	2.4	2.1	1.5	3.9	2.3	1.8	2.1	2.1	2.4
Median	1	1	0.5	0	3	0.5	1	0	0	1
Minimum	0	0	0	0	0	0	0	0	0	0
Maximum	10	10	10	26	23	10	35	49	46	49