


How should I manage immunosuppression in a kidney transplant patient with COVID-19? An ERA-EDTA DESCARTES expert opinion

Umberto Maggiore ¹, Daniel Abramowicz², Marta Crespo³, Christophe Mariat⁴, Geir Mjoen⁵, Licia Peruzzi⁶, Mehmet Sükrü Sever⁷, Gabriel C. Oniscu⁸, Luuk Hilbrands ⁹, and Bruno Watschinger¹⁰; on behalf of the DESCARTES Working Group of the ERA-EDTA

¹Dipartimento di Medicina e Chirurgia, Università di Parma, UO Nefrologia, Azienda Ospedaliero-Universitaria di Parma, Parma, Italy, ²Department of Nephrology, Antwerp University Hospital, Antwerp University, Antwerp, Belgium, ³Department of Nephrology, Hospital del Mar Barcelona, Barcelona, Spain, ⁴Department of Nephrology, Dialysis, and Renal Transplantation, University North Hospital, Saint Etienne, France, ⁵Department of Transplant Medicine, Oslo University Hospital, Oslo, Norway, ⁶Pediatric Nephrology Unit, Regina Margherita Children's Hospital, Turin, Italy, ⁷Division of Nephrology, Department of Internal Medicine, Istanbul Faculty of Medicine, Istanbul University, Istanbul, Turkey, ⁸Transplant Unit, Royal Infirmary of Edinburgh, Edinburgh, UK, ⁹Department of Nephrology, Radboud University Medical Center, Nijmegen, The Netherlands and ¹⁰Department of Nephrology, Medical University of Vienna, Vienna, Austria

Correspondence to: Umberto Maggiore; E-mail: umberto.maggiore@unipr.it; Twitter handle: @UMaggiore, @ERAEDTA, @gabriel_oniscu, @MCrespoBarrio



INTRODUCTION

After severe acute respiratory syndrome coronavirus (SARS-CoV) in 2003 and Middle East respiratory syndrome-related coronavirus (MERS-CoV) in 2012, the world is now facing a third rapidly spreading coronavirus outbreak, caused by SARS-CoV-2 [1–3]. Coronavirus disease 2019 (COVID-19) was declared a global pandemic by the World Health Organization [4] and has already led to >3 million reported cases and >230 000 deaths worldwide [5].

Viral infections pose an important risk of morbidity and mortality in patients after organ transplantation [6]. Indeed, immunosuppression, which is crucial for preventing alloimmune reactions, impairs host defense mechanisms. Respiratory virus infections, such as respiratory syncytial virus, may progress more rapidly to pneumonia in organ transplant recipients, with a tendency to more severe disease and prolonged shedding of potentially infectious virus. Immunosuppression also increases the severity of both adeno- and influenza viruses (reviewed in Razonable [7] and Manuel *et al.* [8]).

With regard to COVID-19, all kidney transplant recipients (KTRs) must strictly follow the general hygiene measures recommended for the general population, including frequent hand

washing, social distancing measures and wearing of masks in crowded places [9]. COVID-19 induces variable clinical courses in normal hosts but seems to progress more rapidly in immunocompromised hosts, with greater rates of intensive care unit admissions and death [10]. The largest series reported to date from Europe and the USA are shown in Table 1 [11–17]. Although there might be a bias towards selection of the most severe cases (apart from one study in which only a minority of patients had severe disease [17]), the mortality rates were between 23% and 28%, which is close to the rate of 21% reported from a cohort of 5700 hospitalized patients in New York [18], but certainly higher than the ≤5% usually reported for COVID-19-infected patients [11]. These articles describe various strategies for the management of immunosuppressive therapy, based on a stepwise reduction of immunosuppression according to the severity of the disease. Due to the heterogeneity of the reported strategies, it is of course impossible to draw generalizable conclusions about the best practice to follow.

A European initiative, promoted by ERA-EDTA and the DESCARTES working group (WG) has recently started and is aiming to rapidly collect data about treatments and outcomes of COVID-19 disease in KTRs [9]. In the meantime, how to deal with immunosuppression among KTRs is left to clinical judgement and common sense, taking into consideration the risk of a serious, potentially fatal disease along with the risk of acute rejection and possibly graft loss. Interestingly, none of the series has reported acute rejection and graft loss as a consequence of immunosuppression reduction (Table 1), but this

Table 1. Largest case series reported of kidney transplant recipients with COVID-19

Study	Location	Time frame (2020)	No. of KT ^a s	Hospitalized (%)	Immunosuppressive strategy	High-dose steroids (%)	Anti-IL-6R Ab; other monoclonal Abs	Follow-up days	Mortality (%)	Rejection or graft failure
Akalin <i>et al.</i> [11]	Montefiore, USA	16 March–1 April	36	78	Withdrawal of MPA/AZA, Tac withheld in severely ill patients	2	2% tocilizumab; 21% leronlimab	21 (14–28)	28	Not reported
Pereira <i>et al.</i> [12] ^a	Columbia University, USA	13 March–3 April	46	76	Moderately decrease the overall amount of immunosuppression with a particular emphasis on decreasing or stopping MPA/AZA	24	21% tocilizumab	20 (14–24)	23	Not reported
Columbia University KT program [13]	Columbia University, USA	Up to 27 March	15	100	Stop MPA/AZA while continuing tacrolimus (4–7 ng/mL) and prednisone	7	7% tocilizumab	7 (3–11)	Incomplete follow-up	Not reported
Fernández-Ruiz <i>et al.</i> [14] ^a	Madrid, Spain	5 March–23 March	8	100	Stop CNI and mTORi upon initiation of LPV/r (given in 50% of the pts) Target Tac 5–10 ng/mL Steroids reduced by 50% MPA/AZA decreased	11	6% tocilizumab	18 (14–28)	28	Not reported
Alberici <i>et al.</i> [15]	Brescia, Italy	Up to 24 March	20	100	Stop all immunosuppressive treatment LPV/r, DRV/r given in 95% of the pts Increased dose of steroids	55	30% tocilizumab	Median follow-up 7 days N.A.	25	Not reported
Banerjee <i>et al.</i> [16]	London, UK	1 March–31 March	7	71	CNI stopped in ventilated patients	0	0%	N.A.	Incomplete follow-up	Not reported
Lubetzky <i>et al.</i> [17]	WCM, USA	13 March–20 April	54	72	MPA stopped (61%) in hospitalized patients Tacrolimus reduced (46%) in hospitalized patients	9	4%	21 (5–43)	13	Not reported

Follow-up (days) is reported as median (range) unless otherwise specified.

^aApart from the number of KT^as, reported data from Pereira *et al.* [12] refer to 90 solid organ transplants combined and from Fernández-Ruiz *et al.* [14] to 18 solid organ transplants combined.

KT, kidney transplantation; Ab, antibody; LPV/r, lopinavir/ritonavir; DRV/r, darunavir/ritonavir; MPA, mycophenolate sodium or mofetil; AZA, azathioprine; tocilizumab, anti-IL-6 mAb; leronlimab, CCR5 antagonist; N.A., not available.

Table 2. Management of immunosuppression in patients who are beyond 3–6 months after transplantation

1. Asymptomatic patients: no knowledge of COVID-19 status (ambulatory, stable patients)	
No pre-emptive/proactive change of immunosuppressive medications	
2. Asymptomatic patients, swab pos for COVID-19	
If it is a high-risk patient: age ≥ 70 years, or comorbidities or risk factors (diabetes, cardiac or pulmonary disease, heavy smoking, BMI >30 kg/m ² , eGFR <30 mL/min/1.73 m ² , lymphocyte depletion therapy within previous 3–6 months): consider reducing/stopping AZA/MPA/mTORi if on triple therapy	
3. Mild disease: the patient is alert, has only mild upper respiratory and/or gastrointestinal symptoms, temperature $<38^{\circ}\text{C}$ and does not refer symptoms suggestive of COVID-19 pneumonia such as dyspnoea, persistent chest pain and intensive cough; if available, oxygen saturation in room air is $>95\%$, respiratory rate <25/min; no evidence of pneumonia on either chest X-ray or CT; no need for hospitalization	
If patient is on:	
Triple therapy	Stop MPA/AZA/mTORi Maintain CNI + steroids
Dual therapy (including steroids)	Continue dual therapy
Dual therapy (steroid-free)	
CNI + MPA	Consider replacing MPA with low-dose steroids
CNI + mTORi	Consider replacing mTORi with low-dose steroids
MPA + mTORi	Consider replacing MPA or mTORi with low-dose steroids
<ul style="list-style-type: none"> • Consider CNI dose reduction (to the lower bound of the therapeutic range according to the immunological risk) if there is no clear improvement over the first 3–5 days • Cautiously restart previous immunosuppression 3–7 days after symptoms have cleared 	
4. Evidence of mild COVID-19 pneumonia: oxygen saturation 94–95% in room air; respiratory rate 25–29/min; or suspect lesions on chest X-ray or CT scan	
a. High-risk patient: age ≥ 70 years, or comorbidities or risk factors (diabetes, cardiac or pulmonary disease, heavy smoking, BMI >30 kg/m², eGFR <30 mL/min/1.73 m², lymphocyte depletion therapy within previous 3–6 months)	
Stop MPA/AZA/mTORi, Stop CNI Increase (or start) steroids 15–25 mg/day	
<ul style="list-style-type: none"> • Cautiously restart previous immunosuppression (CNI first) 5–10 days after symptoms have cleared 	
b. No high-risk patient (as defined above)	
Stop MPA/AZA/mTORi Maintain on dual therapy CNI-steroids Reduce CNI trough levels to target CsA: 50 ± 15 ng/mL, Tac: 3 ± 1 ng/mL Continue steroids in maintenance dose	
<ul style="list-style-type: none"> • In patients starting antiretroviral treatment: stop CNI and monitor as detailed in the text • Cautiously restart previous immunosuppression 5–10 days after symptoms have cleared 	
5. More severe COVID-19 pneumonia: oxygen saturation $<94\%$ in room air, respiratory rate ≥ 30/min, unstable or deteriorating course or requiring non-invasive ventilation or transfer to the intensive care unit (with or without mechanical ventilation)	
Discontinue all immunosuppressive drugs Increase/start steroids at 15–25 mg/day (or higher according to local practice).	
<ul style="list-style-type: none"> • Consider continuing with low-dose CNI in patients with higher risk of rejection (e.g. <1 year after transplantation and/or highly immunized) • Cautiously restart previous immunosuppression (CNI first) 5–15 days after symptoms have cleared 	

When available, risk stratification may additionally benefit from the results of lab parameters indicating severe inflammatory disease at risk of rapid progression, such as a high level of C-reactive protein, IL-6, ferritin and D-dimer.

MPA, mycophenolate mofetil, mycophenolic acid; AZA, azathioprine; CsA, cyclosporine; Tac, tacrolimus.

might be due to a too-short follow-up period. Furthermore, with KTRs amounting to only 0.1% of the general population, it is unlikely that evidence-based medicine will ever be produced for KTRs infected with COVID-19. Indeed, while >1000 studies about COVID-19 are registered in ClinicalTrials.gov (accessed 1 May 2020), none is devoted specifically to treatment of KTRs. While *in vitro* experiments suggest that coronavirus may require intact immunophilin pathways with a role for tacrolimus and cyclosporine to inhibit the growth of human coronaviruses [19, 20], the translation of these experimental findings in clinics remains to be seen. There is also the fear that complete withdrawal of immunosuppressive drugs may exacerbate the hyperinflammatory response that may occur in the late stages of COVID-19. After reading the expert opinions published by single centres (Table 1) and societies (French [21], Spanish [22], British [23], American [24]), and after extensive

discussions between its members, the DESCARTES WG formulated suggestions for COVID-19-infected KTRs who are beyond 3–6 months after kidney transplantation (Table 2).

MANAGEMENT OF DRUG-TO-DRUG INTERACTIONS, USE IN PATIENTS WITH RENAL FAILURE AND OTHER ISSUES RELEVANT TO THE USE OF ANTIVIRAL AND ANTI-INFLAMMATORY TREATMENT IN KTRs

Treatment of COVID-19 is based on antiviral drugs that inhibit SARS-CoV-2 proliferation and on immunomodulatory drugs that inhibit the hyperinflammatory syndrome that may cause acute respiratory distress syndrome (ARDS) and life-threatening respiratory failure [25, 26]. In theory, antiviral drugs could be effective in restraining viral infection when given in the early phase of the infection, whereas immunomodulatory

drugs may show the strongest benefit when administered during the late hyperinflammatory phase of the disease. Herein we do not formulate any advice on the use of antiviral drugs, antibiotics or anti-inflammatory drugs in KTRs with COVID-19. For these issues, please refer to your infectiologist and local and/or national guidelines. Below, we discuss anti-COVID-19 therapies in the context of renal impairment and their potential effects on the exposure of concomitant immunosuppressive agents.

ANTIVIRAL DRUGS FOR COVID-19

Remdesivir

Remdesivir, which is administered intravenously, has a favourable clinical profile and no known drug-to-drug interactions (DDIs). However, the drug has not been studied in patients with kidney failure [estimated glomerular filtration rate (eGFR) <30 mL/min/1.73 m²], which has been an exclusion criterion so far [27].

Hydroxychloroquine

Although a much-cited source [28] indicates that there is a possible increase in the exposure of cyclosporine, tacrolimus and mammalian target of rapamycin inhibitors (mTORis) during simultaneous use of chloroquine/hydroxychloroquine, a thorough PubMed search retrieved only two cases of increased cyclosporine levels. As chloroquine/hydroxychloroquine, which have a long half-life, are given for only 5–7 days, it is preferable but not mandatory to follow the trough levels of cyclosporine, tacrolimus and mTORis in this setting. No dose adjustment is mandatory in patients with kidney failure (eGFR <30 mL/min/1.73 m²), although it is recommended to use with caution in this setting.

Lopinavir–ritonavir and darunavir–cobicistat

Complete withdrawal of calcineurin inhibitor (CNI) and mTORi should be considered in every patient undergoing treatment with ritonavir- or cobicistat-boosted antiviral drugs. In fact, tacrolimus should be reduced to 1/20–1/50 in patients receiving ritonavir (and to 1/10 in patients receiving cobicistat) of the initial dose just to maintain constant blood levels [29, 30]. Therefore the dose should be reduced even further (i.e. even <1/50 of the initial dose) if the therapeutic plan is to ‘reduce’ CNIs rather than stop them. If the treatment with ritonavir-/cobicistat-based drugs is planned for only 5–7 days, then CNIs (and mTORis) should be withdrawn altogether and restarted no earlier than 24–48 h after ritonavir/cobicistat discontinuation at low dose. For longer ritonavir/cobicistat treatments (e.g. 2 weeks [31]), tacrolimus blood levels should be monitored daily and oral tacrolimus 0.5 mg/day administered only after blood levels have dropped below the desired lower bound (e.g. <5 ng/mL). It should also be noted that the hepatic clearance of tacrolimus may be further reduced in the course of multiple organ failure, making tacrolimus dose adjustments even more challenging. DDIs are milder for cyclosporine compared with tacrolimus, the dose reduction usually being 1/5 to maintain constant blood levels. No dose adjustment of lopinavir–

ritonavir, darunavir–cobicistat is needed for renal failure. Nonetheless, it should be mentioned that a randomized trial on the use of lopinavir–ritonavir failed to show a clear benefit [31]; therefore, given the high risk of the DDIs in this category of patients, we recommend against their routine use in KTRs.

ANTI-INFLAMMATORY DRUGS FOR COVID-19

Besides counteracting severe inflammation causing inflammatory lung injury and thrombotic complications in COVID-19, anti-inflammatory drugs in transplant recipients may have the additional benefit of protecting against rejection in patients who have withdrawn CNIs because of severe disease. However, they should be used with caution since suppression of inflammatory responses in the absence of effective antiviral therapy may cause uncontrolled infection [10].

High-dose steroids

A randomized controlled trial is testing the safety/efficacy of steroids (NCT04273321). Until results are available, broad use of high-dose steroids is discouraged [32]. However, for the treatment of a hyperinflammatory state causing abrupt respiratory compromise, and in the absence of viable alternatives, their use may be considered on a case-by-case basis [33–35], even in KTRs.

Tocilizumab and other anti-interleukin (IL)-6/IL-6R monoclonal antibodies (e.g. sarilumab)

After the preliminary successful experience of its use in China [36], and pending results from various clinical trials, tocilizumab is currently the most popular treatment used to counteract hyperinflammatory syndrome causing impending or ongoing respiratory compromise. In this severe clinical setting, usually transplant recipients had already withdrawn mycophenolate/azathioprine/mTORi along with CNIs before they were considered for tocilizumab administration. Tocilizumab should be used with caution if the white blood cell count is <1000/mm³. Its elimination is not influenced by renal dysfunction.

Intravenous immunoglobulins

High-dose intravenous immunoglobulins have been proposed for the use in patients with COVID-19 and deteriorating conditions to counteract inflammation and endothelial activation [37].

INVESTIGATIONAL TREATMENT STRATEGIES

At the time of writing, clinical trials are ongoing for the evaluation of other promising antiviral drugs and anti-inflammatory drugs such as favipiravir [25, 26], umifenovir (arbidol) [25, 26, nelfinavir [25], ivermectin [38], baricitinib [a Janus kinase (JAK) inhibitor] [38], anakinra (an IL-1 receptor antagonist) [39] and colchicine [40]. Favipiravir and arbidol do not need to be adjusted for renal function. CYP3A4 is the major isoform involved in arbidol metabolism, therefore there may be potential

interaction, especially in patients taking cyclosporine and mTORi. In contrast, favipiravir does not have DDIs with CNIs and mTORi [26]. Baricitinib [25] is contraindicated with kidney failure (eGFR <30 mL/min/1.73 m²) and may strongly increase the risk of infections in association with CNIs. Anakinra does not have DDIs with CNIs and mTORis and should be given every other day in patients with kidney failure (eGFR <30 mL/min/1.73 m²). Colchicine is better avoided while the patient is receiving concurrent treatment with ritonavir–cobicistat-based antiviral therapy and its dose should be halved in patients with kidney failure (eGFR <30 mL/min/1.73 m²). DDIs with tacrolimus are likely to be negligible, whereas concomitant administration of cyclosporine may warrant monitoring.

Convalescent plasma therapy and hyperimmune immunoglobulins

Convalescent plasma therapy and hyperimmune immunoglobulins, namely the transfer of passive immunity from convalescent human plasma, offer the most promising novel therapeutic approaches for the treatment of COVID-19 [41, 42].

CONFLICT OF INTEREST STATEMENT

None declared.

REFERENCES

1. Tsang KW, Ho PL, Ooi GC *et al.* A cluster of cases of severe acute respiratory syndrome in Hong Kong. *N Engl J Med* 2003; 348: 1977–1985
2. Zaki AM, van Boheemen S, Bestebroer TM *et al.* Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia. *N Engl J Med* 2012; 367: 1814–1820
3. Zhu N, Zhang D, Wang W *et al.* A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med* 2020; 382: 727–733
4. Cucinotta D, Vanelli M. WHO declares COVID-19 a pandemic. *Acta Biomed* 2020; 91: 157–160
5. Johns Hopkins University. Corona Virus Resource Center. <https://coronavirus.jhu.edu/> (1 May 2020, date last accessed)
6. Patel R, Paya CV. Infections in solid-organ transplant recipients. *Clin Microbiol Rev* 1997; 10: 86–124
7. Razonable RR. Management of viral infections in solid organ transplant recipients. *Expert Rev Anti Infect Ther* 2011; 9: 685–700
8. Manuel O, Lopez-Medrano F, Keiser L *et al.* Influenza and other respiratory virus infections in solid organ transplant recipients. *Clin Microbiol Infect* 2014; 20: 102–108
9. COVID-19 News and Information. 2020. <https://www.era-edta.org/en/covid-19-news-and-information/#toggle-id-6> (1 May 2020, date last accessed)
10. Fishman JA, Grossi PA. Novel coronavirus-19 (COVID-19) in the immunocompromised transplant recipient: #Flatteningthecurve. *Am J Transplant* 2020; doi: 10.1111/ajt.15890
11. Akalin E, Azzi Y, Bartash R *et al.* Covid-19 and kidney transplantation. *N Engl J Med* 2020; doi: 10.1056/NEJMc2011117
12. Pereira MR, Mohan S, Cohen DJ *et al.* COVID-19 in solid organ transplant recipients: initial report from the US epicenter. *Am J Transplant* 2020; doi: 10.1111/ajt.15941
13. Columbia University Kidney Transplant Program. Early description of coronavirus 2019 disease in kidney transplant recipients in New York. *J Am Soc Nephrol* 2020; doi: 10.1681/ASN.2020030375
14. Fernandez-Ruiz M, Andres A, Loinaz C *et al.* COVID-19 in solid organ transplant recipients: a single-center case series from Spain. *Am J Transplant* 2020; doi: 10.1111/ajt.15929
15. Alberici FD, Delbarba E, Manenti C *et al.* A single center observational study of the clinical characteristics and short-term outcome of 20 kidney

- transplant patients admitted for SARS-CoV2 pneumonia. *Kidney Int* 2020; doi: 10.1016/j.kint.2020.04.002
16. Banerjee DP, Shah S, Ster IC *et al.* COVID-19 infection in kidney transplant recipients. *Kidney Int* 2020; doi: 10.1016/j.kint.2020.03.018
17. Lubetzky M, Aull M, Shapiro-Craig R *et al.* Kidney allograft recipients diagnosed with coronavirus disease-2019: a single center report. *Nephrol Dial Transplant* 2020. In press.
18. Richardson S, Hirsch JS, Narasimhan M *et al.* Presenting characteristics, comorbidities, and outcomes among 5700 patients hospitalized with COVID-19 in the New York City area. *JAMA* 2020; doi: 10.1001/jama.2020.6775
19. Carbajo-Lozoya J, Muller MA, Kallies S *et al.* Replication of human coronaviruses SARS-CoV, HCoV-NL63 and HCoV-229E is inhibited by the drug FK506. *Virus Res* 2012; 165: 112–117
20. Ma-Lauer Y, Zheng Y, Malesevic M *et al.* Influences of cyclosporin A and non-immunosuppressive derivatives on cellular cyclophilins and viral nucleocapsid protein during human coronavirus 229E replication. *Antiviral Res* 2020; 173: 104620
21. Société Francophone de Transplantation (SFT); Société Francophone de Néphrologie, Dialyse et Transplantation (SFNDT) Groupe Infection et Immunodépression; Société de pathologie infectieuse de langue française (SPILF). Prise en charge de l'infection due au SARS-CoV-2 chez les patients adultes transplantés d'organe solide 2020. <https://www.sfndt.org/sites/www.sfndt.org/files/medias/documents/Recos%20COVID19%20guide%20pratique%20SFT%20SFNDT%20SPILF%2003%2004%2020.pdf> (1 May 2020, date last accessed)
22. Lopez V, Vazquez T, Alonso-Titos J *et al.* Recommendations on management of the SARS-CoV-2 coronavirus pandemic (Covid-19) in kidney transplant patients. *Nefrologia* 2020; doi: 10.1016/j.nefro.2020.03.002
23. British Transplant Society. Guidance on the management of transplant recipients diagnosed with or suspected of having COVID19. https://bts.org.uk/wp-content/uploads/2020/03/Clinical_management_transplant_recipients.pdf (1 May 2020, date last accessed)
24. American Society of Transplantation. 2019-nCoV (Coronavirus): FAQs for organ donation and transplantation. <https://www.myast.org/sites/default/files/COVID19%20FAQ%20Tx%20Centers%2004.06.2020%20%282%29.pdf> (1 May 2020, date last accessed)
25. Barlow A, Landolf KM, Barlow B *et al.* Review of emerging pharmacotherapy for the treatment of coronavirus disease 2019. *Pharmacotherapy* 2020; doi: 10.1002/phar.2398
26. Sanders JM, Monogue ML, Jodlowski TZ *et al.* Pharmacologic treatments for coronavirus disease 2019 (COVID-19): a review. *JAMA* 2020; doi: 10.1001/jama.2020.6019
27. Grein J, Ohmagari N, Shin D *et al.* Compassionate use of remdesivir for patients with severe COVID-19. *N Engl J Med* 2020; doi: 10.1056/NEJMoa2007016
28. University of Liverpool. COVID-19 drug interactions. <https://www.covid19-druginteractions.org>
29. van Maarseveen EM, Rogers CC, Trofe-Clark J *et al.* Drug-drug interactions between antiretroviral and immunosuppressive agents in HIV-infected patients after solid organ transplantation: a review. *AIDS Patient Care STDS* 2012; 26: 568–581
30. Patel SJ, Kuten SA, Musick WL *et al.* Combination drug products for HIV—A word of caution for the transplant clinician. *Am J Transplant* 2016; 16: 2479–2482
31. Cao B, Wang Y, Wen D *et al.* A trial of lopinavir-ritonavir in adults hospitalized with severe Covid-19. *N Engl J Med* 2020; doi: 10.1056/NEJMoa2001282
32. Russell CD, Millar JE, Baillie JK. Clinical evidence does not support corticosteroid treatment for 2019-nCoV lung injury. *Lancet* 2020; 395: 473–475
33. Mehta P, McAuley DF, Brown M *et al.* COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet* 2020; 395: 1033–1034
34. Shang L, Zhao J, Hu Y *et al.* On the use of corticosteroids for 2019-nCoV pneumonia. *Lancet* 2020; 395: 683–684
35. Wu C, Chen X, Cai Y *et al.* Risk factors associated with acute respiratory distress syndrome and death in patients with coronavirus disease 2019 pneumonia in Wuhan, China. *JAMA Intern Med* 2020; doi: 10.1001/jamainternmed.2020.0994

36. Zhang C, Wu Z, Li JW *et al*. The cytokine release syndrome (CRS) of severe COVID-19 and Interleukin-6 receptor (IL-6R) antagonist tocilizumab may be the key to reduce the mortality. *Int J Antimicrob Agents* 2020; doi: 10.1016/j.ijantimicag.2020.105954
37. Cao W, Liu X, Bai T *et al*. High-dose intravenous immunoglobulin as a therapeutic option for deteriorating patients with coronavirus disease 2019. *Open Forum Infect Dis* 2020; 7: ofaa102
38. Caly L, Druce JD, Catton MG *et al*. The FDA-approved drug ivermectin inhibits the replication of SARS-CoV-2 in vitro. *Antiviral Res* 2020; 178: 104787
39. REMAP-CAP response to the novel COVID-19 pandemic. <https://www.remapcap.org/coronavirus>
40. Gandolfini I, Delsante M, Fiaccadori E *et al*. COVID-19 in kidney transplant recipients. *Am J Transplant* 2020; doi: 10.1111/ajt.15891
41. Chen L, Xiong J, Bao L *et al*. Convalescent plasma as a potential therapy for COVID-19. *Lancet Infect Dis* 2020; 20: 398–400
42. Shen C, Wang Z, Zhao F *et al*. Treatment of 5 critically ill patients with COVID-19 with convalescent plasma. *JAMA* 2020; 323: 1582

Received: 15.4.2020; Editorial decision: 4.5.2020

Nephrol Dial Transplant (2020) 35: 904–907
doi: 10.1093/ndt/gfaa094
Advance Access publication 17 April 2020

The aftermath of coronavirus disease 2019: devastation or a new dawn for nephrology?

Rajiv Agarwal

Division of Nephrology, Department of Medicine, Indiana University School of Medicine and Richard L. Roudebush Veterans Administration Medical Center, Indianapolis, IN, USA

Correspondence to: Rajiv Agarwal; E-mail: ragarwal@iu.edu

ADDITIONAL CONTENT

An author video to accompany this article is available at: <https://academic.oup.com/ndt/>.

The acute crisis with the coronavirus disease of 2019 (COVID-19) caused by the novel coronavirus Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-COV-2) is the largest biomedical catastrophe of our lifetimes. Like so many other disasters in the past, such as war, famine, social unrest and economic calamities, this too shall pass, but it will undoubtedly leave the world changed. Some of the changes are already evident, but some are inevitable once we get over this pandemic. In this perspective, I provide examples of how the virus is already inducing change in the practice of medicine at large and for nephrology in particular. As is true for many changes, some persist after the emergency is over, so I speculate on how nephrology may change once we surmount this predicament (Figure 1).

COLLABORATION

In the worldwide fight against an invisible enemy, the world has learned incredible new ways to collaborate. Within days after the epidemic unfolded in North America, two young physicians, Matthew Sparks and Swapnil Hiremath, curated an annotated resource for use by nephrologists [1]. With a group of colleagues, they were among the first to address the question of whether angiotensin-converting enzyme inhibitors or




FIGURE 1: Coronavirus—devastation or a new dawn for nephrology?

angiotensin receptor blockers should be stopped in patients who are taking these drugs. Their answer—which was to continue to take these medications—was later supported both by a statement from the European Society of Cardiology [2] and a scholarly review in the *New England Journal of Medicine* [3]. In the context of the epidemic, under the umbrella of the World