

Rising prevalence of BMI ≥ 40 kg/m²: A high-demand epidemic needing better documentation

Kath Williamson^{1,3}  | Amy Nimegeer² | Michael Lean¹

¹School of Medicine, Dentistry and Nursing, University of Glasgow, Glasgow, UK

²MRC/CSO Social and Public Health Sciences Unit, University of Glasgow, Glasgow, UK

³NHS Lothian, Edinburgh, UK

Correspondence

Michael Lean, School of Medicine, Dentistry and Nursing, University of Glasgow, Level 2, New Lister Building, Royal Infirmary, 10-16 Alexandra Parade, Glasgow, Edinburgh, UK. Email: mike.lean@glasgow.ac.uk

Funding information

NHS Lothian Research Futures Award; NHS Research Scotland Career Researcher Fellowship; Royal College of Nursing Foundation, UK; University of Glasgow; Chief Scientist Office, Grant/Award Number: SPHSU15; Medical Research Council, Grant/Award Number: MC_UU_12017/15

Summary

Whilst previously rare, some surveys indicate substantial increases in the population with body mass index (BMI) ≥ 40 kg/m² since the 1980s. Clinicians report emerging care challenges for this population, often with high resource demands. Accurate prevalence data, gathered using reliable methods, are needed to inform health care practice, planning, and research. We searched digitally for English language sources with measured prevalence data on adult BMI ≥ 40 collected since 2010. The search strategy included sources identified from recent work by NCD-RisC (2017), grey sources, a literature search to find current sources, and digital snowball searching. Eighteen countries, across five continents, reported BMI ≥ 40 prevalence data in surveys since 2010: 12% of eligible national surveys examined. Prevalence of BMI ≥ 40 ranged from 1.3% (Spain) to 7.7% (USA) for all adults, 0.7% (Serbia) to 5.6% (USA) for men, and 1.8% (Poland) to 9.7% (USA) for women. Limited trend data covering recent decades support significant growth of BMI ≥ 40 population. Methodological limitations include small samples and data collection methods likely to exclude people with very high BMIs. BMI ≥ 40 data are not routinely reported in international surveys. Lack of data impairs surveillance of population trends, understanding of causation, and societal provision for individuals living with higher weights.

KEYWORDS

BMI ≥ 40 , data quality, prevalence, severe obesity

1 | INTRODUCTION

Prior to the 1970s, body mass index (BMI, kg/m²) ≥ 40 was rare.¹ It was regarded as a pathological condition, deemed to indicate genetic, endocrine, or other vulnerability, affecting a tiny, fixed population, with numbers too small for confident analysis.² Whilst prevalence can be minimal in low- to middle-income countries, contemporary studies suggest rising prevalence, with a current estimated global prevalence of 0.64% in men and 1.6% in women.³ It is predicted that by 2025, the global numbers of underweight women will be surpassed by those with BMI ≥ 35 .³

Although numbers may appear relatively small, compared with BMI ≥ 30 , frontline health professionals report increasing challenges in providing safe and effective overall health care for people with very high BMIs.⁴ Emerging issues relate to providing for basic care needs such as appropriate positioning and handling,^{5,6} continence and skincare,⁷ but also cover medical problems such as appropriate dosage of medicines,⁸ difficulty in performing medical imaging⁹ and complex psychological issues, including stigma, which impact treatment adherence.¹⁰ Evidence from several countries indicates that professional guidance or training about severe obesity for care givers is minimal,¹¹⁻¹⁵ threatening quality of care for this population,¹⁶ largely

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. Obesity Reviews published by John Wiley & Sons Ltd on behalf of World Obesity Federation

due to lack of awareness and limited evidence base. This problem is compounded by stigmatization over size and weight by the media, within society and among health professionals.¹⁷

Furthermore, conventional behavioural weight management interventions have minimal impact for BMI ≥ 40 .¹⁸ Only a minority of individuals access bariatric surgery,¹⁹ and effective nonsurgical interventions are not yet widely available.²⁰ Thus, once individuals reach a very high BMI, the potential for sustained weight loss to improve quality of life and reduce secondary medical complications is limited. The severity and impact of muscular-skeletal complications reduce physical activity and mobility, increasing dependence on others and putting weight loss further out of reach. Consequently, individuals with BMI ≥ 40 face reduced life expectancy,²¹ multimorbidity,²² disability,²³ and reduced quality of life.²⁴ In turn, this disease burden produces multifaceted demands on health and social care services, raising direct and indirect costs.²⁵ Until recently, evidence on direct costs of BMI ≥ 40 have been limited.²⁵ Total health care costs in the United Kingdom rise linearly, and double, as BMI increases²⁶ from 20 to 40. A recent systematic review of international health care costs and BMI found costs for people with BMI ≥ 40 to be 50% greater than for people with BMI 18.5 to 24.9.²⁷ Costing studies typically exclude underresearched wider care costs, such as social care²⁸ and nursing home usage related to functional disability and often long-term provision,^{28,29} so current estimates are likely to underestimate the full costs. Forecasts indicate that increased resource usage will continue, including costs required to structurally adapt care facilities to the needs of people with BMI ≥ 40 , alongside providing suitable equipment and training for staff, that is currently missing.³⁰

The present scoping review explores the extent of international prevalence data on BMI ≥ 40 . It focusses on measured data, given the potential for error and bias with self-reported anthropometry by individuals who are overweight.^{31,32} We assessed the extent and quality of epidemiological reporting for the BMI ≥ 40 category internationally, with a view to improving the documentation of this emerging high-demand population in future national surveys, to enable development of a reliable evidence base to guide effective care.

2 | METHODS

The primary epidemiological reports being investigated are health surveys, undertaken by governments for population surveillance to inform strategic policy priorities. Health surveys are not usually reported in the academic literature, unless for a secondary analysis focussed on a specific issue or subpopulation, often with considerable time lag. Thus, database search terms for the primary survey would need to be very broad, making identification through academic databases highly resource intensive, with results prone to being incomplete and outdated. For the present scoping review, an alternative search strategy was therefore applied, based on the sources identified by the most recent systematic review of international BMI survey data, with additional searches to update and supplement these sources, outlined below.

Four key approaches were used to identify potential data sources:

1. Building on previous work

The NCD Risk Factor Collaboration (NCD-RisC) 2017 study on global BMI trends was chosen as a basis for the initial search, on the basis of its size and rigour: It used 2416 sources of measured data, collected up to 2016.³³ These sources were compiled from a systematic medical database search, supplemented by a worldwide network of researchers identifying and accessing national measurement surveys via interested parties, including World Health Organization (WHO). Full details are given in the published paper and its appendix.³³ All 2416 sources in the NCD-RisC appendix were screened as per the inclusion criteria in Table 1, with digital snowball searching used to locate individual data sources.

2. Digital searching of current grey literature sites

Searching of key international organizations websites known to compile BMI population survey data was undertaken, focussing on the Organization for Economic Cooperation Development (OECD), the WHO, and the World Obesity Federation (WOF).

3. Systematic database search

To ensure identification of any new sources since 2016 in the academic literature, a modified version of the NCD-RisC literature search was undertaken in Medline and Embase using the search terms

TABLE 1 Source inclusion and exclusion criteria

Included	Excluded
Reports BMI ≥ 40 prevalence	Does not report BMI ≥ 40 prevalence
Measured anthropometric data	Self-reported anthropometric data
Nationally representative data: sampling strategy showing national coverage based on electoral roll/census or similar, with at least stratification for age and sex	Nonnationally representative, subpopulations by age/sex/rural/urban/regional/community
Adults aged 15 years or older	Children and adolescents
Data collected in or since 2010	Data collected pre-2010
Data already compiled in a publicly available report/website ^a	Data requiring registration/searching through raw data ^b
Report in English language	Report not in English language
Not already identified by search	Already identified through alternative source

Abbreviation: BMI, body mass index.

^aFor immediate use by decision makers.

^bMay be available to academics, but requiring analysis and presentation prior to use by decision makers.

in Appendix A. Results from conference proceedings were deliberately included, as they can provide the first exposure of research analysis, highlighting new or updated data sources, whilst full articles commonly take much longer to publication, if indeed they are ever published in full. Sources were searched both for BMI ≥ 40 data, but also to identify any new sources/health surveys not already located through parts 1 and 2 of the search process.

4. Digital snowball searching

Whilst NCD-RisC identified sources, individual sources needed located and accessed, often directly from a website of government agencies or international bodies. This was undertaken digitally, seeking previously unknown sources or articles.

All sources were screened using inclusion criteria in Table 1. Figures 1A and 1B represent the search process, with numbers of studies included and excluded at each stage. Searches took place between June and September 2019. Where successive or annualized surveys were identified, sources were checked for the most current data, with the most recent data retained. If multiple studies for the same country or source were identified, the one with the largest sample, or that provided the most information to data extraction, was retained. These pragmatic methods adopt a systematic approach to exploring the broad evidence landscape in an emerging area, highlighting gaps for future, more detailed research.³⁴

3 | RESULTS

Prevalence data of measured BMI ≥ 40 from 2010 onwards were located for 18 individual countries, on all five continents, comprising 12% of eligible national survey data examined (Table 2) either in its primary form ($n = 10$) or through secondary analysis ($n = 8$). Data were very recent, with half of the countries reporting prevalence from 2015 onwards, and only three back to 2010. None of the original sources located reported BMI data categories higher than BMI ≥ 40 (for example, BMI $\geq 45/\geq 50$), with just one secondary analysis from Spain doing so.⁴¹

Germany,³⁷ Saudi Arabia,⁴³ Seychelles,⁴⁶ and Poland³⁸ report only for men and women separately, with no data for combined-sex adults. In contrast, only combined-sex-adult data were available for Brazil⁴⁸ and Portugal.³⁹ All regions of the world (as defined by WHO) contained a country with rates above 4% for women, and 2% for men, with the exception of Africa, where the maximum was Seychelles with 1.5% for men.⁴⁶ Other than Brunei Darussalam,⁵¹ rates for women were universally higher than for men.

Prevalence rates are presented graphically in Figure 2A-C. The United States has the highest prevalence,⁴⁹ with rates across all adults, men, and women ranging between 5.6% and 9.7%. Other countries with higher rates (5.0%-7.7%) for all adults and women are New Zealand,⁴⁵ Kuwait,⁴² and Barbados,⁴⁷ although rates in men are

markedly lower. Australia,⁴⁴ Canada,⁵⁰ Scotland,³⁶ England,³⁵ and Saudi Arabia all have rates in the region of 2.5% to 5.5% for all adults and/or women, again with men notably lower. Germany, Serbia,⁴⁰ Spain, Portugal, Poland, Brazil, and Malaysia⁵² display the lowest rates between 0.7% and 2.8%. Brunei Darussalam and Seychelles both exhibit disparate prevalence patterns from those above for men and women.

Table 2 is modelled on Foresight's International Evidence Review,⁵³ which when published illustrated the general availability of regionally comparative BMI ≥ 30 prevalence data. The rigorous inclusion criteria applied during searching means that most sources (with potential exception of Spain and Poland) were from surveys done or supported by national government agencies, with primary reporting through grey literature sources. Due to the constraints associated with gathering measured data, total sample sizes tend to be relatively small, with 13 countries being < 8000 , and five above this. With prevalence under 10% of these numbers, wide confidence intervals result for BMI ≥ 40 , as seen in Table 2.

Less than half of the countries included have regular surveillance, ranging from annual to three to four yearly (Table 2: England, Scotland, Canada, New Zealand, Australia, and the United States), significantly limiting analysis of trends over time. Reporting from other countries appears to be more ad hoc and unpredictable, without planning for regular surveillance.

3.1 | International survey data sources

Globally, BMI ≥ 40 data were generally poorly available compared with BMI ≥ 30 data. Seventy-one NCD-RisC sources had BMI $\geq 25/\geq 30$ data available in English, of which only 11 (15%) provided BMI ≥ 40 data. Forty six studies out of the 80 screened for the literature search reported mean BMI or BMI $\geq 25/\geq 30$ data, with just one (2%) reporting BMI ≥ 40 . For Africa, only the island state of Seychelles presented BMI ≥ 40 data.

The OECD annually publishes country by country Health Statistics with measured data sources for 26 out of 44 countries in its 2019 report.⁵⁴ Many major European countries are missing, as the European Health Interview Survey (EHIS), which is the primary health data collection tool for Eurostat, the Statistics office of the European Union, uses self-reported height and weight data.⁵⁵ The European Health Examination Survey does include measured data but appears not to have been widely adopted.⁵⁶

Additional grey sources were searched, including WHO website, where the categorizations consistently applied to data were BMI 25.0 to 29.9 and ≥ 30 , with no additional categorization for BMI ≥ 40 . The WOF website supports the Global Obesity Observatory, featuring a searchable interactive map detailing national overweight and obesity rates, with data sources referenced. All 179 available countries were individually searched, with 12 displaying data on BMI ≥ 40 ,⁵⁷ of which four had not been previously identified and fitted the inclusion criteria.




FIGURE 1 Search strategy (A) part 1 and 2: NCD-RisC sources and grey sources, (B) part 3 & 4: literature search

3.2 | Data quality

To enable assessment of data quality, the column headers of Table 2 highlight basic quality parameters appropriate to national health surveillance,⁵⁸ with additional categories particularly relevant to the population with BMI ≥ 40 .

Survey methods for the six countries with regular surveillance in place (England, Scotland, Australia, New Zealand, Canada, and the

United States) are available for scrutiny on public websites. These surveys employ complex sample design, aimed at reducing bias and with weighting to reflect the age/sex stratification of the population. Response rates are difficult to compare, due to heterogeneity of definitions. As ongoing programmes, the surveys measure nonresponse across years, which appears to be a growing challenge, with Australian data showing the nonresponse rate for the BMI module specifically, increasing from 26.8% in 2014 to 2015 to 33.8% in 2017 to 2018.⁵⁹

TABLE 2 International BMI ≥ 40 kg/m² prevalence rates for data collected since 2010

Country	Prevalence %		Source	Year Data Collected	Sample Size	Age, years	Ongoing Data Collection	Data Collection Method	Scale Capacity, kg	
	All	Men								Women
WHO European Region										
England	4 ^a (3.1-4.1)	2 ^a (1.9-3.2)	5 ^a (3.9-5.4)	Health Survey England ³⁵	2017	6530	≥16	Annual	Home visit	200
Scotland	3 ^a (2.8-4.1)	2 ^a (1.5-3.2)	4 ^a (3.5-4.5)	Scottish Health Survey ³⁶	2018	3746	≥16	Annual	Home visit	200 ^b
Germany	—	1.2 (0.8-1.8)	2.8 (2.2-3.5)	German Health Interview and Examination Survey for Adults ³⁷	2008-2011	7116	18-79	No	Community centres ^c	250
Poland	—	1.3 (0.09-1.7)	1.8 (1.3-2.3)	WOBASZ II ³⁸	2013-2014	5417	≥20	No	Home visit and clinics	No data
Portugal	1.8 (1.4-2.2)	—	—	National Health Examination Survey (INSEF) ³⁹	2015	4819	25-74	No	Health centres	200
Serbia	1.4 (1.2-1.6)	0.7 (0.5-0.9)	2.0 (1.7-2.4)	National Health Survey ⁴⁰	2013	13 103	≥15	No	Home visit	No data
Spain	1.3 (1.0-1.8)	1.0 (0.6-1.6)	1.6 (1.1-2.3)	Nutritional Study of Spanish Population (ENPE) ⁴¹	2014-2015	3966	25-64	No	Home visit	150
WHO Eastern Mediterranean Region										
Kuwait	5.5 (4.8-6.3)	3.9 (2.9-4.9) ^d	7.0 (5.9-8.1) ^d	STEPS, Weidpass et al ⁴²	2014	3589	18-69	No	Health centres	No data
Saudi Arabia	—	2.5 (2.3-3.3)	4.7 (3.9-5.5)	Saudi Health Interview Survey ⁴³	2013	10 337	≥15	No	Home visit	No data
WHO Western Pacific Region										
Australia	4.0 (3.6-4.4)	3.3 (2.8-3.8)	4.7 (4.2-5.2)	National Health Survey ⁴⁴	2017-2018	18 656	≥18	3-4 yearly	Home visit	200
New Zealand	5.1 (4.5-5.6)	3.5 (2.9-4.1)	6.6 (5.8-7.4)	New Zealand Health Survey ⁴⁵	2017-2018	13 869	≥15	Biennial	Home visit	200
WHO Africa Region										
Seychelles	—	1.5 (0.5-2.5)	6.7 (4.9-8.5)	Seychelles Heart Study IV ⁴⁶	2013-2014	1240	25-64	No	Health centre	No data
WHO Americas Region										
Barbados	5.0 (3.7-6.7)	1.8 (0.8-4.0)	7.9 (6.0-10.5)	Barbados Health of the Nation Study ⁴⁷	2011-2013	1197	≥25	No	Home visit	No data
Brazil	1.8 (1.6-2.0)	—	—	National Health Survey, Wagner et al ⁴⁸	2013	49 359	20-64	No	Not given	No data
United States	7.7 (6.6-8.9)	5.6 (4.3-7.2)	9.7 (8.4-11.2)	NHANES, Hales et al ⁴⁹	2015-2016	5337	≥20	Biennial	Mobile Examination Centre	272

(Continues)

TABLE 2 (Continued)

Country	Prevalence %		Source	Year Data Collected	Sample Size	Age, years	Ongoing Data Collection	Data Collection Method	Scale Capacity, kg
	All	Men							
Canada	4.0 (2.8-5.8)	2.6 ^c (1.3-5.0)	Canadian Health Measures Survey ⁵⁰	2014-2015	5794	18-79	Biennial	Mobile Examination Centre ^c	272 ^d
WHO South East Asia Region									
Brunei Darussalam	3.0 (2.2-4.0)	3.9 (2.6-5.6)	National Health and Nutrition Status Survey ⁵¹	2010-2011	1524	>19	No	Local clinic ^c	200
Malaysia	1.4 (1.2-1.6)	0.9 (0.7-1.2)	National Health and Morbidity Survey ⁵²	2015	5196	>18	No	Not given	150

Abbreviation: BMI, body mass index; WHO, World Health Organization.

^aReported as rounded, whole numbers only.

^bConfirmed through personal communication with survey team.

^cTransport provided to centre if needed.

^dConfidence intervals calculated using % sample size for men and women, as absolute numbers only provided for all adults.

^eSource advises use with caution.




FIGURE 2 International body mass index (BMI) ≥40 prevalence rates by country: (A) all adults, (B) men, and (C) women

Secondary analyses tended to have less available detail on sampling and methodology, with those for Spain⁴¹ and Brazil⁴⁸ referencing previous publications not available in English.

Data collection methods of included surveys were scrutinized for factors that may affect participation of people with BMI ≥40 (Table 2). Eleven countries specifically stated exclusion criteria, which were very similar, namely, the institutionalized population. This excluded those in hospitals and, apart from New Zealand, people in care homes. No surveys appeared to document an upper BMI limit, but the capacity of scales used has potential to enact this, effectively meaning individuals with weights over the maximum capacity of the instrument are either excluded or estimated self-reports.³⁵ No clear

data on scale capacity could be found for seven countries, two countries appear to have used scales with a maximum capacity of 150 kg, whilst nine had scale capacities of 200 kg or over. Functional mobility limitations may affect participants' ability to stand on scales or attend examination centres outside the home. For two countries, place of anthropometry data collection was unavailable, nine countries used home visits, whilst seven required participants to leave their home, although some of these offered transport if needed.

4 | DISCUSSION

Whilst good quality BMI ≥ 30 prevalence data are now available globally, largely due to the widespread implementation of surveillance tools such as STEPS, there are few robust measured data on BMI ≥ 40 or higher categories. The published surveys generally use similar proven methodologies to obtain population-representative data. However, the methods used to assess BMI ≥ 30 , risk providing inaccurate results for higher BMI categories, for example, if they require mobility of participants or if the scales used have an upper limit of 200 kg. Hence, true prevalence may be higher than stated in Table 2.

The data that are available have limitations, notably wide confidence intervals for higher BMI categories, making it difficult to determine the significance of annual changes. If, as predicted, numbers with BMI ≥ 40 continue to grow, this problem may diminish, although is unlikely to disappear altogether as the cost of gathering measured data limits sample sizes. The limited long-term trend data available help to overcome uncertainties with large confidence intervals for the smaller numbers with BMI ≥ 40 . Since 1995, BMI ≥ 40 prevalence in Scotland has trebled for women aged 16 to 64 years.⁶⁰ Australia saw similar increases of 2.9-fold for men and 2.0-fold for women between 1995 and 2011 to 2012.⁶¹ Prevalence of BMI ≥ 40 in the United States quadrupled between 1976 and 2004, substantially surpassing the rise in BMI ≥ 30 .⁶² Prevalence in men in England aged 16 years or older experienced an eight-fold increase since 1995,³⁵ but Brazil reported the most dramatic increase of nearly 20-fold between 1974-1975 and 2013.⁴⁸ Overall population distributions have thus consistently shifted upwards, as illustrated in Figure 3, comparing waves of US NHANES data from 1967-1980 to 2005-2006.⁶³

Comparison with analyses of self-report surveys from the United States and Canada, which allow for much larger samples, all show disproportionately larger growth in higher BMI categories, with increases of up to 10-fold in the BMI ≥ 40 category.⁶⁴⁻⁶⁶ This is despite suggestions that the underestimation of weight for self-report data is likely to be greater with higher BMI.³¹ Thus, despite wide confidence intervals in individual survey years and some variations in methodologies, measured trend data evidence from several countries, together with larger-scale self-report datasets, all support a long-term rise in BMI ≥ 40 .

Two key sources of data for NCD-RisC and WOF were STEPS reports and Demographic Health Survey (DHS) country reports.

The STEPwise approach, featuring three different level of "steps" of key risk factor assessments for NCDs, was developed by WHO, to aid countries increase their surveillance capacity.⁶⁷ One hundred and thirteen country reports or data sheets were publicly available on the WHO STEPS site at the time of searching.⁶⁸ DHS is supported by the US Agency for International Development (USAID), to inform planning particularly in relation to maternal and child health, with over 93 standard DHS reports since 2010 on its website.⁶⁹ Both of these are easily accessible population surveillance tools, widely used by low- to middle-income countries, with standard methodologies including collecting measured height and weight in a household setting. Current standard reporting practice for STEPS and DHS focusses on BMI 25.0 to 29.9 and ≥ 30 , with no additional categorization for BMI ≥ 40 .

International health surveillance provides reliable health information that is comparable over time and between populations. It has allowed documentation of a nutrition transition that is rooted in the impact of large-scale social changes such as reduction in physical activity, increased urbanization, and greater consumption of processed food.⁷⁰ Whilst regional variation remains, the major global concern has moved away from underweight as the primary nutrition issue, towards excess weight as the leading cause of morbidity and mortality through secondary noncommunicable diseases.²³ This has already occurred to such a degree that BMI ≥ 35 in women now surpasses underweight in 165 countries for women and 113 for men.³ Yet current national documentation of BMI distribution does not reflect this shift, with the focus still on reporting categories < 18.5 to ≥ 30 , and/or mean BMI, as the DHS and STEPS reports evidence. This historical bias is obscuring the significant




FIGURE 3 Changes in the distribution of body mass index (BMI) between 1976-1980 and 2005-2006, adults aged 20 to 74 years: United States.⁶³ Data are age adjusted by the direct method to the year US Census 2000 estimates using age groups 20 to 39, 40 to 59, and 60 to 74. Overweight is BMI of 25.0 to 29.9 kg/m²; obesity is BMI at or above 30.0 kg/m²; and severe obesity is BMI at or above 40.0 kg/m². Pregnant women are excluded from the analysis. Sources: NCHS, National Health Examination Survey, and National Health and Nutrition Examination Surveys

changes in higher BMI categories.^{42,66} Without characterizing this progression, policymakers and planners are unable to respond effectively to the needs of the population or evaluate the effectiveness of policies.

The example of Kuwait in this review provides a case in point. The original 2014 STEPS report on the WHO website documents the mean BMI and four separate BMI categories <18.5, 18.5 to 24.9, 25.0 to 29.9, and ≥ 30 , without ≥ 40 , as a distinct category.⁶⁸ The BMI ≥ 40 data were later reported by the analysis of Weiderpass et al of the original STEPS dataset, illustrating that the data had been collected but gone unreported.⁴² Table 2 shows Kuwait's prevalence rates as second highest only to the United States, providing valuable information regarding BMI population distribution, in a region where no other sources of all adult prevalence were found.

One practical solution would be for WHO and similar agencies to call for data on the BMI ≥ 40 population to be included when reporting all anthropometry surveys. The National Child Measurement Programme in England took an equivalent step in 2018, adding severe obesity as a reporting category.⁷¹ Given that the prevalence of BMI ≥ 50 is now similar to that of the BMI ≥ 40 category about 20 years ago, with some real-world datasets including categories of 50/60/70,⁷² it may be wise also to include reporting BMI ≥ 50 , to map future trends. This would hugely increase the amount of BMI ≥ 40 data available globally, with little extra cost, given that the data are already collected. For countries with small numbers to report, the need for caution in interpretation would be dealt with in the same way for low numbers in any category, for example, underweight.

4.1 | Causation

The lack of data on BMI ≥ 40 trajectories by region, nation, age, sex, and class makes it difficult to explore causation. Improved international data, ideally from longitudinal studies, would promote comparison between countries, taking into account their differing social and economic contexts.⁷³ Together with the emergence of large-scale genome studies looking at the inherited susceptibility to BMI ≥ 40 ,⁷⁴ reasons for the escalation of high body weight may be more accurately sought. There are some indications of associations with lower socio-economic status in some populations,^{75,76} but these patterns require further study. Concerningly, some countries report increasing rates of the highest BMI groups growing for children and adolescents,^{71,77} with potential for excess weight to track through into adulthood. This would differ from current patterns, when rates are lower in early adulthood, peaking in middle age.^{35,49,78} Unusually, the survey data from Brunei Darussalam showed 19- to 29-year-olds having some of the highest rates of BMI ≥ 40 across the age trajectory.⁵¹

A lack of prevalence data keeps the population hidden, preventing development of appropriate weight management services to treat this population group, along with comparative

analysis of different treatment models and health care systems.²⁵ BMI ≥ 35 with comorbidities or BMI ≥ 40 with or without comorbidities is a commonly applied threshold for bariatric surgery, yet access to surgery is often very limited.^{19,79} Evidence on effective alternatives to surgery or prevention is needed, whilst access to traditional services can be difficult for people with BMI ≥ 40 due to functional disability.⁸⁰ Improved global prevalence data would facilitate work on economic costing of treatment and prevention for the population with BMI ≥ 40 .

4.2 | Consequences of rise in prevalence

Whilst numbers may appear small in terms of proportion of the whole population, given that these are at national scale, they translate into significant absolute numbers with a large real-life impact on care provision.

4.2.1 | Health risk and comorbidities

It is well recognized that BMI can be a poor proxy for body fat, and in many studies, waist circumference or evidence-based predictive equations for total body fat are better than BMI for assessing health risk, particularly cardiac and metabolic health outcomes.⁸¹⁻⁸⁴ Consequently, surveys from the United States, England, Scotland, New Zealand, and Australia all measure waist circumference. However, people with BMI ≥ 35 exceed these cut points, often with a large abdominal fat apron, making methods which incorporate waist circumference unlikely to be reliable for very high body weights. Thus, BMI remains the best available simple estimate of body fat at the highest levels. Surveys and studies need to consider data collection methods, particularly scales capable of weighing at least 200 kg, with easily accessible wide and low platforms, ideally offering home visits to facilitate accurate data collection from people with high BMIs. Additionally, a review is needed regarding the treatment of what have historically been seen as biologically implausible values (BIVs), the majority of which have been found to be accurate.⁸⁵ The definition of BIVs and use of upper thresholds for weight and BMI in research studies requires reexamination in view of the documented shift in population distribution towards heavier BMIs.

Elevated health risk translates into increased prevalence of multimorbidity (co-occurrence of ≥ 2 conditions) compared with those of normal weight by nearly two⁸⁶ to seven times,⁴⁰ particularly for cardiometabolic multimorbidity.²² Alongside physical diseases, risk of depression increases with BMI,⁸⁷ together with functional disability,⁸⁸ with increasing numbers of people with BMI ≥ 40 living in care homes.²⁹ As such care of these multiple obesity-related consequences amplifies health care costs considerably.⁸⁹ Given the rise in prevalence and size of these costs, in addition to older costing studies being limited by BMI thresholds that are now too low, this is likely to become of increasing concern globally.

4.2.2 | Planning

As an emerging population, people with BMI ≥ 40 , especially those with BMI ≥ 50 and ≥ 60 , have needs that are currently often unaddressed by service providers.⁴ Increasingly, changes to care environments are being needed to accommodate larger body sizes, requiring adjustments such as widened doorways, reinforced floors, suitable seats, and larger rooms. Problems featured in the health literature demonstrate the scope of challenges, such as evacuation planning,⁹⁰ diagnostic scanning,⁹ and positioning during surgery in theatre.⁹¹ Documented requirements include staff training,¹¹ specialist equipment provision,⁹¹ and development of specialist clinical protocols covering essential areas, such as tissue viability guidance.⁹² However, issues are not restricted to health care but affect all aspects of life, including specialist housing,⁹³ adapted workplace design,⁹⁴ larger sized fashionable and safety clothing,⁹⁵ and barriers to travelling.⁹⁶ Projections of future severe obesity prevalence using different datasets agree in predicting continued rises for the foreseeable future.¹ Current analyses suggest five million living with a BMI ≥ 40 in the United Kingdom by 2035, with rates up to 20% for Welsh women aged 55 to 64 years⁹⁷ and over 20 million people affected in the United States.³⁰ Better characterization of the population with BMI ≥ 40 is required to support organizational and societal readiness for this population. Adaptations are expensive in time and money, particularly when made retrospectively, underlining the need for accurate planning to happen now.

4.3 | Strengths and limitations

This review has concentrated on robustly measured data to establish the international prevalence rates for BMI ≥ 40 . Whilst the rationale for exclusion of self-report data is sound, in that it commonly underestimates BMI, the exclusion also acts as a limitation, for example, by excluding EHIS data, which covers many European countries. The sample sizes possible with measured data are reduced by the need for resources to make measurements, and there is potential bias against including very heavy individuals whose mobility is impaired. In some cases, the upper limit of scales excluded the heaviest individuals. These limitations would tend to underestimate the true prevalence of the highest BMI categories, not overestimate.

Applying a lower threshold of BMI ≥ 35 would have broadened the available data, whilst potentially weakening the focus on the highest BMI category, where costs and clinical complexity is greatest. Some studies report the lower threshold of BMI ≥ 35 particularly those examining Asian populations where different BMI cut-offs relating to overweight and obesity are often applied.⁹⁸

Limiting the review to the English language prevented examination of some original data sources, which could only be located in their native language, for example, Spanish for Chile and Mexico. It was not possible to locate English versions of these, and resources did not allow for translation. The OECD reports these original sources

in English in its database, but only at BMI thresholds of 25.0 to 29.9 and BMI ≥ 30 , with prevalence for Chile and Mexico the highest in the world, above even those of the United States.⁹⁹ Thus, they are likely to have significant BMI ≥ 40 prevalence. Additionally, whilst the search processes were broad, encompassing a variety of sources and used systematic methods, they were not exhaustive, as might be expected from a formal systematic literature review or meta-analysis. We believe that they represent a reliable summary of the current evidence base on BMI ≥ 40 , as it is available to decision makers. Some individual sources have not been included, notably those from non-English publications, but they are unlikely to alter the very consistent conclusions.

5 | CONCLUSION

This review highlights the poor availability of robust international data available on the emerging issue of BMI ≥ 40 prevalence. The measured data available suggest significant prevalence on all five continents, with proportionally large rises in recent decades. Given the multiple care challenges, high resource needs and poor current evidence base for this population, routinely reporting BMI ≥ 40 and higher categories in national surveys, would be valuable, with appropriate caveats for interpreting the still small numbers in individual surveys. Accurate characterization of the subpopulations with BMI ≥ 40 and higher categories requires consideration of measurement equipment and the mobility limitations of individuals with high BMI.

ACKNOWLEDGEMENTS

Thanks to Welcome Wami and Andy Peters for providing statistical support.

CONFLICT OF INTEREST

The authors declare no competing financial interests in relation to this work.

FUNDING INFORMATION

Kath Williamson was supported by an NHS Research Scotland Career Researcher Fellowship, an RCN Professional Foundation Bursary, and an NHS Lothian Research Futures award for this work.

Amy Nimegeer is funded by the Medical Research Council (MRC) and the Chief Scientist Office of the Scottish Government Health Directorates (CSO) as part of the MRC/CSO Social and Public Health Sciences Unit's Informing Healthy Public Policy programme (MC_UU_12017/15 and SPHSU15).

Michael Lean is funded by the University of Glasgow.

ORCID

Kath Williamson  <https://orcid.org/0000-0001-8606-8935>

REFERENCES

1. Finkelstein EA, Khavjou OA, Thompson H, et al. Obesity and severe obesity forecasts through 2030. *Am J Prev Med.* 2012;42(6): 563-570.
2. Sturm R. Increases in morbid obesity in the USA: 2000-2005. *Public Health.* 2007;121(7):492-496.
3. NCD-Risk Factor Collaboration (NCD-RisC). Trends in adult body-mass index in 200 countries from 1975 to 2014: a pooled analysis of 1698 population based measurement studies with 19.2 million participants. *Lancet* 2016;387(10026):1377-1396.
4. Lumley E, Homer CV, Palfreyman S, Shackley P, Tod AM. A qualitative study to explore the attitude of clinical staff to the challenges of caring for obese patients. *J Clin Nurs.* 2015;Dec;24(23-24): 3594-3604.
5. Wiggerman N, Smith K, Kumbar D. What bed size does a patient need? The relationship between body mass index and space required to turn in bed. *Nurs Res.* 2017;66(6):483-489.
6. Gardner L-A. Caring for Class III obese patients. *Am J Nurs.* 2013;113 (11):66-70.
7. Rose MA, Pokorny M, Drake DJ. Preventing pressure ulcers in the morbidly obese: in search of an evidence base. *Bariatric Nurs Surg Patient Care.* 2009;4(3):221-227.
8. Polso AK, Lassiter JL, Nagel JL. Impact of hospital guideline for weight-based antimicrobial dosing in morbidly obese adults and comprehensive literature review. *J Clin Pharm Ther.* 2014;39(6):584-608.
9. Wiles R, Meredith SM, Mullany JP, Wiles T. Are English CT departments and radiographers prepared for the morbidly obese patient? *Radiography.* 2017;23(3):187-190.
10. Thille P. Managing anti-fat stigma in primary care: an observational study. *Health Commun.* 2019;34(8):892-903.
11. Royal College of Physicians. In: *Action on Obesity: Comprehensive Care for all. Report of a working party.* London, UK: Royal College of Physicians; 2013.
12. Rinne C, Orschel C, Semkowich B, et al. Knowledge, attitudes, and practice of in-patient physiotherapists in Ontario regarding patients who are super-morbidly obese. *Physiother Can.* 2018;70(2): 102-112.
13. Poitou C, Chakhtoura Z, Uzan C, Nikpayam M, Oppert M. Just the tip of the ice-berg: difficulties in assessing and managing extreme obesity in routine clinical care. *Eur J Clin Nutr.* 2018;72(3):452-454.
14. Hales C, de Vries K, Coombs M. Managing social awkwardness when caring for morbidly obese patients in intensive care: a focused ethnography. *Int J Nurs Stud.* 2016;58:82-89.
15. Pokorny ME, Scott E, Rose MA, et al. Challenges in caring for morbidly obese patients. *Home Healthc Nurse.* 2009;27(1):43-52.
16. Zhang N, Li Y, Temkin-Greener H. Prevalence of obesity in New York nursing homes: associations with facility characteristics. *Gerontologist.* 2013;53(4):567-581.
17. Teachman BA, Brownell KD. Implicit anti-fat bias among health professionals: is anyone immune? *Int J Obes Relat Metab Disord.* 2001;25 (10):1525-1531.
18. Fildes A, Charlton J, Rudisill C, Littlejohns P, Prevost AT, Gulliford MC. Probability of an obese person attaining normal body weight: cohort study using electronic health records. *Am J Public Health.* 2015;105(9): e54-e59. <https://doi.org/10.2105/ajph.2015.302773>
19. Gulliford MC, Charlton J, Prevost T, et al. Costs and outcomes of increasing access to bariatric surgery: cohort study and cost-effectiveness analysis using electronic health records. *Value Health.* 2017;20(1):85-92. <https://doi.org/10.1016/j.jval.2016.08.734>
20. McCombie L, Brosnahan N, Ross H, Bell-Higgs A, Govan L, Lean MEJ. Filling the intervention gap: service evaluation of an intensive non-surgical weight management programme for severe and complex obesity. *J Hum Nutr Diet.* 2019;32(3):329-337.
21. Global BMI Mortality Collaboration. Body-mass index and all-cause mortality: individual-participant-data meta-analysis of 239 prospective studies in four continents. *Lancet.* 2016;388(10046):776-786.
22. Kivimaki M, Kuosma E, Ferrie JE, et al. Overweight, obesity, and risk of cardiometabolic multimorbidity: pooled analysis of individual-level data for 120 813 adults from 16 cohort studies from the USA and Europe. *Lancet Public Health.* 2017;2(6):e277-e285. [https://doi.org/10.1016/s2468-2667\(17\)30074-9](https://doi.org/10.1016/s2468-2667(17)30074-9)
23. Global Burden of Disease 2015 Obesity Collaborators. Health effects of overweight and obesity in 195 countries over 25 years. *N Engl J Med.* 2017;377(1):13-27.
24. Gupta S, Richard L, Forsythe A. The humanistic and economic burden associated with increasing body mass index in the EU5. *Diabetes Metab Syndr Obes.* 2015;8:327-338.
25. Grieve E, Fenwick E, Yang HC, Lean M. The disproportionate economic burden associated with severe and complicated obesity: a systematic review. *Obes Rev.* 2013;14(11):883-894.
26. Tigbe WW, Briggs AH, Lean MEJ. A patient-centred approach to estimate total annual healthcare cost by body mass index in the UK counterweight programme. *Int J Obes (Lond).* 2013;37(8): 1135-1139.
27. Kent S, Gray A, Jebb SA, Cairns BJ, Mihaylova B. Body mass index and healthcare costs: a systematic literature review of individual participant data studies. *Obes Rev.* 2017;18(8):869-879.
28. Copley VR, Cavill N, Wolstenholme J, Fordham R, Rutter H. Estimating the variation in need for community-based social care by body mass index in England and associated cost: population-based cross-sectional study. *BMC Public Health.* 2017;17(1):667. <https://doi.org/10.1186/s12889-017-4665-1>
29. Zhang N, Field T, Mazor K, Zhou Y, Lapane KL, Gurwitz JH. The increasing prevalence of obesity in residents of U.S. nursing homes: 2005-2015. *J Gerontol A Biol Sci Med Sci.* 2019;74(12):1929-1936. <https://doi.org/10.1093/gerona/gly265>
30. Cecchini M. Use of healthcare services and expenditure in the US in 2025: the effect of obesity and morbid obesity. *PLoS ONE.* 2018;13 (11):e0206703. <https://doi.org/10.1371/journal.pone.0206703>
31. Luo J, Thomson CA, Hendryx M, et al. Accuracy of self-reported weight in the Women's health initiative. *Public Health Nutr.* 2019;22 (6):1019-1028.
32. Stommel M, Schoenborn CA. Accuracy and usefulness of BMI measures based on self-reported weight and height: findings from the NHANES & NHIS 2001-2006. *BMC Public Health.* 2009;9(1):1-10.
33. NCD-Risk Factor Collaboration (NCD-RisC). Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *Lancet.* 2017;390(10113):2627-2642.
34. Peters MDJ, Godfrey C, McInerney P, Baldini Soares C, Khalil H, Parker D. Chapter 11: scoping reviews. In: Aromateris E, Munn Z, eds. *Joanna Briggs Institute Reviewer's Manual.* The Joanna Briggs Institute: Available from <https://reviewersmanual.joannabriggs.org/>; 2017.
35. Health Survey for England 2017. NHS Digital Web site. <https://digital.nhs.uk/data-and-information/publications/statistical/health-survey-for-england/2017>. Published 2018. Accessed June 29th, 2019.
36. Scottish Health Survey 2018. Scottish Government Web site. <https://www2.gov.scot/scottishhealthsurvey>. Published September 24th, 2019. Accessed 25th September, 2019.
37. Mensink GBM, Schienkiewitz A, Haftenberger M, Lampert T, Ziese T, Scheidt-Nave C. Overweight and obesity in Germany: results of the German health interview and examination survey for adults, English version of "Übergewicht und Adipositas in Deutschland. Ergebnisse der Studie zur Gesundheit Erwachsener in Deutschland (DEGS1)" in Supplementary Material. *Bundesgesundheitsblatt - Gesundheitsforschung - Gesundheitsschutz.* 2013;56:786-794.

38. Stepaniak U, Micek A, Waskiewicz A, et al. Prevalence of general and abdominal obesity and overweight among adult in Poland. Result of the WOBASZ II study (2013-14) and comparison with the WOBASZ study (2003-5). *Pol Arch Med Wewn.* 2016;126(9):662-671.
39. Gaio V, Antunes L, Namorado S, et al. Prevalence of overweight and obesity in Portugal: results from the first Portuguese Health Examination Survey (INSEF 2015). *Ob Res Clin Pract.* 2018;12(1):40-50.
40. Jovic D, Marinkovic J, Vukovic D. Association between body mass index and prevalence of multimorbidity: a cross-sectional study. *Public Health.* 2016;139:103-111.
41. Aranceta-Bartrina J, Perez-Rodrigo C, Alberdi-Aresti G, Ramos-Carrera N, Lazaro-Masedo S. Prevalence of general obesity and abdominal obesity in the Spanish adult population (aged 25-64 years) 2014-2015: The ENPE study. *Rev Esp Cardiol (Engl Ed).* 2016;69(6):579-587.
42. Weiderpass E, Botter E, Longenecker JC, et al. The prevalence of overweight and obesity in an adult Kuwaiti population in 2014. *Front Endocrinol (Lausanne)* 2019(10):449. <https://doi.org/10.3389/fendo.2019.00449>
43. Ministry of Health Saudi Arabia. Saudi Arabia Health Interview Survey Results 2013. Institute for health metrics & evaluation website. <http://www.healthdata.org/sites/default/files/files/Projects/KSA/Saudi-Health-Interview-Survey-Results.pdf>. Accessed August 16th, 2019.
44. National Health Survey: First Results, 2017-18. Australian Bureau of Statistics Web site. <https://www.abs.gov.au/AUSSTATS/abs@.nsf/DetailsPage/4364.0.55.0012017-18?OpenDocument>. Published December 12th 2018, updated 28th May 2019. Accessed July 1st, 2019.
45. Annual Update of Key Results 2017/8: New Zealand Health Survey. New Zealand Ministry of Health Web site. https://minhealthnz.shinyapps.io/nz-health-survey-2017-18-annual-data-explorer/_w_a04772ad/_w_b1a87660/#/explore-indicators. Published April 5th, 2019. Accessed June 4th, 2019.
46. National Survey of Noncommunicable Diseases in Seychelles 2013-14 (Seychelles Heart Study IV): methods and main findings. Ministry of Health Seychelles Web site. <http://www.health.gov.sc/wp-content/uploads/National-Survey-of-Noncommunicable-Diseases-in-Seychelles-2013-2014.pdf>. Published June 28th, 2015. Accessed August 25th, 2019.
47. Unwin N, Rose AMC, George KS, Hambleton IR, Howitt C. *The Barbados Health of the Nation Survey: Core Findings*. St Thomas, Barbados: Chronic Disease Research Centre. The University of the West Indies and the Barbados Ministry of Health; 2015.
48. Wagner KJP, Boing AF, Cembranel F, da Silva Boing AC, Subramanian SV. Change in the distribution of body mass index in Brazil: analysing the interindividual inequality between 1974 and 2013. *J Epidemiol Community Health.* 2019;73(6):544-548.
49. Hales C, Fryar C, Carroll M, Freedman D, Ogden C. Trends in obesity & severe obesity prevalence in US youth and adults by sex and age, 2007-2008 to 2015-2016. *JAMA.* 2018;319(16):1723-1725.
50. Canadian Health Measures Survey: adult body mass index—Health Canada Classification. Table 13-10-0323-01. Statistics Canada Web site. <https://www150.statcan.gc.ca/t1/tbl1/en/tv.action?pid=1310032301>. Published 2017. Accessed October 1st, 2019.
51. The Second Health and Nutritional Status Survey (NHANSS) 2014. Ministry of Health, Brunei Darussalam Web site. <http://www.ppkk.gov.bn/SitePages/NHANSS.aspx>. Published October 2015. Accessed September 25th, 2019.
52. Malaysian Institute for Public Health. *National Health and Morbidity Survey 2015 (NHMS 2015). Vol.II: Non-Communicable Diseases, Risk Factors, & Other Health Problems*. Kuala Lumpur, Malaysia: Ministry of Health; 2015.
53. Lobstein T, Jackson LR. *Tackling Obesities: Future Choices-International Comparisons of Obesity Trends, Determinants and Responses-Evidence Review (Part 1-Adults)*. Government Office for Science: London, UK; 2007.
54. OECD Health Statistics 2019, Definitions, Sources and Methods. Overweight or Obese Population—Measured data (aged 15 +). Organisation for Economic Cooperation and Development Web site. <http://www.oecd.org/els/health-systems/Table-of-Content-Metadata-OECD-Health-Statistics-2018.pdf>. Published 2019. Accessed July 2nd, 2019.
55. Eurostat, European Commission. *European Health Interview Survey (EHIS-Wave 2) Methodological Manual*. Luxembourg: Publications Office of the European Union; 2013.
56. European Health Examination Survey. <http://www.ehes.info>. Published 2019. Accessed September 25th, 2019.
57. Global Obesity Observatory, Obesity Prevalence Worldwide—Adults. World Obesity Federation Web site. <https://www.worldobesity.org/data/map/overview-adults>. Published 2018, updated 2019. Accessed September 27th, 2019.
58. Pineda E, Sanchez-Romero L-M, Brown M, et al. Forecasting future trends in obesity across Europe: The value of improving surveillance. *Obes Facts.* 2018;11(5):360-371.
59. National Health Survey: First Results, 2017-18. Appendix 2: Physical measurements in the 2017-18 National Health Survey. Australian Bureau of Statistics, Web site. <https://www.abs.gov.au/AUSSTATS/abs@.nsf/Latestproducts/4364.0.55.001Appendix22017-18?opendocument&tabname=Notes&prodno=4364.0.55.001&issue=2017-18&num=&view=>. Published March 26th, 2019. Accessed October 24th, 2019.
60. Scottish Health Survey 2017: Trend Tables. Scottish Government Web site. <https://www2.gov.scot/Topics/Statistics/Browse/Health/scottish-health-survey/Publications/Trend2017>. Published 2018. Accessed June 30th, 2019.
61. Keating C, Backholer K, Gearon E, et al. Prevalence of Class-I, Class-II and Class-III obesity in Australian adults between 1995 and 2011-12. *Obes Res Clin Pract.* 2015;9(6):553-562.
62. Ruhm CJ. Current and future prevalence of obesity and severe obesity in the United States. *Forum Health Econ Policy.* 2007;10(2):1-26.
63. Ogden CL, Carroll MD, McDowell MA, Flegal KM. Obesity Among Adults in the U.S.—No Statistically Significant Change Since 2003-2004. *NCHS Data Brief No 1*. Hyattsville, MD, USA. 2007.
64. Sturm R, Hattori A. Morbid obesity rates continue to rise rapidly in the United States. *Int J Obes (Lond).* 2013;37(6):889-891.
65. Krishna A, Razak F, Lebel A, Davey Smith G, Subramanian SV. Trends in group inequalities and interindividual inequalities in BMI in the United States, 1993-2012. *Am J Clin Nutr.* 2015;101(3):598-605.
66. Lebel A, Subramanian SV, Hamel D, Gagnon P, Razak F. Population-level trends in the distribution of body mass index in Canada, 2000-2014. *Can J Public Health.* 2018;109(4):539-548.
67. NCDs-STEPwise approach to noncommunicable disease risk factor surveillance (STEPS). World Health Organisation Web site. <http://www.who.int/ncds/surveillance/steps/riskfactor/en/>. Published 2017. Accessed October 8th, 2018.
68. NCDs-STEPS Country Reports. World Health Organisation Web site. <http://www.who.int/ncds/surveillance/steps/reports/en/>. Published 2019. Accessed October 10th, 2019.
69. The DHS Programme-Survey Search. Demographic Health Survey Web site. <https://dhsprogram.com/What-We-Do/survey-search.cfm?pgtype=main&SrvtP=country>. Published 2019. Accessed October 10th, 2019.

70. Popkin B, Adair LS, Ng SW. Global nutrition transition and the pandemic of obesity in developing countries. *Nutr Rev.* 2012;70(1):3-21.
71. National Child Measurement Programme, England 2017/18 school year. NHS Digital Web site. <https://files.digital.nhs.uk/9F/22AF4D/nati-chil-meas-prog-eng-2017-2018-rep.pdf>. Published 2018. Accessed October 7th, 2019.
72. Moussa OM, Erridge S, Chidambaram S, Ziprin P, Darzi A, Purkayastha S. Mortality of the severely obese: a population study. *Ann Surg.* 2019;269(6):1087-1091.
73. Inoue Y, Qin B, Poti J, Sokol R, Gordon-Larsen P. Epidemiology of obesity in adults: latest trends. *Curr Obes Rep.* 2018;7(4):276-288.
74. Khera AV, Chaffin M, Wade KH, et al. Polygenic prediction of weight and obesity trajectories from birth to adulthood. *Cell.* 2019;177(3):587-596.
75. Keating CL, Moodie ML, Bulfone L, Swinburn BA, Stevenson CE, Peeters A. Healthcare utilization and costs in severely obese subjects before bariatric surgery. *Obesity.* 2012;20(12):2412-2419.
76. Booth HP, Charlton J, Gulliford MC. Socioeconomic inequality in morbid obesity with body mass index more than 40 kg/m² in the United States and England. *Soc Sci Med Pop Health.* 2017;Dec;3:172-178.
77. Skinner AC, Ravanbakht SN, Skelton JA, Perrin EM, Armstrong SC. Prevalence of obesity and severe obesity in US children, 1999–2016. *Pediatrics.* 2018;141(3):e20173459.
78. Scottish Health Survey 2017: Trend Tables. Scottish Government Web site. <https://www2.gov.scot/Topics/Statistics/Browse/Health/scottish-health-survey/Publications/Trend2017>. Published 2018. Accessed 30th June, 2019.
79. National Institutes of Health. Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults: The Evidence Report. NIH Publication No. 98-4083. https://www.nhlbi.nih.gov/files/docs/guidelines/ob_gdlns.pdf. Published 1998. Accessed October 6th, 2019.
80. Skea ZC, Aceves-Martins M, Robertson C, De Bruin M, Avenell A. Acceptability and feasibility of weight management programmes for adults with severe obesity: a qualitative systematic review. *BMJ Open.* 2019;9(9):e029473.
81. Tanamas SK, Lean MEJ, Combet E, Vlassopoulos A, Zimmet PZ, Peeters A. Changing guards: time to move beyond body mass index for population monitoring of excess adiposity. *QJM.* 2016;109(7):443-446.
82. Han TS, Al-Gindan YY, Govan L, Hankey CR, Lean MEJ. Associations of BMI, waist circumference, body fat, and skeletal muscle with type 2 diabetes in adults. *Acta Diabetol.* 2019;56(8):947-954.
83. Han TS, Al-Gindan YY, Govan L, Hankey CR, Lean MEJ. Associations of body fat and skeletal muscle with hypertension. *J Clin Hypertens.* 2019;21(2):230-238.
84. Al-Gindan YY, Hankey C, Govan L, Gallagher D, Heymsfield SB, Lean MEJ. Derivation and validation of simple equations to predict total muscle mass from simple anthropometric and demographic data. *Am J Clin Nutr.* 2014;100(4):1041-1051.
85. Freedman DS, Lawman HG, Skinner AS, McGuire LC, Allison DB, Ogden CL. Validity of the WHO cutoffs for biologically implausible values of weight, height, and BMI in children and adolescents in NHANES from 1999 through 2012. *Am J Clin Nutr.* 2015;102(5):1000-1006.
86. Booth H, Prevost AT, Gulliford MC. Impact of body mass index on prevalence of multimorbidity in primary care: cohort study. *Fam Pract.* 2014;31(1):38-43.
87. Moussa OM, Ardissino M, Kulatilake P, et al. Effect of body mass index on depression in a UK cohort of 363 037 obese patients: a longitudinal analysis of transition. *Clin Obes.* 2019;9(3):e12305. <https://doi.org/10.1111/cob.12305>
88. Hajek A, Konig H-H. The curvilinear effect of BMI on functional health—the evidence of the long-running German aging survey. *Obes Facts.* 2017;10(3):252-260.
89. Tremmel M, Gerdtham U-G, Nilsson P, Saha S. Economic Burden of Obesity: A Systematic Literature Review. *Int J Environ Res Public Health.* 2017;14(4):435. <https://doi.org/10.3390/ijerph1404035>
90. Gray L, MacDonald C. Morbid obesity in disasters: bringing the “conspicuously invisible” into Focus. *Int J Environ Res Public Health.* 2016;13(10):E1029. <https://doi.org/10.3390/ijerph13101029>
91. Rosenfeld HE, Limb R, Chan P, Fitzgerald M, Bradley WPL, Rosenfeld JV. Challenges in the surgical management of spine trauma in the morbidly obese patient: a case series. *J Neurosurg Spine.* 2013;19(1):101-109.
92. Cowdell F, Radley K. What do we know about skin-hygiene care for patients with bariatric needs? Implications for nursing practice. *J Adv Nurs.* 2014;70(3):543-552.
93. Aberdeen City Council. Local Housing Strategy 2018-2023. <https://www.aberdeencity.gov.uk/services/housing/local-housing-strategy>. Published 2018. Accessed September 17th, 2019.
94. Gyi D, Masson A, Hignett S. Plus size and obese workers: anthropometry estimates to promote inclusive design. *Ergonomics.* 2019;62(9):1234-1242.
95. Peters LD. You are what you wear: how plus-size fashion figures in fat identity formation. *Fash Theory.* 2015;18(1):45-71.
96. Flaherty GT, Geoghegan R, Brown IG, Finucane FM. Severe obesity as a barrier to international travel: a qualitative analysis. *J Travel Med.* 2019;26(3):taz018. <https://doi.org/10.1093/jtm/taz018>
97. Keaver L, Xu B, Jaccard A, Webber L. Morbid obesity in the UK: a modelling projection study to 2035. *Scand J Public Health.* 2018 Aug 30;140349481879481. <https://doi.org/10.1177/1403494818794814>
98. Wu F, Guo Y, Chatterji S, et al. Common risk factors for chronic non-communicable diseases among older adults in China, Ghana, Mexico, India, Russia and South Africa: the study on global AGEing and adult health (SAGE) wave 1. *BMC Public Health.* 2015;15(1):88. <https://doi.org/10.1186/s12889-015-1407-0>
99. OECD: Overweight or obese population (indicator). <https://data.oecd.org/healthrisk/overweight-or-obese-population.htm>. Published 2019. Accessed October 22nd, 2019.

How to cite this article: Williamson K, Nimegeer A, Lean M. Rising prevalence of BMI ≥ 40 kg/m²: A high-demand epidemic needing better documentation. *Obesity Reviews.* 2020;21:e12986. <https://doi.org/10.1111/obr.12986>

APPENDIX A

Search strategy and terms, last search undertaken on 28 September 2019.

Medline (Ovid) and Embase

(National health survey or national population surveillance or national prevalence or epidemiological survey).tw

(BMI or body mass index or obese*).tw

Limited to humans, (all adults (19 plus years)), English language, 2016 – current, All types of publication

(Child or children or adolescents or baby). Tw records remove