



Similar Replicative Fitness Is Shared by the Subtype B and Unique BF Recombinant HIV-1 Isolates that Dominate the Epidemic in Argentina

Andrea E. Rubio¹, Awet Abraha², Crystal A. Carpenter², Ryan M. Troyer², Ángel L. Reyes-Rodríguez², Horacio Salomon¹, Eric J. Arts², Denis M. Tebit^{2*}✉

1 Centro Nacional de Referencia para el SIDA, Departamento de Microbiología, Facultad de Medicina Universidad de Buenos Aires, Buenos Aires, Argentina, **2** Division of Infectious Diseases and HIV Medicine, Case Western Reserve University, Cleveland, Ohio, United States of America

Abstract

The HIV-1 epidemic in South America is dominated by pure subtypes (mostly B and C) and more than 7 BF and BC recombinant forms. In Argentina, circulating recombinant forms (CRFs) comprised of subtypes B and F make up more than 50% of HIV infections. For this study, 28 HIV-1 primary isolates were obtained from patients in Buenos Aires, Argentina and initially classified into subtype B (n=9, 32.1%), C (n=1, 3.6%), and CRFs (n=18, 64.3%) using partial *pol* and *vpu-env* sequences, which proved to be inconsistent and inaccurate for these phylogenetic analyses. Near full length genome sequences of these primary HIV-1 isolates revealed that nearly all intersubtype BF recombination sites were unique and countered previous “CRF” B/F classifications. The majority of these Argentinean HIV-1 isolates were CCR5-using but 4 had a dual/mixed tropism as predicted by both phenotypic and genotypic assays. Comparison of the replicative fitness of these BF primary HIV-1 isolates to circulating B, F, and C HIV-1 using pairwise competitions in peripheral blood mononuclear cells (PBMCs) indicated a similarity in fitness of these BF recombinants to subtypes B and F HIV-1 (of the same co-receptor usage) whereas subtype C HIV-1 was significantly less fit than all as previously reported. These results suggest that the multitude of BF HIV-1 strains present within the Argentinean population do not appear to have gained replicative fitness following recent B and F recombination events.

Citation: Rubio AE, Abraha A, Carpenter CA, Troyer RM, Reyes-Rodríguez AL, et al. (2014) Similar Replicative Fitness Is Shared by the Subtype B and Unique BF Recombinant HIV-1 Isolates that Dominate the Epidemic in Argentina. PLoS ONE 9(4): e92084. doi:10.1371/journal.pone.0092084

Editor: Jean-Pierre Vartanian, Institut Pasteur, France

Received: November 29, 2013; **Accepted:** February 18, 2014; **Published:** April 11, 2014

Copyright: © 2014 Rubio et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by NIAID, NIH grant AI49170, Grant PICT-PRH 2008-120 from the National Scientific and Technical Research Agency (Argentina), and PIP 11420090100254 from CONICET. All virus work was performed in the biosafety level 2 and 3 facilities of the Case Western Reserve University Hospitals Center for AIDS Research (AI36219). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: dmt6u@virginia.edu

✉ Current address: Myles Thaler Center for AIDS and Human Retrovirus Research, Department of Microbiology, Immunology and Cancer Biology, University of Virginia, Charlottesville, Virginia, United States of America

Introduction

Human immunodeficiency virus type 1 (HIV-1) evolved from the simian immunodeficiency virus (SIVcpz) predominant in the subspecies of chimpanzee, *Pan troglodytes troglodytes* (Ptt) [1–4]. The transmission events from apes to humans occurred on at least four separate occasions which is represented by phylogenetic distinct groups termed M (major), O (outlier), N (non-M, non-O) and P. HIV-1 group M is responsible for the global pandemic shown clearly by its divergence into nine “pure” clades or subtypes (A–D, F–H, J and K), 63 circulating recombinant forms (CRF) (www.hiv.lanl.gov) and a myriad of unique recombinant forms (URF). Globally, CRFs account for about 10–20% of all new HIV-1 infections while URFs are responsible for over 30% of infections in regions where two or more subtypes co-circulate [2,3]. Intersubtype recombinants are clearly contributing to HIV-1 evolution and may ultimately result in complete disappearance of the “pure” HIV-1 subtypes [5], which in fact originated from earlier SIV/HIV recombination events [6].

The prevalence of HIV-1 subtypes and CRFs in the human population varies by geographical region and can be influenced by various socio-cultural factors as well as virus-host genetics (founder effects and host restrictive factors) [7]. Although subtype B is dominant in developed countries and is the most widely studied clade, non-subtype B HIV-1 are responsible for >90% of HIV infections [3,4]. Subtype C dominates in prevalence (>50% of infections worldwide) due in part to epidemics in India, China, and southern African countries (South Africa, Zimbabwe, Botswana, Zambia, Namibia, Malawi) [3,4]. In South America, the HIV-1 epidemic has been well characterized with subtype B HIV-1 found in the northern Pacific and Caribbean coastal countries (Venezuela, Colombia, Ecuador, Peru) whereas in the Southern cone (i.e. Argentina, Chile, Paraguay, and Uruguay) the epidemic is more diverse with a higher HIV-1 prevalence. In southern Brazil and Argentina [8,9] subtype B, C, and F as well as B/C and B/F recombinants are the most prevalent (Figure 1).

Intersubtype BF recombinants were first reported as a CRF (CRF12_BF) in Argentina and Uruguay in 2001 [10], followed by other novel BF CRFs in Brazil, Uruguay and Chile (i.e.




Figure 1. Changes in the prevalence of HIV-1 subtypes and recombinant forms during the past ten years in Buenos Aires, Argentina (white) and Rio Grande do Sol, Brazil (gray). Rio do Sol is located in the most southern part of Brazil while Buenos Aires is in north eastern Argentina. Both regions are located close to each other and share a common characteristic in that they harbor dominant and evolving recombinant forms. The prevalence numbers shown were obtained by combining the subtype distribution from published reports originating in these 2 regions. Only samples with a known sampling date were included in the analyses. doi:10.1371/journal.pone.0092084.g001

CRF17_BF, [10]; CRF28_BF, CRF29_BF [11]; CRF38_BF [12]; CRF39_BF, CRF40_BF [13], CRF44_BF; CRF46_BF [14]. CRF12_BF, CRF38_BF, and related BF recombinants appear to have originated in the late 1970s and early 1980s, dates which coincide with the beginning of the local epidemic [15,16]. In the HIV-infected population of Brazil (with the exception of southern Brazil; Figure 1), subtype B (71%) dominates over a lower prevalence of intersubtype BF recombinants (12%), subtype C (10%), subtype F (5%) and other subtypes/intersubtypes (2%) [17–19]. In contrast, intersubtype BF recombinant (56.7%) are found at high prevalence in HIV-infected Argentinians with subtype B (34.5%), F and other subtypes/intersubtypes (<5%) observed at low frequency [20–23] (Figure 1). About 50% of these Argentinian intersubtype BF-like viruses are related to, but distinct from

CRF12_BF [10,20,22]. Notably, most BF recombinants appear unique, but share a number of breakpoints with CRF12_BF [22]. Thus, previous or continuous super-infections involving CRF12_BF and/or other B, F, or BF recombinant viruses results in circulation of multiple BF recombinants which may be better defined as unique recombinant forms (URFs) rather than CRFs. This subtype/recombinant diversity and dynamics is quite unique to the Argentinean epidemic. In this study, the full genome sequences revealed that all of the 18 BF recombinant HIV-1 isolates from Buenos Aires, Argentina were URFs and not CRFs.

Based on the dominance of BFs in the Argentinean epidemic, Sanabani et al. (2006) [24] have proposed that these recombinants may have higher replicative fitness than “pure” subtypes B and F. However, aside from the fitness studies of CRF02_AG versus

subtype A or G, there are no other comparisons of replicative fitness between recombinant HIV-1 and their parental isolates. Two studies have examined potential phenotypic differences among BF viruses [25,26] and suggested that HIV-1 BF strains might have a higher replicative fitness over subtypes B and F HIV-1 isolates. In an effort to understand the high prevalence of intersubtype BF recombinants in Argentina, we compared *ex vivo* 'pathogenic' or replicative fitness of primary HIV-1 isolates of subtype B, F, C, and intersubtype BF recombinants derived from chronically infected patients. Fitness of these primary HIV-1 isolates was determined by direct head-to-head virus competitions in human peripheral blood cells (PBMCs) of HIV-negative donors. We discovered that the BF recombinants were of equal or lower replicative fitness compared with Argentinean B and reference F HIV-1 isolates.

Materials and Methods

Ethics Statement

Confidential one-on-one interviews were conducted onsite by health care workers. During these encounters, the study was explained and subjects were invited to participate. Only those subjects who were willing to participate were provided with a written informed consent, enrolled, and sampled. All procedures were approved by the Independent Research Ethics Committee, School of Medicine, University of Buenos Aires (CIEI-FM-UBA), Office for Human Research Protection (OHRP) reference numbers IORG#0004063 and IRB#00004817.

Patients, Sample Collection, Viral Load Determination and CD4 Lymphocytes Count

Patients were sampled from September 2002 to July 2004 from two medical facilities in Buenos Aires: namely the *Hospital de Agudos José María Ramos Mejía*, (n = 15), a voluntary counseling and testing center where routine HIV testing is performed and the "National Reference Center for AIDS, School of Medicine, University of Buenos Aires" (n = 13) where patients are monitored routinely for viral load and CD4 cell count as standard-of-care. Subjects attending these centers were from the city of Buenos Aires and nearest suburbs. Clinical records were reviewed to confirm that these patients were antiretroviral naïve.

Whole blood samples were collected and an aliquot was used for CD4 determination. Briefly, absolute count of CD4⁺ T-lymphocytes (cells/ul) from peripheral whole blood was determined in each HIV-1-infected patient by flow cytometry using a Coulter XL (Beckman) instrument (**Table 1**). Remaining blood was separated into plasma and buffy coat. Viral load in plasma was assessed by the branched DNA technology [VERSANT[®] HIV-1 RNA 3.0 Assay (b-DNA)] with a detection range of 50–500,000 HIV-1 RNA copies/ml (**Table 1**).

Virus Isolation, Co-receptor Usage and Titration

PBMCs isolated from HIV infected patients were co-cultivated with uninfected PBMCs from a healthy donor, as described previously [27]. Briefly, patient and uninfected cells were separately stimulated with 2 µg/ml phytohemagglutinin (PHA; Gibco, BRL) in RPMI medium for 2 to 3 days, and further maintained in complete RPMI medium containing 10% fetal bovine serum, 1 ng of recombinant interleukin (IL-2 invitrogen), 100 U/ml penicillin, and 100 µg/ml streptomycin (Cellgro)]. Subsequently, 1.5×10^6 stimulated cells from uninfected donors were co-cultured, in 24-well tissue culture plates, with 1.5×10^6 PBMC from HIV-1-infected patients. On day 4, 7, 11, 14 and 18, supernatant was harvested from each well to assess radioactive

reverse transcriptase (RT) activity. On day 7 and 14, 1.5×10^6 PHA-stimulated PBMC from the HIV-negative donor were thawed from frozen aliquots (derived from the same donor and blood draw) and added to each well/co-cultivation. Cell-free supernatants were harvested at peak viremia, as measured by a RT assay, centrifuged and stored at -80°C until subsequently needed. Twenty eight primary HIV-1 isolates from Argentina were obtained from this process. Two subtype F reference strains CA20 and VI850 were kindly provided by Dr. Guido Vanham (Institute of Tropical Medicine in Antwerp, Belgium). The other subtype reference strains (C-97ZA003, C-96USNG58, C-93MW959, F1-093BR020_1) were obtained from the National Institutes of Health AIDS Research and Reference Reagent Program (Table 1).

To determine co-receptor usage, U87 cell lines (human glioma cell line) expressing CD4 and either CCR5 or CXCR4 chemokine receptor (obtained from D. Littman and the AIDS Research and Reagent Program) were maintained in Dulbecco Modified Eagle Medium (DMEM; Cellgro) supplemented with 15% FBS, 100 U/ml penicillin, 100 mg of streptomycin, and 300 µg/ml geneticin (G418; Gibco-BRL) and 1 µg/ml puromycin (Life Technologies, Inc.) to retain receptor and co-receptor expression, respectively. {-Deng, 1996 5706/id} Co-receptor usage was then assessed by infecting 2.5×10^3 U87.CD4 cells expressing either CCR5 or CXCR4, in parallel, with each viral stock, in 48-well plates, as described previously [28]. Virus production was determined by assaying RT activity in the cell-free supernatants at days 3, 5, 7 and 9. Finally, CXCR4-usage was confirmed by infection of the HTLV-1-transformed MT2 cell line as described [29] (**Tables 1 and 2**).

The tissue culture infectious dose for 50% infectivity (TCID₅₀) values were determined by serially diluting each stock of HIV-1 isolate in triplicate and infecting 10^5 stimulated PBMCs in a 96-well flat-bottom plate. Infectivity in each well was tested by the radiolabeled RT assay [28], and the TCID₅₀ assay end point was determined on day 10. TCID₅₀ values was calculated as infectious units per milliliter (IU/ml) using the Reed and Muench method [30].

DNA Extraction, PCR, Sequencing and Phylogenetic Analysis

DNA was extracted from PBMC of each HIV-1-infected patient with the QIAamp DNA Blood Mini Kit (Qiagen). Nested PCR was performed on extracted DNA to obtain several non-contiguous fragments spanning various regions of the HIV-1 genome or the full length genome of isolates. Primers used for these amplifications are available upon request. HIV-1 PCR products were purified using the Qiaquick DNA purification kit (Qiagen) and then sequenced in both directions using an ABI PRISM 3100 Genetic Analyzer (Applied Biosystems). Chromatograms were manually edited for alignment by using Sequencher 4.10.1 Software (Gene Codes) and Vector NTI suite.

Nucleotide sequences were aligned using Clustal X Multiple Alignment neighbor joining protocol version 7.0.5.3. Alignments included a representative set of known HIV-1 subtype reference sequences from the Los Alamos HIV Sequence Database. Neighbor-joining (NJ) trees were constructed under the Kimura 2-parameter model with the MEGA3 program. The codon alignment was performed using Gene Cutter tool available at the Los Alamos National Laboratory website (www.hiv.lanl.gov/content/sequence/GENE_CUTTER/cutter.html). Each HIV sequence was analyzed for recombination patterns using Simplot version 3.5.1 [31] with a bootscan sliding window of 300 nucleotides in 20 nucleotide steps (100 bootstrap replicates 50%

Table 1. Characteristics of Argentinean HIV-1 primary isolates and reference strains.

Isolate name	Subtype Pol	Subtype Env	Age (years)	Viral load (copies/mL)	CD4 (cells/ μ L)	Coreceptor usage	Phenotype (NSI/SI/DM)	Country of Origin
B271	B	B	59	175,113	121	R5	NSI	Argentina
B524	B	B	36	409,639	87	R5	NSI	Argentina
B542	B	B	34	123,704	nd	R5	NSI	Argentina
B552	B	B	40	nd	171	X4	SI	Argentina
B563	B	B	34	46,719	nd	R5	NSI	Argentina
B735	B	B	27	63,237	238	R5	NSI	Argentina
B775	B	B	37	2,802	nd	X4	SI	Argentina
B872	B	B	55	>500,000	nd	R5	NSI	Argentina
B958	B	B	27	28,838	318	R5	NSI	Argentina
B F027	BF	BF	36	127,093	150	R5×4	DM	Argentina
BF116	BF	BF	38	112,521	236	R5	NSI	Argentina
BF118	BF	BF	42	344,685	255	R5	NSI	Argentina
BF119	BF	BF	36	26,921	nd	R5	NSI	Argentina
BF134	BF	BF	25	18,384	nd	R5×4	DM	Argentina
BF303	BF	BF	42	199,019	280	R5	NSI	Argentina
BF329	B	BF	48	239,552	217	R5	NSI	Argentina
BF333	BF	BF	51	339,539	78	R5	NSI	Argentina
BF456	BF	BF	34	7,485	nd	R5	NSI	Argentina
BF461	BF	BF	36	68,750	nd	R5	NSI	Argentina
BF549	BF	F	34	68,421	nd	R5	NSI	Argentina
BF555	BF	BF	36	176,805	nd	R5×4	DM	Argentina
BF559	B	BF	38	>500,000	nd	X4	SI	Argentina
BF640	BF	BF	29	130,651	228	R5×4	DM	Argentina
BF714	B	BF	34	3,825	nd	R5	NSI	Argentina
BF819	BF	BF	46	27,317	nd	X4	SI	Argentina
BF955	BF	BF	30	>500,000	47	R5	NSI	Argentina
BF992	BF	BF	32	451,371	519	R5	NSI	Argentina
C674	C	C	51	nd	nd	R5	NSI	Argentina
97ZA003	C	C	na	nd	nd	R5	NSI	South Africa
96USNG58	C	C	na	na	na	R5	NSI	Nigeria
93MMW959	C	C	na	na	na	R5	NSI	Malawi
VI850	F1	F1	na	na	na	R5	NSI	DRC
CA20	F2	F2	na	na	na	R5	NSI	Cameroon
93BR020_1	F1	F1	na	na	na	X4	SI	Brazil

DRC: Democratic Republic of Congo; na: not applicable; nd: not done.
doi:10.1371/journal.pone.0092084.t001

region [53]. Nested PCR products of the *env* gene were analyzed by the heteroduplex tracking assay (HTA) to determine the amount of virus produced in the dual infection/competition experiments [39]. The same genomic region (*env* C2-V3) was PCR-amplified from different subtype-specific HIV-1 *env* clones [subtype A RW020 and SF170, subtype C BR025, subtype E TH22 and CAR7] [41] for use as DNA probes. However, in this case, the E80 primer was radiolabeled with T4 polynucleotide kinase and 2uCi of 5'-[γ^{32} P] ATP, prior to amplification, as described [39]. At least two DNA probes in separate HTAs were used to determine relative virus production in each dual infection. Probes and PCR products were mixed, denatured and annealed on a 6% polyacrylamide gel. Heteroduplexes corresponding to each virus in a given competition were quantified using a Molecular Imager FX (Bio-Rad) imager. The final ratio of the two viruses produced in a dual infection was estimated by comparing the virus production in the competition to the virus production in the mono-infection. Production of individual HIV-1 isolates in a dual infection (f_0) divided by its initial proportion in the inoculum (i_0) is referred to as the relative fitness ($w = f_0/i_0$). Thus, the ratio of the relative fitness values of each HIV-1 variant in the competition is a measure of the fitness difference (W_D) between both HIV-1 strains ($W_D = W_M/W_L$), where W_M and W_L correspond to the relative fitness of the more and less fit viruses, respectively [39].

Statistical Analysis

Statistical analyses were performed using GraphPad Prism 4 (GraphPad Software, USA). Two-tailed Student's *t* tests were used to compare patients' mean age. Two-tailed Mann-Whitney tests were used to compare viral load values and CD4 cell counts. Two-sided Fisher's exact tests were used to compare winner/loser

proportions in competitions between two groups of viral isolates. All tests were considered significant when $p < 0.05$.

Results

Patient Clinical Characteristics

Twenty-eight HIV isolates obtained from 28 patients at various stages of HIV disease were used in this study. The mean age of subjects infected with HIV-1 subtype B and BF-like recombinants was similar (mean age 38.7 and 37.7 years, respectively, $p = 0.1669$, Student's *t* test, unpaired, two-tailed) while the mean age for all 28 subjects was 38.1 (Table 1). Viral load values were also similar for both groups; subtype B and BF strains (1.2×10^5 and 1.4×10^5 copies/ml respectively) (Table 1). Comparison of CD4 cell counts between subtype B and BF-like suggested all subjects were in late stages of disease (187 and 230 cells/ μ l respectively, $p = 0.9399$, Mann-Whitney test, two-tailed) (Table 1). Furthermore, no significant association could be established between patient viral load values (VL) and viral tropism (VL: R5 vs. X4; see below), nor between VL and subtype (data not shown).

Because drug resistance mutations affect replicative fitness of viral isolates [61], all primary isolates were screened for the presence of drug resistance mutations in the *pol* gene using the Stanford Drug Resistance Database tool (<http://cpr.stanford.edu/cpr/>). Two of the 28 HIV-1 isolates harbored mutations associated with high level resistance to protease inhibitors (PIs) and to reverse transcriptase inhibitors (RTIs). Namely, isolates B524 carried mutations V82A, I84V, L90M in PR, and K103N, P225H in RT while isolate BF819 had mutations D30N, N88D in PR and E101E/K and M184V in RT (Table S1).




Figure 2. Genotypic Characterization of HIV-1 primary isolates from Argentina. Phylogenetic trees were constructed for (a) partial *pol* (2253–3749, HXB2) and (b) *vpu* to *env* (5968–9092, HXB2) of Argentinean primary isolates. Sequences were aligned using Clustal X and trees constructed by neighbor joining method under the Kimura 2-parameter model. (c) Sequences were aligned and bootscanned as described in materials and methods to determine recombination sites. Sequences in green represent the subtype F1 regions and in blue-subtype B. Regions in the genome that were not sequenced are represented by empty boxes. The gray shading represents those sequences used for the phylogenetic trees in 2a and b respectively.
doi:10.1371/journal.pone.0092084.g002

Phylogenetic Characterization of Argentinean HIV-1 Isolates

Sequencing of two genomic fragments from a single virus facilitates subtype classification, detection of recombination sites within the HIV genome, and mapping of these recombination sites to the common breakpoints found in the BF CRF. Neighbor-joining (NJ) phylogenetic trees of partial *pol* (2253–3749, HXB2 numbering) and *vpu-env* (5968–9092, HXB2 numbering) sequences of these 28 Argentinean HIV-1 isolates predicted 9 (32.1%) subtype Bs, 18 (64.3%) intersubtype BF recombinants, and 1 (3.2%) subtype C (Fig. 2a, b and shaded regions of Fig. 2c; **Table 1**). Six samples (BF329, BF549, BF559, BF640, BF714, BF992) had discordant phylogenetic clustering patterns in the *pol* compared to the *vpu-env* sequences suggesting that a recombination event occurred between or within these two genes (Fig. 2a and b). Considering earlier reports of diverse BF strains circulating in Argentina [22,42], we performed bootscan analyses on the *pol* and *vpu-env* sequences (see Materials and Methods). These algorithms confirmed the initial classification into B, F, and BF subtypes/recombinant forms as well as identified probable recombination sites, almost all of which were unique to the known BF CRFs (CRF12,17, 38, 39, 40, 42, 44, 46, and 47).

Near full length sequencing of 18 of these 28 samples confirmed results from partial *pol* and *env* sequencing, i.e. the BF recombinant patterns were different from the two main BF recombinants CRF12_BF and 17 (Fig. 2c) as well as the other BF CRFs that circulate in Argentina or other regions of South America (not shown in Fig. 2c). Only one BF recombinant, BF549, had a “pure” subtype F sequence for the *vpu* to *env* region whereas the other BF isolates had a mosaic *rev* and/or *env* composed of both subtype B and F sequences (Fig. 2c).

Isolation of Primary Isolates, Co-receptor Usage and V3-loop Properties

Twenty-eight primary HIV-1 strains were obtained by PBMC co-cultivation using blood samples derived from HIV-1 positive patients in Buenos Aires, Argentina (Table 1). Co-receptor usage was assessed by infecting U87.CD4 cells expressing either CCR5 or CXCR4. Of the 28 primary HIV-1 isolates, 20 could only replicate in U87.CD4.CCR5 cells (R5-tropic), 4 only infected U87CD4.CXCR4 cells, and 4 were dual/mixed tropic (R5/X4) (**Table 1 and 2**). Additionally, only B553, B775, BF559, BF819, BF027, BF134, BF555, and BF640 replicated in MT-2 cells, i.e. the same viruses designated X4 or R5/X4 using the U87.CD4 co-receptor usage assay. The X4 and R5/X4 HIV-1 isolates were derived from late stage disease and typically had viral loads ($>10^5$ copies/ml) and low CD4 cell counts (<200 cells/ml) (values not available for all patients at the time of sample collection). Two X4 isolates were subtype B and two, were BF recombinants whereas all four dual tropic viruses were BF isolates (**Table 1 and 2**).

The co-receptor usage of these 28 HIV-1 Argentinean isolates was confirmed by inputting the Env V3 amino acid loop sequence into various algorithms: the net amino acid charge, the 11/25 rule, Position-Specific Scoring Matrix (PSSM), and geno2pheno. It is important to note that PSSM and geno2pheno provide the highest predictive values for co-receptor usage with subtype B HIV-1 sequences as compared with other HIV-1 subtypes. The X4 and R5/X4 HIV-1 isolates had the highest positive net amino acid charge in the V3 loop when compared to that of the R5 viral isolates (mean X4 net charge = 6.75 vs. mean R5 net charge = 4.69, $p = 0.0006$, Student's *t* test, unpaired, two-tailed). Three of eight CXCR4-using viruses (B775, and BF559, and BF555) had a positively charged amino acid at position 11 or 25




Figure 3. Comparing the relative replicative fitness difference (W_D) of HIV-1 subtype B and BF recombinants from Argentina in direct competitions. Subtype B strains a) B271 b) B524, c) B542 d) B563 e) B735 f) B872 g) B958 were competed against CRF-BF-like isolates (BF116, BF118, BF119, BF303, BF329, BF333, BF456, BF461, BF549, BF714, BF955, BF992) in PHA-stimulated, IL2-treated PBMCs. The fitness difference represented on the Y axis ranges from 0.01 to 100. Bars represent the fitness of the isolates shown on the Y-axis. doi:10.1371/journal.pone.0092084.g003

(306 and 322 in HXB2 Env amino acid numbering), (Table 2). In subtype B, over 90% of the X4 or dual tropic HIV-1 isolates have a positively charged amino acid at positions 11 and 25. Furthermore, the R5 isolate B563 had an Arginine at position 25, which predicts CXCR4 usage with 80–90% specificity [35]. Finally, PSSM and geno2pheno correctly predicted co-receptor usage for 26 and 28 HIV-1 isolates. Only R5 BF992 and X4 BF819 were incorrectly predicted as X4 and R5 by PSSM (respectively) whereas R5-B563 and X4-BF819 were wrongly




Figure 4. Relative fitness difference (W_D) between Argentinean subtype B strains a) B271, b) B563 c) B735 d) B872, e) B958 and Subtype C strains C674 (Argentina), C5, C8, C9 (reference strains). Competitions were performed in PHA/IL2 treated PBMCs as in Figure 3. The fitness difference shown on the Y axis ranges from 0.01 to 100. Bars represent the fitness of the isolates shown on the Y-axis. doi:10.1371/journal.pone.0092084.g004

predicted as dual and X4 tropic (respectively) by geno2pheno (**Table 2**). None of these “subtype B” algorithms were absolute in predicting co-receptor usage as determined by phenotypic assays. Limitations in predicting co-receptor usage may relate to the subtype F V3 loop sequences bore by most BF isolates (Fig. 2b and c). An antigenically distinct subtype B variant has been described in Brazil, which has a GWGR rather than GPGR signature motif at the tip of the V3-loop. This rare variant is found in ~50% of Brazilian subtype B isolates and has been associated with slower disease progression [43]. Considering the overlap between the Argentinean and Brazilian HIV epidemics, we analyzed the diversity of the V3-loop. Only one of the Argentinean HIV sequence (BF134) had a GWGR motif, which was also the only BF strain with a subtype B Env gp120 coding region. The subtype F V3 sequences of the BF strains had either a GPGR or GPGQ sequence, the latter being more common for subtype F but infrequent for subtype B [44].

The N-linked glycosylation site, N301, is part of a highly conserved NNTR sequence found in the V3 loop of CCR5 tropic primary isolates [45]. All of the subtype B, F, and BF sequences had NNTR sequence with the exception of two X4 isolates, which harbored either NTIKQ (B552, with an isoleucine insertion) and

NKTQ (B775) (**Table 2**). Aside from N-linked sites at amino acid position 301, no significant differences were observed in the sequence lengths of any variable loop or in the number of N-linked glycosylation patterns in the V1–V5 regions with the Argentinean subtype B and BF isolates.

Relative Replicative Fitness of Primary Subtype B, C, F and Intersubtype BF Recombinant HIV-1 Isolates of Different Phenotypes

In order to determine the *ex vivo* pathogenic fitness of primary R5 and X4 subtype B, C and intersubtype BF recombinant isolates from Argentina, nearly 700 dual infection/competitions experiments were performed in PHA stimulated human PBMCs using the 28 primary HIV-1 isolates (described above) and 6 reference strains of subtypes C (n = 3); F1 (n = 2), and F2 (n = 1) (**Table 1**). Full pairwise competition experiments were performed with the 25 primary R5 isolates and the 9 primary X4- and X4/R5-tropic isolates (Figures 3–7). The fitness difference, W_D is defined as the fitness of a B isolate over the fitness of BF isolate or C isolate in PBMC competitions (Fig. 3 and 4, respectively). In Fig. 3, we show that the replicative fitness of each R5 subtype B isolate (y axis of panels A–G) is, on average, similar to each of the 12 BF isolates in




Figure 5. Comparing the relative replicative fitness difference (W_D) of HIV-1 sub-subtype F1 reference strains; (a) CA20 and (b) V1850 and Argentinean subtype B (B271, B524, B542, B563, B735 and B872) in direct competitions using PHA/IL2 PBMCs. Similarly, subtype F (c) CA20 and (d) V1850 were competed against BFs (BF116, BF118, BF119, BF303, BF456, BF549, BF714 and BF955). Bars represent the relative fitness of the viruses shown on the Y-axis.
doi:10.1371/journal.pone.0092084.g005

these pairwise competitions. Only one BF isolate (BF549) is consistently more fit than all the B isolates (Fig. 3a–g) and other BF strains (Fig. S1a). Intersubtype BF recombinants and subtype B isolates had nearly equal number of “wins” in direct head-to-head competitions (BF wins: 55.95%, $n=47$ while B wins: 44.05%, $n=37$, $p=0.1647$). However, as shown in Fig. 4, the subtype B and BF isolates clearly out-compete the subtype C HIV-1 isolates with C viruses winning only one of 22 head-to-head competitions ($p<0.0001$, two-sided Fisher’s exact test) confirming previous observations that subtype C strains are less fit than most HIV-1 group M strains [36,40,46]. The only Argentinean subtype C isolate in our panel, C674 was the least fit compared to the other 3 C HIV-1 isolates derived from South Africa, Nigeria and Malawi (Fig. 4).

HIV-1 subtype F1 is a parental subtype of BF recombinants. However, “pure” subtype F HIV-1 is now rare in Argentina as reflected by its absence in our set of Argentinean primary isolates and recent molecular epidemiology studies [16,42] (Fig. 1). We therefore included a CCR5-using subtype F1 primary isolate, CA20 from Cameroon and subtype F2, V1850 from the

Democratic Republic of Congo (DRC) for competitions against 6 Argentinean subtype B (Fig. 5a and b) and 8 BF strains (Fig. 5c and d). The sub-subtype F1 is most related to South American F isolates and the F segments in BF recombinant genomes. The F1 isolate, CA20 was of equal or slightly greater fitness than the subtype B and BF isolates whereas the F2 isolate, V1850 was slightly less fit and lost the majority of competitions against the subtype B and BF isolates from Argentina. Nonetheless, there was no significant difference in replicative fitness between any of these subtype B, F, or BF HIV-1 isolates.

The mean intra- and intersubtype relative values were plotted for each of the B and BF isolates (Fig. 6) to obtain a clear representation of relative replicative fitness in human PBMCs. In Fig. 6, the mean intra-subtype/-recombinant relative fitness values (y axis) represents the average fitness of each B (or BF) isolate against each of the other B (or BF) isolates. Inter-subtype/-recombinant (x-axis) represents the average fitness of each B (or BF) isolate against each of the BF (or B) isolate. When comparing the intra- and inter-group M and subtype C fitness of 29 primary HIV-1 isolates in a previous study, we clearly showed that the




Figure 6. Comparison of intra- and intersubtype fitness of R5 HIV-1 subtype B and various BF recombinants from Argentina. Mean relative fitness values were determined for each HIV-1 isolate from pairwise competitions between isolates of the same subtype (mean intrasubtype relative fitness) and of different subtypes (mean intersubtype relative fitness). The mean intrasubtype and intersubtype fitness values for each HIV-1 isolate were then plotted as the x and y coordinates. Subtype B viruses are shown as black filled squares while BF recombinants are indicated by a red filled square. The mean fitness of subtype B and BF are very similar and also indicated as red and black filled squares respectively. The insert figure represents the fitness of HIV-1 subtype C relative to group M strains as described in a previous study (Abraha et al, 2009). Subtype C strains (red squares and triangles) have low mean relative fitness and cluster together (orange oval shade) while group M strains (green dots and tetragon) are more fit and cluster together as well (green oval shade). doi:10.1371/journal.pone.0092084.g006

subtype C isolates were on average less fit than any group M isolate (e.g. subtype A, B, D, and CRF01_AE) (see insert panel A of Fig. 6). In contrast, the replicative fitness of the subtype B and

BF isolates from the inter-subtype/-recombinant comparisons overlapped with the intra-subtype fitness comparisons. In other words, some isolates such as BF549 had both high intra- and inter-




Figure 7. Comparing the relative replicative fitness difference (W_D) of syncytium inducing (SI) HIV-1 Argentinean subtype B (a) B552, (b) B775 and (c) reference subtype F, 93BR020 against BF Argentinean recombinants (BF134; BF559; BF819, BF027, BF555 and BF640) in direct competitions using PHA/IL2 PBMCs. Bars represent the fitness of the viruses shown on the Y-axis. doi:10.1371/journal.pone.0092084.g007

subtype/-recombinant fitness but on average the BF isolates were not more fit than B isolates or vice versa (see mean R5 B fitness vs mean R5 BF fitness in Fig. 6).

A more limited set of CXCR4-using subtype B (B552, B775) and BF (BF134, BF027, BF555, BF640) HIV-1 isolates were available from Argentinean patients for these fitness studies. These viruses as well as one subtype F X4 reference strain (93BR020) were used for pairwise competitions in PBMCs (Fig. 7, Fig. S1b). As described in Fig. 7, it appears that X4 B isolates had higher fitness than the BF HIV-1 isolates in direct head-to-head competitions (B wins 10 of 12 competitions, 83%, $p=0.0033$). However, the BF819 isolate is an X4 isolate carrying drug resistance mutations D30N and N88G in PR and M184V in RT. The lamivudine-resistant M184V mutation is associated with a fitness cost [47]. When excluding the BF819 isolate from these analyses, there was no statistical difference in replicative fitness between these CXCR4-using subtype B and BF isolates. The sole subtype F X4 isolate was more fit than all the BF recombinant and B isolates (not shown). In summary, a total of 40 competitions between the other two subtype F isolates, 9 subtype B isolates, and 18 BF recombinants show similar replicative fitness ($p=0.129$) and suggest that this X4 subtype F isolate, 93BR020 and the R5 BF549 isolates were outliers with high replicative fitness.

Regardless of subtype, X4 HIV-1 isolates usually replicate faster and to higher levels than R5 HIV-1 isolates in PBMC cultures [39,46]. However, as previously reported [36,39,46]), these dual-infection experiments in PBMC cultures do not represent “true” competitions as R5 and X4 isolates do not compete for the same susceptible cells. CCR5 and CXCR4 are typically expressed on different CD4+ T-cells (predominantly on memory and naïve, respectively) [48] or at different levels in macrophages (higher CCR5 over CXCR4 levels). In this study, two R5 subtype B and one R5 intersubtype BF isolate were competed against three X4 primary isolates (1 subtype B and 2 intersubtype BFs isolates). All X4 isolates out-produced R5 variants, in a subtype independent manner, further supporting the higher pathogenic fitness of X4 isolates over the R5 HIV-1 isolates in PHA-activated, IL2-maintained PBMC cultures (Data not shown).

Discussion

In many epidemics around the world, generation and transmission of intersubtype HIV-1 recombinants appears to outpace the expansion of the “pure” parental subtypes. For example, BC recombinants are now increasing in prevalence as compared to subtype B or C HIV-1 isolates in Rio Grande do Sol province in Brazil [49–51] (Fig. 1). BC recombinant forms (specifically CRF07 and 08) dominate over subtype B or C in China [52]. Finally, the HIV-1 epidemic in Argentina involves a complex mixture of BF recombinants expanding faster than subtype B infections [42] (Fig. 1). This increasing prevalence of recombinant forms might infer greater replicative fitness over the parental HIV-1 subtype in the epidemic. However, “fitness” in the HIV-1 epidemic is not necessarily comparable to *ex vivo* replicative fitness in PBMCs whereas virulence (i.e. rate of disease progression within a patient) is a direct correlate of replicative fitness [53]. With the exception of studies comparing the relative fitness of subtype A, G, and CRF02_AG HIV-1 isolates [54], [55] *ex vivo* replicative fitness of intersubtype HIV-1 recombinants have not been compared to their parental HIV-1 subtypes in primary human cells.

To obtain a preliminary understanding on how HIV-1 recombination in the epidemic impacts replicative fitness (and possibly pathogenesis), we compared the *ex vivo* fitness of reference subtype F, primary isolates of subtype B, and recombinant BF

HIV-1 from Argentina where a significant emergence of BF recombinants is recent (mid 1990s–early 2000s) and complex [42,56], (Fig. 1). Initial V3 loop genotypic analyses of HIV-1 in Argentina suggested that the epidemic was similar to that in North America where subtype B was and still is highly prevalent [57]. However, we now know that subtype F HIV-1 may have a longer history in Argentina, having been introduced between 1975–1980 [56]. Recent studies have shown that the HIV epidemic in Argentina is dominated by pure subtype B, various BF recombinant forms, and a rare occurrence of subtypes C and F. Accordingly, we were unable to identify a “pure” subtype F isolate among 28 HIV infected individuals which is consistent with recent reports of rare subtype F HIV-1 infections in Argentina [42,58]. Early in the Argentinean epidemic, subtype B was most common in men who have sex with men (MSM) and subtype BF among IDUs and the heterosexually-active population [59,60] but differential prevalence of subtypes/recombinants in these transmission groups has eroded over the past decade. Despite attempts to classify the BFs as CRFs, our analyses of 18 full and near full length BF genomes revealed that most BF breakpoints did not match the recombination sites used to classify CRF12_17, 38, 39, 40, 42, 44, 46, and 47. In the past, a majority of BF recombinants isolated from HIV-infected Argentineans have been classified as CRF12_BF and CRF17_BF based on partial pol sequences [17,22,42]. We are now analyzing all of the full length BF sequences in the Los Alamos HIV-1 sequence database ($n > 40$) and preliminary results suggest that the vast majority of the BF strains classified as a CRF might instead be URFs of recent B+F or BF+B co-infections, the latter being the most likely. We observed little evidence of an epidemic with stable BF CRFs in Buenos Aires. These findings are consistent with recent studies showing a decrease in CRF12_BF prevalence (69% to 46% from 1986–93 to 2001–2008) [42] and an increased prevalence of unique BF recombinant forms in Argentina (up to 50% infections by URF_BF) [17,42] (summarized in Fig. 1). The later introduction of subtype C in Argentina from Brazil has led to the identification of BC recombinants (which may have originated in Brazil) and of a complex B/C/F recombinant in one patient [61].

Does the predominance of BF recombinants suggest a better fitness of these strains? Factors such as founder effects, host restrictive factors as well as cultural/behavioral factors may affect differential global spread of HIV variants in the human population [3,4,62]. We are also adopting infectious disease models to understand HIV spread in the epidemic [63]. Lower virulence (related to reduced replicative fitness in PBMCs [64]) leads to longer asymptomatic infection and increased opportunity for transmission. Aside from an initial founder event(s), the other key factor in these models is transmission efficiency related to the amount of transferred virus (e.g. viral load in donor [65,66]) and the ability to establish new host infections (e.g. relative replicative capacity and transmission through mucosal/genital tissue [36,38,40]). The dominance of BF-like recombinants over subtypes B, C and F in Argentina may be related to some biological advantage such as better transmission efficiency coupled with weaker virulence over the parental “pure” subtypes B and F. A second but not exclusive factor relates to the timing of founder events into the Argentinean population, and possibly into different transmission groups [49]. As described above, with the continuous emergence of new BF recombinants, these different transmission groups (e.g. IV drug users, heterosexuals, and MSMs) have now mixed and the BF epidemic in Argentina is now driven by heterosexual transmissions. This study only suggests that the similar replicative fitness of B, F, or BF HIV-1 isolates may relate to similar rates of disease progression (as determined by increasing

viral loads and decreasing CD4 cell counts). Thus, we propose that the increased spread of BF isolates in the Argentinean epidemic may be related to better transmission opportunity and/or efficiency of BF over B or F HIV-1 isolates. We are currently testing transmission efficiency in our ex vivo transmission models involving infections of human penile, cervical, vaginal, and rectal explants.

After >700 head-to-head competitions between 9 B, 18 BF, 3 F, and 4 C HIV-1 isolates (28 from Argentina), it was clear that subtype B and BF HIV-1 isolates had similar replicative fitness in human PBMC cultures. Subtype C HIV-1 isolates were less fit than all of the 27 B, BF, and F HIV-1 isolates, which remained consistent with multiple studies showing poor replicative fitness of subtype C HIV-1 compared to any other group M HIV-1 isolate (derived from >2000 pairwise competitions employing over 50 primary HIV-1 isolates) [36,40,46]. Our studies and those of others have suggested that HIV-1 subtype F isolates may be of slightly higher replicative fitness than the majority of group M isolates. The proposed higher replicative capacity of F and BF HIV-1 strains is based on specific gene sequence or other genetic factors. One study suggested that the subtype F long terminal repeat (LTR) showed greater transcriptional activation than the LTRs of subtype B [25], possibly leading to higher levels of virus production. Aulicino et al (2007) [58] examined 40 Argentinean *vpu* sequences of BF strains and revealed a high substitution rate ($\sim 11 \times 10^{-3}$ substitution per site per year) than that observed among subtype B strains [67]. These studies were limited to one or few B, BF, or F sequences and did not examine the effects of these gene regions/sequences in the context of the entire virus isolate. Nonetheless, both studies provide some evidence for increased subtype F fitness over subtype B HIV-1 isolates.

Based on the cumulative data from over 5000 pairwise competitions in PBMCs [36,40,46], we now rank replicative fitness of subtypes and dominant CRFs as follows: $B = D = F > CRF02_AG \geq A = CRF01_AE >> C$. In addition there is a strong correlation between relative replicative fitness in PBMCs and virulence in humans. Interestingly, the subtypes of highest replicative fitness (B, D, and F) are also the viruses expanding the slowest in numerous regional pandemics resulting in apparent shifts in the dominant subtypes, e.g. CRF01_AE or URFs over subtype B in Thailand, subtype C-containing URF and subtype C over subtype B in Brazil, and BF URFs over subtype B and F in Argentina. We suspect that these subtype/CRFs/URFs of lower virulence are expanding due to greater opportunity for transmission and retention of high transmission efficiency. Preliminary studies suggest that all HIV-1 subtypes in group M have similar transmission fitness when employing human genital tissue as a model for primary infection. As with the near extinction of HIV-1 subtype F in the Argentinean HIV-1 epidemic over the past 40 years, the dominance of subtype F HIV-1 over other subtypes has also decreased in Romania [68] and is almost absent in Cameroon (5% in 1999); [27] despite the moderate frequency of subtype F1 (17% of all HIV infections) during the late 80's and early 90's [69]. Increase in prevalence and new infections by a specific subtype may simply be due to founder events in a specific transmission group. Nonetheless, we consistently observe that HIV-1 subtypes

(D, F, G, and CRF01_AE) that have decreased in prevalence (in Uganda, Romania/Argentina, Cameroon, and Thailand, respectively) also have the highest ex vivo replicative fitness.

Conclusion

The present study suggests that BF recombinants circulating in Buenos Aires, Argentina have multiple recombinant structures which most likely resulted from continuous dual/superinfection between B and BF. Further, similar replicative fitness between subtypes B and BF HIV-1 isolates implies that recombination of the parental HIV-1 isolates does not necessarily increase replicative fitness. Initial dual infection or superinfection may result in higher fitness based on immune escape, host adaptation, and/or increase replicative fitness. Subsequent transmission of the recombinant form (e.g. BF) may select for factors associated with high transmission efficiency, which appear to be different than those factors governing replicative fitness in PBMCs. It is worth noting that CRF02_AG, another dominant recombinant was found to be more fit than its parental subtypes A and G, [54,55] suggesting that this higher replicative fitness may be associated with its dominance in the African HIV epidemic. However, the origin of CRF02_AG (or sub-subtype A3) dates back to the start of the HIV-1 pandemic in the Congo basin whereas our BF strains may have originated from very recent dual or super infections in Argentina [15]. Following approximately 50 years of evolution, the replicative fitness of CRF02_AG may reflect the changes derived from the original recombination as well as the accumulation of stable substitutions within the genome. With the analyses of the BF recombinants, evolution may be limited to just one or few transmission events.

Supporting Information

Figure S1 Fitness difference for pairwise competition with R5 and X4 HIV-1 isolates. Fitness differences (WD) for pairwise competition with A) R5 and B) X4 HIV-1 isolates are shown as the relative fitness of the “column” isolate over the “row” isolate. Gray boxes represent intra-subtype fitness differences while unshaded boxes represent inter-subtype fitness differences among HIV-1 patient's isolates and reference strains from subtype B (bold italic), intersubtype BF recombinants (bold), subtype C (italic) and subtype F (normal). (XLSX)

Table S1 Drug resistance mutations: calibrated population resistance tool from Stanford University was used to evaluate the presence of drug resistance mutations in *pol* gene sequence from each viral isolate (2253–3749, HXB2 numbering). (XLSX)

Author Contributions

Conceived and designed the experiments: HS EJA. Performed the experiments: AER AA CAC RMT ÁLRR DMT. Analyzed the data: AER DMT EJA. Contributed reagents/materials/analysis tools: HS EJA. Wrote the paper: AER EJA DMT.

References

- Gao F, Bailes E, Robertson DL, Chen Y, Rodenburg CM, et al. (1999) Origin of HIV-1 in the chimpanzee Pan troglodytes troglodytes. *Nature* 397: 436–441. 10.1038/17130 [doi].
- Keele BF, Van HF, Li Y, Bailes E, Takehisa J, et al. (2006) Chimpanzee reservoirs of pandemic and nonpandemic HIV-1. *Science* 313: 523–526. 1126531 [pii];10.1126/science.1126531 [doi].
- Tebit DM, Arts EJ (2011) Tracking a century of global expansion and evolution of HIV to drive understanding and to combat disease. *Lancet Infect Dis* 11: 45–56. S1473-3099(10) 70186-9 [pii];10.1016/S1473-3099(10)70186-9 [doi].
- Hemelaar J (2012) The origin and diversity of the HIV-1 pandemic. *Trends Mol Med* 18: 182–192. S1471-4914(11)00210-3 [pii];10.1016/j.molmed.2011.12.001 [doi].

5. Quinones-Mateu ME, Gao Y, Ball SC, Marozsan AJ, Abraha A, et al. (2002) In vitro intersubtype recombinants of human immunodeficiency virus type 1: comparison to recent and circulating in vivo recombinant forms. *J Virol* 76: 9600–9613.
6. Bailes E, Gao F, Bibollet-Ruche F, Courgnaud V, Peeters M, et al. (2003) Hybrid origin of HIV in chimpanzees. *Science* 300: 1713. 10.1126/science.1080657 [doi];300/5626/1713 [pii].
7. Tebit DM, Nankya I, Arts EJ, Gao Y (2007) HIV diversity, recombination and disease progression: how does fitness “fit” into the puzzle? *AIDS Rev* 9: 75–87.
8. Guimaraes ML, Dos Santos MA, Loureiro R, Galvao-Castro B, Morgado MG (2002) High frequency of recombinant genomes in HIV type 1 samples from Brazilian southeastern and southern regions. *AIDS Res Hum Retroviruses* 18: 1261–1269. 10.1089/088922202320886307 [doi].
9. Montano SM, Sanchez JL, Laguna-Torres A, Cuchi P, Avila MM, et al. (2005) Prevalences, genotypes, and risk factors for HIV transmission in South America. *J Acquir Immune Defic Syndr* 40: 57–64. 00126334-200509010-00010 [pii].
10. Carr JK, Avila M, Gomez CM, Salomon H, Hierholzer J, et al. (2001) Diverse BF recombinants have spread widely since the introduction of HIV-1 into South America. *AIDS* 15: F41–F47.
11. De Sa Filho DJ, Sucupira MC, Caseiro MM, Sabino EC, Diaz RS, et al. (2006) Identification of two HIV type 1 circulating recombinant forms in Brazil. *AIDS Res Hum Retroviruses* 22: 1–13. 10.1089/aid.2006.22.1 [doi].
12. Ruchansky D, Casado C, Russi JC, Arbiza JR, Lopez-Galindez C (2009) Identification of a new HIV Type 1 circulating recombinant form (CRF38_BF1) in Uruguay. *AIDS Res Hum Retroviruses* 25: 351–356. 10.1089/aid.2008.0248 [doi].
13. Guimaraes ML, Eyer-Silva WA, Couto-Fernandez JC, Morgado MG (2008) Identification of two new CRF_BF in Rio de Janeiro State, Brazil. *AIDS* 22: 433–435. 10.1097/QAD.0b013e3282f74d0 [doi];00002030-200801300-00015 [pii].
14. Sanabani SS, Pastena ER, Neto WK, Martinez VP, Sabino EC (2010) Characterization and frequency of a newly identified HIV-1 BF1 intersubtype circulating recombinant form in Sao Paulo, Brazil. *Virology* 74: 1743–1748. 10.1016/j.virus.2010.07.014 [doi].
15. Dilemnia DA, Jones LR, Pando MA, Rabinovich RD, Damilano GD, et al. (2011) Analysis of HIV type 1 BF recombinant sequences from South America dates the origin of CRF12_BF to a recombination event in the 1970s. *AIDS Res Hum Retroviruses* 27: 569–578. 10.1089/AID.2010.0118 [doi].
16. Gomez CM, Avila M, Hierholzer J, Pando M, Martinez PL, M et al. (2002) Mother-to-child HIV type 1 transmission in Argentina: BF recombinants have predominated in infected children since the mid-1980s. *AIDS Res Hum Retroviruses* 18: 477–483. 10.1089/088922202317406619 [doi].
17. Gomez-Carrillo M, Pampuro S, Duran A, Losso M, Harris DR, et al. (2006) Analysis of HIV type 1 diversity in pregnant women from four Latin American and Caribbean countries. *AIDS Res Hum Retroviruses* 22: 1186–1191. 10.1089/aid.2006.22.1186 [doi].
18. de Souza AC, de Oliveira CM, Rodrigues CL, Silva SA, Levi JE (2008) Short communication: Molecular characterization of HIV type 1 BF pol recombinants from Sao Paulo, Brazil. *AIDS Res Hum Retroviruses* 24: 1521–1525. 10.1089/aid.2008.0089 [doi].
19. de Sa-Filho DJ, Soares MS, Candido V, Gagliani LH, Cavaliere E, et al. (2008) HIV type 1 pol gene diversity and antiretroviral drug resistance mutations in Santos, Brazil. *AIDS Res Hum Retroviruses* 24: 347–353. 10.1089/aid.2007.0203 [doi].
20. Thomson MM, Villahermosa ML, Vazquez-de-Parga E, Cuevas MT, Delgado E, et al. (2000) Widespread circulation of a B/F intersubtype recombinant form among HIV-1-infected individuals in Buenos Aires, Argentina. *AIDS* 14: 897–899.
21. Dilemnia DA, Gomez AM, Lourtau L, Marone R, Losso MH, et al. (2007) HIV type 1 genetic diversity surveillance among newly diagnosed individuals from 2003 to 2005 in Buenos Aires, Argentina. *AIDS Res Hum Retroviruses* 23: 1201–1207. 10.1089/aid.2007.0068 [doi].
22. Quarleri JF, Rubio A, Carobene M, Turk G, Vignoles M, et al. (2004) HIV type 1 BF recombinant strains exhibit different pol gene mosaic patterns: descriptive analysis from 284 patients under treatment failure. *AIDS Res Hum Retroviruses* 20: 1100–1107. 10.1089/aid.2004.20.1100 [doi].
23. Pando MA, Gomez-Carrillo M, Vignoles M, Rubio AE, dos Ramos Farias MS, et al. (2011) Incidence of HIV type 1 infection, antiretroviral drug resistance, and molecular characterization in newly diagnosed individuals in Argentina: A Global Fund Project. *AIDS Res Hum Retroviruses* 27: 17–23. 10.1089/aid.2010.0013 [doi].
24. Sanabani S, Kleine NW, Kalmar EM, Diaz RS, Janini LM, et al. (2006) Analysis of the near full length genomes of HIV-1 subtypes B, F and BF recombinant from a cohort of 14 patients in Sao Paulo, Brazil. *Infect Genet Evol* 6: 368–377. S1567-1348(06) 00005-0 [pii];10.1016/j.meegid.2006.01.003 [doi].
25. Turk G, Carobene M, Monczor A, Rubio AE, Gomez-Carrillo M, et al. (2006) Higher transactivation activity associated with LTR and Tat elements from HIV-1 BF intersubtype recombinant variants. *Retrovirology* 3: 14. 1742-4690-3-14 [pii];10.1186/1742-4690-3-14 [doi].
26. Carobene MG, Rodriguez RC, De Candia CA, Turk G, Salomon H (2009) In vitro dynamics of HIV-1 BF Intersubtype Recombinants genomic regions involved in the regulation of gene expression. *Virology* 6: 107. 1743-422X-6-107 [pii];10.1186/1743-422X-6-107 [doi].
27. Tebit DM, Zekeng L, Kaptue L, Salminen M, Krausslich HG, et al. (2002) Genotypic and phenotypic analysis of HIV type 1 primary isolates from western Cameroon. *AIDS Res Hum Retroviruses* 18: 39–48. 10.1089/088922202573394709 [doi].
28. Torre VS, Marozsan AJ, Albright JL, Collins KR, Hartley O, et al. (2000) Variable sensitivity of CCR5-tropic human immunodeficiency virus type 1 isolates to inhibition by RANTES analogs. *J Virol* 74: 4868–4876.
29. Karlsson A, Parsmyr K, Aperia K, Sandstrom E, Fenyo EM, et al. (1994) MT-2 cell tropism of human immunodeficiency virus type 1 isolates as a marker for response to treatment and development of drug resistance. *J Infect Dis* 170: 1367–1375.
30. Salzman AJ, Fraundorf E, Abraha A, Baird H, Moore D, et al. (2004) Relationships between infectious titer, capsid protein levels, and reverse transcriptase activities of diverse human immunodeficiency virus type 1 isolates. *J Virol* 78: 11130–11141. 10.1128/JVI.78.20.11130-11141.2004 [doi];78/20/11130 [pii].
31. Lole KS, Bollinger RC, Paranjape RS, Gadkari D, Kulkarni SS, et al. (1999) Full-length human immunodeficiency virus type 1 genomes from subtype C-infected seroconverters in India, with evidence of intersubtype recombination. *J Virol* 73: 152–160.
32. Salminen MO, Carr JK, Burke DS, McCutchan FE (1995) Identification of breakpoints in intergenotypic recombinants of HIV type 1 by bootscanning. *AIDS Res Hum Retroviruses* 11: 1423–1425.
33. Rozanov M, Plikat U, Chappay C, Kochergin A, Tatusova T (2004) A web-based genotyping resource for viral sequences. *Nucleic Acids Res* 32: W654–W659. 10.1093/nar/gkh419 [doi];32/suppl_2/W654 [pii].
34. Schultz AK, Zhang M, Leitner T, Kuiken C, Korber B, et al. (2006) A jumping profile Hidden Markov Model and applications to recombination sites in HIV and HCV genomes. *BMC Bioinformatics* 7: 265. 1471-2105-7-265 [pii];10.1186/1471-2105-7-265 [doi].
35. Low AJ, Swenson LC, Harrigan PR (2008) HIV coreceptor phenotyping in the clinical setting. *AIDS Rev* 10: 143–151.
36. Abraha A, Nankya IL, Gibson R, Demers K, Tebit DM, et al. (2009) CCR5- and CXCR4-tropic subtype C human immunodeficiency virus Type 1 isolates have a lower level pathogenic fitness than other dominant group M subtypes: Implications for the Epidemic. *J Virol* 83: 5592–5605. JVI.02051-08 [pii];10.1128/JVI.02051-08 [doi].
37. Tebit DM, Lobritz M, Lalonde M, Immonen T, Singh K, et al. (2010) Divergent evolution in reverse transcriptase (RT) of HIV-1 group O and M lineages: impact on structure, fitness, and sensitivity to non-nucleoside RT inhibitors. *J Virol* 84: 10099-10 [pii];10.1128/JVI.00991-10 [doi].
38. Arien KK, Troyer RM, Gali Y, Colebunders RL, Arts EJ, et al. (2005) Replicative fitness of historical and recent HIV-1 isolates suggests HIV-1 attenuation over time. *AIDS* 19: 1555–1564. 00002030-200510140-00001 [pii].
39. Quinones-Mateu ME, Ball SC, Marozsan AJ, Torre VS, Albright JL, et al. (2000) A dual infection/competition assay shows a correlation between ex vivo human immunodeficiency virus type 1 fitness and disease progression. *J Virol* 74: 9222–9233.
40. Ball SC, Abraha A, Collins KR, Marozsan AJ, Baird H, et al. (2003) Comparing the ex vivo fitness of CCR5-tropic human immunodeficiency virus type 1 isolates of subtypes B and C. *J Virol* 77: 1021–1038.
41. Delwart EL, Sheppard HW, Walker BD, Goudsmit J, Mullins JI (1994) Human immunodeficiency virus type 1 evolution in vivo tracked by DNA heteroduplex mobility assays. *J Virol* 68: 6672–6683.
42. Aulicino PC, Bello G, Guimaraes ML, Ruchansky D, Rocco C, et al. (2011) Longitudinal analysis of HIV-1 BF1 recombinant strains in vertically infected children from Argentina reveals a decrease in CRF12_BF pol gene mosaic patterns and high diversity of BF unique recombinant forms. *Infect Genet Evol* 11: 349–357. S1567-1348(10) 00309-6 [pii];10.1016/j.meegid.2010.11.008 [doi].
43. Diaz RS, Leal E, Sanabani S, Sucupira MC, Tanuri A, et al. (2008) Selective regimes and evolutionary rates of HIV-1 subtype B V3 variants in the Brazilian epidemic. *Virology* 381: 184–193. S0042-6822(08)00530-8 [pii];10.1016/j.virus.2008.08.014 [doi].
44. Coetzer M, Cilliers T, Ping LH, Swanstrom R, Morris L (2006) Genetic characteristics of the V3 region associated with CXCR4 usage in HIV-1 subtype C isolates. *Virology* 356: 95–105. S0042-6822(06)00507-1 [pii];10.1016/j.virus.2006.07.030 [doi].
45. Clevestig P, Pramanik L, Leitner T, Ehrnst A (2006) CCR5 use by human immunodeficiency virus type 1 is associated closely with the gp120 V3 loop N-linked glycosylation site. *J Gen Virol* 87: 607–612. 87/3/607 [pii];10.1099/vir.0.81510-0 [doi].
46. Arien KK, Abraha A, Quinones-Mateu ME, Kestens L, Vanham G, et al. (2005) The replicative fitness of primary human immunodeficiency virus type 1 (HIV-1) group M, HIV-1 group O, and HIV-2 isolates. *J Virol* 79: 8979–8990. 79/14/8979 [pii];10.1128/JVI.79.14.8979-8990.2005 [doi].
47. Paredes R, Sagar M, Marconi VC, Hoh R, Martin JN, Parkin NT, et al. (2009) In vivo fitness cost of the M184V mutation in multidrug-resistant human immunodeficiency virus type 1 in the absence of lamivudine. *J Virol* 83: 2038–2043. JVI.02154-08 [pii];10.1128/JVI.02154-08 [doi].
48. Bleul CC, Wu L, Hoxie JA, Springer TA, Mackay CR (1997) The HIV coreceptors CXCR4 and CCR5 are differentially expressed and regulated on human T lymphocytes. *Proc Natl Acad Sci U S A* 94: 1925–1930.

49. Santos AF, Schrago CG, Martínez AM, Mendoza-Sassi R, Silveira J, et al. (2007) Epidemiologic and evolutionary trends of HIV-1 CRF31_BC-related strains in southern Brazil. *J Acquir Immune Defic Syndr* 45: 328–333. 10.1097/QAI.0b013e3180690d6a [doi].
50. Soares MA, de Oliveira T, Brindeiro RM, Díaz RS, Sabino EC, et al. (2003) A specific subtype C of human immunodeficiency virus type 1 circulates in Brazil. *AIDS* 17: 11–21. 10.1097/01.aids.0000042595.93174.a4 [doi].
51. Brígido LF, Nunes CC, Oliveira CM, Knoll RK, Ferreira JL, et al. (2007) HIV type 1 subtype C and CB Pol recombinants prevail at the cities with the highest AIDS prevalence rate in Brazil. *AIDS Res Hum Retroviruses* 23: 1579–1586. 10.1089/aid.2007.0102 [doi].
52. Wang W, Jiang S, Li S, Yang K, Ma L, et al. (2008) Identification of subtype B, multiple circulating recombinant forms and unique recombinants of HIV type 1 in an MSM cohort in China. *AIDS Res Hum Retroviruses* 24: 1245–1254. 10.1089/aid.2008.0095 [doi].
53. Arts EJ, Quinones-Mateu ME (2003) Sorting out the complexities of HIV-1 fitness. *AIDS* 17: 780–781. 10.1097/01.aids.0000050881.72891.32 [doi].
54. Konings FA, Burda ST, Urbanski MM, Zhong P, Nadas A, et al. (2006) Human immunodeficiency virus type 1 (HIV-1) circulating recombinant form 02_AG (CRF02_AG) has a higher in vitro replicative capacity than its parental subtypes A and G. *J Med Virol* 78: 523–534. 10.1002/jmv.20572 [doi].
55. Njai HF, Gali Y, Vanham G, Clybergh C, Jennes W, et al. (2006) The predominance of Human Immunodeficiency Virus type 1 (HIV-1) circulating recombinant form 02 (CRF02_AG) in West Central Africa may be related to its replicative fitness. *Retrovirology* 3: 40. 1742-4690-3-40 [pii];10.1186/1742-4690-3-40 [doi].
56. Bello G, Aulicino PC, Ruchansky D, Guimaraes ML, Lopez-Galindez C, et al. (2010) Phylodynamics of HIV-1 circulating recombinant forms 12_BF and 38_BF in Argentina and Uruguay. *Retrovirology* 7: 22. 1742-4690-7-22 [pii];10.1186/1742-4690-7-22 [doi].
57. Gomez CM, Piccardo C, Libonatti O (1992) [Molecular analysis of the principal neutralization epitope (V3 loop) of human immunodeficiency virus type 1 in Argentina]. *Rev Argent Microbiol* 24: 91–101.
58. Aulicino PC, Bello G, Rocco C, Romero H, Mangano A, et al. (2007) Description of the first full-length HIV type 1 subtype F1 strain in Argentina: implications for the origin and dispersion of this subtype in South America. *AIDS Res Hum Retroviruses* 23: 1176–1182. 10.1089/aid.2007.0038 [doi].
59. Avila MM, Pando MA, Carrion G, Peralta LM, Salomon H, et al. (2002) Two HIV-1 epidemics in Argentina: different genetic subtypes associated with different risk groups. *J Acquir Immune Defic Syndr* 29: 422–426.
60. Espinosa A, Vignoles M, Carrillo MG, Sheppard H, Donovan R, et al. (2004) Intersubtype BF recombinants of HIV-1 in a population of injecting drug users in Argentina. *J Acquir Immune Defic Syndr* 36: 630–636. 00126334-200405010-00012 [pii].
61. Pando MA, Eyzaguirre LM, Segura M, Bautista CT, Marone R, et al. (2006) First report of an HIV-1 triple recombinant of subtypes B, C and F in Buenos Aires, Argentina. *Retrovirology* 3: 59. 1742-4690-3-59 [pii];10.1186/1742-4690-3-59 [doi].
62. Arien KK, Vanham G, Arts EJ (2007) Is HIV-1 evolving to a less virulent form in humans? *Nat Rev Microbiol* 5: 141–151. nrmicro1594 [pii];10.1038/nrmicro1594 [doi].
63. Anderson RM, May RM (1992) *Infectious diseases of humans: dynamics and control*. Oxford; New York: Oxford University Press.
64. Troyer RM, Collins KR, Abraha A, Fraundorf E, Moore DM, et al. (2005) Changes in human immunodeficiency virus type 1 fitness and genetic diversity during disease progression. *J Virol* 79: 9006–9018. 79/14/9006 [pii];10.1128/JVI.79.14.9006-9018.2005 [doi].
65. Gray RH, Wawer MJ, Brookmeyer R, Sewankambo NK, Serwadda, et al. (2001) Probability of HIV-1 transmission per coital act in monogamous, heterosexual, HIV-1-discordant couples in Rakai, Uganda. *Lancet* 357: 1149–1153. S0140-6736(00)04331-2 [pii];10.1016/S0140-6736(00)04331-2 [doi].
66. Reynolds SJ, Makumbi F, Nakigozi G, Kagaayi J, Gray RH, et al. (2011) HIV-1 transmission among HIV-1 discordant couples before and after the introduction of antiretroviral therapy. *AIDS* 25: 473–477. 10.1097/QAD.0b013e3283437c2b [doi].
67. Wolfs TF, de Jong JJ, Van den Berg H, Tijnagel JM, Krone WJ, et al. (1990) Evolution of sequences encoding the principal neutralization epitope of human immunodeficiency virus 1 is host dependent, rapid, and continuous. *Proc Natl Acad Sci U S A* 87: 9938–9942.
68. Temereanca A, Ene L, Mehta S, Manolescu L, Duiculescu D, et al. (2013) Transmitted HIV drug resistance in treatment-naïve Romanian patients. *J Med Virol*. 10.1002/jmv.23572 [doi].
69. Nkengasong JN, Janssens W, Heyndrickx L, Franssen K, Ndumbe PM, et al. (1994) Genotypic subtypes of HIV-1 in Cameroon. *AIDS* 8: 1405–1412.