

Contents lists available at [ScienceDirect](#)

Data in Brief

journal homepage: www.elsevier.com/locate/dib

Data Article

Data set of dissolved major and trace elements from the lacustrine systems of Clearwater Mesa, Antarctica

Karina L. Lecomte^{a,b,*}, Cecilia V. Echegoyen^{a,b}, Paula A. Vignoni^{b,c,d},
 Kateřina Kopalová^e, Tyler J. Kohler^{e,1}, Silvia H. Coria^f,
 Juan M. Lirio^f

^a Centro de Investigaciones en Ciencias de la Tierra (CICTERRA), CONICET/Universidad Nacional de Córdoba, Av. Vélez Sarsfield, 1611, X5016CGA Córdoba, Argentina

^b Facultad de Ciencias Exactas Físicas y Naturales, Universidad Nacional de Córdoba, Av. Vélez Sarsfield, 1611, X5016CGA Córdoba, Argentina

^c Institute of Geosciences, Potsdam University, Karl-Liebknecht-Straße. 24-25, 14476, Potsdam-Golm, Germany

^d Climate Dynamics and Landscape Evolution, German Research Centre for Geoscience GFZ, Telegrafenberg, 14473, Potsdam, Germany

^e Department of Ecology, Faculty of Science, Charles University, Viničná 7, 128 44, Praha 2, Czech Republic

^f Instituto Antártico Argentino, 25 de Mayo 1143, San Martín, Prov. Buenos Aires, Argentina

ARTICLE INFO

Article history:

Received 19 November 2019

Revised 4 March 2020

Accepted 10 March 2020

Available online 19 March 2020

Keywords:

Water chemistry

Pristine environments

Dissolved trace elements

High latitude lakes

ABSTRACT

This article presents analytical observations on physicochemical parameters and major and trace element concentrations of water, ice, and sediment samples from the lake systems of Clearwater Mesa (CWM), northeast Antarctic Peninsula. Geochemical analyses include inductively coupled plasma mass spectrometry (ICP-MS) for cations and trace elements and ion chromatography for anions. Some figures are included (i.e. Piper and Gibbs diagrams) which indicate water classification type and rock-water interactions in CWM, respectively. It also contains PHREEQC software output, listing the chemical speciation for dissolved elements, Saturation Indexes (SI), and modelling outputs. Each lake SI are also illustrated in a figure. Finally, total organic and inorganic carbon (TOC and TIC, respectively) were determined for bottom lake sediments

* Corresponding author.

E-mail address: karina.lecomte@unc.edu.ar (K.L. Lecomte).

¹ Current Address: Stream Biofilm and Ecosystem Research Laboratory, School of Architecture, Civil and Environmental Engineering, École Polytechnique Fédérale de Lausanne, CH-1015, Lausanne, Switzerland

and marginal salt samples. This information will be useful for future research assessing the impacts of anthropogenic pollution and the effects of climate change, providing insights into naturally occurring geochemical processes in a pristine environment, and evaluating geochemical behaviour of dissolved elements in high-latitude hydrological systems. These data correspond to the research article "Dissolved major and trace geochemical dynamics in Antarctic Lacustrine Systems" [1].

© 2020 The Authors. Published by Elsevier Inc.
This is an open access article under the CC BY license.
(<http://creativecommons.org/licenses/by/4.0/>)

Specifications table

Subject	Earth and Planetary Sciences
Specific subject area	Earth-Surface Geochemical Processes
Type of data	Table Figure Other (Software Output)
How data were acquired	Major cations and trace elements were determined by inductively coupled plasma mass spectrometry (PerkinElmer Sciex ELAN 9000 ICP/MS, PerkinElmer Nexion, Thermo icapQ or Agilent 7700, ActLabs, Canada). Chloride and sulphate were determined by chemically suppressed ion chromatography with conductivity detection (Thermo, model Constametric 3500, with Dionex suppressor and IonPac AS22 Dionex column -4 × 250 mm- for anions, CICTERRA-UNC, Argentina). <i>In situ</i> measurements include: alkalinity (volumetric methods), pH and Eh (Hach digital detector), electrical conductivity, temperature, and total dissolved solids (TDS, Hach conductivitymeter). TIC and TOC values were determined in sediment samples following Loss on ignition method (LOI, Heiri et al. 2001). Chemical information was processed with PHREEQC Interactive software (U.S. Geological Survey) [2]. Sediment texture was determined with a particle analyser (Horiba LA-950, LabGEO, CICTERRA-UNC, Argentina).
Data format	Raw analyzed Other (Software Output)
Parameters for data collection	Samples were collected following standard protocols [3]. Parameters for data collection are: 1-Temperature, pH, redox potential, alkalinity, electrical conductivity, and TDS measured <i>in situ</i> . 2- Dissolved major and trace element concentrations, in filtered and acidified samples. 3-TIC and TOC in lyophilized sediments. 4- Grain size characterization in sediments with removed carbonates and organic matter.
Description of sample collection	72 samples were collected (2 ice, 35 lake water, 34 lake sediment, and one salt sample) from 35 different lakes and ponds during January 2015. Clean polythene bottles were used to collect and transport samples to each specific laboratory. Water samples were collected and stabilized (filtered in the field through a 0.22 µm pore size filter and acidified to pH<2) according to established protocols, and stored in a refrigerator (<4 °C) until analysis. Sediment samples were collected from the margins of lakes using a standard scoop sampler and stored in plastic bags.
Data source location	Clearwater Mesa (James Ross Island, northeast Antarctic Peninsula).
Data accessibility	Data are presented with this article.
Related research article	Karina L. Lecomte, Paula A. Vignoni, Cecilia V. Echegoyen, Pia Santolaya, Kateřina Kopalová, Tyler J. Kohler, Matěj Roman, Silvia H. Coria, Juan M. Lirio, Dissolved major and trace geochemical dynamics in Antarctic lacustrine systems, <i>Chemosphere</i> , (2020) 240, 124,938 DOI: https://doi.org/10.1016/j.chemosphere.2019.124938




Fig. 1. a) pH-Eh diagram; b) Piper diagram from Clearwater Mesa lake and ice water samples.

Value of the data

- Given the logistical constraints associated with collecting information from this unique and scarcely known high latitude environment, these observations present an opportunity for hydrological and geochemical research without the need for resampling.
- This information can be helpful for multidisciplinary researchers, such as environmentalists, biologists, physics, and geologists, and can be readily incorporated into meta-analyses.
- This information can be used to identify and analyse processes controlling high latitude lake chemistry through geochemical diagrams and models, and compare them with other environments.
- The present information contributes to a growing database that can serve as a baseline in environmental change and pollution studies (physicochemical and trace element) for Antarctic systems.

1. Data description

Water, ice, and sediment samples from 35 different lakes and ponds were collected during the austral summer (15–29 January 2015) on Clearwater Mesa (CWM), located on James Ross Island, northeast Antarctic Peninsula [1]. Fig. 1a shows the different hydrogeochemical environ-




Fig. 2. Gibbs Diagram of Clearwater Mesa samples.

ments defined according to the relationship between pH vs. Eh (adapted from [4]). Water ionic classification is shown in Fig. 1b with a Piper diagram [5].

Physico-chemical parameters and major ion concentrations were determined for 37 lake and ice samples, and are presented in Table 1. This table also indicates lake area; Mg^{2+}/Ca^{2+} and $Na^+/(Na^+ + Ca^{2+})$ ratios; and sediment TOC and TIC. Fig. 2 shows TDS values vs. $Na^+/(Na^+ + Ca^{2+})$ ratios in a Gibbs diagram [6]. Table 2 displays the hydrochemical statistical values for each hydrogeological system.

Table 3 presents dissolved trace element concentrations. They were normalized to the Upper Continental Crust [7] and a spidergram is shown in Fig. 3, where World average [8] and the mean James Ross Island Volcanic Group geochemical composition have been indicated [9]. Table 4 shows statistical values for differentiated hydrogeological system's trace elements.

PHREEQC models are presented in the Software Output, which is included in the *Supplementary Material*. It comprises ion speciation, Saturation Indexes, and geochemical models (i.e. mixing modelling between Blancmange Glacier and Cecilia samples and inverse modelling between Blancmange Glacier and Florencia samples). Finally, Fig. 4 illustrates those minerals that are prone to precipitate due to their positive Saturation Index (SI).

2. Experimental design, materials, and methods

2.1. Sampling and analyses

Water and sediment samples were collected from 35 different lakes and ponds on Clearwater Mesa during the 2015 Antarctic summer field campaign following standard protocols [3]. Two

Table 1

Sampling points' location and lake's physical and chemical characteristics. It is included TIC and TOC data for CWM lakes.

System	Lake	Jan 2015	Surface m ²	pH	Eh mV	T °C	Conduc $\mu\text{S cm}^{-1}$	TDS Mg L ⁻¹	T Alk	Mg ⁺²	Ca ⁺²	Na ⁺	K ⁺	HCO ₃ ⁻	CO ₃ ⁻²	Cl ⁻	SO ₄ ⁻²	Balance	TOC %	TIC	Na/ Na+Ca meq L ⁻¹	Mg/Ca	
Isolated closed lakes	Adriana	23	18,010	8.66	315.1	10.5	4490.0	2240	156.2	94.9	12.3	178.4	3.6	156.2	0.0	455.3	37.4	0.3	2.4	0.4	0.9	12.9	
	Aldru	20	1840	9.00	369.9	12.6	260.0	131.1	43.4	7.1	6.5	27.1	5.2	42.0	1.4	45.0	6.4	2	2.0	0.3	0.8	1.8	
	Andrea	18	20,630	7.69	394.9	7.7	935.0	470.0	156.5	24.7	29.7	79.5	2.6	156.5	0.0	152.5	7.4	0.4	4.8	0.3	0.7	1.4	
	Joanna	24	5260	8.52	341.0	16.5	312.0	156.9	42.3	9.6	6.7	23.1	1.3	42.3	0.0	44.2	8.9	0.6	2.0	0.2	0.7	2.4	
	Juanita	23	5680	9.14	346.2	9.4	654.0	327.0	209.5	17.6	17.7	59.6	1.3	186.7	22.8	62.0	5.5	-6.8	1.5	0.6	0.7	1.7	
	María	24	2510	8.34	361.3	15.2	267.0	135.5	50.4	7.3	5.6	23.2	1.0	50.4	0.0	34.3	4.4	0.5	3.0	0.3	0.8	2.2	
	Martina	24	820	8.60	360.9	16.6	2490.0	1240	121.4	90.5	38.4	190.0	4.2	121.4	0.0	521.3	44.9	0.3	1.2	0.5	0.8	3.9	
	Natasha	17	13,300	8.58	410.4	1.4	1126.0	564.0	144.4	49.5	14.2	79.1	2.9	140.8	3.6	204.4	7.0	-0.3	1.3	0.3	0.8	5.8	
	Natasha Ice	18	-	6.52	455.4	-	609.0	305.0	250.5	37.9	8.8	59.9	1.5	250.5	0.0	71.2	2.0	0.3	-	-	-	-	
	Soledad	24	5030	8.72	310.8	15.7	263.0	129.3	28.3	4.9	3.2	17.5	0.8	28.3	0.0	27.5	4.2	0.4	2.6	0.5	0.8	2.6	
Katerina	Tamara	23	21,600	7.90	394.0	8.6	300.0	150.0	41.1	9.0	6.2	29.1	0.8	41.1	0.0	52.5	7.8	0.4	1.9	0.3	0.8	2.4	
	Adela	25	18,010	8.84	353.0	15.5	392.0	197.2	151.2	12.7	7.8	34.4	1.3	133.2	18.0	20.9	7.5	-8.8	-	-	0.8	2.7	
	Alejandra	27	27,480	8.82	277.0	15.5	367.0	184.6	100.8	12.9	9.5	30.1	1.0	88.8	12.0	44.3	6.8	-6.2	1.6	0.5	0.7	2.3	
	Claudina	27	30,520	8.76	304.7	16.8	255.0	128.3	101.6	8.2	6.4	23.6	0.8	98.0	3.6	12.5	2.8	-2.6	2.0	0.3	0.8	2.1	
	Graciela	25	30,070	8.74	308.9	18.5	603.0	302.0	58.3	24.1	15.0	46.0	1.7	45.1	13.2	128.6	17.6	-4.1	2.2	0.2	0.7	2.7	
	Ileana	21	6630	7.77	385.2	14.3	255.0	128.5	34.6	7.4	5.3	18.1	1.0	34.6	0.0	36.0	3.7	0.7	2.3	0.3	0.7	2.3	
	Karina	22	20,760	7.92	399.3	11.3	328.0	164.0	142.5	15.8	7.3	23.8	1.0	142.5	0.0	11.8	1.4	0.4	2.4	0.2	0.7	3.6	
	Katerina	21	126,950	8.37	342.4	14.1	496.0	248.0	191.1	25.5	11.9	32.8	1.2	191.1	0.0	31.2	5.1	0.3	2.5	0.4	0.7	3.6	
																			4.4	0.4	14.3	0.4	
		Linda	25	20,550	8.73	305.9	18.5	651.0	326.0	157.7	28.2	18.3	49.0	2.0	142.1	15.6	95.7	16.2	-4.2	2.3	0.3	0.7	2.6
Trinidad-Tatana	Ludmila	25	10,000	8.67	335.2	15.3	203.0	102.3	65.4	5.8	4.9	23.3	0.7	65.4	0.0	22.0	1.8	0.3	2.3	0.2	0.8	2.0	
	Marta	28	2740	8.78	274.9	18.5	415.0	208.0	103.8	14.7	9.3	36.1	1.4	95.4	8.4	53.9	7.6	-3.8	2.2	0.4	0.8	2.6	
	Nora	28	65,720	8.75	275.6	15.9	465.0	233.0	131.2	22.2	9.6	35.0	1.3	120.4	10.8	60.0	7.8	-4.1	1.8	0.3	0.8	3.9	
	Paula	28	14,890	9.21	263.3	17.2	403.0	201.0	150.3	16.4	12.7	37.1	1.8	132.3	18.0	45.3	7.3	-7.1	1.8	0.3	0.7	2.2	
	Sandra	28	1830	8.04	300.6	12.0	291.0	111.6	59.5	5.5	4.1	24.4	0.8	59.5	0.0	24.0	3.3	0.4	-	-	0.8	2.2	
	Sara	28	6630	9.10	262.8	17.6	533.0	267.0	127.1	25.9	12.5	43.6	1.7	117.5	9.6	88.7	10.7	-2.9	2.3	0.4	0.8	3.5	
	Silvia	21	37,350	8.75	341.5	13.0	630.0	315.0	197.7	29.4	17.8	46.4	1.8	197.7	0.0	64.0	13.4	0.4	1.6	0.2	0.7	2.8	
	Susan	22	20,130	7.92	398.3	11.3	181.0	91.2	69.2	7.7	4.1	13.6	0.8	69.2	0.0	9.1	1.7	0.4	2.4	0.3	0.7	3.1	
	Valentina	28	3910	9.22	263.3	19.1	506.0	253.0	180.9	18.4	12.6	43.5	1.9	170.1	10.8	39.5	6.3	-3.8	2.3	0.3	0.8	2.4	
	Esther	29	14,080	9.43	271.2	16.3	302.0	152.1	61.5	6.7	7.7	27.1	0.8	51.9	9.6	41.5	4.6	-6.9	2.7	0.3	0.8	1.5	
Norma	Tatana	29	5350	9.02	275.5	16.0	1427.0	715.0	304.7	44.8	32.4	145.0	2.6	295.1	9.6	229.8	14.1	-1.1	0.9	0.6	0.8	2.3	
	Argentina	29	16,820	9.32	279.2	13.3	642.0	321.0	42.2	15.7	10.7	50.1	1.1	31.4	10.8	118.0	7.8	-4	-	-	0.8	2.4	
	Trinidad	29	65,370	8.85	277.5	13.0	1823.0	912.0	775.9	87.6	20.0	154.0	3.0	763.9	12.0	83.1	1.8	-1.1	1.6	0.3	0.9	7.3	
Florencia	Joaquina	15	2870	8.33	270.5	8.1	367.0	184.7	109.9	13.3	9.6	39.9	1.4	109.9	0.0	46.9	8.9	0.4	1.7	0.3	0.8	2.3	
	Norma	15	610	7.31	412.9	6.8	248.0	125.5	57.2	8.8	5.9	24.2	0.8	57.2	0.0	36.9	4.5	0.3	0.8	0.1	0.8	2.5	
	Cecilia	18	6860	8.47	378.3	7.4	556.0	278.0	6.8	8.4	8.8	53.3	2.3	6.8	0.0	116.0	8.5	-0.8	2.2	0.3	0.8	1.6	
	Blancmange Glacier	18	-	6.40	458.8	-	11.7	5.9	54.9	1.7	2.7	16.5	0.6	54.9	0.0	2.7	0.6	0.3	-	-	0.8	1.0	
	Florencia	18	~132,000	7.22	446.8	2.0	152.9	76.7	3.4	0.2	13.0	1.5	0.3	3.4	0.0	22.2	2.4	-0.5	2.2	0.3	0.1	0.0	

K.L. Leconte, C.V. Echegoyen and P.A. Vignoni et al./Data in Brief 30 (2020) 105438

Table 2
Statistical values for each hydrogeological system's physical and chemical characteristics.

System		pH	Eh m V	T °C	Conduc. $\mu\text{S cm}^{-1}$	TDS mg L ⁻¹	T Alk	Mg ⁺²	Ca ⁺²	Na ⁺	K ⁺	HCO ₃ ⁻	CO ₃ ⁻²	Cl ⁻	SO ₄ ⁻²	TOC %	TIC	Na/(Na+Ca) meq L ⁻¹	Mg/Ca
Isolated closed lakes	Maximum	9.1	455	16.6	4490	2240	250.5	94.9	38.4	190	5.2	250.5	22.8	521	44.9	4.8	0.6	0.93	12.9
	Minimum	6.5	311	1.4	260	129.3	28.3	4.9	3.2	17.5	0.8	28.3	0.0	27.5	2.0	1.2	0.2	0.70	1.4
	Mean	8.3	369	11.4	1064	531.7	113.1	32.1	13.6	69.7	2.3	110.6	2.5	152	12.4	2.3	0.4	0.80	3.7
	Median	8.6	361	11.6	609	305.0	121.4	17.6	8.8	59.6	1.5	121.4	0.0	62.0	7.0	2.0	0.3	0.79	2.4
Katerina	Maximum	9.2	399	19.1	651	326.0	197.7	29.4	18.3	49.0	2.0	197.7	18.0	129	17.6	14.3	0.5	0.84	3.9
	Minimum	7.8	263	11.3	181	91.2	34.6	5.5	4.1	13.6	0.7	34.6	0.0	9.1	1.4	1.6	0.2	0.69	2.0
	Mean	8.6	317	15.6	410	203.6	119.0	16.5	9.9	33.0	1.3	111.9	7.1	46.3	7.1	3.0	0.3	0.75	2.7
Trinidad-Tatana	Maximum	8.8	306	15.5	403	201.0	127.1	15.8	9.5	34.4	1.3	117.5	8.4	39.5	6.8	2.3	0.3	0.75	2.6
	Minimum	9.4	279	16.3	1823	912.0	775.9	87.6	32.4	154	3.0	763.9	12.0	230	14.1	2.7	0.6	0.87	7.3
	Mean	8.9	271	13.0	302	152.1	42.2	6.7	7.7	27.1	0.8	31.4	9.6	41.5	1.8	0.9	0.3	0.75	1.5
Norma	Maximum	9.2	277	14.7	1035	518.0	183.1	30.3	15.4	97.6	1.9	173.5	10.2	101	6.2	1.6	0.3	0.80	2.4
	Minimum	8.3	413	8.1	367	184.7	109.9	13.3	9.6	39.9	1.4	109.9	0.0	46.9	8.9	1.7	0.3	0.78	2.5
	Mean	7.3	271	6.8	248	125.5	57.2	8.8	5.9	24.2	0.8	57.2	0.0	36.9	4.5	0.8	0.1	0.78	2.3
	Median	7.8	342	7.5	308	155.1	83.6	11.1	7.8	32.1	1.1	83.6	0.0	41.9	6.7	1.3	0.2	0.78	2.4
Florenxia	Maximum	7.8	342	7.5	308	155.1	83.6	11.1	7.8	32.1	1.1	83.6	0.0	41.9	6.7	1.3	0.2	0.78	2.4
	Minimum	8.5	459	7.4	556	278.0	54.9	8.4	13.0	53.3	2.3	54.9	0.0	116	8.5	2.2	0.3	0.84	1.6
	Mean	6.4	378	2.0	12	5.9	3.4	0.2	2.7	1.5	0.3	3.4	0.0	2.7	0.6	2.2	0.3	0.09	0.0
	Median	7.4	428	4.7	240	120.2	21.7	3.4	8.2	23.8	1.1	21.7	0.0	47.0	3.8	2.2	0.3	0.59	0.9
CWM lakes	Maximum	7.2	447	4.7	153	76.7	6.8	1.7	8.8	16.5	0.6	6.8	0.0	22.2	2.4	2.2	0.3	0.84	1.0
	Minimum	9.4	459	19.1	4490	2240	775.9	94.9	38.4	190	5.2	763.9	22.8	521	44.9	14.3	0.6	0.93	12.9
	Mean	6.4	263	1.4	12	5.9	3.4	0.2	2.7	1.5	0.3	3.4	0.0	2.7	0.6	0.8	0.1	0.09	0.0
	Median	8.4	338	13.2	654	326.5	126.6	22.2	11.6	49.7	1.6	121.4	5.1	85.3	8.4	2.5	0.3	0.76	2.9
	Maximum	8.7	341	14.3	403	201.0	103.8	14.7	9.5	35.0	1.3	98.0	0.0	45.3	6.8	2.2	0.3	0.77	2.4

Table 3

Trace element determined in the 2015 sampling campaign and JRIVG mean values (in ppm) are from concentrations reported by Košler et al., 2009.

Sample	Si mg L ⁻¹	Al	Fe	Ti μg L ⁻¹	Mn	Ba	Sr	Zr	Rb	Zn	V	Cr	Cu
Detection limit (μg L ⁻¹)	200	2	10	0.1	0.1	0.1	0.04	0.01	0.005	0.5	0.1	0.5	0.2
Adela	4.80	0.10	0.21	5.60	17.40	2.80	5.21	0.21	0.63	8.70	1.30	2.70	5.50
Adriana	0.20	0.01	0.03	0.70	1.70	0.50	20.40	0.08	0.35	0.50	1.30	4.00	3.90
Adru	3.60	0.06	0.03	8.20	3.60	0.20	2.79	0.09	0.81	16.20	1.60	0.50	2.00
Alejandra	4.30	0.01	0.02	1.70	3.60	0.30	2.89	0.04	0.21	6.20	0.70	0.50	1.70
Andrea	4.10	0.00	0.04	2.20	5.60	0.20	22.30	0.16	0.47	6.10	7.30	1.40	1.60
Argentina	3.80	0.12	0.19	7.00	9.50	1.40	4.04	0.19	0.50	11.00	2.60	0.50	3.60
Cecilia	2.60	0.03	0.02	1.50	3.20	0.30	15.00	0.73	0.43	8.90	5.70	0.60	1.10
Claudina	2.50	0.01	0.02	1.00	2.80	0.10	2.45	0.03	0.14	2.20	0.50	0.50	1.60
Ester	3.40	0.09	0.12	3.60	10.10	1.60	3.99	0.08	0.37	8.60	2.20	0.70	3.50
Blancmange Glacier	1.60	0.04	0.04	4.50	2.40	0.20	1.45	0.05	0.19	5.50	6.00	0.50	1.10
Florencia	0.40	0.19	0.15	20.90	9.10	0.90	1.81	0.44	0.21	12.10	0.50	0.50	2.00
Graciela	3.30	0.13	0.19	8.20	11.70	2.60	6.55	0.16	0.50	12.10	1.00	0.70	3.40
Ileana	2.80	0.10	0.26	7.60	7.70	1.50	2.87	0.10	0.30	6.10	0.60	0.60	2.40
Joanna	4.20	0.08	0.12	5.10	13.60	1.70	4.67	0.37	0.64	21.30	0.60	1.20	2.80
Joaquina	6.00	0.03	0.07	3.10	5.60	0.60	4.35	0.07	0.17	10.90	1.60	1.00	1.90
Juanita	5.70	0.01	0.02	2.00	4.30	0.20	5.40	0.06	0.25	6.80	2.70	0.60	1.30
Karina	2.80	0.07	0.13	3.80	7.90	4.32	3.59	0.10	0.21	5.80	0.50	0.60	2.90
Katerina	3.00	0.10	0.19	6.10	14.00	2.60	7.21	0.11	0.26	6.90	0.70	0.70	2.50
Linda	4.70	0.32	0.59	22.50	15.60	2.10	7.60	0.40	0.66	11.10	1.70	1.00	4.40
Ludmila	3.60	0.05	0.07	2.50	7.70	1.10	1.63	0.03	0.26	9.90	0.80	0.80	3.10
Maria	3.80	0.07	0.06	3.00	11.70	2.10	3.19	0.11	0.42	12.90	0.50	0.80	2.70
Marta	4.40	0.09	0.09	6.60	5.80	1.30	5.80	0.13	0.26	4.00	2.00	0.80	1.90
Martina	9.70	0.18	0.11	5.80	18.10	4.40	21.10	0.22	0.91	14.00	1.70	0.50	4.20
Natasha	1.90	0.03	0.05	1.80	10.40	0.70	7.98	0.11	0.28	37.00	0.70	3.00	6.30
Natasha Ice	3.80	0.22	0.45	20.50	21.70	1.50	12.90	0.42	0.53	14.70	1.70	0.90	4.00
Nora	3.00	0.09	0.14	6.30	8.50	1.50	7.12	0.11	0.30	2.90	1.30	0.50	2.00
Norma	2.60	0.03	0.13	2.70	11.80	0.90	2.61	0.03	0.12	11.10	0.70	0.50	2.20
Paula	4.40	0.40	0.62	30.50	23.60	2.30	9.96	0.53	0.78	91.50	3.20	1.00	4.10
Sandra	3.50	0.06	0.03	2.40	8.40	2.00	2.96	0.04	0.31	8.50	0.70	0.60	2.00
Sara	4.10	0.14	0.16	5.70	17.70	4.60	9.64	0.09	0.43	13.80	1.60	0.80	4.00
Silvia	2.90	0.02	0.03	2.10	5.30	0.30	7.36	0.03	0.29	4.90	0.80	0.60	2.50
Soledad	2.70	0.17	0.25	12.90	11.90	1.80	2.39	0.16	0.39	10.90	1.00	0.60	2.70
Susan	1.80	0.20	0.14	3.90	11.20	3.20	3.04	0.06	0.34	11.50	0.40	0.50	2.70
Tamara	2.40	0.02	0.03	1.50	7.40	0.80	3.88	0.03	0.16	6.30	0.60	0.80	1.70
Tatana	12.60	0.10	0.11	7.00	8.80	1.60	17.90	0.16	0.61	9.00	7.80	1.30	5.60
Trinidad	2.70	0.03	0.06	3.20	8.30	0.80	15.80	0.18	0.71	15.20	1.80	1.10	3.50
Valentina	4.70	0.16	0.18	10.20	12.80	2.20	8.85	0.20	0.41	4.40	3.40	0.60	2.30
Basalts mean values	22.22	4.06	7.40	917.80	139.41	126.88	505.63	163.50	13.60	-	160.38	458.13	43.39
Sample	Y μg L ⁻¹	Li	Ni	Pb	Sc	Th	Co	Hf	Cs	U	As	Mo	Sb
Detection limit (μg L ⁻¹)	0.003	1	0.3	0.01	1	0.001	0.005	0.001	0.001	0.001	0.03	0.1	0.01
Adela	0.06	7.00	2.10	0.22	1.00	0.00	0.55	0.00	0.10	0.08	0.60	0.30	0.04
Adriana	0.01	5.00	1.10	0.01	1.00	0.00	0.04	0.00	0.21	0.44	2.80	1.40	0.05
Adru	0.00	3.00	0.90	0.34	2.00	0.00	0.14	0.00	0.03	0.01	0.47	0.10	0.98
Alejandra	0.00	2.00	1.40	0.45	2.00	0.00	0.10	0.00	0.20	0.06	0.44	0.30	0.02
Andrea	0.01	2.00	1.60	0.51	2.00	0.00	0.16	0.00	0.10	0.14	0.55	0.30	0.02
Argentina	0.03	2.00	1.90	0.38	2.00	0.00	0.32	0.01	0.09	0.04	0.42	0.20	0.08
Cecilia	0.04	1.00	0.70	0.50	1.00	0.01	0.07	0.03	0.05	0.05	0.28	0.20	0.02
Claudina	0.00	2.00	1.10	0.16	1.00	0.00	0.08	0.00	0.02	0.04	0.40	0.20	0.01
Ester	0.01	3.00	2.70	0.40	2.00	0.00	0.43	0.00	0.02	0.03	0.27	0.10	0.02
Blancmange Glacier	0.00	1.00	0.50	0.40	1.00	0.00	0.07	0.00	0.02	0.04	0.28	0.20	0.01

(continued on next page)

Table 3
(continued)

Sample	Y $\mu\text{g L}^{-1}$	Li	Ni	Pb	Sc	Th	Co	Hf	Cs	U	As	Mo	Sb
Florencia	0.04	9.00	0.70	0.58	1.00	0.00	0.17	0.01	0.02	0.01	0.04	0.10	0.02
Graciela	0.03	9.00	1.90	0.69	2.00	0.01	0.38	0.01	0.02	0.15	0.58	0.30	0.02
Ileana	0.04	2.00	2.00	0.53	1.00	0.00	0.24	0.00	0.02	0.04	0.45	0.10	0.02
Joanna	0.02	11.00	1.90	0.45	2.00	0.00	0.40	0.00	0.02	0.02	0.28	0.20	0.04
Joaquina	0.01	2.00	2.10	0.49	3.00	0.00	0.22	0.00	0.06	0.07	0.51	0.40	0.03
Juanita	0.00	6.00	0.80	0.20	3.00	0.00	0.06	0.00	0.01	0.06	0.80	0.30	0.02
Karina	0.01	4.00	1.40	0.28	1.00	0.00	0.26	0.00	0.02	0.06	0.33	0.10	0.01
Katerina	0.02	5.00	1.80	0.26	1.00	0.00	0.36	0.00	0.03	0.10	0.60	0.20	0.03
Linda	0.10	10.00	2.50	0.52	2.00	0.01	0.40	0.01	0.03	0.23	0.79	0.30	0.04
Ludmila	0.00	5.00	1.10	0.37	2.00	0.00	0.23	0.00	0.01	0.01	0.17	0.10	0.01
Maria	0.00	13.00	1.20	0.35	2.00	0.00	0.34	0.00	0.03	0.02	0.22	0.10	0.02
Marta	0.01	3.00	1.30	0.23	2.00	0.01	0.27	0.00	0.02	0.06	0.60	0.30	0.03
Martina	0.03	13.00	2.80	0.36	5.00	0.00	0.81	0.01	0.07	0.73	1.37	1.20	0.05
Natasha	0.01	5.00	2.40	1.57	5.00	0.01	0.17	0.01	0.05	0.07	0.60	0.50	0.05
Natasha Ice	0.10	6.00	3.60	0.66	2.00	0.01	0.33	0.01	0.13	0.27	1.17	0.40	0.05
Nora	0.01	5.00	1.50	0.25	1.00	0.01	0.28	0.00	0.01	0.08	0.50	0.20	0.03
Norma	0.00	2.00	1.10	0.23	1.00	0.00	0.21	0.00	0.01	0.02	0.36	0.20	0.02
Paula	0.10	8.00	2.20	0.77	2.00	0.02	0.60	0.01	0.03	0.17	1.17	0.30	0.06
Sandra	0.00	3.00	1.00	0.24	2.00	0.00	0.37	0.00	0.01	0.03	0.25	0.10	0.03
Sara	0.11	5.00	2.30	0.55	2.00	0.00	0.36	0.00	0.02	0.12	0.83	0.40	0.04
Silvia	0.00	7.00	1.80	0.22	1.00	0.00	0.08	0.00	0.07	0.26	0.73	0.30	0.04
Soledad	0.04	12.00	1.30	0.45	1.00	0.00	0.55	0.00	0.37	0.02	0.26	0.10	0.04
Susan	0.02	4.00	4.00	0.55	1.00	0.01	0.30	0.00	0.09	0.02	0.23	0.10	0.03
Tamara	0.00	5.00	1.50	0.51	1.00	0.00	0.14	0.00	0.02	0.00	0.48	0.20	0.02
Tatana	0.01	3.00	2.60	0.56	6.00	0.00	0.33	0.00	0.09	0.14	1.09	0.90	0.05
Trinidad	0.01	5.00	2.90	0.38	2.00	0.00	0.17	0.00	0.05	0.13	0.97	0.40	0.04
Valentina	0.03	6.00	1.60	0.28	2.00	0.01	0.44	0.01	0.01	0.27	1.27	0.50	0.06
Basalts mean values	23.94	6.19	108.38	2.94	0.00	22.78	2.58	43.78	3.54	0.13	0.97	–	–

ice samples (i.e., Blancmange Glacier and Lake Natasha-ice), were also collected, along with a precipitated salt sample from the margin of Lake Andrea.

Water temperature, pH, redox potential, electrical conductivity, TDS, and alkalinity were measured at each waterbody *in situ*. Redox potential and pH were measured with a Hach digital detector, while temperature, TDS, and electrical conductivity were measured using a digital Hach conductivitymeter. Alkalinity was measured as CaCO_3 by end point titration in the field, and using a 0.16 N H_2SO_4 solution until $\text{pH} = 4.5$. For the determination of anions, major cations, and trace elements, samples were vacuum-filtered in the field with 0.22 μm pore size cellulose filters (HA-type, Millipore Corp.). An aliquot was stored in polyethylene bottles at 4 °C for the determination of chloride and sulphate by chemically suppressed ion chromatography with conductivity detection. The other aliquot was acidified ($\text{pH} < 2$) with concentrated, redistilled and ultrapure HNO_3 (Sigma-Aldrich) for the analytical determination of major and trace elements by inductively coupled plasma-mass spectrometry (ICP-MS, Activation Laboratories Ltd., Ancaster, Ontario, Canada). Major and trace elements concentrations were validated using NIST (National Institute of Standards and Technology) 1640 and Riverine Water Reference Materials for Trace Metals certified by the National Research Council of Canada (SRLS-4), and detection limits are reported in the corresponding tables. Statistical parameters (i.e. mean and median) were calculated for each hydrogeological system's trace elements and physical and chemical characteristics.

2.2. Sediment analysis

Bottom lake sediment samples were collected from 32 shallow lakes using a standard scoop sampler. Lake Katerina was sampled 3 times in order to characterize spatial variability in this

Table 4
Statistical values for each hydrogeological system's trace elements.

System		Si Mg L ⁻¹	Al	Fe	Ti µg L ⁻¹	Mn	Ba	Sr	Zr	Rb	Zn	V	Cr	Cu	Y	Li	Ni	Pb	Sc	Th	Co	Hf	Cs	U	As	Mo	Sb
Isolated closed lakes	Max	9.7	0.2	0.3	12.9	18.1	4.4	22.3	0.4	0.9	37.0	7.3	4.0	6.3	0.042	13.0	2.8	1.6	5.0	0.008	0.8	0.006	0.37	0.73	2.80	1.4	0.98
	Min	0.2	0.0	0.0	0.70	1.7	0.2	2.4	0.0	0.2	0.5	0.5	0.5	1.3	0.003	2.0	0.8	0.0	1.0	0.001	0.0	0.001	0.01	0.00	0.22	0.1	0.02
	Mean	3.8	0.1	0.1	4.32	8.8	1.3	9.4	0.1	0.5	13.2	1.8	1.3	2.9	0.013	7.5	1.6	0.5	2.4	0.003	0.3	0.003	0.09	0.15	0.78	0.4	0.13
Katerina	Median	3.7	0.0	0.0	2.60	8.9	0.8	5.0	0.1	0.4	11.9	1.2	0.8	2.7	0.008	5.5	1.4	0.4	2.0	0.002	0.2	0.003	0.04	0.04	0.52	0.3	0.04
	Max	4.8	0.4	0.6	30.5	23.6	4.6	10.0	0.5	0.8	91.5	3.4	2.7	5.5	0.109	10.0	4.0	0.8	2.0	0.019	0.6	0.013	0.20	0.27	1.27	0.5	0.06
	Min	1.8	0.0	0.0	1.00	2.8	0.1	1.6	0.0	0.1	2.2	0.4	0.5	1.6	0.003	2.0	1.0	0.2	1.0	0.001	0.1	0.001	0.01	0.01	0.17	0.1	0.01
Trinidad-Tatana	Mean	3.6	0.1	0.2	7.45	10.7	2.0	5.6	0.1	0.4	12.4	1.2	0.8	2.9	0.033	5.1	1.8	0.4	1.5	0.005	0.3	0.004	0.04	0.10	0.58	0.2	0.03
	Median	3.5	0.1	0.1	5.70	8.5	2.1	5.8	0.1	0.3	6.9	0.8	0.6	2.5	0.020	5.0	1.8	0.3	2.0	0.004	0.3	0.003	0.02	0.08	0.58	0.3	0.03
	Max	12.6	0.1	0.2	7.00	10.1	1.6	17.9	0.2	0.7	15.2	7.8	1.3	5.6	0.033	5.0	2.9	0.6	6.0	0.004	0.4	0.005	0.09	0.14	1.09	0.9	0.08
Norma	Min	2.7	0.0	0.1	3.20	8.3	0.8	4.0	0.1	0.4	8.6	1.8	0.5	3.5	0.006	2.0	1.9	0.4	2.0	0.001	0.2	0.002	0.02	0.03	0.27	0.1	0.02
	Mean	5.6	0.1	0.1	5.20	9.2	1.4	10.4	0.2	0.5	11.0	3.6	0.9	4.1	0.014	3.3	2.5	0.4	3.0	0.003	0.3	0.004	0.06	0.08	0.69	0.4	0.05
	Median	3.6	0.1	0.1	5.30	9.2	1.5	9.9	0.2	0.6	10.0	2.4	0.9	3.6	0.008	3.0	2.7	0.4	2.0	0.003	0.3	0.004	0.07	0.08	0.70	0.3	0.05
Florenca	Max	6.0	0.0	0.1	3.10	11.8	0.9	4.4	0.1	0.2	11.1	1.6	1.0	2.2	0.006	2.0	2.1	0.5	3.0	0.002	0.2	0.003	0.06	0.07	0.51	0.4	0.03
	Min	2.6	0.0	0.1	2.70	5.6	0.6	2.6	0.0	0.1	10.9	0.7	0.5	1.9	0.003	2.0	1.1	0.2	1.0	0.002	0.2	0.002	0.01	0.02	0.36	0.2	0.02
	Mean	4.3	0.0	0.1	2.90	8.7	0.8	3.5	0.1	0.1	11.0	1.2	0.8	2.1	0.005	2.0	1.6	0.4	2.0	0.002	0.2	0.003	0.03	0.04	0.44	0.3	0.03
CWM lakes	Median	4.3	0.0	0.1	2.90	8.7	0.8	3.5	0.1	0.1	11.0	1.2	0.8	2.1	0.005	2.0	1.6	0.4	2.0	0.002	0.2	0.003	0.03	0.04	0.44	0.3	0.03
	Max	2.6	0.2	0.2	20.9	9.1	0.9	15.0	0.7	0.4	12.1	6.0	0.6	2.0	0.042	9.0	0.7	0.6	1.0	0.005	0.2	0.033	0.05	0.05	0.28	0.2	0.02
	Min	0.4	0.0	0.0	1.50	2.4	0.2	1.5	0.1	0.2	5.5	0.5	0.5	1.1	0.003	1.0	0.5	0.4	1.0	0.001	0.1	0.003	0.02	0.01	0.04	0.1	0.01
CWM lakes	Mean	1.5	0.1	0.1	8.97	4.9	0.5	6.1	0.4	0.3	8.8	4.1	0.5	1.4	0.027	3.7	0.6	0.5	1.0	0.002	0.1	0.016	0.03	0.03	0.20	0.2	0.02
	Median	1.6	0.0	0.0	4.50	3.2	0.3	1.8	0.4	0.2	8.9	5.7	0.5	1.1	0.037	1.0	0.7	0.5	1.0	0.001	0.1	0.011	0.02	0.04	0.28	0.2	0.02
	Max	12.6	0.4	0.6	30.5	23.6	4.6	22.3	0.7	0.9	91.5	7.8	4.0	6.3	0.109	13.0	4.0	1.6	6.0	0.019	0.8	0.033	0.37	0.73	2.80	1.4	0.98
CWM lakes	Min	0.2	0.0	0.0	0.70	1.7	0.1	1.5	0.0	0.1	0.5	0.4	0.5	1.1	0.003	1.0	0.5	0.0	1.0	0.001	0.0	0.001	0.01	0.00	0.04	0.1	0.01
	Mean	3.7	0.1	0.1	6.59	9.7	1.5	7.3	0.2	0.4	12.1	1.9	0.9	2.9	0.025	5.3	1.8	0.4	1.9	0.004	0.3	0.005	0.06	0.11	0.63	0.3	0.06
	Median	3.5	0.1	0.1	4.50	8.8	1.5	5.2	0.1	0.4	9.0	1.3	0.7	2.7	0.013	5.0	1.6	0.4	2.0	0.002	0.3	0.003	0.03	0.06	0.50	0.2	0.03




Fig. 3. Upper Continental Crust normalized spidergram. World average [8] and James Ross Island Volcanic Group average [9] are added for comparison.

large lake. Particular attention was paid to collect the sediment/water interface. Samples were skimmed from the benthic surface (top 2–3 cm), stored in plastic bags, and refrigerated until analysis.

TOC and TIC was determined in sediment samples using the loss on ignition method (LOI; [10]). Prior to their analysis by LOI, they were partitioned and lyophilized in order to remove water by sublimation. To prepare the samples, sediment was first ground to obtain a homogeneous sample. A quantity of 2 g was then weighed on a high-precision analytical balance (Precisa XR 2058- DR) and placed in numbered crucibles. Each sample was weighed to obtain the initial weight. The organic matter content was determined by calculating the difference in mass between sediment samples dried to a constant weight at 105 °C, and after furnacing at 550 °C for 4 h. Carbonate content was calculated as the mass loss after burning the LOI residue at 950 °C for 2 h [10].

2.3. Geochemical modelling

Chemical information was processed with PHREEQC [2] constructed using the AQUACHEM PHREEQC interface. These programs were used to calculate the SI for different mineral phases, element speciation in the given conditions (i.e. the element distribution in all the possible dissolved chemical species that can be found in the samples), and geochemical weathering and mixing models. Software outputs are presented in the *Supplementary Material*. The SI is the ratio between the ion activity product for the given material and the reaction constant at a given temperature. When $SI > 0$, it means that the solution is supersaturated with respect to the mineral phase and may therefore precipitate, whereas when $SI < 0$ the solution is below saturation of the specified mineral.




Fig. 4. Mineral Saturation Index (SI) calculated with PHREEQC.

Geochemical modelling uses the interaction of cations and anions as a function of temperature, redox potential, pH, and ionic strength. Inverse modelling was performed to quantify weathering processes occurring, first between connected lakes Esther and Tatana, and second between Blancmange Glacier and Lake Florencia. Inverse modelling begins with known initial and final solutions, and available mineral phases which attempts to quantify the processes that lead to the final solution chemistry (e.g., [11]). In the process, all possible combinations of dissolution and/or precipitation reactions explaining chemical changes observed between the two solutions and the mineral phases are reconstructed [12]. Models were performed under equilibrium with $O_2(g)$ given that these are surficial hydrological systems. Finally, three mixing models with different proportions of Blancmange Glacier and Lake Cecilia solutions (i.e., one model where both the glacier and lake contribute 50% of the water, one where 20% of the water is from the glacier and 80% is from Lake Cecilia, and one where 80% of the water is from the glacier and 20% is from Lake Cecilia) were made to verify eventual mixing processes.

Acknowledgments

The authors wish to thank Dirección Nacional del Antártico (DNA) and Instituto Antártico Argentino (IIA) for the financial and logistical support in Antarctica. This work was supported by Agencia Nacional de Promoción Científica y Tecnológica (ANPCYT projects PICTO-2010-0096 and PICT 2017e2026), by the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET, Argentina, projects PIP 11220170100088CO and PUE-CICTERRA 2016), and by the Universidad Nacional de Córdoba (SeCyT, project 336-20180100385-CB). Authors TJK and KK were further supported by Charles University Research Centre program No. 204069.

Conflict of Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Supplementary material

Supplementary material associated with this article can be found, in the online version, at doi:10.1016/j.dib.2020.105438.

References

- [1] K.L. Lecomte, P.A. Vignoni, C.V. Echegoyen, P. Santolaya, K. Kopalová, T.J. Kohler, M. Roman, S.H. Coria, J.M. Lirio, Dissolved major and trace geochemical dynamics in Antarctic lacustrine systems, *Chemosphere* 240 (2020) 124938 <https://doi.org/10.1016/j.chemosphere.2019.124938>.
- [2] D.L. Parkhurst, User's guide to PHREEQC-A computer program for speciation, reaction-path, advective-transport, and inverse geochemical calculations, U.S. Geol. Surv. Water Res. Invest. Rep. 95 (1995) 4227 Lakewood, Colorado.
- [3] E.W. Rice, R.B. Baird, A.D. Eaton, L.S. Clesceri, *Standard Methods for the Examination of Water and Wastewater*, 22nd Ed., American Water Works Assn, Washington DC, 2012.
- [4] L.G. Baas Becking, I.R. Kaplan, D. Moore, Limits of the natural environment in terms of pH and oxidation-reduction potentials, *J. Geol.* 68 (1960) 243–284.
- [5] A.M. Piper, A graphic procedure in the geochemical interpretation of water analyses, *Trans. Am. Geophys. Union* 25 (1944) 914–923.
- [6] R.J. Gibbs, Mechanisms controlling world water chemistry, *Science* 170 (1970) 1088–1090.
- [7] S.M. McLennan, Relationships between the trace element composition of sedimentary rocks and upper continental crust, *Geochem. Geophys. Geosyst.* (2001) <https://doi.org/10.1029/2000GC000109>.
- [8] J. Gaillardet, J. Viers, B. Dupré, Trace elements in river waters, in: H.D. Holland, K.K. Turekian (Eds.), *Treatise on Geochemistry*, 2nd edn., Elsevier, Oxford, 2014, pp. 195–235.
- [9] J. Košler, T. Magna, B. Mlcoch, P. Mixa, D. Nyíl, F.V. Holub, Combined Sr, Nd, Pb and Li isotope geochemistry of alkaline lavas from northern James Ross Island (Antarctic Peninsula) and implications for back-arc magma formation, *Chem. Geol.* 258 (2009) 207–218 <https://doi.org/10.1016/j.chemgeo.2008.10.006>.
- [10] O. Heiri, A.F. Lotter, G. Lemcke, Loss on ignition as a method for estimating organic and carbonate content in sediments: reproducibility and comparability of results, *J. Paleolimnol.* 25 (2001) 101–110.
- [11] K.L. Lecomte, A.I. Pasquini, P.J. Depetris, Mineral weathering in a semiarid mountain river: its assessment through PHREEQC inverse modeling, *Aquat. Geochem.* 11 (2005) 173–194.
- [12] D.L. Parkhurst, C.A. Apello, User's guide to PHREEQC (version 2): a computer code program for speciation, batch-reaction, one-dimensional transport and inverse geochemical calculations, U.S. Geol. Surv. Water Res. Invest. Rep. 99 (1999) Washington DC.