

Morroniside protects against cerebral ischemia/reperfusion injury by inhibiting neuron apoptosis and MMP2/9 expression

GUOYONG ZENG¹, WEIJIANG DING², YIN LI², MEIYING SUN³ and LIYING DENG²

¹Department of Neurology, The Affiliated Ganzhou Hospital of Nanchang University, Ganzhou, Jiangxi 341000;

²Department of Neurology, The Second Affiliated Hospital of Nanchang University, Nanchang, Jiangxi 330006;

³Department of Neurology, The People's Hospital of Fuzhou, Fuzhou, Jiangxi 344000, P.R. China

Received November 3, 2017; Accepted March 9, 2018

DOI: 10.3892/etm.2018.6457

Abstract. The aim of the present study was to investigate the effect of morroniside against matrix metalloproteinase (MMP)2/9 and focal cerebral ischemia/reperfusion (I/R) injury in rats. A rat model of focal cerebral I/R injury rats was established and rats were administered with 30, 90 or 270 mg/kg/day morroniside for 7 days. The expression of MMP2/9 and neuronal apoptosis were assessed. In addition, the expression of active caspase-3, B-cell lymphoma 2 (Bcl-2) and Bcl-2-associated X protein (Bax) were measured. The results revealed that MMP2 and MMP9 expression was upregulated and the percentage of apoptotic neurons was increased in rats with focal cerebral I/R injury compared with the control. However, treatment with morroniside significantly inhibited I/R-induced MMP2/9 expression and neuron apoptosis compared with the untreated I/R injury group. Morroniside administration also decreased the expression of active caspase-3 and increased the Bcl-2/Bax ratio compared with untreated rats with focal cerebral I/R injury. The inhibitory effect of morroniside on MMP2/9 expression and neuron apoptosis was dose dependent. In summary, the results of the present study suggest that morroniside is able to protect against cerebral I/R injury in the brain and may have potential as a therapeutic treatment for patients who have suffered a stroke.

Introduction

Ischemia/reperfusion (I/R) causes severe damage to vital organs, including the brain, heart, lungs and kidneys (1-4). I/R injury encompasses stroke, which is characterized by a sudden

focal or global neurological impairment and is one of the most prevalent causes of mortality and disability worldwide (5-7).

In principle, cerebral ischemia is caused by cerebral blood flow reduction leading to a series of pathological changes, including ionic homeostasis loss, energy failure, increased oxidative stress, apoptosis, irreversible tissue/organ damage and neurological and behavioral deficits (8,9). I/R injury-induced damage is associated with dysregulation of complex interactions, inflammatory responses and extracellular matrix (ECM) remodeling (10). The ECM forms a key component of the basement membrane, and is essential for the permeability and integrity of blood-brain barrier (BBB), while matrix metalloproteinases (MMPs) can exert proteolytic activity to degrade all ECM components (1,11).

Abnormal degradation of ECM components in rats with cerebral I/R injury reduces the integrity and permeability of the BBB, which may lead to leakage of fluids from the blood to the brain and subsequent brain edema (1). Type IV collagen is a key component of the ECM complex and a major proteolytic substrate of MMP2 and MMP9 (1). Overexpression of MMP2 and MMP9 has previously been reported in I/R injured tissues, including the heart, liver and kidneys (1,3,12). Furthermore, breakdown of the BBB in rats may be induced by upregulated proteolytic activity of MMP2 and MMP9, which subsequently results in over-degradation of type IV collagen to reduce BBB integrity and promote BBB permeability (1).

At present, typical pharmacological agents used to treat cerebral I/R injury therapy are neuroprotective, antioxidative, antiapoptotic or anti-inflammatory agents (13,14). Previous studies have reported the protective effect of morroniside, an active extract of *Cornus officinalis*, against neuronal apoptosis, oxidative stress and inflammation (8,15-17). Furthermore, the neuroprotective effect of morroniside has been reported in rats with focal cerebral ischemia (8). However, to the best of our knowledge, the effect of morroniside on MMP2/9 expression and neuron apoptosis in cerebral I/R injury has yet to be reported.

The aim of the present study was to elucidate the effect of morroniside on MMP2/9 expression and neuron apoptosis in rats with cerebral I/R injury. Rats with focal cerebral I/R injury were administered with morroniside (30, 90 and 270 mg/kg) for 7 days, following which MMP2/9 expression and neuronal apoptosis were detected and comparatively analyzed.

Correspondence to: Dr Liying Deng, Department of Neurology, The Second Affiliated Hospital of Nanchang University, 1 Minde Road, Nanchang, Jiangxi 330006, P.R. China
E-mail: dengliying_1@tom.com

Key words: morroniside, cerebral ischemia/reperfusion injury, matrix metalloproteinase-2, matrix metalloproteinase-9

Materials and methods

Animal model and treatment. A total of 50 adult male Sprague Dawley rats (age, 7-8 weeks; weight, 260-280 g) were purchased from Beijing Vital River Laboratory Animal Technology Co., Ltd. (Beijing, China). All rats were housed at 25±2°C and 50% humidity, with a 12-h light/dark cycle, and food and water *ad libitum*. Ethical approval was granted by the Medical Ethics Committee of the Affiliated Ganzhou Hospital of Nanchang University (Nanchang, China).

Rats were randomly assigned into five groups (n=10 in each). Rats in the control group underwent sham surgery. All other rats underwent suture-occluded surgery as described by Longa *et al.* (18), with a 0.26 mm nylon monofilament (Beijing Shandong Biotech Co., Ltd., Beijing, China) inserted through the right common carotid artery and were divided into groups as follows: The cerebral I/R injury model group (model), no treatment; low dose group, 30 mg/kg/day morroniside (Sinopharm Chemical Reagent Co., Ltd., Shanghai, China) by gavage; moderate dose group, 90 mg/kg/day morroniside by gavage; high dose group, 270 mg/kg/day morroniside by gavage. Rats in the control and model groups received an equal volume of normal saline. Longa's five-grade scale methods were used to score neurological deficit following surgery (18) and rats with a score of 0 or 4 were excluded from the current study.

Immunofluorescence assay. Following 7 days of morroniside treatment, rats were anesthetized intravenously (30 mg/kg 5% pentobarbital sodium; Sigma Aldrich; Merck KGaA, Darmstadt, Germany) and brains were harvested. Brain tissues were mounted in optimal cutting temperature medium (Sakura Finetek USA, Inc., Torrance, CA, USA), frozen at -20°C in a Leica cryostat device (Leica Microsystems GmbH, Wetzlar, Germany) and cut into 4-μm sections. Sections were treated with 3% H₂O₂ solutions for 20 min at 37°C, blocked with normal goat serum (Sigma Aldrich; Merck KGaA) for 30 min at 37°C, and incubated with anti-MMP2 (cat. no. sc-13594) and anti-MMP9 (cat. no. sc-21733; 1:200; Santa Cruz Biotechnology, Inc., Dallas, TX, USA) antibodies at 4°C for 20 h. Sections were subsequently incubated with Cy3[®]-labelled goat anti-mouse immunoglobulin (Ig)G (cat. no. ab97035; 1:100; Abcam, Cambridge, MA, USA) in the dark at 37°C for 2 h. DAPI solution was used for nuclear staining at 37°C for 10 min. Slides were observed using a fluorescence microscope (magnification, x200) and assessed using Image-Pro Plus 6 software (Media Cybernetics, Inc., Rockville, MD USA).

Apoptotic cell analysis. Apoptotic cells in brain tissues were detected using a TUNEL assay. Briefly, ischemic penumbra cortex area was resected and the frozen sections (4-μm-thick) were fixed using 4% polyoxymethylene at room temperature for 20 min. and a colorimetric TUNEL kit (EMD Millipore, Billerica, MA, USA) was used according to the manufacturer's protocol for *in situ* apoptosis detection. The sections were washed by PBS and then incubated with prepared TUNEL solution for 1 h at 37°C in a dark chamber. A 50 μl DAB solution (2.5 μl DAB, 0.5 μl 30% H₂O₂ and 47 μl PBS) was added for nuclear staining at 37°C for 20 min. The TUNEL-positive

cells were observed and counted under a light microscope at a magnification of x200. Apoptotic cells were stained brown and the mean was calculated from 10 independent fields.

Western blotting. Brain tissues were homogenized and lysed in radioimmunoprecipitation assay buffer (Sigma Aldrich; Merck KGaA). Proteins were quantified using BCA method and equal amounts (50 μg) were separated by 8% SDS-PAGE. Proteins were transferred to a polyvinylidene fluoride membrane, blocked with 5% non-fat milk at 37°C for 1 h, and then was subsequently incubated with specific primary antibodies against MMP2 (cat. no. sc-13594), MMP9 (cat. no. sc-21733), active Caspase-3 (cat. no. sc-56052), B-cell lymphoma 2 (Bcl-2; cat. no. sc-56015) and Bcl-2-associated X protein (Bax; cat. no. sc-70407) and GAPDH (cat. no. sc-47724; all 1:1,000; Santa Cruz Biotechnology, Inc.) at 4°C overnight. The membrane was incubated with horseradish peroxidase-conjugated goat anti-rabbit IgG (cat. no. sc-2004; 1:1,000, Santa Cruz Biotechnology, Inc.) for 2 h at 37°C followed by enhanced chemiluminescence reagents (GE Healthcare Life Sciences, Little Chalfont, UK). Quantity One 4.4.1 software (Bio-Rad Laboratories, Inc., Hercules, CA, USA) was used to quantify bands with GAPDH as an internal control.

Reverse transcription-quantitative polymerase chain reaction (RT-qPCR). RT-qPCR was performed to measure the relative expression of MMP2 and MMP9 mRNA. Total RNA was extracted from brain tissues using TRIzol (Invitrogen; Thermo Fisher Scientific, Inc., Waltham, MA, USA) and RT was performed to synthesize cDNA. RNA was converted into cDNA using a High Capacity cDNA Reverse Transcription kit (Applied Biosystems; Thermo Fisher Scientific, Inc.) according to the manufacturer's protocol. MMP2 and MMP9 mRNA expression was detected on an ABI 7500 fast real time PCR platform (Applied Biosystems; Thermo Fisher Scientific, Inc.) using a SYBR Premix Ex Taq[™] kit (Takara Bio, Inc., Otsu, Japan) and the following primers: MMP2, forward 5'-CTGGGC CAGCCATCGCTGC-3' and reverse 5'-GCTTGCGGGGAA AGAAGTTG-3'; MMP9, forward 5'-GGCAGCCCCTGCTCC TGGTG-3' and reverse 5'-CCTTTAGTGTCTCGCTGTCC-3'; GAPDH, forward 5'-CTCCTCCTGGCCTCGCTGT-3' and reverse 5'-GCTGTCACCTTACCGTTCC-3'. Temperature protocols were as follows: 95°C for 4 min followed by 40 cycles of 95°C for 30 sec, 60°C for 35 sec and 72°C for 20 sec and a final extension step at 65°C for 6 min. The relative mRNA expression level was analyzed using the 2^{-ΔΔC_q} method with GAPDH as the internal reference gene (19). Each reaction was run in triplicate.

Statistical analysis. All data are expressed as the mean ± standard error of the mean. SPSS 17.0 software was used for statistical analyses (SPSS, Inc., Chicago, IL, USA). One-way analysis of variance followed by Scheffe's test was used to assess differences between groups. P<0.05 was considered to indicate a statistically significant difference.

Results

Morroniside reduces the expression of MMP2 and MMP9 in an I/R injury model. The results of western blotting and RT-qPCR revealed that MMP2 and MMP9 protein (Fig. 1A)




Figure 1. Western blotting and reverse transcription-quantitative polymerase chain reaction were performed to measure MMP2 and MMP9 (A) protein and (B) mRNA, respectively, in rat brain tissues (n=10). **P<0.01 vs. Con, &P<0.05 and &&P<0.01 vs. Model. MMP, matrix metalloproteinase; Con, control; Low, 30 mg/kg morroniside; Mod, 90 mg/kg morroniside; High, 270 mg/kg morroniside.




Figure 2. An immunofluorescence assay was performed to assess the expression of (A) MMP2 and (B) MMP9 in rat brain tissues (n=10). Magnification, x200. MMP, matrix metalloproteinase; Con, control; Low, 30 mg/kg morroniside; Mod, 90 mg/kg morroniside; High, 270 mg/kg morroniside.

and mRNA (P<0.01; Fig. 1B) expression was increased in the model group compared with the control group. However, treatment with morroniside reduced the expression of MMP2 and MMP9 mRNA (Fig. 1A) and protein (low dose, P<0.05; moderate and high doses, P<0.01; Fig. 1B) in a dose dependent manner compared with the model group. Immunofluorescence results revealed that the pattern of MMP2 and MMP9 positive cells in cortical infarctions followed a similar trend to protein expression (Fig. 2). The MMP2 and MMP9 fluorescence

intensity in the brains of rats was increased in the model group compared with the control (Fig. 2). However, treatment with morroniside markedly reduced MMP2 and MMP9 fluorescence compared with the model group (Fig. 2).

Morroniside inhibits neuron apoptosis. Apoptotic neurons on the sections from ischemic penumbra cortex were detected using a TUNEL assay. The percentage of apoptotic neurons was significantly increased in the model group compared




Figure 3. (A) A TUNEL assay was performed to quantify apoptotic neurons in a cerebral ischemia/reperfusion injury model (n=10 rats). (B) Western blotting was performed to measure the expression of active caspase-3, Bcl-2 and Bax protein. Magnification, $\times 200$. ** $P < 0.01$ vs. Con, & $P < 0.05$ and && $P < 0.01$ vs. Model. Bcl-2, B-cell lymphoma 2; Bax, Bcl-2-associated X protein; Con, control; Low, 30 mg/kg morroniside; Mod, 90 mg/kg morroniside; High, 270 mg/kg morroniside.

with the control ($P < 0.01$; Fig. 3A). However, treatment with morroniside significantly reduced I/R-associated neuron apoptosis in a dose dependent manner (low and moderate doses, $P < 0.05$; high dose, $P < 0.01$; Fig. 3A). The expression of apoptosis-associated proteins was assessed using western blotting. The results demonstrated that active caspase-3 and Bax were significantly upregulated in the model group compared with the control group ($P < 0.01$; Fig. 3B), while Bcl-2 was significantly downregulated ($P < 0.01$; Fig. 3B). The expression of active caspase-3 and Bax was significantly downregulated by morroniside treatment in a dose-dependent manner (low dose, $P < 0.05$; moderate and high doses, $P < 0.01$; Fig. 3B), while the expression of Bcl-2 was significantly upregulated (low dose, $P < 0.05$; moderate and high doses, $P < 0.01$; Fig. 3B).

Discussion

The upregulation of MMPs, in particular MMP2 and MMP9, results in the degradation of type IV collagen, which in

turn impairs BBB integrity and enhances I/R injuries in the brain (1). The present study is the first to demonstrate the effect of morroniside treatment on MMP2/9 expression and neuron apoptosis and in I/R. The results revealed that MMP2 and MMP9 are upregulated in the brains of rats with cerebral I/R injury, whereas the administration of morroniside significantly ameliorated this effect. These results suggest that morroniside may convey a protective effect in the brains of rats with I/R injury.

Increased MMP2 and MMP9 expression has been reported in I/R injuries of the lung, heart, skeletal muscle, kidneys and other organs (1-4). Previous studies have revealed that MMP2/9 overexpression disrupts the BBB and results in edema (20,21). It has also been reported that the inhibition of MMP2/9 may protect against cerebral ischemia and improve organ function (12,22-25). However, MMP2 knockout does not protect against acute focal ischemia injury in brain (26). Kato *et al* (10) demonstrated that MMP2 deletion was able to impair poly ADP-ribose polymerase 1

degradation, enhance MMP9 activity and exacerbate hepatic I/R injury in mice (10). It has also been reported that whereas MMP9 deficiency may protect against hepatic I/R injury (12). Together, these reports suggest that effect of MMP2/9 in I/R injury is complex and, while MMP2/9 inhibition may be beneficial, it is not necessarily required to improve I/R injury. However, the results of the present study demonstrated that morroniside administration inhibits MMP2 and MMP9 in I/R injured brains, revealing morroniside was beneficial to protect against cerebral I/R injury.

The results of the present study revealed that morroniside significantly reduces the percentage of apoptotic neurons, inhibits caspase-3 activation and increases the Bcl-2/Bax protein ratio in rats with cerebral I/R injury. Cerebral I/R injury results in neurological and behavioral deficits, which are accompanied by neuron apoptosis (9,27,28). Wang *et al* (16) reported that morroniside was able to reduce H₂O₂-induced SH-SY5Y cell apoptosis, intracellular Ca²⁺ accumulation and mitochondrial membrane potential, while also inhibiting the H₂O₂-induced decrease in superoxide dismutase activity (16). The protective effect of morroniside on cell apoptosis and other diseases has been widely reported (8,29,30). The results of the present study suggest that morroniside treatment has a protective effect against I/R-induced neuron apoptosis; future studies should investigate the effect of MMP2 or MMP9 in rats with I/R injury to confirm the mechanisms of morroniside.

In conclusion, the present study revealed that protective effect of morroniside in rats with cerebral I/R injury. Morroniside inhibits the I/R-induced upregulation of MMP2/9 and neuron apoptosis, suggesting that morroniside may promote behavioral deficit repair following cerebral I/R injury. However, the underlying mechanism responsible for these effects remains to be elucidated.

Acknowledgements

Not applicable.

Funding

No funding was received.

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Authors' contributions

GZ wrote the manuscript and conducted the majority of the experiments. WD, YL and MS collected and analyzed the data. LD designed the study and approved the manuscript for submission.

Ethics approval and consent to participate

Ethical approval was granted by the Medical Ethics Committee of the affiliated Ganzhou Hospital of Nanchang University (Nanchang, China).

Consent for publication

Not applicable.

Competing interests

All of the authors have no conflict of interest in this research.

References

1. Roach DM, Fitrige RA, Laws PE, Millard SH, Varelias A and Cowled PA: Up-regulation of MMP-2 and MMP-9 leads to degradation of type IV collagen during skeletal muscle reperfusion injury; protection by the MMP inhibitor, doxycycline. *Eur J Vasc Endovasc Surg* 23: 260-269, 2002.
2. Yano M, Omoto Y, Yamakawa Y, Nakashima Y, Kiriya M, Saito Y and Fujii Y: Increased matrix metalloproteinase 9 activity and mRNA expression in lung ischemia-reperfusion injury. *J Heart Lung Transplant* 20: 679-686, 2001.
3. Cheung PY, Sawicki G, Wozniak M, Wang W, Radomski MW and Schulz R: Matrix metalloproteinase-2 contributes to ischemia-reperfusion injury in the heart. *Circulation* 101: 1833-1839, 2000.
4. Zhang X, Sakamoto T, Hata Y, Kubota T, Hisatomi T, Murata T, Ishibashi T and Inomata H: Expression of matrix metalloproteinases and their inhibitors in experimental retinal ischemia-reperfusion injury in rats. *Exp Eye Res* 74: 577-584, 2002.
5. Navarro-Sobrinho M, Rosell A, Hernández-Guillamon M, Penalba A, Boada C, Domingues-Montanari S, Ribó M, Alvarez-Sabín J and Montaner J: A large screening of angiogenesis biomarkers and their association with neurological outcome after ischemic stroke. *Atherosclerosis* 216: 205-211, 2011.
6. Feigin VL, Forouzanfar MH, Krishnamurthi R, Mensah GA, Connor M, Bennett DA, Moran AE, Sacco RL, Anderson L, Truelsen T, *et al*: Global and regional burden of stroke during 1990-2010: Findings from the Global burden of disease study 2010. *Lancet* 383: 245-254, 2014.
7. Chen L, Xiang Y, Kong L, Zhang X, Sun B, Wei X and Liu H: Hydroxysafflor yellow A protects against cerebral ischemia-reperfusion injury by anti-apoptotic effect through PI3K/Akt/GSK3 β pathway in rat. *Neurochem Res* 38: 2268-2275, 2013.
8. Wang W, Xu J, Li L, Wang P, Ji X, Ai H, Zhang L and Li L: Neuroprotective effect of morroniside on focal cerebral ischemia in rats. *Brain Res Bull* 83: 196-201, 2010.
9. Wang Q, Sun AY, Simonyi A, Jensen MD, Shelat PB, Rottinghaus GE, MacDonald RS, Miller DK, Lubahn DE, Weisman GA and Sun GY: Neuroprotective mechanisms of curcumin against cerebral ischemia-induced neuronal apoptosis and behavioral deficits. *J Neurosci Res* 82: 138-148, 2005.
10. Kato H, Duarte S, Liu D, Busuttill RW and Coito AJ: Matrix metalloproteinase-2 (MMP-2) gene deletion enhances MMP-9 activity, impairs PARP-1 degradation, and exacerbates hepatic ischemia and reperfusion injury in mice. *PLoS One* 10: e0137642, 2015.
11. Rosenberg GA, Estrada E, Kelley RO and Kornfeld M: Bacterial collagenase disrupts extracellular matrix and opens blood-brain barrier in rat. *Neurosci Lett* 160: 117-119, 1993.
12. Hamada T, Fondevila C, Busuttill RW and Coito AJ: Metalloproteinase-9 deficiency protects against hepatic ischemia/reperfusion injury. *Hepatology* 47: 186-198, 2008.
13. Ai J, Wan H, Shu M, Zhou H, Zhao T, Fu W and He Y: Guhong injection protects against focal cerebral ischemia-reperfusion injury via anti-inflammatory effects in rats. *Arch Pharm Res* 40: 610-622, 2017.
14. Zhao P, Zhou R, Zhu XY, Hao YJ, Li N, Wang J, Niu Y, Sun T, Li YX and Yu JQ: Matrine attenuates focal cerebral ischemic injury by improving antioxidant activity and inhibiting apoptosis in mice. *Int J Mol Med* 36: 633-644, 2015.
15. Xu HQ, Hao HP, Zhang X and Pan Y: Morroniside protects cultured human umbilical vein endothelial cells from damage by high ambient glucose. *Acta Pharmacol Sin* 25: 412-415, 2004.
16. Wang W, Sun F, An Y, Ai H, Zhang L, Huang W and Li L: Morroniside protects human neuroblastoma SH-SY5Y cells against hydrogen peroxide-induced cytotoxicity. *Eur J Pharmacol* 613: 19-23, 2009.

17. Park CH, Noh JS, Kim JH, Tanaka T, Zhao Q, Matsumoto K, Shibahara N and Yokozawa T: Evaluation of morroniside, iridoid glycoside from *Corni Fructus*, on diabetes-induced alterations such as oxidative stress, inflammation, and apoptosis in the liver of type 2 diabetic db/db mice. *Biol Pharm Bull* 34: 1559-1565, 2011.
18. Longa EZ, Weinstein PR, Carlson S and Cummins R: Reversible middle cerebral artery occlusion without craniectomy in rats. *Stroke* 20: 84-91, 1989.
19. Livak KJ and Schmittgen TD: Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) method. *Methods* 25: 402-408, 2001.
20. Pérez-Hernández M, Fernández-Valle ME, Rubio-Araiz A, Vidal R, Gutiérrez-López MD, O'shea E and Colado MI: 3,4-Methylenedioxymethamphetamine (MDMA, ecstasy) produces edema due to BBB disruption induced by MMP-9 activation in rat hippocampus. *Neuropharmacology* 118: 157-166, 2017.
21. Tasaki A, Shimizu F, Sano Y, Fujisawa M, Takahashi T, Haruki H, Abe M, Koga M and Kanda T: Autocrine MMP-2/9 secretion increases the BBB permeability in neuromyelitis optica. *J Neurol Neurosurg Psychiatry* 85: 419-430, 2014.
22. Lin HB, Cadete VJ, Sra B, Sawicka J, Chen Z, Bekar LK, Cayabyab F and Sawicki G: Inhibition of MMP-2 expression with siRNA increases baseline cardiomyocyte contractility and protects against simulated ischemic reperfusion injury. *Biomed Res Int* 2014: 810371, 2014.
23. Yan P, Chen KJ, Wu J, Sun L, Sung HW, Weisel RD, Xie J and Li RK: The use of MMP2 antibody-conjugated cationic microbubble to target the ischemic myocardium, enhance Timp3 gene transfection and improve cardiac function. *Biomaterials* 35: 1063-1073, 2014.
24. Lin H, Sra B, Cadete VJ, Sawicka J, Cayabyab FS and Sawicki G: 774 MMP-2 siRNA Protects Against Contractile Dysfunction in Cardiomyocytes Subjected to Ischemia/Reperfusion. *Canadian J Cardiology* 28: S396-S397, 2012.
25. Zhang HT, Zhang P, Gao Y, Li CL, Wang HJ, Chen LC, Feng Y, Li RY, Li YL and Jiang CL: Early VEGF inhibition attenuates blood-brain barrier disruption in ischemic rat brains by regulating the expression of MMPs. *Mol Med Rep* 15: 57-64, 2017.
26. Asahi M, Sumii T, Fini ME, Itohara S and Lo EH: Matrix metalloproteinase 2 gene knockout has no effect on acute brain injury after focal ischemia. *Neuroreport* 12: 3003-3007, 2001.
27. Hua F, Ma J, Ha T, Xia Y, Kelley J, Williams DL, Kao RL, Browder IW, Schweitzer JB, Kalbfleisch JH and Li C: Activation of toll-like receptor 4 signaling contributes to hippocampal neuronal death following global cerebral ischemia/reperfusion. *J Neuroimmunol* 190: 101-111, 2007.
28. Liang K, Ye Y, Wang Y, Zhang J and Li C: Formononetin mediates neuroprotection against cerebral ischemia/reperfusion in rats via downregulation of the Bax/Bcl-2 ratio and upregulation PI3K/Akt signaling pathway. *J Neurol Sci* 344: 100-104, 2014.
29. Wang W, Huang W, Li L, Ai H, Sun F, Liu C and An Y: Morroniside prevents peroxide-induced apoptosis by induction of endogenous glutathione in human neuroblastoma cells. *Cell Mol Neurobiol* 28: 293-305, 2008.
30. Yokozawa T, Yamabe N, Kim HY, Kang KS, Hur JM, Park CH and Tanaka T: Protective effects of morroniside isolated from *Corni Fructus* against renal damage in streptozotocin-induced diabetic rats. *Biol Pharm Bull* 31: 1422-1428, 2008.



This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) License.