

Activity-dependent changes in MAPK activation in the Angelman Syndrome mouse model

Irina Filonova, Justin H. Trotter, Jessica L. Banko, and Edwin J. Weeber¹

Department of Molecular Pharmacology and Physiology, Morsani College of Medicine, University of South Florida, Byrd Alzheimer's Institute, Tampa, Florida 33613, USA

Angelman Syndrome (AS) is a devastating neurological disorder caused by disruption of the maternal *UBE3A* gene. Ube3a protein is identified as an E3 ubiquitin ligase that shows neuron-specific imprinting. Despite extensive research evaluating the localization and basal expression profiles of Ube3a in mouse models, the molecular mechanisms whereby Ube3a deficiency results in AS are enigmatic. Using in vitro and in vivo systems we show dramatic changes in the expression of Ube3a following synaptic activation. In primary neuronal culture, neuronal depolarization was found to increase both nuclear and cytoplasmic Ube3a levels. Analogous up-regulation in maternal and paternal Ube3a expression was observed in Ube3a-YFP reporter mice following fear conditioning. Absence of Ube3a led to deficits in the activity-dependent increases in ERK1/2 phosphorylation, which may contribute to reported deficits in synaptic plasticity and cognitive function in AS mice. Taken together, our findings provide novel insight into the regulation of Ube3a by synaptic activity and its potential role in kinase regulation.

Angelman Syndrome (AS) is a neurological disorder that affects ~1:12,000 children (Steffenburg et al. 1996) and is characterized by severe developmental delay, cognitive disruption, an absence of speech, contagiously happy demeanor, ataxia, and a greater propensity for seizures (Pelc et al. 2008; Dagli et al. 2011). Absence, mutation, or disruption of the maternal *UBE3A* allele results in a near complete absence of protein expression in the central nervous system (CNS). *UBE3A* encodes an E3 ubiquitin ligase (E6-AP) that shows neuron-specific paternal imprinting in humans and mice (Jiang et al. 1998; Gustin et al. 2010; Daily et al. 2012). Ube3a has few identified biochemical targets including activity-regulated cytoskeleton-associated protein (Arc) and Ephexin 5 (Greer et al. 2010; Margolis et al. 2010); however, Ube3a is likely to play multiple roles in the neuron and no single target of Ube3a fully explains the unique phenotype associated with the disorder.

A recent study by Greer et al. (2010) demonstrated dynamic expression of Ube3a during synaptic activity. Specifically, Ube3a mRNA was significantly increased in response to membrane depolarization or glutamate receptor activation in cultured primary neurons. In addition, Ube3a expression was up-regulated in the mouse brain during kainate-induced seizures and in response to enhanced environmental stimuli during the novel object exploration test (Greer et al. 2010). Stringent control of Ube3a expression is necessary for neuronal functioning. A prime example of this is human *UBE3A* gene dosage. Absence of *UBE3A* results in AS, while the interstitial duplication of *UBE3A* is associated with autism spectrum disorders (ASD) (Schanen 2006). Supportive observations are made in mice lacking Ube3a and those with a Ube3a duplication, both of which partially phenocopy aspects of AS and ASD, respectively. Maternal Ube3a-deficient animals show impaired locomotor activity, enhanced seizure propensity, and deficits in associative and spatial learning and memory (Jiang et al. 1998). Conversely, duplication of Ube3a results in poor social interaction, behavioral inflexibility, and anxiety (Smith et al. 2011). Taken together, these data suggest that there is a critical ho-

meostatic level of Ube3a expression that must be maintained for normal neuronal function.

In the present study, we evaluate changes in Ube3a expression in response to neuronal depolarization in vitro and neuronal activity in vivo. We present evidence that control of Ube3a expression is critical for activity-dependent kinase activation. These results better define the role of Ube3a in normal synaptic function in the CNS, expand our knowledge about the paternal Ube3a protein expression, and examine consequences of Ube3a absence in the context of neuronal activity.

Results

Ube3a expression changes in primary neurons immediately after neuronal depolarization

Maternal Ube3a protein is present in the neuronal nucleus, soma, and dendrites, and is enriched in the postsynaptic compartment (Dindot et al. 2008; Gustin et al. 2010). Subcellular fractionation was used to isolate soluble and nonsoluble nuclear, membrane, and cytosolic compartments (Fig. 1A). To verify the specificity of the Ube3a antibody, hippocampi from Ube3a-deficient mice (Ube3a m⁻/p⁻) and littermate controls were fractionated. The antibody shows specificity to Ube3a (~110 kDa) with a complete absence of reactivity in Ube3a m⁻/p⁻ mice. We observed a single unidentified protein band of a slightly higher molecular weight detected in the nuclear and chromatin-bound fractions. We confirmed quality of fractions by detection of HSP90, ApoER2, NeuN, and histone H3 to the cytosolic, membrane, nuclear, and chromatin-associated fractions, respectively (Fig. 1A). Consistent with previous reports (Gustin et al. 2010), we find that Ube3a is enriched in the cytosol with lower detection in the nucleus and membrane compartments. Next, we determined if synaptic

¹Corresponding author
E-mail eweeber@health.usf.edu

Article is online at <http://www.learnmem.org/cgi/doi/10.1101/lm.032375.113>.

© 2014 Filonova et al. This article is distributed exclusively by Cold Spring Harbor Laboratory Press for the first 12 months after the full-issue publication date (see <http://learnmem.cshlp.org/site/misc/terms.xhtml>). After 12 months, it is available under a Creative Commons License (Attribution-NonCommercial 3.0 Unported), as described at <http://creativecommons.org/licenses/by-nc/3.0/>.




Figure 1. Ube3a expression changes in cytosol and nucleus following 55 mM KCl application. (A) Subcellular fractionation technique was used to isolate cellular fractions with minimal cross-contamination. (CEB) cytosolic, (MEB) membrane, (NEB) nuclear soluble, (CRB) chromatin bound. Representative Western blot of Ube3a total protein increase in cytosolic and nuclear fractions at 5 min (B) and 30 min (C) following 55 mM KCl stimulation in primary neuronal cultures (DIV 8-10). Ube3a is shown in two exposures (low [upper panel], high [lower panel]). (NT) Nontreated control. (D-F) Quantification of Western blot analysis ($n = 5$, treatments standardized to NT group, one-way ANOVA, post hoc Tukey's test, [*] $P < 0.05$, [**] $P < 0.01$).

depolarization could modulate Ube3a protein expression in neuronal culture. Mixed primary cultures (DIV 8-10) prepared from E17-18 wild-type brains were depolarized with 55 mM KCl for 5 and 30 min. Western blot analysis revealed that Ube3a expression following KCl treatment is significantly increased in the nucleus and cytosol as early as 5 min (nuclear soluble, $201 \pm 21.42\%$, $P < 0.05$; cytosol, $153.5 \pm 16.59\%$, $P < 0.05$) and remained elevated up to 30 min (nucleus, $182.4 \pm 34.78\%$, $P < 0.05$; cytosol, $167.3 \pm 17.51\%$, $P < 0.01$) (Fig. 1B-F). Ube3a was slightly down-regulated in the membrane fraction at 30 min; however, the decrease failed to reach statistical significance. These data demonstrate that KCl-induced depolarization affects Ube3a protein levels resulting in increased Ube3a expression in the cytosol and the nucleus.

Consequences of Ube3a absence on depolarization-induced kinase activation

The wide variety and severity of AS symptoms suggests that the loss of Ube3a causes a global disruption of synaptic function. This is supported in the AS mouse model with identified synaptic disruption in the visual cortex, cerebellum, and hippocampus (Jiang et al. 1998; Heck et al. 2008; Yashiro et al. 2009). A multitude

of Ca^{2+} -dependent signal transduction pathways are stimulated by synaptic depolarization. Previously, basal levels of several kinases important for LTP induction and maintenance in the hippocampus of AS mice were examined in the attempt to identify direct or indirect targets of Ube3a ubiquitination. In this study, we sought to revisit the p44/p42 extracellular signal-regulated kinase (ERK1/2) and evaluate its response to neuronal depolarization in wild-type (Ube3a m+/p+) and Ube3a-deficient mice (Ube3a m-/p+). The necessity of ERK p44/p42 (ERK1/2) is well established for synaptic plasticity and memory formation (English and Sweatt 1997; Atkins et al. 1998; Thomas and Haganir 2004).

Acute hippocampal slices from Ube3a m-/p+ mice and wild-type littermates were treated with 55 mM KCl for 5 or 30 min. Following Western blotting, the levels of total and phosphorylated ERK1/2 (p-ERK, Thr202/Tyr204) and mitogen-activated kinase 1/2 (MEK1/2, p-MEK [Ser217/221]) were analyzed. Under basal conditions, Ube3a m+/p+ and Ube3a m-/p+ groups ($n = 6-7$ slices, three animals per genotype) had similar levels of total ERK1/2 and p-ERK (Fig. 2A,B). ERK1/2 activation was determined by same blot p-ERK to total ERK ratio and standardized to nontreated controls within genotype. All treated groups show elevated p-ERK after depolarization; however, p-ERK levels in Ube3a m-/p+ slices (p-ERK1, $311 \pm 20.83\%$; p-ERK2, $207 \pm 16.98\%$; $P < 0.01$, two-way ANOVA) failed to reach wild-type levels (p-ERK1, 477.2 ± 57.65 ; p-ERK2, $354 \pm 80\%$) at 5 min post depolarization (Fig. 2C,D). Interestingly, levels were elevated from basal levels in

both groups 30 min post depolarization, but were not significantly different ($n = 6$ per genotype). Basal expression, phosphorylation, and activation of MEK were not affected by the absence of Ube3a (Fig. 2E,F). These data suggest that after neuronal depolarization only the immediate activation of ERK is impaired.

Spatial and temporal properties of Ube3a expression following fear conditioning training

Next, we wanted to determine if similarities in the observed changes exist for Ube3a and ERK1/2 in vivo. Recent reports suggest a dynamic regulation of Ube3a and a close integration of this regulation with synaptic plasticity in wild-type mouse brain (Greer et al. 2010). The Ube3a allele is believed to show biallelic expression in all cell types, with the exception of neurons. However, maternal deletion shows a near 100% absence of Ube3a detection in the CNS (Fig. 3). Increased Western blot exposure reveals Ube3a protein that cannot be detected in the Ube3a m-/p- mouse brain (Fig. 3). This observation raises the question of whether dynamic regulation of the paternal allele is occurring, albeit at a reduced expression, or if the detection of protein is from nonneuronal sources. Therefore, we sought to determine if both maternal and paternal copies respond to neuronal activation.




Figure 2. Reduced KCl depolarization-dependent p-ERK phosphorylation in acute hippocampal slices of Ube3a m⁻/p⁺ mice. (A) Representative immunoblot. (B,E) Levels of total ERK and total MEK1/2 are unchanged in m⁻/p⁺ mice. (C,D) Quantitative Western blot analysis shows significantly decreased levels in pERK1/2 activation in hippocampal slices of Ube3a m⁻/p⁺ mice at 5 min but not 30 min after 55 mM KCl application ($n = 5$, treatments standardized to NT group within genotype, two-way ANOVA, post hoc Tukey's test, [*] $P < 0.05$, [**] $P < 0.01$). (F) KCl-induced phosphorylation of MEK1/2 is normal in both genotypes.

The heterozygous Ube3a-Yellow Fluorescent Protein (Ube3a-YFP) fusion protein reporter mouse allows the ability to distinguish paternal from maternal protein expression by increasing the molecular weight of a single allele with the addition of the YFP. Figure 4A shows that paternal Ube3a-YFP is resolved higher than maternal due to the presence of the 20 kD YFP tag. Using this strategy, we next sought to determine if Ube3a protein changes could be detected with *in vivo* synaptic activity following fear conditioned associative memory formation. Fear conditioning (FC) was performed using a three foot-shock paradigm. To verify that this training paradigm leads to robust memory formation, a separate group of paternal Ube3a-YFP mice was shown to have a significant increase in freezing behavior compared to a no shock, context only group (Fig. 4B). Western blot analysis of the whole hippocampus and cortex collected at 5-min, 15-min, 1-h, 3-h, and 6-h intervals post training revealed significant changes in maternal Ube3a expression in the hippocampus ($149.6 \pm 14.68\%$, $P < 0.05$) and cortex ($378.4 \pm 78.15\%$, $P < 0.001$) at 6 h after FC when compared to control group (Fig. 4C). Remarkably, paternal Ube3a was significantly elevated only in hippocampus (1 h, $185.6 \pm 15.82\%$; 3 h, $196 \pm 14.05\%$; 6 h, $210 \pm 19.65\%$; $P < 0.01$) of pUbe3a-YFP mice starting at the 1-h time point; however, no similar response was observed in cortical samples (Fig. 4D). To evaluate if exposure to the context could result in altered Ube3a ex-

pression, a group of naïve mice (handled only) was examined and showed no difference (data not shown). To determine if paternal Ube3a-YFP is affected by the inclusion of YFP, maternal Ube3a-YFP animals were generated and Ube3a-YFP was compared to naïve Ube3a. We found reduced maternal ($86.6 \pm 3.38\%$, $P < 0.05$) or paternal Ube3a ($56 \pm 13.56\%$, $P < 0.01$) expression of the Ube3a-YFP fusion protein compared to naïve wild-type protein (Fig. 4E,F).

Taken together our results suggest that both maternal and paternal alleles are activated by synaptic activity. It appears that paternal Ube3a expression has a specific temporal expression profile that is similar to the maternal activation in hippocampus, but not in cortex. Paternal Ube3a comprises ~3%–5% of total Ube3a in neurons; thus, the amount of paternal protein increase is negligible. Nevertheless, it is critical to consider and further characterize the paternal Ube3a allele, especially given the growing interest in gene-based therapy approaches.

Activity-dependent p-ERK dysregulation in Ube3a deficient mouse model

Reduced ERK activation in response to neuronal stimulation may contribute to the cognitive disruption present in Ube3a maternal deficient mice and human AS. Numerous studies report that reduction in ERK activity disrupts associative fear conditioned memory formation (Blum et al. 1999; Selcher et al. 2003). Next we determined if hippocampal-dependent associative learning in adult Ube3a m⁻/p⁺ showed reduced

p-ERK activation in area CA1 following a three foot-shock FC paradigm. This experimental approach has been previously shown increased ERK activation in area CA1 of the hippocampus (Chwang et al. 2006). We subjected both Ube3a m⁺/p⁺ ($n = 5$ control, $n = 6$ experimental) and Ube3a m⁻/p⁺ ($n = 5$ for control, $n = 4$ experimental) 3- to 4-mo-old mice to FC and measured p-ERK levels in CA1 60 min after to FC training. No significant change in total ERK1/2 or p-ERK1/2 was observed at baseline (Fig. 5B). Significant differences in p-ERK activation became apparent when the samples were compared at 60 min post training. Consistent with published data (Chwang et al. 2006) area CA1 of Ube3a m⁺/p⁺ hippocampi showed a significant increase in p-ERK2 levels ($143.4 \pm 14.19\%$ of Ube3a m⁺/p⁺, $P < 0.05$) compared to control group. By comparison, p-ERK2 activation was blunted in Ube3a m⁻/p⁺ animals ($82.10 \pm 4.6\%$ of Ube3a m⁻/p⁺ controls). Analysis of CA3 and DG areas did not reveal changes in total or p-ERK in either wild-type or in transgenic littermates (data not shown).

Discussion

A preponderance of imprinted genes is associated with placental regulation and represents a genetically unique method of regulating protein expression. However, maternal Ube3a levels




Figure 3. Ube3a expression in Ube3a m+/p+, Ube3a m-/p+, and Ube3a m-/p- mouse brains. (A) Paternal Ube3a is expressed at low levels throughout in the Ube3a m-/p+ mouse brain. (HIP) Hippocampus, (PFC) prefrontal cortex, (COR) cortex, (STR) striatum, (CER) cerebellum. Immunostaining (B) and subsequent analysis (C) of coronal sections of mouse brain demonstrated near absence of paternal Ube3a expression in Ube3a m-/p+ mice ($n = 3$) and no Ube3a expression in Ube3a m-/p- ($n = 2$) compared to wild-type controls ($n = 3$) (one-way ANOVA, post hoc Tukey's test, [*] $P < 0.05$).

can undergo further regulation beyond paternal allele silencing. For example, Ube3a is a target of itself, suggesting a negative feedback for regulating protein level. This form of biochemical regulation may be in response to the recent discovery of altered neuronal Ube3a expression following glutamatergic activation and neuronal depolarization. A wide variety of activators, including *N*-methyl-D-aspartate (NMDA), α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid (AMPA), and brain-derived neurotrophic factor (BDNF), can initiate an increase of Ube3a mRNA within 5 h in primary cultures (Greer et al. 2010). In light of direct and indirect alterations of Arc and CaMKII, both major players in synaptic plasticity, changes in Ube3a expression suggest a role in synaptic function that supersedes the traditional roles of an E3 ubiquitin ligase. We sought to determine if Ube3a is altered in response to neuronal depolarization and in vivo learning and memory. Furthermore, we wanted to investigate if other signaling proteins modulated by calcium influx were likewise altered during synaptic activity in the Angelman Syndrome mouse model.

Although we did not specifically delineate isoform expression, we found increased total Ube3a in response to KCl treatment at 5 and 30 min post application in both cytosolic and nuclear fractions isolated from primary neurons, suggesting that this is not an isoform-dependent phenomenon. Although ubiquitously expressed, Ube3a localizes to the nucleus where it serves as a transcriptional coactivator and plays a role in the chromosome segregation (Nawaz et al. 1999; Singhmar and Kumar 2011). Nuclear association is also expected due to other Ube3a targets, including p53, p27, or HHR23A (Jiang et al. 1998; Kumar et al. 1999; Mishra and Jana 2008; Mishra et al. 2009). However, the potential role for up-regulation of nuclear associated Ube3a is yet to be de-

termined. Greer et al. (2010) reported that Ube3a mRNA is up-regulated in mouse hippocampus 1 h following novel environment exploration task. In our study we utilized the three foot-shock fear conditioning paradigm to examine spatial and temporal Ube3a changes in the paternal Ube3a-YFP reporter mouse brain. This particular model allows quantification of both maternal and paternal proteins due to the differential molecular weight of the paternal-derived Ube3a-YFP protein. Our results demonstrate an increase in both maternal and paternal Ube3a expression starting at 1 h post training in hippocampus, while only maternal Ube3a expression is increased in the cortex. The spatial differences in expression profiles may reflect the behavioral paradigm, as fear conditioning is a hippocampal-dependent behavioral task requiring extensive coordination of hippocampal pyramidal cell activity. Alternatively, the cortex may represent a brain region where the imprinting center at 15q11 is much more tightly regulated and the silencing mechanism is more stringent or insensitive to fear conditioning. However, it is intriguing to find that neuronal activity might be sufficient to overcome epigenetic silencing and allow, albeit modest, production of the paternal Ube3a protein.

Identifying activity-dependent regulation of Ube3a protein raises the possibility that Ube3a is involved in activities other than housekeeping protein degradation. It stands to reason that Ube3a may be involved in modulating pathways that are involved in synaptic plasticity, and this may explain why our earlier work investigating kinase alterations in the Ube3a maternal deficient mouse only revealed an alteration in α -CaMKII phosphorylation (Weeber et al. 2003). Those previous experiments were performed during basal conditions. The use of KCl-induced neuronal depolarization should effectively activate multiple signaling pathways leading to ERK activation. Interestingly, we demonstrate reduced phosphorylation of ERK in response to KCl depolarization in the Ube3a m-/p+ mouse, suggesting that Ube3a is involved in a signaling cascade that targets ERK phosphorylation. Curiously, we did not find a concurrent difference in either total MEK or p-MEK levels, the kinase responsible for phosphorylation and activation of ERK. This result may reflect a direct or indirect action of Ube3a on MEK's ability to phosphorylate ERK or an Ube3a-dependent regulation of phosphatases. The exact mechanism for this difference in ERK activation is currently under investigation.

There has been extensive investigation into the regulation of ERK in mice performing various behavioral tasks. Depending on the learning paradigm, the p-ERK levels are shown to increase as early as 1 min post training and stay elevated for up to 9 h (Blum et al. 1999; Chwang et al. 2006; Trifilieff et al. 2006, 2007). Both memory formation and hippocampal synaptic plasticity are sensitive to inhibition of ERK activation and reduced p-ERK levels correlate nicely with deficits in memory retention (Blum et al. 1999; Selcher et al. 1999). Our finding that p-ERK levels are reduced following contextual fear conditioning in Ube3a m-/p+ mice provides an additional rationale to explain




Figure 4. Changes in maternal and paternal Ube3a expression following contextual fear conditioning. (A) YFP-tagged Ube3a has higher molecular weight than wild-type Ube3a. Depending on the paternal transmission, animals can carry an imprinted paternal Ube3a-YFP gene and wild-type maternal gene (pUbe3a-YFP) or imprinted paternal wild-type Ube3a and maternal Ube3a-YFP (mUbe3a-YFP). Addition of YFP affects basal expression of Ube3a. Expressions of mUbe3a-YFP and pUbe3a proteins are significantly reduced compared to wild-type mUbe3a (C) and pUbe3a (D), respectively (mUbe3a-YFP, $n = 6$; pUbe3a-YFP, $n = 4$; Student's t -test, [*] $P < 0.05$, [**] $P < 0.01$). (B) pUbe3a-YFP animals that received three shocks freeze significantly more than littermate pUbe3a-YFP controls 24 h later ($n = 5$ per group; Student t -test, $df = 8$, [***] $P < 0.001$). (E, F) mUbe3a expression is elevated starting at 1 h and significantly increased in the hippocampus (C) and cortex (D) of pUbe3a-YFP animals at 6 h. pUbe3a-YFP is only increased in hippocampus. All the time points are normalized and compared to "context only" control group (ctr) ($n = 3$ per group, one-way ANOVA, Tukey's post hoc test, [*] $P < 0.05$, [**] $P < 0.01$, [***] $P < 0.001$).

the well-established associative learning deficit exhibited by these mice (Jiang et al. 1998; van Woerden et al. 2007; Huang et al. 2013).

In conclusion, the present study suggests that Ube3a protein levels can change significantly in response to neuronal depolarization and, importantly, during the processes of associative learning and memory. Increased levels in both maternal and paternal Ube3a are intriguing and raise important questions about regulatory mechanisms of both maternal and paternal alleles. Furthermore, subtle changes in the paternal Ube3a allele, its distribution, isoform production, and function, should be addressed before considering it as a potential therapeutic target.

Materials and Methods

Animals and reagents

All animals utilized in the experiments were handled in accordance with protocols approved by the Institutional Animal Care and Use Committee of the University of South Florida. To generate paternally imprinted Ube3a-YFP, C57 females were bred to Ube3a-YFP males. Maternal Ube3a-YFP mutants were obtained by the opposite crosses. pUbe3a-YFP males and females that were 3–3.5 mo old were used in fear conditioning experiments to establish Ube3a expression profiles. 129SV/C57 hybrid Ube3a m^+/p^- and wild-type littermates 2–4 mo old were used to assess p-ERK activation. Antibodies: E6-AP (Bethyl lab); actin, ERK1/2, p-ERK1/2, MEK1/2, p-MEK1/2, Histone H3, and HSP90 (Cell signaling); NeuN (Millipore); ApoER2 (Abcam). Chemicals: KCl (50 mM, Sigma, P9549).

Fear conditioning and tissue collection

Paternal Ube3a-YFP mice were handled 5 min/d for 5 d in the room where fear conditioning training took place. On the day of experiment, each animal was placed in a sound attenuation chamber (SAC) for 4–5 min; after 3-min habituation three 0.5-mA shocks with 30-sec intervals were applied to a floor grid. Mice were euthanized with isoflurine at 5 min, 15 min, 1 h, 3 h, and 6 h following conditioning (three animals per group, males and females). Mouse brains were removed and cortex and hippocampus were dissected in ice-cold PBS containing Halt protease and phosphatase inhibitors cocktail (Thermo Scientific, 78440). Dissected tissue was frozen on dry ice and stored at -80°C until further processing. A control group of mice was subjected to fear conditioning training and was tested 24 h later. Freezing behavior was recorded and scored by AnyMaze software.

Western blot

Frozen tissue and primary neuronal cultures were homogenized by sonication in ice-cold RIPA buffer (50 mM Tris-HCl, pH 7.4, 150 mM NaCl, 2 mM EDTA, 0.1% SDS, 0.5% sodium deoxycholate, 1% triton $\times 100$) containing a Halt protease and phosphatase inhibitors cocktail. The lysates were clarified at 22,000g for 30 min. The protein concentrations were determined using the BCA Protein Analysis Kit (Pierce). Samples were resolved by 10% TGX gels (Bio-Rad) and transferred to PVDF membrane. Membranes were blocked with 5% milk in TBST (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 0.05% Tween-20) for 1 h and incubated with primary antibodies at 4°C overnight. After three 10-min washes with TBST, secondary antibodies were applied for 1 h at room temperature. Blots were detected with film for chemiluminescence using Pierce ECL. Developed images were analyzed using ImageJ software. Statistical analysis (Student t -test, ANOVA) was performed using GraphPad Software.




Figure 5. (A,C) Disregulation of p-ERK activation in Ube3a m⁻/p⁺ mouse brain following fear conditioning. Ube3a m⁻/p⁺ and their littermate controls were subjected to a three shocks conditioning paradigm. ERK1/2 phosphorylation was determined by densitometry. p-ERK2 expression was significantly increased in wild-type CA1 hippocampal region ($n = 4-5$ per group, one-way ANOVA, Tukey's post hoc test, [*] $P < 0.05$, [**] $P < 0.01$). Fear conditioning failed to induce the same changes in Ube3a m⁻/p⁺ mice. (B,D) There was no change in p-ERK1/2 or total ERK1/2 expression between Ube3a deficient and littermate control groups.

Subcellular fractionation

Mouse brains and murine mixed primary neuronal cultures were fractionated using the Subcellular Protein Fractionation Kit for Cultured Cells (Thermo Scientific, 78840) according to manufacture manual.

Primary neuronal culture

Mixed (cortical and hippocampal) cultures were used for Western blot analysis and subcellular fractionation. Brains from E16-E18 wild-type 129Sv embryos were removed and dissected under microscope in ice-cold isotonic buffer (137 mM NaCl, 5.4 mM KCl, 170 μ M Na₂HPO₄, 148 μ M K₂HPO₄, 5.5 mM glucose, 58.4 mM sucrose). Dissected tissue was pooled together in 15-mL tube and incubated with 1 mL 0.25% trypsin for 5 min at 37°C. Trypsin was removed and 5 mL dissociation media was added (DMEM, high glucose, 10% heat inactivate FBS, 1 \times antimycotic/antibacterial solution). Hippocampi were dissociated by slowly pipetting through glass pipettes. After brief centrifugation, dissociation media was replaced by neurobasal medium supplemented with 2% B27 and 0.5 mM glutamax. Cells were plated in six well (2,500,000/well) plates coated with poly-L-lysine. Cultures were treated with 55 mM KCl on DIV 7-10.

Nonfluorescent immunohistochemistry

Mice were transcardially perfused with saline solution (0.9% NaCl in water) followed by 4% PFA diluted in 0.1 M PBS. Removed brains were dehydrated via sucrose gradient (10%, 20%, 30%) and sectioned on a cryostat at 30 μ m. Free-floating sections were blocked in 4% goat serum, 0.2% lysine in 0.1 M PBS and incubated with Ube3a primary antibody overnight (1:1000). Secondary antibody was added for 2 h and the avidin-biotin-HRP complex was formed with VECTASTAIN ABC kit (Vector labs). DAB was used to develop the stain. Sections were imaged with a Zeiss Mirax Scan 150 microscope. Identification of positive staining and percent area of positive stain was performed using Image Analysis software (created by Andrew Lesniak, Zeiss).

Acute hippocampal slice preparation and treatment

Two-month-old m⁻/p⁺ and m⁺/p⁺ littermates were sacrificed by decapitation, the brains were rapidly removed and briefly submerged in ice-cold cutting solution (110 mM sucrose, 60 mM NaCl, 3 mM KCl, 1.25 mM NaH₂PO₄, 28 mM NaHCO₃, 0.5 mM CaCl₂, 5 mM glucose, 0.6 mM ascorbate) saturated with 95% O₂ and 5% CO₂. Hippocampal slices (400 μ m) were cut on a vibratome and allowed to equilibrate in artificial cerebrospinal fluid solution (125 mM NaCl, 2.5 mM KCl, 1.24 mM NaH₂PO₄, 25 mM NaHCO₃, 10 mM glucose, 2 mM CaCl₂, 1 mM MgCl₂) at 32°C for 2 h. Slices were treated with 55 mM KCl for 5 and 30 min, snap frozen, and stored at -80°C for further processing.

Acknowledgments

This work was supported by The Foundation for Angelman Syndrome Therapeutics (FAST) and NICHD R21-HD67249. We thank Sharon Claridge for intellectual contribution and direction and Dr. April Lussier for data and manuscript editing.

References

- Atkins CM, Selcher JC, Petraitis JJ, Trzaskos JM, Sweatt JD. 1998. The MAPK cascade is required for mammalian associative learning. *Nat Neurosci* **1**: 602–609.
- Blum S, Moore AN, Adams F, Dash PK. 1999. A mitogen-activated protein kinase cascade in the CA1/CA2 subfield of the dorsal hippocampus is essential for long-term spatial memory. *J Neurosci* **19**: 3535–3544.
- Chwang WB, O'Riordan KJ, Levenson JM, Sweatt JD. 2006. ERK/MAPK regulates hippocampal histone phosphorylation following contextual fear conditioning. *Learn Mem* **13**: 322–328.
- Dagli A, Buiting K, Williams C. 2011. Molecular and clinical aspects of Angelman syndrome. *Mol Syndromol* **2**: 100–112.
- Daily J, Smith AG, Weeber EJ. 2012. Spatial and temporal silencing of the human maternal *UBE3A* gene. *Eur J Paediatr Neurol* **15**: 587–591.
- Dindot SV, Antalffy BA, Bhattacharjee MB, Beaudet AL. 2008. The Angelman syndrome ubiquitin ligase localizes to the synapse and nucleus, and maternal deficiency results in abnormal dendritic spine morphology. *Hum Mol Genet* **17**: 111–118.
- English JD, Sweatt JD. 1997. A requirement for the mitogen-activated protein kinase cascade in hippocampal long term potentiation. *J Biol Chem* **272**: 19103–19106.
- Greer PL, Hanayama R, Bloodgood BL, Mardinly AR, Lipton DM, Flavell SW, Kim T-K, Griffith EC, Waldon Z, Maehr R. 2010. The Angelman Syndrome protein Ube3A regulates synapse development by ubiquitinating arc. *Cell* **140**: 704–716.
- Gustin RM, Bichell TJ, Bubser M, Daily J, Filonova I, Mrelashvili D, Deutch AY, Colbran RJ, Weeber EJ, Haas KF. 2010. Tissue-specific variation of Ube3a protein expression in rodents and in a mouse model of Angelman syndrome. *Neurobiol Dis* **39**: 283–291.
- Heck DH, Zhao Y, Roy S, LeDoux MS, Reiter LT. 2008. Analysis of cerebellar function in Ube3a-deficient mice reveals novel genotype-specific behaviors. *Hum Mol Genet* **17**: 2181–2189.
- Huang H-S, Burns AJ, Nonneman RJ, Baker LK, Riddick NV, Nikolova VD, Riday TT, Yashiro K, Philpot BD, Moy SS. 2013. Behavioral deficits in an Angelman syndrome model: Effects of genetic background and age. *Behav Brain Res* **243**: 79–90.
- Jiang YH, Armstrong D, Albrecht U, Atkins CM, Noebels JL, Eichele G, Sweatt JD, Beaudet AL. 1998. Mutation of the Angelman ubiquitin ligase in mice causes increased cytoplasmic p53 and deficits of contextual learning and long-term potentiation. *Neuron* **21**: 799–811.
- Kumar S, Talis AL, Howley PM. 1999. Identification of HHR23A as a substrate for E6-associated protein-mediated ubiquitination. *J Biol Chem* **274**: 18785–18792.
- Margolis SS, Salogiannis J, Lipton DM, Mandel-Brehm C, Wills ZP, Mardinly AR, Hu L, Greer PL, Bikoff JB, Ho HY, et al. 2010.

- EphB-mediated degradation of the RhoA GEF Ephexin5 relieves a developmental brake on excitatory synapse formation. *Cell* **143**: 442–455.
- Mishra A, Jana NR. 2008. Regulation of turnover of tumor suppressor p53 and cell growth by E6-AP, a ubiquitin protein ligase mutated in Angelman mental retardation syndrome. *Cell Mol Life Sci* **65**: 656–666.
- Mishra A, Godavarthi SK, Jana NR. 2009. UBE3A/E6-AP regulates cell proliferation by promoting proteasomal degradation of p27. *Neurobiol Dis* **36**: 26–34.
- Nawaz Z, Lonard DM, Smith CL, Lev-Lehman E, Tsai SY, Tsai M-J, O'Malley BW. 1999. The Angelman syndrome-associated protein, E6-AP, is a coactivator for the nuclear hormone receptor superfamily. *Mol Cell Biol* **19**: 1182–1189.
- Pelc K, Cheron G, Dan B. 2008. Behavior and neuropsychiatric manifestations in Angelman syndrome. *Neuropsychiatr Dis Treat* **4**: 577–584.
- Schanen NC. 2006. Epigenetics of autism spectrum disorders. *Hum Mol Genet* **15**: R138–R150.
- Selcher JC, Atkins CM, Trzaskos JM, Paylor R, Sweatt JD. 1999. A necessity for MAP kinase activation in mammalian spatial learning. *Learn Mem* **6**: 478–490.
- Selcher JC, Weeber EJ, Christian J, Nekrasova T, Landreth GE, Sweatt JD. 2003. A role for ERK MAP kinase in physiologic temporal integration in hippocampal area CA1. *Learn Mem* **10**: 26–39.
- Singhmar P, Kumar A. 2011. Angelman syndrome protein UBE3A interacts with primary microcephaly protein ASPM, localizes to centrosomes and regulates chromosome segregation. *PLoS One* **6**: e20397.
- Smith SE, Zhou Y-D, Zhang G, Jin Z, Stoppel DC, Anderson MP. 2011. Increased gene dosage of Ube3a results in autism traits and decreased glutamate synaptic transmission in mice. *Sci Transl Med* **3**: 103ra197.
- Steffenburg S, Gillberg CL, Steffenburg U, Kyllerman M. 1996. Autism in Angelman syndrome: A population-based study. *Pediatr Neurol* **14**: 131–136.
- Thomas GM, Haganir RL. 2004. MAPK cascade signalling and synaptic plasticity. *Nat Rev Neurosci* **5**: 173–183.
- Trifilieff P, Herry C, Vanhoutte P, Caboche J, Desmedt A, Riedel G, Mons N, Micheau J. 2006. Foreground contextual fear memory consolidation requires two independent phases of hippocampal ERK/CREB activation. *Learn Mem* **13**: 349–358.
- Trifilieff P, Calandrea L, Herry C, Mons N, Micheau J. 2007. Biphasic ERK1/2 activation in both the hippocampus and amygdala may reveal a system consolidation of contextual fear memory. *Neurobiol Learn Mem* **88**: 424–434.
- van Woerden GM, Harris KD, Hojjati MR, Gustin RM, Qiu S, de Avila Freire R, Jiang Y-H, Elgersma Y, Weeber EJ. 2007. Rescue of neurological deficits in a mouse model for Angelman syndrome by reduction of α CaMKII inhibitory phosphorylation. *Nat Neurosci* **10**: 280–282.
- Weeber EJ, Jiang YH, Elgersma Y, Varga AW, Carrasquillo Y, Brown SE, Christian JM, Mirnikjoo B, Silva A, Beaudet AL, et al. 2003. Derangements of hippocampal calcium/calmodulin-dependent protein kinase II in a mouse model for Angelman mental retardation syndrome. *J Neurosci* **23**: 2634–2644.
- Yashiro K, Riday TT, Condon KH, Roberts AC, Bernardo DR, Prakash R, Weinberg RJ, Ehlers MD, Philpot BD. 2009. Ube3a is required for experience-dependent maturation of the neocortex. *Nat Neurosci* **12**: 777–783.

Received July 10, 2013; accepted in revised form October 24, 2013.