



# Further Mechanisms and Locations in Which Antisporozoite Antibodies Neutralize Malaria Sporozoites

Jerome P. Vanderberg (Retired)<sup>a</sup>

<sup>a</sup>New York University School of Medicine, New York, New York, USA

**KEYWORDS** antibodies, malaria, preerythrocytic, skin, sporozoites, vaccine, immune complexes, antisporozoite antibodies, circumsporozoite protein, immune complexes, *Plasmodium*, sporozoite

Having worked in this area for many years, I read with much interest the recent article “Antibody-Mediated Protection against *Plasmodium* Sporozoites Begins at the Dermal Inoculation Site” by Flores-Garcia et al. (1). It is now universally recognized, as described by the authors, that mosquitoes inject most of their sporozoites into extravascular skin tissue rather than directly into blood vessels (2, 3). From there, they migrate through the dermis to enter blood vessels (3) and are carried to the liver, where they invade hepatocytes to continue their development. The authors further note that sporozoites spend much of their extracellular time at the skin inoculation site, thus rendering them vulnerable to antibody-mediated destruction at this site. Such antibody-mediated immobilization or destruction of sporozoites has been shown both *in vitro* and *in vivo* (3–5). I am gratified that the authors have corroborated these prior findings made by us and several other authors and have expanded them with new data.

The authors concluded that antibodies targeting the migratory sporozoites exert a large proportion of their protective effect at the inoculation site but that the mechanisms by and location in which they neutralize parasites have not been fully elucidated (1). I would like, however, to call attention to the evidence we presented in our 2009 publication (5) in which we elucidated an entirely new and complementary way in which sporozoites can be neutralized by host antibodies.

It is known that living sporozoites release large quantities of soluble circumsporozoite protein (CSP) into their environment both *in vitro* and *in vivo* and that soluble CSP is found within the saliva of malaria-infected mosquitoes (6, 7). Thus, infected mosquitoes introduce into the skin of immunized hosts not only CSP-covered sporozoites but also soluble CSP, both of which encounter and interact with homologous anti-CSP antibodies *in situ* within avascular tissue of the host dermis. We presented evidence that many of the sporozoites are trapped there within apparent immune complexes, as determined by confocal microscopy and specific staining with fluorescein isothiocyanate (FITC)-conjugated protein A and A/C. Thus, sporozoites were not only immobilized by CS antibodies as has previously been shown but were additionally entrapped by being encased within these immune complexes. I respectfully suggest, in disagreement with the conclusions of the authors (1), that the mechanisms by and location in which antisporozoite antibodies neutralize parasites have indeed been further elucidated, as demonstrated in our 2009 paper (5).

**Citation** Vanderberg JP. 2019. Further mechanisms and locations in which antisporozoite antibodies neutralize malaria sporozoites. *mBio* 10:e01588-19. <https://doi.org/10.1128/mBio.01588-19>.

**Invited Editor** Carole A. Long, NIAID/NIH

**Editor** Patricia J. Johnson, University of California Los Angeles

**Copyright** © 2019 Vanderberg. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to [Vandej01@nyumc.org](mailto:Vandej01@nyumc.org).

For the author reply, see <https://doi.org/10.1128/mBio.02108-19>.

**Published** 10 September 2019

## REFERENCES

1. Flores-Garcia Y, Nasir G, Hopp CS, Munoz C, Balaban AE, Zavala F, Sinnis P. 2018. Antibody-mediated protection against *Plasmodium* sporozoites begins at the dermal inoculation site. *mBio* 9:e02194-18. <https://doi.org/10.1128/mBio.02194-18>.
2. Sidjanski S, Vanderberg JP. 1997. Delayed migration of *Plasmodium* sporozoites from the mosquito bite site to the blood. *Am J Trop Med Hyg* 57:426–429. <https://doi.org/10.4269/ajtmh.1997.57.426>.
3. Vanderberg JP, Frevert U. 2004. Intravital microscopy demonstrating

- antibody-mediated immobilisation of *Plasmodium berghei* sporozoites injected into skin by mosquitoes. *Int J Parasitol* 34:991–996. <https://doi.org/10.1016/j.ijpara.2004.05.005>.
4. Stewart MJ, Nawrot R, Schulman S, Vanderberg JP. 1986. *Plasmodium berghei* sporozoite invasion is blocked in vitro by sporozoite-immobilizing antibodies. *Infect Immun* 51:859–864.
  5. Kebaier C, Voza T, Vanderberg JP. 2009. Kinetics of mosquito-injected *Plasmodium* sporozoites in mice: fewer sporozoites are injected into sporozoite-immunized mice. *PLoS Pathog* 5:e1000399. <https://doi.org/10.1371/journal.ppat.1000399>.
  6. Stewart MJ, Vanderberg JP. 1991. Malaria sporozoites release circumsporozoite protein from their apical end and translocate it along their surface. *J Protozool* 38:411–421. <https://doi.org/10.1111/j.1550-7408.1991.tb01379.x>.
  7. Beier JC, Vaughan JA, Madani A, Noden BH. 1992. *Plasmodium falciparum*: release of circumsporozoite protein by sporozoites in the mosquito vector. *Exp Parasitol* 75:248–256. [https://doi.org/10.1016/0014-4894\(92\)90185-D](https://doi.org/10.1016/0014-4894(92)90185-D).