

RESEARCH ARTICLE

Open Access

Deep sequencing and expression of microRNAs from early honeybee (*Apis mellifera*) embryos reveals a role in regulating early embryonic patterning

Lisa Zondag^{1,2}, Peter K Dearden¹ and Megan J Wilson^{1,2*}

Abstract

Background: Recent evidence supports the proposal that the observed diversity of animal body plans has been produced through alterations to the complexity of the regulatory genome rather than increases in the protein-coding content of a genome. One significant form of gene regulation is the contribution made by the non-coding content of the genome. Non-coding RNAs play roles in embryonic development of animals and these functions might be expected to evolve rapidly. Using next-generation sequencing and *in situ* hybridization, we have examined the miRNA content of early honeybee embryos.

Results: Through small RNA sequencing we found that 28% of known miRNAs are expressed in the early embryo. We also identified developmentally expressed microRNAs that are unique to the Apoidea clade. Examination of expression patterns implied these miRNAs have roles in patterning the anterior-posterior and dorso-ventral axes as well as the extraembryonic membranes. Knockdown of Dicer, a key component of miRNA processing, confirmed that miRNAs are likely to have a role in patterning these tissues.

Conclusions: Examination of the expression patterns of novel miRNAs, some unique to the *Apis* group, indicated that they are likely to play a role in early honeybee development. Known miRNAs that are deeply conserved in animal phyla display differences in expression pattern between honeybee and *Drosophila*, particularly at early stages of development. This may indicate miRNAs play a rapidly evolving role in regulating developmental pathways, most likely through changes to the way their expression is regulated.

Background

A major component of the transcriptome of animals consists of non-protein coding RNAs [1-3]. Micro-RNAs (miRNAs) are a subset of small non-coding RNAs that are 18-24 nucleotides long and have a key role in regulating gene expression in eukaryotes. They are produced from a primary full-length transcript (pri-miRNA), which is cleaved to form hairpin structures around 70 nucleotides in length. These are called precursor miRNAs (pre-miRNAs) and are exported to the cytoplasm

to be processed further to functional mature miRNAs by the ribonuclease Dicer [4,5]. Once assembled into the RNA-induced silencing complex (RISC), the miRNA acts on its target by binding to complementary sequences present in the 3' untranslated regions (UTR) of the target mRNA [6]. This results in either translational repression or mRNA cleavage, thus providing another level of gene regulation [7].

There is accruing evidence to suggest that miRNAs play a role in regulating multiple developmental pathways, including fundamental developmental processes of animal development such as axis formation and organ morphogenesis. Many miRNAs are expressed in developmentally restricted patterns [8,9], for example, examination of miRNA expression in *Drosophila* during embryogenesis found that many are expressed in

* Correspondence: meganj.wilson@otago.ac.nz

¹Laboratory for Evolution and Development, Genetics Otago and the National Research Centre for Growth and Development, Department of Biochemistry, University of Otago, P.O. Box 56, Dunedin 9054, New Zealand

²Developmental Biology Laboratory, Department of Anatomy, University of Otago, P.O. Box 56, Dunedin 9054, New Zealand

Table 1 Summary of small RNA sequencing results

	Sample 1	Sample 2
Total number of reads	20120856	21480569
Clean reads	19563073	20873296
Number of reads mapped back to genome	13200933	14892414
Number of reads map to annotated pre-miRNA	21450	18469

restricted patterns along the anterior-posterior and dorso-ventral axes and in specific tissues. Their expression is developmentally regulated, often by their own promoter and regulatory elements, similar to developmental protein coding factors [10]. Loss of miRNA function often results in defective development and patterning [11-14], indicating an essential role in animal development. MiRNAs are also proposed to provide developmental stability particularly under times of environment stress, acting to buffer developmental pathways [15].

Many miRNA families are ancient and can be traced to more basal animals. Expression analysis and functional studies indicate that their roles are often conserved [8]. As single miRNAs can bind to many different mRNAs to regulate their expression, they can potentially impact on several regulatory pathways [16]. Thus any changes to the way miRNAs are expressed are likely to impact multiple developmental processes. Many miRNAs have been found to be specific to particular phylogenetic groups, some found only in particular lineages. Over 40 miRNA families arose early in the lineage leading to the vertebrates, and it has been suggested that these contributed to the evolution of vertebrate complexity [17].

Here we have profiled the miRNAs expressed during early embryogenesis in the honeybee (*Apis mellifera*) to determine if they are likely to play a significant role in honeybee embryogenesis. The expression and function of miRNAs in insect development has to date only been investigated in *Tribolium* and *Drosophila* [8,18]. Like *Drosophila*, *Apis* development begins with a syncytial blastoderm stage prior to cellularisation, where much of the body patterning information is established [19]. Both *Apis* and *Drosophila* are considered long germ band insects where segmentation occurs across the whole body [20]. However there are some significant differences, notably in patterning of the extraembryonic membranes. In *Drosophila* the extraembryonic membranes are patterned as one tissue, the amino-serosa, a process regulated by the transcription factor zen [21]. In honeybee, these membranes are patterned separately, although both tissues still require zen [22]. There are also significant differences in the nature of the regulatory networks required to pattern the anterior-posterior axis [22-24]. The honeybee genome has been sequenced and 168

Table 2 Profile of known miRNAs in present in 24-30 hour embryos

miRNA	Sample 1	Sample 2
ame-miR-100	101	73
ame-miR-1	94	5
ame-miR-71	269	88
ame-miR-3759	5	107
ame-miR-184	5294	4431
ame-miR-927	21	22
ame-miR-275	22	15
ame-miR-7	158	73
ame-miR-279	23	8
ame-miR-8	9	8
ame-miR-92b	103	149
ame-miR-283	8	16
ame-miR-3756	362	264
ame-miR-2	167	261
ame-miR-263b	6	16
ame-miR-3785	13	14
ame-miR-11	15	26
ame-miR-3747a	606	143
ame-miR-279c	544	310
ame-miR-315	4	13
ame-miR-13a	10	3
ame-miR-12	8	20
ame-miR-2944	2452	1177
ame-miR-263	39	141
ame-miR-375	6	6
ame-miR-279b	335	318
ame-miR-34	9	6
ame-miR-750	8	7
ame-miR-989	8	4
ame-miR-9c	89	69
ame-miR-3477	211	199
ame-miR-3715	15	35
ame-miR-10	3923	3561
ame-miR-125	49	87
ame-miR-3747b	63	23
ame-miR-996	17	3
ame-miR-9a	2214	1977
ame-miR-92a	175	214
ame-let-7	135	185
ame-miR-3786	31	32
ame-miR-252	14	10
ame-bantam	13	6
ame-miR-317	360	354
ame-miR-306	608	619
ame-miR-3791	592	707

A *Ame-mir-10 (sample 1)*

```

CCCAGTTAATGCTCTACATCTACCCCTGTAGATCCGAATTTGTTTGATAAGAGGCCGACAAATTCCGGTTCCTAGAGAGGTTTGTGTGGTGATACAGAGCTAC
...(((((((...(((((((...(((((((...(((((((...(((((((...(((((((...(((((((...(((((((...(((((((...(((((((...(((((((...
*****ACCCTGTAGATCCGAATTTGT*****length read count
.....CTACCCCTGTAGATCCGAATTTGT.....23 1
.....TACCCCTGTAGATCCGAATTTGT.....22 3511
.....TACCCCTGTAGATCCGAATTTG.....21 187
.....TACCCCTGTAGATCCGAAT.....18 4
.....TACCCCTGTAGATCCGAATT.....19 4
.....TACCCCTGTAGATCCGAATTT.....20 3
.....TACCCCTGTAGATCCGAATTTGTTT.....24 1
.....ACCCTGTAGATCCGAATTTGT.....21 185
.....ACCCTGTAGATCCGAATTTGTT.....22 14
.....ACCCTGTAGATCCGAATTTG.....20 7
.....ACCCTGTAGATCCGAATTTGTTT.....23 1
.....ACCCTGTAGATCCGAATTTGTTTGATA.....27 1
.....ACCCTGTAGATCCGAATTT.....19 1
.....ACCCTGTAGATCCGAATT.....18 1
.....CCTGTAGATCCGAATTTGT.....19 2
.....CAAATTCGGTTCCTAGAGAGGTTT.....23 91
.....CAAATTCGGTTCCTAGAGAGGTTG.....24 37
.....CAAATTCGGTTCCTAGAGAGGTT.....22 5
.....CAAATTCGGTTCCTAGAGAGGT.....21 2
.....CAAATTCGGTTCCTAGAGAGG.....20 1
.....CAAATTCGGTTCCTAGAGA.....18 1
.....AAATTCGGTTCCTAGAGAGGTTTG.....23 264
.....AAATTCGGTTCCTAGAGAGGTTT.....22 22
.....AAATTCGGTTCCTAGAGAGGT.....20 1
.....AAATTCGGTTCCTAGAGAGTT.....21 1
    
```

Ame-mir-1

```

CCGGGCGATGCTGTTCCGTGCTTCCTACTTCCCATAGTGGATGCGACGTA TGGAAATGTAAGAAGTATGGAGCTGCGCCCGG
(((((((...(((((((...(((((((...(((((((...(((((((...(((((((...(((((((...(((((((...(((((((...(((((((...
*****TGGAAATGTAAGAAGTATGGAG***** length read count
.....ATGGAATGTAAGAAGTATGGAG.....23 1
.....TGGAAATGTAAGAAGTATGGAG.....22 67
.....TGGAAATGTAAGAAGTATGGA.....21 22
.....TGGAAATGTAAGAAGTATGG.....20 3
.....TGGAAATGTAAGAAGTATG.....19 1
    
```





Figure 1 Alignment of sequence reads back to the miRNA precursors for mir-184 and mir-1. (A) An example of mapping sequence reads back to the pre-miRNA precursor. Bracket-notation for secondary structure is shown below the precursor and the mature miRNA sequence (miRBase) is flanked by asterisks. The number of reads giving rise to a particular sequence is noted beside each sequence. Highlighted in red is the most abundant small RNA read. **(B)** Venn diagram of the distribution of known miRNAs from the honeybee embryo small RNA libraries between *Drosophila*, other arthropods and vertebrates.

miRNAs have so far been predicted. The *Drosophila* genome encodes at least 430 miRNAs (miRBase). We examined the expression of honeybee miRNAs during early development, by deep sequencing and *in situ* hybridization. This included developmental stages at which the anterior-posterior and dorso-ventral axes have been established, patterning including segmentation is underway, just prior to gastrulation. As the pattern of miRNA expression is often reflective of their function in a developmental process or tissue patterning, we

examined the expression of eight miRNAs identified in our study. Additionally, RNAi knockdown of *Dicer* during early embryogenesis indicated that small RNAs are likely to contribute to early honeybee embryo development.

Results and discussion

Abundance and expression of previously known miRNAs
 Total RNA was extracted from honeybee embryos aged from 24 to 30 hours old (stages 4 to 5). Approximately



20 million reads were generated per sample, corresponding to 3 million unique reads (Table 1). Almost 70% of unique small RNA tags were mapped back to the *Apis mellifera* genome (Table 1). The number of reads for known (i.e. present in miRBase) *Apis mellifera* miRNAs are shown in Table 2 and an example of an alignment is shown in Figure 1A. This provides further experimental evidence that these are transcribed miRNAs in the honeybee genome. Of the 168 mature miRNA in miRBase, 45 were represented in both samples. This indicates that almost 28% of known miRNAs are expressed in the early honeybee embryo, implying a

possible role for these in regulating developmental pathways. An example of read alignments to precursor miRNA are shown Figure 1A.

We determined if these miRNAs are also present in the genomes of other phyla. Of the miRNAs we isolated from honeybee embryos, 36 are also present in the genomes of other arthropods. 15 of these are present in the vertebrate lineage, indicating these are likely to be ancestral (Figure 1B). Less conservation was observed between *Apis* and *Drosophila* miRNA content which has also been noted previously between *Tribolium* and *Drosophila* [18], indicating that Arthropod groups

Table 3 Novel miRNA profile and presence (+) or absence (-) in the genomes of other insects

Candidate miRNA	Reads Sample 1	Reads Sample 2	<i>Bombus</i>	<i>Nasonia</i>	<i>Atta cephalotes</i>
mir-0002	197	144	+	-	-
mir-0003	106	82	+	-	-
mir-0004	7	8	+	-	-
mir-0005	1642	2263	+	+	+
mir-0006	30	35	-	-	-
mir-0007	5688	5036	+	-	-
mir-0008	59	29	+	-	+
mir-0009	6	9	+	-	-
mir-0010	44	30	-	-	-
mir-0011	6	14	-	-	-
mir-0012	2095	1152	-	-	-
mir-0013	10	17	-	-	-
mir-0014	405	206	-	-	-
mir-0017	405	206	-	-	-
mir-0018	37	10	-	-	-
mir-0020	87	88	+	-	-

(other than Diptera) have more miRNA content in common to each other than when compared separately to Diptera.

To determine if the developmental expression patterns of miRNAs with orthologues in both honeybee and *Drosophila* were conserved, we examined the expression of four miRNAs detected in our study, with expression and functional data of their orthologue in *Drosophila*.

In situ hybridisation was performed using RNA probes designed to bind to the pri-miRNA, the longer transcript from which the pre-miRNA is produced, prior to export from the nucleus. This strategy has been successfully used in previous studies illustrating that it reflects the mature miRNA expression (when detected using LNA probes) [25]. The probes used detect nascent transcripts before processing by Drosha RNase III enzyme and thus are expected to detect nuclear dots rather than the cytoplasmic staining produced with probes against an mRNA transcript (for example see Additional file 1: Figure S3).

Mir-10 is a widely conserved miRNA in both sequence and genomic location in both invertebrates and vertebrate *Hox* gene clusters. It is located within the *Hox* complex in *Drosophila* between the *deformed* (*dfd*) and *sex-combs reduced* (*Scr*) *Hox* genes and has been predicted to directly regulate mRNA translation of nearby *Hox* genes that contain mir-10 binding sites in their 3'UTRs [26]. However ectopic expression of *Dme-mir-10* had no significant effect on the expression of predicted *Hox* targets (*Scr* and *Abd-B*) [25], suggesting that they

may not be biologically relevant targets for mir-10 in a laboratory setting. *Apis mellifera mir-10* (*Ame-mir-10*) is also located in the *Hox* gene complex between *Am-dfd* and *Am-Scr* (Additional file 2: Figure S4). We detected expression of this miRNA during embryogenesis (Table 1 and Figure 1A). *In situ* hybridisation analysis revealed expression of *pri-Ame-mir-10* at the anterior pole of the embryo (Figure 2A and B). Later *pri-Ame-mir-10* RNA expression (stage 9; approximately 48 hours old) was detected throughout the posterior two-thirds of the embryo, in the posterior terminal segment and the underlying mesoderm (Figure 2D). This is consistent with expression of a *Hox* regulator, as it shows limited expression along the anterior-posterior axis. In comparison, *Drosophila pri-Dme-mir-10* expressed in a broad band across the middle of blastoderm embryos and, following germ band extension, it is expressed in the posterior half of the embryo, in the anal pad, ventral neuroectoderm and hindgut [25]. This indicates that while there are similarities in late embryonic expression of *mir-10* between *Drosophila* and honeybee embryos, *mir-10* expression in early embryos is quite different. This may indicate a shift in *mir-10* function in early embryogenesis between Diptera and hymenoptera.

Previous deep sequencing of *Drosophila* embryonic RNA revealed that most of the reads correspond to the 3' arm of the mir-10 precursor [27], although RNA expression patterns of both mir-10-5p and mir-10-3p is similar in *Drosophila* embryos [25]. We found that the 90% of sequence reads are from the 5' arm of the pre-miRNA in honeybee embryos, indicating that mir-10-5p is responsible for the majority of mir-10 function (Figure 1). This has also been found for mir-10 in *Tribolium* [18]. Changes to which part of the pre-miRNA strand provides the dominant or functional miRNA sequence (arm switching) is proposed to be one mechanism of miRNA evolution to drive miRNA diversification [28]. Our results and those from *Tribolium* [18] would indicate that the ancestral dominant arm was the mir-10-5p (producing the mature miRNA) and that this has switched during *Drosophila* evolution to the mir-10-3p arm. However, while only 10% of the total reads for pre-mir-10 were from the 3p arm in honeybee embryos (Figure 1), it was still a significant number (425) and more abundant than some of the other miRNAs detected (Table 2), indicating that the mature miRNA from this arm of the mir-10 hairpin (mir-10-3p) may have a distinct role during honeybee development.

Mir-1 is a highly conserved miRNA that has been suggested to play a role in myogenesis in *Drosophila* and vertebrates [29-31]. We detected low numbers of reads for *Ame-mir-1* in both our samples (Table 1), but because of its conservation between vertebrates and invertebrates, we

group or have been lost from other lineages. We examined the expression patterns of four of these newly identified miRNAs during honeybee embryogenesis.

The novel miRNA *Ame-mir-0002* is located within an intron of the *vesicular glutamate transporter 2.1* gene on chromosome 11 (Figure 3C). *Pri-Ame-mir-0002* RNA was expressed throughout the ectoderm of the stage 4 embryo (Figure 3D), by late stage 5 RNA expression is no longer detected at the anterior-dorsal region of the embryo, or along the dorsal surface (Figure 3E), this includes areas where the extraembryonic membranes form [22]. Later *pri-Ame-mir-0002* RNA expression was found throughout the neuroectoderm (Figure 3F). These expression patterns imply a possible role for mir-0002 in patterning the ectoderm and neuroectoderm during honeybee development. Many species-specific miRNAs are located in the introns of protein coding genes compared to those are proposed to be more ancient which tend to be intergenic [39]. Most are often transcribed from the host-gene promoter, although some intronic miRNAs have their own promoter [40,41]. Further work

is required to determine if mir-0002 expression is driven from the host-gene promoter or if it has its own promoter region.

The novel miRNA mir-0004 is on chromosome 15, located between *smad4* and a hypothetical gene, and about 5.6 kbp from *mir-252* (Figure 4C). Expression of *pri-Ame-mir0004* RNA was detected in ectoderm cell nuclei at the anterior and posterior half of the embryo, but weaker within the central region of the embryo (Figure 4D). Embryo staining produced high background, probably as a result of longer staining times required to visualize the RNA expression; read count from the small RNA libraries indicated mir-0004 was present at low levels.

mir-0007 is located in the intergenic space between two predicted genes (Figure 5A). *Pri-mir-0007* RNA was detected in the dorsal-posterior ectoderm (Figure 5D), a part of the embryo where the amnion arises [22]. By stage 9, *pri-mir-0007* RNA was detected in the posterior half of the amnion (Figure 5E).

Mir-0005 was detected with high numbers of reads (1642 and 2263 reads per sample) in our small RNA




Figure 5 Expression and sequence of novel *Ame-mir-0007*. Embryos are positioned anterior to the left, dorsal side up. (A) Sequence of the mir-0007 precursor with aligned sequence reads and read counts. Highlighted in red is the sequence with the highest read counts. Bracket-notation of secondary structure is shown below the precursor. Mature miRNA and miRNA* sequences are shown surrounded by asterisks. (B) The predicted mir-0007 hairpin structure, the most abundant read sequence highlighted in red. (C) Genomic location of the pre-mir-0007; mir-0007 is located between the predicted genes LOC726303 and LOC410888 on chromosome 2. (D) Expression during embryogenesis at stage 4. (E) Expression of *pri-mir0007* at stage 9 is though the posterior of the extraembryonic membrane.




Figure 7 Genomic location of pre-mir-92a and pre-mir-92b in arthropods. *Mir-92a* and *b* are intragenic positioned closely in hemimetabola, hymenoptera and coleopteran genomes and are predicted to be expressed as part of the same pri-RNA transcript. However in Diptera, *mir92a* and *b* are now separated by considerable distances and may no longer be part of the same polycistronic transcript.

to assess morphology. Approximately 20% survived through to stage 9, in line with previous survival ratios from similar procedures due to the nature of the injection procedure [22,23]. Posterior and anterior regions of the embryos were malformed with loss of terminal patterning (Figure 8B, C, F and H). In some embryos there was a significant reduction in the amnion that normally covers the yolk sac, and the dorsal regions of the embryo extend further towards the ventral side of the embryo (Figure 8B and H). This phenotype is very similar to that of *Am-zen* RNAi knockdown [22], suggesting problems with dorso-ventral patterning. Abdominal tracheal pits were present but segments appeared to be fused together, indicating a defect in anterior-posterior patterning (Figure 8B and C).


Conclusions

We examined the miRNA content of early honeybee embryos by deep sequencing followed by determination of the expression patterns of eight of these miRNAs.

Consistent with both miRNA expression patterns and target prediction, Dicer siRNA knockdown embryos had defects in extraembryonic membrane formation, anterior-posterior and dorso-ventral patterning, suggesting that miRNAs may have functions in regulating these patterning pathways or their gene target(s) have roles in multiple pathways.

Many miRNAs that are expressed during embryo development are deeply conserved throughout metazoans indicating a more ancient origin. However, several are unique to the arthropod group, and some restricted just to the *Apis* lineage. Given the developmental expression of these miRNAs, they may have taken on roles particular to the development of the honeybee embryo. Previous studies have hypothesised that miRNAs are continuously added to the metazoan genomes, are stabilized once added, and are rarely lost [27,42,43].

Interestingly, some of the highly conserved miRNAs identified in our study had very different expression



patterns in the honeybee to those documented in other animals. *Mir-1* expression differed between *Apis* and other animals, which may result from loss of a regulatory element or binding sites for cis-regulatory proteins [44]. Early embryonic expression of *mir-10* differed between *Drosophila* and *Apis*, but later expression was similar and is consistent with regulation by separate cis-regulatory elements (one controlling early expression, one controlling later expression) as suggested previously for *Drosophila* *mir-10* [8]. This suggests that earlier expression regulatory elements are evolving more rapidly. A similar pattern of more labile expression in early development has previously observed for protein-coding genes [24,45-47]. These changes or shifts in miRNA expression imply the regulatory regions controlling miRNA expression are also rapidly evolving. This indicates the importance that changes to gene-regulatory sequences contribute to the evolution of developmental pathways extends to also changes associated with regulatory elements that control miRNA genes.

Methods

Sample collection and preparation

A queen honeybee was caged with an empty area of an Eziqueen queen rearing frame and placed back into the hive. After 5 hours the frame was removed and the queen released back into the hive. The eggs were removed and while still attached to the black strips of the Eziqueen frame incubated for 24 hours. Eggs were collected and total RNA extracted using TRIzol (Life Technologies). Total RNA concentration and purity determined was using a Nanodrop spectrometer (Thermo Scientific). 10 µg of purified total RNA was sent to Beijing Genomics Institute (BGI) for sequencing on an Illumina HiSeq 2000 sequencer. Low quality reads, reads without the adaptors, reads with polyA sequences and reads without the insert tag were removed. Also discarded were any tRNA, rRNA, snRNA and snoRNA sequences. Sequence reads were mapped to the genome using the programme SOAP [44]. Small RNA tags were aligned to known miRNA *Apis mellifera* precursors

(miRBASE). The two small RNA libraries shared 98.06% of sequences. Analyses of the length distribution of cleaned reads showed enrichment of small RNAs from 22 to 31 nucleotides (Additional file 7: Figure S1); we would expect about a length of 22nt for miRNAs. Examination of the first base of the 22 nucleotide sequences revealed most show a first base bias to uridine as predicted from previous deep sequencing miRNA studies (Additional file 8: Figure S2).

Amplification of pri-miRNA fragments

Oligonucleotide primers were designed to amplify 500-800 bp regions using genomic template. In each amplicon, the precursor miRNA resides in the centre of the sequence. PCR fragments were cloned into the vector pBluescript II KS (+/-) and sequenced. Cloned fragments were used to produce RNA sense and antisense *in situ* hybridisation probes. Oligonucleotide primer pairs were as follows: mir-10 5'ACAAATGGACGACGAAGAGG3' and 5'GCGGCACGTACGTTACTTTA3', mir-1 5'GCCACGTACGTTTCGAAAAC3' and 5'TTCGCAAGACGATAACATCA3', mir-184 5'GCCTCGGGTTTCGAGCGTT3' and 5'AGGAGAAGGGAAGAATGTGCAGAGA3', mir-9a 5'CCGATTTCTCCGTCTTTTCTG3' and 5'CCGATTTCTCCGTCTTTTCTG3', mir-0002 TGTA CGGGCAGTACTGGG and TCTTGATGATGCGTCTTG, mir-0004 5'CAACGATGCGTTTCGACTTA3' and 5'GTACCCACGAGTCGTAC3', mir-0005 5'TCGATA TTCGAAACGCAACA3' and 5'TGGATTTGAATTCGTGTATGAAA3', mir-0007 5'ACGAGGATACACGGATGGAC3' and CAATTCACTTCCTTTTCACCTCA3'.

In situ hybridization on honeybee embryos

Performed as per Osborne [48] with the following modifications. Incubations with pri-miRNA anti-sense and sense probes were carried out at 60°C with rotation for 48 hours before post-hybridisation wash steps to remove excess probe. Embryos were incubated overnight at 4°C with anti-dioxygenin-alkaline phosphatase antibody with rotation before post-antibody wash steps and colour reaction.

Dicer siRNA knockdown

Two siRNAs were designed against Am-dicer; GGACGAAGAGUUAGAGUUUU and UGAAACAGCUAGUGAUUUUU. The two siRNAs were injected together at a final concentration of 5 µg/ml into freshly laid eggs attached to plastic strips from an Eziqueen frame [49]. As a control, a non-targeting siRNA (D-001810-01-05, Dharmacon) was also injected at the same concentration. Following injection, embryos were placed in a humidified incubator at 35°C for 48 hours. After incubation they were fixed with heptane/formaldehyde in PBS overnight, rocking at room temperature.

After fixation, embryos were washed with PBS and stained with DAPI before visualization on an Olympus BX61 microscope with a DP71 camera. Embryos were staged as per DuPraw [19].

Additional files

Additional file 1: Figure S3. (A) Pseudocoloured image showing DAPI (blue) and *Am-eve* RNA staining in red. *Am-eve* RNA is detected in the cytoplasm of embryonic cells. (B) Pseudocoloured image of *pri-mir-0002* RNA staining (red) overlaid with DAPI (blue) staining. *Pri-mir-0002* RNA is detected in the nucleus of embryonic cells.

Additional file 2: Figure S4. Genome location and read count for *Ame-mir-10*. Genome location of *Ame-mir-10* in the *Hox* complex, between *deformed (dfd)* and *sex-combed reduced (Scr)*.

Additional file 3: Figure S5. Clusterdraw analysis of the upstream regions of *Dme-mir-1* and *Ame-mir-1*. Cluster of two binding sites using the clusterdraw programme [50] with background model either site at *D. melanogaster* (A) or *A. mellifera* (B). This programme has successfully identified cis-regulatory elements in *Apis* and *Drosophila* previously [22,50-52]. P values cut off on the Y-axis and position in the sequence along the X-axis.

Additional file 4: Figure S6. Alignment of mir-0008 and mir-0005/mir-92b pre-miRNAs. Abbreviations: *Apis mellifera*, *Bombus impatiens*, *Atta cephalotes*, *Nasonia vitripennis*, *Drosophila melanogaster*. Boxed are the mature miRNA sequences.

Additional file 5: Figure S7. Alignment of sequence reads to *Apis* mir-92a, mir-92b-1 and mir-0005/mir-92b pre-miRNAs.

Additional file 6: Table S1. Phenotype of surviving larvae (at 72 hours) following siRNA injections.

Additional file 7: Figure S1. Length distribution in both samples of clean small RNA reads.

Additional file 8: Figure S2. miRNA nucleotide bias at each position.

Competing interests

The authors declare they have no competing interests.

Authors' contributions

MJW performed most of the *in situ* hybridisations, siRNA experiments and designed the project and wrote the publication. LZ cloned and synthesized the *in situ* probes, performed some of the honeybee *in situ* hybridisations and was involved in the writing of the manuscript. PKD discussed the data and took part in writing the manuscript. All authors read and approved the final manuscript.

Acknowledgments

We would like to thank James Smith for critical reading of this manuscript. Otto Hyink for help with the honeybee hives. Cris Anderson for technical assistance. This work was supported by a University of Otago Research Grant to MJW.

Received: 7 August 2012 Accepted: 22 October 2012

Published: 2 November 2012

References

1. Birney E, et al: Identification and analysis of functional elements in 1% of the human genome by the ENCODE pilot project. *Nature* 2007, **447**:799-816. doi:10.1038/nature05874.
2. Graveley BR, et al: The developmental transcriptome of *Drosophila melanogaster*. *Nature* 2011, **471**:473-479. doi:10.1038/nature09715.
3. Jacquier A: The complex eukaryotic transcriptome: unexpected pervasive transcription and novel small RNAs. *Nat Rev Genet* 2009, **10**:833-844. doi:10.1038/nrg2683.
4. Lee Y, Jeon K, Lee JT, Kim S, Kim VN: MicroRNA maturation: stepwise processing and subcellular localization. *EMBO J* 2002, **21**:4663-4670.

5. Lee YS, et al: Distinct roles for *Drosophila* Dicer-1 and Dicer-2 in the siRNA/miRNA silencing pathways. *Cell* 2004, **117**:69–81.
6. Hutvagner G, Zamore PD: A microRNA in a multiple-turnover RNAi enzyme complex. *Science* 2002, **297**:2056–2060. doi:10.1126/science.1073827.
7. Cannell IG, Kong YW, Bushell M: How do microRNAs regulate gene expression? *Biochem Soc Trans* 2008, **36**:1224–1231. doi:10.1042/BST0361224.
8. Aboobaker AA, Tomancak P, Patel N, Rubin GM, Lai EC: *Drosophila* microRNAs exhibit diverse spatial expression patterns during embryonic development. *Proc Natl Acad Sci U S A* 2005, **102**:18017–18022. doi:0508823102 [pii] 10.1073/pnas.0508823102.
9. He X, He X, Yan YL, DeLaurier A, Postlethwait JH: Observation of miRNA gene expression in zebrafish embryos by *in situ* hybridization to microRNA primary transcripts. *Zebrafish* 2011, **8**:1–8. doi:10.1089/zeb.2010.0680.
10. Lau NC, Lim LP, Weinstein EG, Bartel DP: An abundant class of tiny RNAs with probable regulatory roles in *Caenorhabditis elegans*. *Science* 2001, **294**:858–862. doi:10.1126/science.1065062.
11. He X, et al: miR-196 regulates axial patterning and pectoral appendage initiation. *Dev Biol* 2011, **357**:463–477. doi:10.1016/j.ydbio.2011.07.014.
12. Chen Z, et al: DiGeorge syndrome critical region 8(DGCR8) mediated miRNA biogenesis is essential for vascular smooth muscle cell development in mice. *J Biol Chem* 2012. doi:10.1074/jbc.M112.351791.
13. Zhao Z, et al: A negative regulatory loop between microRNA and Hox gene controls posterior identities in *Caenorhabditis elegans*. *PLoS Genet* 2010, **6**. doi:10.1371/journal.pgen.1001089.
14. Shibata M, Nakao H, Kiyonari H, Abe T, Aizawa S: MicroRNA-9 regulates neurogenesis in mouse telencephalon by targeting multiple transcription factors. *J Neurosci* 2011, **31**:3407–3422. doi:10.1523/JNEUROSCI.5085-10.2011.
15. Shomron N: MicroRNAs and developmental robustness: a new layer is revealed. *PLoS Biol* 2010, **8**:e1000397. doi:10.1371/journal.pbio.1000397.
16. Krek A, et al: Combinatorial microRNA target predictions. *Nat Genet* 2005, **37**:495–500. doi:10.1038/ng1536.
17. Heimberg AM, Sempere LF, Moy VN, Donoghue PC, Peterson KJ: MicroRNAs and the advent of vertebrate morphological complexity. *Proc Natl Acad Sci U S A* 2008, **105**:2946–2950. doi:10.1073/pnas.0712259105.
18. Marco A, Hui JH, Ronshaugen M, Griffiths-Jones S: Functional shifts in insect microRNA evolution. *Genome Biol Evol* 2010, **2**:686–696. doi:10.1093/gbe/evq053.
19. DuPraw EJ: In *Methods in developmental biology*. Edited by Wilt FH, Wessells NK, Thomas Y. New York, USA: Crowell Company; 1967:183–218.
20. Osborne PW, Dearden PK: Expression of Pax group III genes in the honeybee (*Apis mellifera*). *Dev Genes Evol* 2005, **215**:499–508.
21. Rushlow C, Levine M: Role of the *zerknüllt* gene in dorsal-ventral pattern formation in *Drosophila*. *Adv Genet* 1990, **27**:277–307.
22. Wilson MJ, Dearden PK: Diversity in insect axis formation: two orthodenticle genes and hunchback act in anterior patterning and influence dorsoventral organization in the honeybee (*Apis mellifera*). *Development* 2011, **138**:3497–3507. doi:10.1242/dev.067926.
23. Wilson MJ, Dearden PK: Tailless patterning functions are conserved in the honeybee even in the absence of Torso signaling. *Dev Biol* 2009, **335**:276–287. doi:S0012-1606(09)01174-9 [pii] 10.1016/j.ydbio.2009.09.002.
24. Wilson MJ, Havler M, Dearden PK: Giant, Kruppel, and caudal act as gap genes with extensive roles in patterning the honeybee embryo. *Dev Biol* 2010, **339**:200–211. doi:S0012-1606(09)01405-5 [pii] 10.1016/j.ydbio.2009.12.015.
25. Lemons D, Pare A, McGinnis W: Three *Drosophila* Hox complex microRNAs do not have major effects on expression of evolutionarily conserved Hox gene targets during embryogenesis. *PLoS One* 2012, **7**:e31365. doi:10.1371/journal.pone.0031365.
26. Enright AJ, et al: MicroRNA targets in *Drosophila*. *Genome Biol* 2003, **5**:R1. doi:10.1186/gb-2003-5-1-r1.
27. Ruby JG, et al: Evolution, biogenesis, expression, and target predictions of a substantially expanded set of *Drosophila* microRNAs. *Genome Res* 2007, **17**:1850–1864. doi:10.1101/gr.6597907.
28. Griffiths-Jones S, Hui JH, Marco A, Ronshaugen M: MicroRNA evolution by arm switching. *EMBO Rep* 2011, **12**:172–177. doi:10.1038/embor.2010.191.
29. Sokol NS, Ambros V: Mesodermally expressed *Drosophila* microRNA-1 is regulated by Twist and is required in muscles during larval growth. *Genes Dev* 2005, **19**:2343–2354. doi:10.1101/gad.1356105 [pii] 10.1101/gad.1356105.
30. Tang Y, et al: MicroRNA-1 regulates cardiomyocyte apoptosis by targeting Bcl-2. *Int Hear J* 2009, **50**:377–387.
31. Chen JF, et al: The role of microRNA-1 and microRNA-133 in skeletal muscle proliferation and differentiation. *Nat Genet* 2006, **38**:228–233. doi:10.1038/ng1725.
32. Biemar F, et al: Comprehensive identification of *Drosophila* dorsal-ventral patterning genes using a whole-genome tiling array. *Proc Natl Acad Sci U S A* 2006, **103**:12763–12768. doi:10.1073/pnas.0604484103.
33. Kwon C, Han Z, Olson EN, Srivastava D: MicroRNA1 influences cardiac differentiation in *Drosophila* and regulates Notch signaling. *Proc Natl Acad Sci U S A* 2005, **102**:18986–18991. doi:10.1073/pnas.0509535102.
34. Li Y, Wang F, Lee JA, Gao FB: MicroRNA-9a ensures the precise specification of sensory organ precursors in *Drosophila*. *Genes Dev* 2006, **20**:2793–2805. doi:10.1101/gad.1466306.
35. Leucht C, et al: MicroRNA-9 directs late organizer activity of the midbrain-hindbrain boundary. *Nat Neurosci* 2008, **11**:641–648. doi:10.1038/nn.2115.
36. Deo M, Yu JY, Chung KH, Tippens M, Turner DL: Detection of mammalian microRNA expression by *in situ* hybridization with RNA oligonucleotides. *Dev Dyn* 2006, **235**:2538–2548. doi:10.1002/dvdy.20847.
37. Iovino N, Pane A, Gaul U: miR-184 has multiple roles in *Drosophila* female germline development. *Dev Cell* 2009, **17**:123–133. doi:10.1016/j.devcel.2009.06.008.
38. Chen, et al: Identification and Characterization of novel amphioxus microRNAs by Solexa sequencing. *Genome Biology* 2009, **10**:R78. doi:10.1186/gb-2009-10-7-r78. http://sourceforge.net/projects/mireap/.
39. Campo-Paysaa F, Semon M, Cameron RA, Peterson KJ, Schubert M: microRNA complements in deuterostomes: origin and evolution of microRNAs. *Evol Dev* 2011, **13**:15–27. doi:10.1111/j.1525-142X.2010.00452.x.
40. Rodriguez A, Griffiths-Jones S, Ashurst JL, Bradley A: Identification of mammalian microRNA host genes and transcription units. *Genome Res* 2004, **14**:1902–1910. doi:10.1101/gr.2722704.
41. Martinez NJ, et al: Genome-scale spatiotemporal analysis of *Caenorhabditis elegans* microRNA promoter activity. *Genome Res* 2008, **18**:2005–2015. doi:10.1101/gr.083055.108.
42. Sempere LF, Martinez P, Cole C, Baguna J, Peterson KJ: Phylogenetic distribution of microRNAs supports the basal position of aceol flatworms and the polyphyly of Platyhelminthes. *Evol Dev* 2007, **9**:409–415. doi:10.1111/j.1525-142X.2007.00180.x.
43. Wheeler BM, et al: The deep evolution of metazoan microRNAs. *Evol Dev* 2009, **11**:50–68. doi:10.1111/j.1525-142X.2008.00302.x.
44. Li R, et al: SOAP2: an improved ultrafast tool for short read alignment. *Bioinformatics* 2009, **25**:1966–1967. doi:10.1093/bioinformatics/btp336. 1966-1967.
45. Dearden PK, et al: Patterns of conservation and change in honey bee developmental genes. *Genome Res* 2006, **16**:1376–1384.
46. Wilkins A: *The Evolution of Developmental Pathways*. Sunderland, MA, USA: Sinauer Associates; 2002.
47. Davidson EH: *The regulatory genome: gene regulatory networks in development and evolution*. Burlington, MA, USA: Academic Press, Elsevier; 2006.
48. Osborne PW, Dearden PK: Non-radioactive *in-situ* hybridisation to honeybee embryos and ovaries. *Apidologie* 2005, **36**:113–118.
49. Dearden PK, Duncan EJ, Wilson MJ: RNA interference (RNAi) in honeybee (*Apis mellifera*) embryos. *Cold Spring Harb Protoc* 2009, doi:2009/6/pdb.prot5228 [pii] 10.1101/pdb.prot5228. pdb prot5228.
50. Papatsenko D: ClusterDraw web server: a tool to identify and visualize clusters of binding motifs for transcription factors. *Bioinformatics* 2007, **23**:1032–1034. doi:10.1093/bioinformatics/btm047.
51. Papatsenko D, Goltsev Y, Levine M: Organization of developmental enhancers in the *Drosophila* embryo. *Nucleic Acids Res* 2009, **37**:5665–5677. doi:10.1093/nar/gkp619.
52. Perry MW, Boettiger AN, Levine M: Multiple enhancers ensure precision of gap gene-expression patterns in the *Drosophila* embryo. *Proc Natl Acad Sci U S A* 2011, **108**:13570–13575. doi:10.1073/pnas.1109873108.

doi:10.1186/1471-2148-12-211

Cite this article as: Zondag et al: Deep sequencing and expression of microRNAs from early honeybee (*Apis mellifera*) embryos reveals a role in regulating early embryonic patterning. *BMC Evolutionary Biology* 2012 **12**:211.