



HHS Public Access

Author manuscript

Int J Obes (Lond). Author manuscript; available in PMC 2021 May 18.

Published in final edited form as:

Int J Obes (Lond). 2021 January ; 45(1): 25–35. doi:10.1038/s41366-020-00717-x.

Gestational Perfluoroalkyl Substance Exposure and Body Mass Index Trajectories over the First 12 Years of Life

Joseph M. Braun¹, Melissa Eliot¹, George D. Papandonatos², Jessie P. Buckley³, Kim M. Cecil⁴, Heidi J. Kalkwarf⁵, Aimin Chen⁶, Charles B. Eaton^{1,7}, Karl Kelsey¹, Bruce P. Lanphear⁸, Kimberly Yolton⁵

¹Department of Epidemiology, Brown University, Providence, RI

²Department of Biostatistics, Brown University, Providence, RI

³Department of Environmental Health and Engineering, Johns Hopkins University, Baltimore, MD

⁴Department of Radiology, Cincinnati Children's Hospital Medical Center, University of Cincinnati College of Medicine, Cincinnati, OH

⁵Department of Pediatrics, Cincinnati Children's Hospital Medical Center, University of Cincinnati College of Medicine, Cincinnati, OH

⁶Department of Biostatistics, Epidemiology and Informatics, University of Pennsylvania Philadelphia, PA

⁷Department of Family Medicine, Warren Alpert Medical School of Brown University, Providence, RI

⁸Faculty of Health Sciences, Simon Fraser University, Vancouver, BC, Canada

Abstract

Background/Objectives: Gestational exposure to perfluoroalkyl substances (PFAS), a ubiquitous class of persistent endocrine disrupting chemicals, is associated with increased risk of obesity and cardiometabolic disease. However, it is unclear if gestational PFAS exposure is associated with adiposity trajectories related to adult obesity and cardiometabolic health.

Subjects/Methods: We measured perfluorooctanoic acid (PFOA), perfluorooctanesulfonic acid (PFOS), perfluorononanoic acid, and perfluorohexanesulfonic acid (PFHxS) concentrations in maternal serum collected between 16-weeks gestation and delivery in a cohort of 345 mother-child pairs in Cincinnati, OH (enrolled 2003–06). From age 4-weeks to 12 years, we measured weight and length or height up to eight times and calculated child BMI (1,865 repeated measures). Using

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

Corresponding Author: Joseph M. Braun, Department of Epidemiology, Brown University, Box G-S121(2), Providence, RI 02912, 401-863-5397, joseph_braun_1@brown.edu.

Competing Interests: Brown University was compensated for Dr. Braun's services as an expert witness for plaintiffs in litigation related to PFAS-contaminated drinking water; these funds were not paid to JMB directly. Dr. Braun received a honorarium for serving on an advisory board to Quest Diagnostics. Dr. Eaton was compensated as an expert witness for plaintiffs in litigation related to PFAS contaminated drinking water. Dr. Lanphear served as an expert witness in cases related to childhood lead poisoning, but he has not personally received any compensation for these services. The other authors report no conflicts of interest.

covariate-adjusted linear mixed models and splines to account for repeated BMI measures and non-linear BMI patterns, respectively, we estimated the age/magnitude of infancy BMI zenith (~1 year) and childhood BMI nadir (~5 years), BMI accrual from 8–12 years, and BMI at age 12 years by PFAS terciles.

Results: BMI trajectories varied by PFOA concentrations (age × PFOA interaction p -value=0.03). Children born to women with higher PFOA concentrations had lower infancy and early childhood BMI, earlier BMI nadir, accelerating BMI gains in mid-childhood and adolescence, and higher BMI at age 12 years. Some of these associations were non-monotonic. PFOS and PFHxS were not associated with alterations in BMI trajectories, but were monotonically associated with lower BMI across infancy, childhood, and adolescence. Compared to children in the 1st PFOS tercile, those in the 2nd (β :-0.83; 95%CI:-2.11, 0.51 kg/m²) and 3rd (β :-1.41; 95%CI:-2.65, -0.14 kg/m²) had lower BMI at age 12 years.

Conclusions: These results suggest that gestational PFOA exposure may be associated with BMI trajectories related to adult obesity and cardiometabolic disease, while PFOS and PFHxS exposure is associated with lower BMI in the first 12 years of life.

Introduction

In the United States (US), a staggering 32% of children are obese or overweight.(1) Obesity during childhood or adolescence increases the risk of atherosclerosis, cardiovascular disease, type 2 diabetes, hypertension, some cancers, and obesity in adulthood.(2–5) Early life adiposity patterns are strong risk factors for adolescent and adult obesity and cardiometabolic disease.(6–16) Later age and higher magnitude of infancy body mass index (BMI) zenith has been associated with excess adiposity and increased risk of obesity in adolescents and adults.(8, 14, 15) Moreover, earlier age of adiposity rebound is associated with increased risk of obesity and cardiometabolic disease in adolescence and adulthood.(8, 14–16) In one study, a 1.9-year earlier BMI rebound was associated with 3.9-times the odds of metabolic syndrome at age 31 years.(14)

Gestational exposure to obesogenic chemicals has been associated with early life adiposity patterns, suggesting that the risk of adulthood obesity and cardiometabolic disease may be preventable.(17, 18) Perfluoroalkyl substances (PFAS), a class of environmentally persistent anthropogenic chemicals, are suspected obesogens. PFAS are used in oil/water repellant textiles, food packaging, cleaning products, and firefighting foams or as processing aids for fluoropolymer manufacturing.(19) Nearly all adults in the US and internationally, including pregnant women, have detectable levels of PFAS in their serum and many PFAS have biological half-lives on the order of years in humans.(20–25) Gestational PFAS exposure may perturb early life adiposity patterns and increase adipose tissue accrual by re-programming biological pathways regulating growth, adipocyte differentiation or proliferation, neuroendocrine regulation, or energy metabolism.(26–31)

Epidemiological studies suggest that gestational PFAS exposure is associated with decreased fetal growth, (32, 33) alterations in infant or childhood growth,(34, 35) and increased adiposity during infancy, childhood, and adulthood.(35–40) However, we are unaware of studies examining whether gestational PFAS exposure is associated with childhood adiposity

patterns related to later life obesity or cardiometabolic disease, namely higher infancy BMI zenith, earlier BMI nadir, or accelerations in BMI accrual during childhood and adolescence. (7–13) Thus, we used a prospective cohort study to investigate whether maternal serum concentrations of four PFAS were associated with features of BMI patterns from age 4 weeks to 12 years in 345 children from Cincinnati, Ohio.

Methods

Study Participants

Between March 2003 and January 2006, we recruited pregnant women into a longitudinal pregnancy and birth cohort study, the Health Outcomes and Measures of the Environment Study (The HOME Study).^(41, 42) We identified women living in the Cincinnati, OH region who attended one of nine prenatal practices affiliated with three hospitals. Eligibility criteria included: 16±3 weeks gestation, 18 years old, residing in a residence built in or before 1978, not living in a mobile/trailer home, HIV-negative, not taking medications for seizures or thyroid disorders, planning to continue prenatal care and deliver at the collaborating clinics and hospitals, planning to live in the greater Cincinnati area for the next year, English fluency, and no diagnosis of diabetes, bipolar disorder, schizophrenia, or cancer resulting in radiation treatment or chemotherapy. The present analysis includes 345 mothers who gave birth to a singleton child and had gestational serum PFAS measurements, covariate data, and at least one child anthropometry assessment between ages 4 weeks and 12 years.

The Institutional Review Boards of Cincinnati Children's Hospital Medical Center (CCHMC) and all delivery hospitals approved the study protocol. The Centers for Disease Control and Prevention (CDC) deferred to the CCHMC IRB as the IRB of record since the role of the CDC was primarily technical oversight of the PFAS assays. Women provided written informed consent for themselves and their children. Children provided written informed assent at the 12-year visit.

Gestational PFAS Exposure Assessment

We measured serum PFAS concentrations in maternal venous blood samples collected at approximately 16 weeks gestation, 26 weeks gestation, or within 24 hours of delivery. After separating serum from clotted blood, we stored samples at -80°C until they were shipped on dry ice to the CDC laboratories. While most women had a sufficient volume of serum for quantification of PFASs in their 16 week sample (n=294, 85.2%), some women did not and we quantified PFAS concentrations in samples collected at 26 weeks (n=34, 9.9%) or within 24 hours of delivery (n=17, 4.2%). Staff at the CDC laboratory quantified perfluorooctanoic acid (PFOA), perfluorooctanesulfonic acid (PFOS), perfluorononanoic acid (PFNA), and perfluorohexanesulfonic acid (PFHxS) concentrations using on-line solid phase extraction coupled with high performance liquid chromatography-isotope dilution tandem mass spectrometry.⁽⁴³⁾ We detected all four PFASs in every sample, with limits of detection ranging between 0.082–0.2 ng/mL. Quality control materials and reagent blanks were included in each analytic batch with coefficients of variation in repeated quality control materials of approximately 6%.

Infant and Child Anthropometry

We measured children's weight and length or height during home or clinic visits at ages 4 weeks and 1, 2, 3, 4, 5, 8, and 12 years of age. At each visit, we measured weight to the nearest 0.01 kg with the child dressed in undergarments or a dry diaper using a digital scale. At the 4-week and 1-year visit, we measured infant length with a length board to the nearest 0.1 cm. We measured height at later study visits to the nearest 0.1 cm with the child standing straight without shoes or head coverings and heels positioned against the wall using a wall-mounted stadiometer. Each weight, length, and height measure was taken in triplicate and averaged for analysis. We calculated BMI in kg/m^2 . Examiners were blinded to mother and children's PFAS concentrations.

Potential Confounders

Trained research assistants collected sociodemographic covariates including maternal race, age, marital status, and household income using standardized computer-assisted interviews. We assessed perinatal variables including maternal pre-pregnancy weight, height, and parity via standardized chart abstraction forms. We calculated pre-pregnancy body mass index (BMI) using self-reported weight/height for ~66% of women and imputed missing values for the remainder using Super Learner (Supplemental Methods).(45) Finally, we averaged serum cotinine (a biomarker of tobacco smoke exposure) concentrations measured at 16 or 26 weeks gestation or within 48 hours of birth to assess secondhand or active tobacco smoke exposure.(46)

We selected potential confounders that might be associated with both gestational PFAS concentrations and child growth using a directed acyclic graph (Supplemental Figure 1). We did not adjust for potential causal intermediates (e.g., birth weight, breastfeeding) in our primary analyses, as prenatal PFAS concentrations have previously been associated with reduced fetal growth and decreased breastfeeding duration.(32, 44) We derived a minimum sufficient set of variables that included maternal race, maternal age, maternal BMI, parity, household income, and gestational tobacco smoke exposure. We included child sex and child age at BMI measurement as precision variables.

Statistical Analyses

First, we calculated univariate statistics of gestational PFAS concentrations, repeated BMI measures, and covariates. We also computed the Pearson correlation coefficient between \log_2 -transformed PFOA, PFOS, PFNA, and PFHxS concentrations.

Next, we examined the shape of the relation between age and infant/child BMI (kg/m^2) using spaghetti plots. Then, we fit a linear mixed model with fixed effects given by a truncated cubic polynomial spline for age with child-specific random intercepts and linear age slopes to account for the non-linear shape of the age-BMI relation and repeated measures on each child, respectively.(47) We began by using a cubic polynomial spline to characterize the shape of this relation; we then chose the number and position of the knots based on the distribution of the anthropometry measurements (Supplemental Table 1). A forward selection approach was based on adding knots one at a time and choosing their location via profile likelihood estimation, while keeping the location of previously selected

knots fixed. Non-nested model comparison using the Akaike Information Criterion resulted in the single-knot model $BMI(t_{ij}) = (\beta_0 + b_{0i}) + (\beta_1 + b_{1i}) * t_{ij} + \beta_2 * t_{ij}^2 + \beta_3 * t_{ij}^3 + \gamma_3 * (t_{ij} - t_0)^3 * I(t_{ij} > t_0) + \epsilon_{ij}$. Here (b_{0i}, b_{1i}) is a vector of normally-distributed random effects with mean zero and unstructured covariance matrix, ϵ_{ij} denotes a normally-distributed residual of zero mean and homoscedastic variance, and $I(s)$ is an indicator function taking the value 0 for $s \leq 0$ and 1 for $s > 0$. As study visit did not occur on the same date for all children, the j_{th} visit time for the i_{th} child given by t_{ij} is allowed to vary across children. The fixed effects part of this model has a first derivative $BMI^{(1)}(t_{ij}) = \beta_1 + 2 * \beta_2 * t_{ij} + 3 * \beta_3 * t_{ij}^2 + 3 * \gamma_3 * (t_{ij} - t_0)^2 * I(t_{ij} > t_0)$, a truncated quadratic spline whose roots closest to the knot t_0 can also be obtained in closed form and give the zenith and nadir of early childhood BMI.

With adjustment for covariates, we examined several aspects of children's longitudinal BMI measurements by fitting the linear mixed model above using R package lme4 (<https://cran.r-project.org/web/packages/lme4>). First, we determined whether the overall BMI trajectory varied according to PFAS tercile using the age \times PFAS interaction term p-value; we considered BMI trajectories to vary if this p-value was < 0.05 . Then, we estimated five features of average BMI trajectories according to terciles of gestational PFAS concentrations that have been associated with increased risk of later life obesity and cardiometabolic disease.(6–13) These features included: 1) the age (years) of the infancy BMI zenith, 2) the magnitude (kg/m^2) of infancy BMI zenith, 3) the age (years) of early childhood BMI nadir, 4) the magnitude (kg/m^2) of early childhood BMI nadir, and 5) the rate ($kg/m^2/y$) of adolescent BMI change (age 8 to 12 years). These features can be expressed as functions of the fixed effects parameters of the truncated basis spline model given above, when PFAS exposure is coded so that the tercile of interest is the reference group. In addition to PFAS exposure, our model adjusted for potential confounding by maternal race, age, BMI, parity, household income, gestational tobacco smoke exposure, and child sex. All potential confounders were centered at the sample mean, including categorical ones resulting in average BMI trajectories at a particular PFAS tercile for a theoretical sample whose maternal race and child sex composition reflects that of the HOME Study. We used percentile bootstrap approaches to estimate 95% confidence intervals (CIs) for all quantities of interest based on parametric bootstrapping with $B=10,000$ iterations in a parallel processing environment.

Secondary and Sensitivity Analyses

We conducted a secondary analysis by modeling the age-BMI relation separately for boys and girls given that prior studies suggest potential sex-specific associations.(38) Then, we conducted sensitivity analyses to test the robustness of our results to various adjustments and assumptions. First, we examined whether plasma volume expansion during pregnancy, hypertensive disorders during pregnancy, or maternal diet during pregnancy confounded the association between PFAS and BMI trajectories by limiting our analyses to women with serum PFAS measures at 16 weeks gestation, excluding women with pregnancy-induced hypertensive disorders, and adjusting for maternal intake of fish, respectively. We examined hypertensive disorders since reduced renal function during pregnancy may be associated with both higher serum PFAS levels and lower birth weight, with the latter being a predictor of subsequent BMI (Supplemental Figure 2). Using standardized chart abstraction, we

characterized gestational hypertension, preeclampsia, eclampsia, and HELLP (hemolysis, elevated liver enzymes, low platelet count) syndrome as pregnancy induced hypertensive diseases.(48) We assessed the frequency of fin- or shellfish intake during the first half of pregnancy with a standardized question. Finally, we restricted our analyses to children who had three or more BMI measures between ages 4 weeks and 12 years to determine if our results were sensitive to including children with fewer repeated BMI measures.

Results

At baseline, women included in this analysis were predominately non-Hispanic White (62.0%), college-educated (50.1%), nulliparous (42.9%), and non-smokers (89.6%). On average, they were 29.9 years old at delivery and had a median household income of \$55,000/year during pregnancy. Baseline characteristics of women included in this analysis were similar to the full cohort.(42)

Median serum PFAS concentrations ranged from 0.9 (PFNA) to 13.8 (PFOS) ng/mL (Supplemental Table 2). Pearson correlations coefficients between log₂-transformed PFAS concentrations ranged from 0.38 (PFNA-PFHxS) to 0.64 (PFOS-PFHxS) (Supplemental Figure 3). Among women with two or more serum PFAS measures at 16 weeks, 26 weeks, and birth, the intraclass correlation coefficients (ICC) between repeated PFOA (ICC=0.76), PFOS (ICC=0.76), PFNA (ICC=0.68), and PFHxS (ICC=0.78) indicated very good to excellent reproducibility.(49)

Three-hundred and forty-five children (53.3% girls) had 1,865 repeated BMI measures and 83.1% of children (n=287) had at least three BMI measurements (Supplemental Table 3). Fifty-eight percent (n=200) of children had at least one BMI measurement during the early childhood, middle childhood, and adolescence periods (Supplemental Table 4). Those with one BMI measurement (n=63, n=18.3%) during these three periods had them almost exclusively during early childhood (n=57, 90%), while 50% of those with two measures had them during either middle childhood or adolescence. The average BMI rose in the first year of life, declined to a nadir between 4 and 5 years of age, and rose linearly from age 5 to 12 years (Supplemental Table 1).

Using a covariate-adjusted linear mixed model with a spline to flexibly estimate the relation between child age and BMI, BMI trajectories varied non-monotonically according to PFOA tertiles (PFOA × age interaction p-value=0.03) (Figure 1, Supplemental Figure 4). Notably, children in the 2nd PFOA tertile had similar magnitude and age of BMI zenith (Table 2), an earlier nadir (-0.32 years; 95% CI: -0.96, 0.27) (Table 3), a more rapid increase in BMI from age 8 to 12 years (0.35 kg/m²/y; 95% CI: 0.11, 0.39) (Table 4), and higher absolute BMI at age 12 years (1.57 kg/m²; 95% CI: 0.34, 2.83) (Table 5) compared to children in the 1st PFOA tertile. Relative to children in the 1st PFOA tertile, those in the 3rd tertile had lower magnitude of BMI zenith (-0.5 kg/m²; 95% CI: -0.87, -0.15) (Table 2), lower magnitude (-0.52 kg/m²; 95% CI: -1.04, 0) and earlier age (-0.29 years; 95% CI: -0.94, 0.32) of BMI nadir (Table 3), but similar rate of BMI change from age 8 to 12 years and BMI at age 12 years (Tables 4 and 5).

While children's BMI trajectories did not vary according to maternal PFOS (PFOS \times age interaction p-value=0.10) or PFHxS terciles (PFHxS \times age interaction p-value=0.18) (Figure 1, Supplemental Figure 4), PFOS and PFHxS were associated with reduced BMI from 4 weeks to 12 years of age. Compared to children in the 1st tercile, children born to women in the 3rd terciles of PFOS and PFHxS had lower magnitude of infancy BMI zenith (PFOS: -0.40 kg/m²; 95% CI: -0.75, -0.06 and PFHxS: -0.42 kg/m²; 95% CI: -0.80, -0.06), lower magnitude of BMI nadir (PFOS: -0.67 kg/m²; 95% CI: -1.02, -0.19 and PFHxS: -0.60 kg/m²; 95% CI: -1.09, -0.09), and lower BMI at age 12 years (PFOS: -1.41 kg/m²; 95% CI: -2.65, -0.14 and PFHxS: -0.50 kg/m²; 95% CI: -1.78, 0.76) (Tables 2, 3, and 5). These associations were monotonic for PFOS. PFOS and PFHxS levels were also associated with later age at adiposity nadir (~0.4 to 0.6 years), but in a non-monotonic manner (Table 4).

Children's BMI trajectories did not vary according to maternal PFNA terciles (PFNA \times age interaction p-value=0.48) (Figure 1, Supplemental Figure 4). Moreover, PFNA levels were not associated with the childhood BMI features we examined (Tables 2–5).

Secondary and Sensitivity Analyses

There was evidence that the association of PFOA with children's BMI trajectories was modified by child sex (PFOA \times sex \times age interaction p-value=0.013) (Supplemental Figure 4). BMI at the time of early childhood BMI nadir was monotonically lower among boys in the 2nd and 3rd PFOA terciles, whereas the age of early childhood BMI nadir was non-monotonically earlier among girls in the top two terciles (Supplemental Tables 4–6). In both sexes, the association of PFOA with other BMI trajectory features were similar. Child sex did not modify the association of PFOS, PFNA, and PFHxS with BMI trajectories (PFAS \times sex \times age interaction p-values = 0.19) (Supplemental Figure 4).

The pattern of associations was not markedly different when we limited our analysis to women with serum PFAS measures at 16 weeks, excluded women with pregnancy induced hypertensive disorders, adjusted for fish intake during pregnancy, or excluded children with <3 BMI measurements (Supplemental Table 7–10). However, when restricting to 16 week PFAS measurements or adjusting for fish intake during pregnancy, some associations were attenuated towards the null for PFOA (rate of BMI change from age 8 to 12 years for the 2nd vs. 1st tercile) and PFOS (magnitude of BMI zenith and nadir and BMI nadir timing) (Supplemental Tables 7 and 9).

Discussion

In this cohort, maternal serum PFOA, PFOS, and PFHxS concentrations were associated with features of BMI trajectories from birth to age 12 years. The dose-response and direction of these associations varied by the specific PFAS and childhood BMI trajectory feature. We observed non-monotonic associations of PFOA with age at adiposity nadir, BMI change from age 8 to 12 years, and BMI at age 12 years. Children in the 2nd PFOA tercile had an earlier age at adiposity nadir, greater increases in BMI between age 8 and 12 years, and higher BMI at age 12 years. The magnitude and age of early childhood BMI nadir across PFOA terciles differed for boys and girls, but patterns were similar for infancy BMI zenith and rate of BMI gain from age 8 to 12 years. In contrast to our findings for PFOA, PFOS

and PFHxS were associated had lower BMI in the first 12 years of life, and associations were monotonic for PFOS. PFNA was not associated with the BMI features we examined.

In multiple studies, higher serum PFOA concentrations during pregnancy have been associated with lower birth weight and reduced early childhood BMI,(32, 33, 50) altered growth and excess adiposity during childhood and adolescence,(35, 37, 38, 51) and excess adiposity in adulthood.(36) Experimental studies suggest that PFOA exposure can increase adipocyte differentiation and cause elevated body weight and adipocyte lipid accumulation in rodents.(29, 52–55) In a pilot study of HOME Study neonates, we observed that prenatal serum PFOA concentrations were associated with hypomethylation of genes related to fetal programming, growth, and obesity in leukocytes.(56)

We speculate that prenatal exposure to chemicals that restrict fetal growth, like PFOA, may set off a cascade of maladaptive processes that re-program systems involved in growth, energy metabolism, appetite, or adipogenesis.(57, 58) Earlier BMI nadir and BMI accelerations from childhood to adolescence are risk factors for subsequent obesity, cardiometabolic disease, cardiovascular disease, ischemic stroke, and type 2 diabetes.(7–13) Given this, future studies could examine prenatal PFOA concentrations in relation to adolescent or adulthood cardiometabolic or cardiovascular risk factors or disease.

Prenatal PFOS and PFHxS concentrations have not been consistently associated with BMI features across infancy and childhood. In two studies, prenatal PFOS concentrations were associated with lower adiposity in 415 US girls at age 5 months (40) and lower BMI and waist circumference in 811 Danish children at age 7 years.(59) In contrast, prenatal PFOS concentrations were associated with higher childhood adiposity in 811 8-year old girls from the US,(38) higher BMI in 381 Swedish children at ages 3 to 5 years,(51) and higher waist-to-hip ratio in 1,022 5 to 9 year old children from Greenland and Ukraine.(37)

The pattern of associations for individual PFAS from this and other studies do not conclusively indicate that a given PFAS is obesogenic or associated with infancy, childhood, or adolescent BMI trajectories related to later life obesity. Indeed, some studies did not observe an association of prenatal PFAS exposure with excess child or adult adiposity or higher risk of being overweight or obese.(60–62) However, there is consistent evidence in both epidemiological studies and experimental studies of animals that prenatal PFOA exposure is associated with reduced fetal growth.(32, 63) The discrepant findings across cohorts might be due to different levels of maternal serum PFAS concentrations, as well as the method and timing of child adiposity assessment. The timing of assessment could be critical if PFAS exposures are associated with alterations in changes in weight or length/height because these studies have assessed childhood adiposity at times ranging from 1-month of age to adulthood. Finally, it is not clear why we observed that PFOS and PFHxS were associated with lower BMI, and PFOA with a more “obesogenic” BMI profile, but prior studies report that the biological mechanisms of PFOA and PFOS exposure differ, despite subtle differences in chemical structures.(64, 65)

Adolescence may be a period of life to mitigate the potential impact of PFAS exposure on later life risk of adverse cardiometabolic outcomes. Notably, in a study of over 62,000

Danish men, those who went from being overweight to normal weight between age 7 and 13 years or 13 years and early adulthood had reduced risk for type 2 diabetes relative to men who were persistently overweight.(13) Moreover, in over 2,700 participants from the Bogulusa Heart Study, greater rates of BMI change in adolescence were associated with increased risk of adult obesity and the rates of change were stronger predictors than absolute BMI during adolescence.(9)

This study has several strengths and limitations. First, we applied appropriate growth modeling techniques in over 1,800 repeated BMI measurements between infancy and adolescence to determine if PFAS concentrations were associated with features of BMI trajectories related to later life risk of obesity and cardiometabolic disease. Despite having eight repeated BMI measurements, we had a greater density of measurements in early childhood. Thus, caution is warranted in interpreting features at later ages. Relatedly, there was loss to follow-up, particularly at ages 4 and 5 years. Reassuringly, participants who remained in the study were generally similar to the full cohort and the presented results were not substantially different when we limited to participants with 3 or more repeated BMI measurements.(41, 42) Second, while we controlled for variables associated with maternal PFAS concentrations that are also potentially important determinants of childhood growth patterns,(66) there may be residual confounding. Notably, we had a relatively crude measure of maternal diet during pregnancy and we were not able to control for maternal renal function during pregnancy (i.e., glomerular filtration rate [GFR]). Higher GFR has been associated with both higher birth weight and lower serum PFAS concentrations.(67) Thus, not controlling for GFR could bias our results. While our results were similar when we restricted to women with PFAS measures at 16 weeks gestation, the rate of BMI change from age 8 to 12 years for the 2nd PFOA tercile was attenuated with this restriction. Interestingly, a prior study examining prenatal serum PFAS and childhood adiposity did not report substantially different results when controlling for maternal GFR during pregnancy. (38) Finally, we did not estimate the potential aggregate effect of PFAS mixtures on childhood growth patterns. Maternal serum PFAS concentrations were positively correlated, indicating joint exposure to multiple PFAS, although we identified different dose-response patterns between PFOA and PFOS exposures. Future studies could use novel and evolving methods to examine the impact of PFAS mixtures on BMI trajectories and also examine the impact of postnatal PFAS exposures on child BMI trajectories.(68)

In this cohort, maternal serum PFOA concentrations were non-monotonically associated with features of childhood BMI trajectories related to adolescent and adulthood obesity and cardiometabolic disease. PFOS concentrations, and to a lesser extent, PFHxS concentrations, were associated with lower BMI during the first 12 years of life. These results highlight the importance of quantifying the potential effects of chemical obesogens on both absolute levels of adiposity, as well as adiposity trajectories. Future studies should determine if adiposity trajectory features mediate previously observed associations of prenatal PFAS exposure with later life obesity and cardiometabolic disease.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported by National Institute of Environmental Health Sciences grants P01 ES011261, R01 ES014575, R01 ES020349, R01 ES027244, R01 ES025214, and R01 ES030078. We would like to acknowledge the Centers for Disease Control and Prevention (CDC) laboratory staff who performed the PFAS measurements. We are grateful to our participants for the time they have given to our study.

References

1. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of childhood and adult obesity in the United States, 2011–2012. *JAMA*. 2014;311(8):806–14. [PubMed: 24570244]
2. Sahoo K, Sahoo B, Choudhury AK, Sofi NY, Kumar R, Bhadoria AS. Childhood obesity: causes and consequences. *J Family Med Prim Care*. 2015;4(2):187–92. [PubMed: 25949965]
3. Park MH, Falconer C, Viner RM, Kinra S. The impact of childhood obesity on morbidity and mortality in adulthood: a systematic review. *Obesity reviews : an official journal of the International Association for the Study of Obesity*. 2012;13(11):985–1000. [PubMed: 22731928]
4. Ebbeling CB, Pawlak DB, Ludwig DS. Childhood obesity: public-health crisis, common sense cure. *Lancet*. 2002;360(9331):473–82. [PubMed: 12241736]
5. Llewellyn A, Simmonds M, Owen CG, Woolacott N. Childhood obesity as a predictor of morbidity in adulthood: a systematic review and meta-analysis. *Obesity reviews : an official journal of the International Association for the Study of Obesity*. 2015.
6. Di Gravio C, Krishnaveni GV, Somashekara R, Veena SR, Kumaran K, Krishna M, et al. Comparing BMI with skinfolds to estimate age at adiposity rebound and its associations with cardio-metabolic risk markers in adolescence. *International journal of obesity (2005)*. 2019;43(4):683–90. [PubMed: 30006579]
7. Brisbois TD, Farmer AP, McCargar LJ. Early markers of adult obesity: a review. *Obesity reviews : an official journal of the International Association for the Study of Obesity*. 2012;13(4):347–67. [PubMed: 22171945]
8. Aris IM, Rifas-Shiman SL, Li LJ, Kleinman KP, Coull BA, Gold DR, et al. Patterns of body mass index milestones in early life and cardiometabolic risk in early adolescence. *International journal of epidemiology*. 2019.
9. Zhang T, Whelton PK, Xi B, Krousel-Wood M, Bazzano L, He J, et al. Rate of change in body mass index at different ages during childhood and adult obesity risk. *Pediatr Obes*. 2019;14(7):e12513. [PubMed: 30702812]
10. Geserick M, Vogel M, Gausche R, Lipek T, Spielau U, Keller E, et al. Acceleration of BMI in Early Childhood and Risk of Sustained Obesity. *The New England journal of medicine*. 2018;379(14):1303–12. [PubMed: 30281992]
11. Bjerregaard LG, Adelborg K, Baker JL. Change in body mass index from childhood onwards and risk of adult cardiovascular disease(). *Trends Cardiovasc Med*. 2019.
12. Gjaerde LK, Gamborg M, Angquist L, Truelsen TC, Sorensen TIA, Baker JL. Association of Childhood Body Mass Index and Change in Body Mass Index With First Adult Ischemic Stroke. *JAMA Neurol*. 2017;74(11):1312–8. [PubMed: 28828465]
13. Bjerregaard LG, Jensen BW, Angquist L, Osler M, Sorensen TIA, Baker JL. Change in Overweight from Childhood to Early Adulthood and Risk of Type 2 Diabetes. *The New England journal of medicine*. 2018;378(14):1302–12. [PubMed: 29617589]
14. Sovio U, Kaakinen M, Tzoulaki I, Das S, Ruokonen A, Pouta A, et al. How do changes in body mass index in infancy and childhood associate with cardiometabolic profile in adulthood? Findings from the Northern Finland Birth Cohort 1966 Study. *International journal of obesity (2005)*. 2014;38(1):53–9. [PubMed: 24080793]
15. Aris IM, Bernard JY, Chen LW, Tint MT, Pang WW, Lim WY, et al. Infant body mass index peak and early childhood cardio-metabolic risk markers in a multi-ethnic Asian birth cohort. *International journal of epidemiology*. 2017;46(2):513–25. [PubMed: 27649801]

16. Peneau S, Gonzalez-Carrascosa R, Gusto G, Goxe D, Lantieri O, Fezeu L, et al. Age at adiposity rebound: determinants and association with nutritional status and the metabolic syndrome at adulthood. *International journal of obesity* (2005). 2016;40(7):1150–6. [PubMed: 27113489]
17. Aris IM, Rifas-Shiman SL, Li LJ, Kleinman K, Coull BA, Gold DR, et al. Pre-, Perinatal, and Parental Predictors of Body Mass Index Trajectory Milestones. *The Journal of pediatrics*. 2018;201:69–77 e8. [PubMed: 29960766]
18. Heindel JJ, Blumberg B. Environmental Obesogens: Mechanisms and Controversies. *Annu Rev Pharmacol Toxicol*. 2018.
19. Toward a New Comprehensive GLObal Database of Per- and Polyfluoroalkyl Substances (PFAS). Paris, France: Organisation for Economic Co-operation and Development; 2018.
20. Calafat AM, Wong LY, Kuklennyik Z, Reidy JA, Needham LL. Polyfluoroalkyl chemicals in the U.S. population: data from the National Health and Nutrition Examination Survey (NHANES) 2003–2004 and comparisons with NHANES 1999–2000. *Environmental health perspectives*. 2007;115(11):1596–602. [PubMed: 18007991]
21. Woodruff TJ, Zota AR, Schwartz JM. Environmental chemicals in pregnant women in the United States: NHANES 2003–2004. *Environmental health perspectives*. 2011;119(6):878–85. [PubMed: 21233055]
22. Olsen GW, Burris JM, Ehresman DJ, Froehlich JW, Seacat AM, Butenhoff JL, et al. Half-life of serum elimination of perfluorooctanesulfonate, perfluorohexanesulfonate, and perfluorooctanoate in retired fluorochemical production workers. *Environmental health perspectives*. 2007;115(9):1298–305. [PubMed: 17805419]
23. Kashino I, Sasaki S, Okada E, Matsuura H, Goudarzi H, Miyashita C, et al. Prenatal exposure to 11 perfluoroalkyl substances and fetal growth: A large-scale, prospective birth cohort study. *Environment international*. 2020:105355.
24. Fornis J, Verner MA, Iszatt N, Nowack N, Bach CC, Vrijheid M, et al. Early Life Exposure to Perfluoroalkyl Substances (PFAS) and ADHD: A Meta-Analysis of Nine European Population-Based Studies. *Environmental health perspectives*. 2020;128(5):57002. [PubMed: 32378965]
25. Haines DA, Khoury C, Saravanabhavan G, Werry K, Walker M, Malowany M. Human biomonitoring reference values derived for persistent organic pollutants in blood plasma from the Canadian Health Measures Survey 2007–2011. *International journal of hygiene and environmental health*. 2017;220(4):744–56. [PubMed: 28372942]
26. Heindel JJ, Blumberg B, Cave M, Machtiger R, Mantovani A, Mendez MA, et al. Metabolism disrupting chemicals and metabolic disorders. *Reproductive toxicology* (Elmsford, NY). 2017;68:3–33.
27. Ye L, Guo J, Ge RS. Environmental pollutants and hydroxysteroid dehydrogenases. *Vitamins and hormones*. 2014;94:349–90. [PubMed: 24388197]
28. Fletcher T, Galloway TS, Melzer D, Holcroft P, Cipelli R, Pilling LC, et al. Associations between PFOA, PFOS and changes in the expression of genes involved in cholesterol metabolism in humans. *Environment international*. 2013;57–58:2–10.
29. Vanden Heuvel JP, Thompson JT, Frame SR, Gillies PJ. Differential activation of nuclear receptors by perfluorinated fatty acid analogs and natural fatty acids: a comparison of human, mouse, and rat peroxisome proliferator-activated receptor- α , - β , and - γ , liver X receptor- β , and retinoid X receptor- α . *Toxicol Sci*. 2006;92(2):476–89. [PubMed: 16731579]
30. Qi W, Clark JM, Timme-Laragy AR, Park Y. Perfluorobutanesulfonic acid (PFBS) potentiates adipogenesis of 3T3-L1 adipocytes. *Food Chem Toxicol*. 2018;120:340–5. [PubMed: 30031040]
31. Watkins AM, Wood CR, Lin MT, Abbott BD. The effects of perfluorinated chemicals on adipocyte differentiation in vitro. *Molecular and cellular endocrinology*. 2015;400:90–101. [PubMed: 25448844]
32. Johnson PI, Sutton P, Atchley DS, Koustas E, Lam J, Sen S, et al. The Navigation Guide - evidence-based medicine meets environmental health: systematic review of human evidence for PFOA effects on fetal growth. *Environmental health perspectives*. 2014;122(10):1028–39. [PubMed: 24968388]

33. Shoaff J, Papandonatos GD, Calafat AM, Chen A, Lanphear BP, Ehrlich S, et al. Prenatal Exposure to Perfluoroalkyl Substances: Infant Birth Weight and Early Life Growth. *Environ Epidemiol.* 2018;2(2).
34. Maisonet M, Terrell ML, McGeehin MA, Christensen KY, Holmes A, Calafat AM, et al. Maternal concentrations of polyfluoroalkyl compounds during pregnancy and fetal and postnatal growth in British girls. *Environmental health perspectives.* 2012;120(10):1432–7. [PubMed: 22935244]
35. Braun JM, Chen A, Romano ME, Calafat AM, Webster GM, Yolton K, et al. Prenatal perfluoroalkyl substance exposure and child adiposity at 8 years of age: The HOME study. *Obesity (Silver Spring, Md.* 2016;24(1):231–7.
36. Halldorsson TI, Rytter D, Haug LS, Bech BH, Danielsen I, Becher G, et al. Prenatal exposure to perfluorooctanoate and risk of overweight at 20 years of age: a prospective cohort study. *Environmental health perspectives.* 2012;120(5):668–73. [PubMed: 22306490]
37. Hoyer BB, Ramlau-Hansen CH, Vrijheid M, Valvi D, Pedersen HS, Zvezdai V, et al. Anthropometry in 5- to 9-Year-Old Greenlandic and Ukrainian Children in Relation to Prenatal Exposure to Perfluorinated Alkyl Substances. *Environmental health perspectives.* 2015;123(8):841–6. [PubMed: 25809098]
38. Mora AM, Oken E, Rifas-Shiman SL, Webster TF, Gillman MW, Calafat AM, et al. Prenatal Exposure to Perfluoroalkyl Substances and Adiposity in Early and Mid-Childhood. *Environmental health perspectives.* 2016.
39. Chen MH, Ng S, Hsieh CJ, Lin CC, Hsieh WS, Chen PC. The impact of prenatal perfluoroalkyl substances exposure on neonatal and child growth. *The Science of the total environment.* 2017;607–608:669–75.
40. Starling AP, Adgate JL, Hamman RF, Kechris K, Calafat AM, Dabelea D. Prenatal exposure to per- and polyfluoroalkyl substances and infant growth and adiposity: the Healthy Start Study. *Environment international.* 2019;131:104983. [PubMed: 31284113]
41. Braun JM, Buckley JP, Cecil KM, Chen A, Kalkwarf HJ, Lanphear BP, et al. Adolescent follow-up in the Health Outcomes and Measures of the Environment (HOME) Study: cohort profile. *BMJ open.* 2020;10(5):e034838.
42. Braun JM, Kalloo G, Chen A, Dietrich KN, Liddy-Hicks S, Morgan S, et al. Cohort Profile: The Health Outcomes and Measures of the Environment (HOME) study. *International journal of epidemiology.* 2017;46(1):24. [PubMed: 27006352]
43. Kato K, Basden BJ, Needham LL, Calafat AM. Improved selectivity for the analysis of maternal serum and cord serum for polyfluoroalkyl chemicals. *J Chromatogr A.* 2011;1218(15):2133–7. [PubMed: 21084089]
44. Romano ME, Xu Y, Calafat AM, Yolton K, Chen A, Webster GM, et al. Maternal serum perfluoroalkyl substances during pregnancy and duration of breastfeeding. *Environmental research.* 2016.
45. van der Laan MJ, Polley EC, Hubbard AE. Super learner. *Statistical applications in genetics and molecular biology.* 2007;6:Article25.
46. Braun JM, Daniels JL, Poole C, Olshan AF, Hornung R, Bernert JT, et al. A prospective cohort study of biomarkers of prenatal tobacco smoke exposure: the correlation between serum and meconium and their association with infant birth weight. *Environ Health.* 2010;9:53. [PubMed: 20799929]
47. Grajeda LM, Ivanescu A, Saito M, Crainiceanu C, Jaganath D, Gilman RH, et al. Modelling subject-specific childhood growth using linear mixed-effect models with cubic regression splines. *Emerg Themes Epidemiol.* 2016;13:1. [PubMed: 26752996]
48. Werner EF, Braun JM, Yolton K, Khoury JC, Lanphear BP. The association between maternal urinary phthalate concentrations and blood pressure in pregnancy: The HOME Study. *Environ Health.* 2015;14(1):75. [PubMed: 26380974]
49. Rosner B *Fundamentals of Biostatistics.* 4th ed. Pacific Grove, CA: Duxbury; 2000.
50. Tanner EM, Bornehag CG, Gennings C. Dynamic growth metrics for examining prenatal exposure impacts on child growth trajectories: Application to perfluorooctanoic acid (PFOA) and postnatal weight gain. *Environmental research.* 2019;182:109044. [PubMed: 31874421]

51. Gyllenhammar I, Diderholm B, Gustafsson J, Berger U, Ridefelt P, Benskin JP, et al. Perfluoroalkyl acid levels in first-time mothers in relation to offspring weight gain and growth. *Environment international*. 2017;111:191–9. [PubMed: 29223808]
52. Taxvig C, Dreisig K, Boberg J, Nellemann C, Schelde AB, Pedersen D, et al. Differential effects of environmental chemicals and food contaminants on adipogenesis, biomarker release and PPAR γ activation. *Molecular and cellular endocrinology*. 2012;361(1–2):106–15. [PubMed: 22526026]
53. Bastos Sales L, Kamstra JH, Cenijn PH, van Rijt LS, Hamers T, Legler J. Effects of endocrine disrupting chemicals on in vitro global DNA methylation and adipocyte differentiation. *Toxicology in vitro : an international journal published in association with BIBRA*. 2013;27(6):1634–43. [PubMed: 23603478]
54. Hines EP, White SS, Stanko JP, Gibbs-Flournoy EA, Lau C, Fenton SE. Phenotypic dichotomy following developmental exposure to perfluorooctanoic acid (PFOA) in female CD-1 mice: Low doses induce elevated serum leptin and insulin, and overweight in mid-life. *Molecular and cellular endocrinology*. 2009;304(1–2):97–105. [PubMed: 19433254]
55. Lv Z, Li G, Li Y, Ying C, Chen J, Chen T, et al. Glucose and lipid homeostasis in adult rat is impaired by early-life exposure to perfluorooctane sulfonate. *Environ Toxicol*. 2013;28(9):532–42. [PubMed: 23983163]
56. Kingsley SL, Kelsey KT, Butler R, Chen A, Eliot MN, Romano ME, et al. Maternal serum PFOA concentration and DNA methylation in cord blood: A pilot study. *Environmental research*. 2017;158:174–8. [PubMed: 28645023]
57. Ornoy A. Prenatal origin of obesity and their complications: Gestational diabetes, maternal overweight and the paradoxical effects of fetal growth restriction and macrosomia. *Reprod Toxicol*. 2011;32(2):205–12. [PubMed: 21620955]
58. Barker DJP. Developmental origins of chronic disease. *Public Health*. 2012;126(3):185–9. [PubMed: 22325676]
59. Andersen CS, Fei C, Gamborg M, Nohr EA, Sorensen TI, Olsen J. Prenatal exposures to perfluorinated chemicals and anthropometry at 7 years of age. *American journal of epidemiology*. 2013;178(6):921–7. [PubMed: 23825166]
60. Barry V, Darrow LA, Klein M, Winquist A, Steenland K. Early life perfluorooctanoic acid (PFOA) exposure and overweight and obesity risk in adulthood in a community with elevated exposure. *Environmental research*. 2014;132C:62–9.
61. Martinsson M, Nielsen C, Bjork J, Rylander L, Malmqvist E, Lindh C, et al. Intrauterine exposure to perfluorinated compounds and overweight at age 4: A case-control study. *PloS one*. 2020;15(3):e0230137. [PubMed: 32176721]
62. Manzano-Salgado CB, Casas M, Lopez-Espinosa MJ, Ballester F, Iniguez C, Martinez D, et al. Prenatal Exposure to Perfluoroalkyl Substances and Cardiometabolic Risk in Children from the Spanish INMA Birth Cohort Study. *Environmental health perspectives*. 2017;125(9):097018. [PubMed: 28934720]
63. Koustas E, Lam J, Sutton P, Johnson PI, Atchley DS, Sen S, et al. The Navigation Guide-Evidence-Based Medicine Meets Environmental Health: Systematic Review of Nonhuman Evidence for PFOA Effects on Fetal Growth. *Environmental health perspectives*. 2014.
64. Abbott BD. Review of the expression of peroxisome proliferator-activated receptors alpha (PPAR alpha), beta (PPAR beta), and gamma (PPAR gamma) in rodent and human development. *Reproductive toxicology (Elmsford, NY)*. 2009;27(3–4):246–57.
65. Abbott BD, Wolf CJ, Das KP, Zehr RD, Schmid JE, Lindstrom AB, et al. Developmental toxicity of perfluorooctane sulfonate (PFOS) is not dependent on expression of peroxisome proliferator activated receptor-alpha (PPAR alpha) in the mouse. *Reproductive toxicology (Elmsford, NY)*. 2009;27(3–4):258–65.
66. Kingsley SL, Eliot MN, Kelsey KT, Calafat AM, Ehrlich S, Lanphear BP, et al. Variability and predictors of serum perfluoroalkyl substance concentrations during pregnancy and early childhood. *Environmental research*. 2018;165:247–57. [PubMed: 29734025]
67. Verner MA, Luccisano AE, Morken NH, Yoon M, Wu H, McDougall R, et al. Associations of Perfluoroalkyl Substances (PFASs) with Lower Birth Weight: An Evaluation of Potential

- Confounding by Glomerular Filtration Rate Using a Physiologically Based Pharmacokinetic Model (PBPK). *Environmental health perspectives*. 2015.
68. Lazarevic N, Barnett AG, Sly PD, Knibbs LD. Statistical Methodology in Studies of Prenatal Exposure to Mixtures of Endocrine-Disrupting Chemicals: A Review of Existing Approaches and New Alternatives. *Environmental health perspectives*. 2019;127(2):26001. [PubMed: 30720337]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 1:

Adjusted mean child BMI as a function of age according to terciles of maternal PFOA (Panel A), PFOS (Panel B), PFNA (Panel C), and PFHxS (Panel D) concentrations during pregnancy among HOME Study mother-child dyads: Derived using linear mixed models with a spline for the age-BMI relation ($n=345$, 1,865 repeated BMI measures)^a
 a-Adjusted for maternal race (non-Hispanic White, non-Hispanic Black, and other), maternal age (continuous, years), maternal BMI (continuous, kg/m^2), parity (nulliparous, parous: 1, and parous: 2+), household income (continuous, USD/year), gestational tobacco smoke exposure (\log_{10} -transformed serum cotinine concentrations), and child sex.

*-Ranges of 1st, 2nd, and 3rd PFOA concentration terciles were 0.5–4.2, 4.3–6.5, and 6.7–26 ng/mL, respectively. Ranges of 1st, 2nd, and 3rd PFOS concentration terciles were 0.4–10, 11–16, and 16–57 ng/mL, respectively. Ranges of 1st, 2nd, and 3rd PFNA concentration terciles were 0.1–0.8, 0.9–1.0, and 1.1–2.9 ng/mL, respectively. Ranges of 1st, 2nd, and 3rd PFHxS concentration terciles were 0.1–1.0, 1.1–2.0, and 2.1–33 ng/mL, respectively.

**-Fringes along x-axis of each plot indicate age and number of observations

***- PFOA × age interaction p-value=0.03, PFOS × age interaction p-value=0.10, PFNA × age interaction p-value=0.58, and PFHxS × age interaction p-value=0.18.

Table 1:

Baseline sociodemographic and perinatal characteristics of HOME Study women and their infants (n=345)

Characteristic	N (%)
Maternal Race	
Non-Hispanic White	214 (62.0)
Non-Hispanic Black	109 (31.6)
Other	22 (6.4)
Maternal Age at Delivery (years)	
18 – <25	81 (23.5)
25 – <30 years	95 (27.5)
30 – <35 years	114 (33.0)
35 years	55 (15.9)
Household Income (\$ per year)	
<20,000	76 (22.0)
20,000 – <40,000	62 (18.0)
40,000 – <80,000	111 (32.2)
80,000	96 (27.8)
Maternal Education	
Less than high school	4 (1.2)
Some High School or High School	79 (22.9)
Some College or Technical School	89 (25.8)
Bachelor's Degree	173 (50.1)
Parity	
Nulliparous (0)	148 (42.9)
Parous: 1	108 (31.3)
Parous: 2+	89 (25.8)
Tobacco Smoke Exposure^a	
None	109 (31.6)
Secondhand	200 (58.0)
Active	36 (10.4)
Pre-pregnancy BMI	
Underweight/Lean (<25 kg/m ²)	147 (42.6)
Overweight (25-<30 kg/m ²)	115 (33.3)
Obese (>30 kg/m ²)	83 (4.1)
Infant Sex	
Female	184 (53.3)
Male	161 (46.7)

^a-Characterized by maternal serum cotinine concentrations during pregnancy. None: <0.015 ng/mL, secondhand: 0.015-<3 ng/mL, and active: 3 ng/mL.

Table 2:

Adjusted magnitude and age of infancy BMI zenith according to maternal PFAS concentration terciles during pregnancy among HOME Study mother-child dyads: Derived using linear mixed models with a spline for the age-BMI relation (n=345, 1,865 repeated BMI measures)^a

PFAS and Tercile (Range, ng/mL)	Mean BMI (kg/m ²)	BMI Difference (95% CI)	Mean Age (years)	Age Difference (95% CI)
PFOA-T1 (0.5–4.2)	17.2	Ref	1.25	Ref
PFOA-T2 (4.3–6.5)	17.3	0.05 (−0.31, 0.40)	1.27	0.02 (−0.10, 0.07)
PFOA-T3 (6.7–26)	16.7	−0.50 (−0.87, −0.15)	1.23	−0.02 (−0.11, 0.06)
PFOS-T1 (0.4–10)	17.3	Ref	1.27	Ref
PFOS-T2 (11–16)	17.0	−0.24 (−0.60, 0.10)	1.25	−0.02 (−0.06, 0.12)
PFOS-T3 (16–57)	16.9	−0.40 (−0.76, −0.06)	1.25	−0.02 (−0.10, 0.06)
PFNA-T1 (0.1–0.8)	17.2	Ref	1.24	Ref
PFNA-T2 (0.9–1.0)	16.9	−0.32 (−0.48, 0.21)	1.30	0.06 (−0.04, 0.18)
PFNA-T3 (1.1–2.9)	17.0	−0.16 (−0.48, 0.16)	1.25	0.01 (−0.07, 0.08)
PFHxS-T1 (0.1–1.0)	17.2	Ref	1.27	Ref
PFHxS-T2 (1.1–2.0)	17.2	0.06 (−0.29, 0.43)	1.28	0.01 (−0.07, 0.10)
PFHxS-T3 (2.1–33)	16.7	−0.42 (−0.80, −0.06)	1.22	−0.05 (−0.14, 0.03)

^aAdjusted for maternal race (non-Hispanic White, non-Hispanic Black, and other), maternal age (continuous, years), maternal BMI (continuous, kg/m²), parity (nulliparous, parous: 1, and parous: 2+), household income (continuous, USD/year), gestational tobacco smoke exposure (log₁₀-transformed serum cotinine concentrations), and child sex (male and female).

Table 3:

Adjusted magnitude and age of early childhood BMI nadir according to maternal PFAS concentration terciles during pregnancy among HOME Study mother-child dyads: Derived using linear mixed models with a spline for the age-BMI relation (n=345, 1,865 repeated BMI measures)^a

PFAS and Tercile (Range, ng/mL)	Mean BMI (kg/m ²)	BMI Difference (95% CI)	Mean Age (years)	Age Difference (95% CI)
PFOA-T1 (0.5–4.2)	16.1	Ref	4.65	Ref
PFOA-T2 (4.3–6.5)	16.2	0.11 (−0.40, 0.61)	4.33	−0.32 (−0.96, 0.27)
PFOA-T3 (6.7–26)	15.6	−0.52 (−1.04, 0.00)	4.36	−0.29 (−0.94, 0.32)
PFOS-T1 (0.4–10)	16.3	Ref	4.13	Ref
PFOS-T2 (11–16)	15.9	−0.43 (−0.92, 0.07)	4.71	0.58 (0.00, 1.20)
PFOS-T3 (16–57)	15.6	−0.67 (−1.02, −0.19)	4.51	0.38 (−0.17, 0.95)
PFNA-T1 (0.1–0.8)	16.0	Ref	4.51	Ref
PFNA-T2 (0.9–1.0)	16.0	0.00 (−0.54, 0.52)	4.44	−0.07 (−0.77, 0.62)
PFNA-T3 (1.1–2.9)	15.8	−0.17 (−0.64, 0.29)	4.37	−0.14 (−0.65, 0.38)
PFHxS-T1 (0.1–1.0)	16.2	Ref	4.35	Ref
PFHxS-T2 (1.1–2.0)	16.0	−0.12 (−0.64, 0.40)	4.72	0.37 (−0.28, 1.02)
PFHxS-T3 (2.1–33)	15.6	−0.60 (−1.09, −0.09)	4.31	−0.04 (−0.59, 0.51)

^aAdjusted for maternal race (non-Hispanic White, non-Hispanic Black, and other), maternal age (continuous, years), maternal BMI (continuous, kg/m²), parity (nulliparous, parous: 1, and parous: 2+), household income (continuous, USD/year), gestational tobacco smoke exposure (log₁₀-transformed serum cotinine concentrations), and child sex (male and female).

Table 4:

Adjusted rate of BMI change from age 8 to 12 years according to maternal PFAS concentration terciles during pregnancy among HOME Study mother-child dyads: Derived using linear mixed models with a spline for the age-BMI relation (n=345, 1,865 repeated BMI measures)^a

PFAS and Tercile (Range, ng/mL)	Mean Rate of BMI Change (kg/m ² /y)	Difference in BMI Rate (95% CI)
PFOA-T1 (0.5–4.2)	0.65	Ref
PFOA-T2 (4.3–6.5)	0.90	0.35 (0.11, 0.39)
PFOA-T3 (6.7–26)	0.74	0.09 (–0.05, 0.24)
PFOS-T1 (0.4–10)	0.79	Ref
PFOS-T2 (11–16)	0.82	0.03 (–0.12, 0.17)
PFOS-T3 (16–57)	0.70	–0.09 (–0.24, 0.04)
PFNA-T1 (0.1–0.8)	0.76	Ref
PFNA-T2 (0.9–1.0)	0.82	0.06 (–0.11, 0.22)
PFNA-T3 (1.1–2.9)	0.75	–0.01 (–0.15, 0.12)
PFHxS-T1 (0.1–1.0)	0.79	Ref
PFHxS-T2 (1.1–2.0)	0.73	–0.06 (–0.20, 0.09)
PFHxS-T3 (2.1–33)	0.78	–0.01 (–0.15, 0.13)

^aAdjusted for maternal race (non-Hispanic White, non-Hispanic Black, and other), maternal age (continuous, years), maternal BMI (continuous, kg/m²), parity (nulliparous, parous: 1, and parous: 2+), household income (continuous, USD/year), gestational tobacco smoke exposure (log₁₀-transformed serum cotinine concentrations), and child sex (male and female).

Table 5:

Adjusted mean BMI (kg/m^2) and difference in BMI (kg/m^2) from ages 4-weeks to 12-years according to maternal PFAS tercile during pregnancy among HOME Study mother-child dyads: Derived using linear mixed models with a spline for the age-BMI relation ($n=345, 1,865$ repeated BMI measures)^a

Age (years)	Tercile ^{b,c}	PFOA (95% CI)	PFOS (95% CI)	PFNA (95% CI)	PFHxS (95% CI)
0.07	T1	15.0 (14.8, 15.3)	14.9 (14.6, 15.2)	14.9 (14.7, 15.1)	14.9 (14.7, 15.2)
0.07	T2-T1	-0.30 (-0.66, 0.08)	-0.01 (-0.39, 0.38)	-0.17 (-0.56, -0.24)	-0.19 (-0.57, 0.18)
0.07	T3-T1	-0.58 (-0.95, -0.19)	-0.47 (-0.85, -0.08)	-0.37 (-0.72, -0.01)	-0.37 (-0.77, 0.02)
1	T1	17.1 (16.9, 17.4)	17.2 (16.9, 17.4)	17.1 (16.9, 17.3)	17.1 (16.8, 17.3)
1	T2-T1	0.03 (-0.34, 0.38)	-0.23 (-0.59, 0.12)	-0.34 (-0.71, 0.02)	-0.05 (-0.30, 0.41)
1	T3-T1	-0.50 (-0.85, -0.14)	-0.40 (-0.76, -0.04)	-0.17 (-0.50, 0.15)	-0.40 (-0.77, 0.04)
2	T1	16.9 (16.6, 17.1)	16.9 (16.7, 17.2)	16.8 (16.6, 17.0)	16.8 (16.6, 17.1)
2	T2-T1	0.05 (-0.30, 0.38)	-0.24 (-0.58, 0.10)	-0.20 (-0.55, 0.15)	0.05 (-0.29, 0.40)
2	T3-T1	-0.55 (-0.89, -0.20)	-0.46 (-0.80, -0.13)	-0.18 (-0.49, 0.13)	-0.51 (-0.86, -0.16)
3	T1	16.4 (16.1, 16.7)	16.5 (16.2, 16.7)	16.3 (16.0, 16.5)	16.4 (16.1, 16.7)
3	T2-T1	0.04 (-0.35, 0.42)	-0.28 (-0.67, 0.11)	-0.07 (-0.47, 0.34)	-0.01 (-0.40, 0.40)
3	T3-T1	-0.57 (-0.98, -0.18)	-0.55 (-0.93, -0.17)	-0.20 (-0.56, 0.16)	-0.58 (-0.98, -0.18)
4	T1	16.1 (15.8, 16.5)	16.3 (16.0, 16.6)	16.0 (15.7, 16.3)	16.2 (15.8, 16.5)
4	T2-T1	0.07 (-0.39, 0.53)	-0.37 (-0.84, 0.09)	0.00 (-0.49, 0.49)	-0.08 (-0.54, 0.40)
4	T3-T1	-0.55 (-1.03, -0.07)	-0.64 (-1.10, -0.19)	-0.18 (-0.62, 0.25)	-0.60 (-1.07, -0.12)
5	T1	16.1 (15.7, 16.5)	16.4 (16.0, 16.8)	16.0 (15.7, 16.4)	16.2 (15.8, 16.6)
5	T2-T1	0.15 (-0.40, 0.69)	-0.51 (-1.05, 0.03)	0.01 (-0.57, 0.58)	-0.16 (-0.70, 0.40)
5	T3-T1	-0.48 (-1.05, 0.07)	-0.73 (-1.27, -0.20)	-0.15 (-0.66, 0.36)	-0.58 (-1.14, -0.03)
8	T1	17.0 (16.4, 17.6)	17.8 (17.2, 18.4)	17.2 (16.6, 17.7)	17.4 (16.8, 18.0)
8	T2-T1	0.57 (-0.27, 1.43)	-0.94 (-1.80, -0.09)	-0.08 (-0.98, 0.84)	-0.41 (-1.26, 0.46)
8	T3-T1	-0.16 (-1.03, 0.70)	-1.02 (-1.87, -0.17)	-0.02 (-0.82, 0.78)	-0.45 (-1.32, 0.43)
12	T1	19.6 (18.7, 20.5)	21.0 (20.1, 21.9)	20.2 (19.4, 21.0)	20.6 (19.7, 21.5)
12	T2-T1	1.57 (0.34, 2.83)	-0.83 (-2.11, 0.41)	0.13 (-1.23, 1.50)	-0.65 (-1.90, 0.65)
12	T3-T1	0.23 (-1.06, 1.53)	-1.41 (-2.65, -0.14)	-0.07 (-1.26, 1.09)	-0.50 (-1.78, 0.76)

^a- Adjusted for maternal race (non-Hispanic White, non-Hispanic Black, and other), maternal age (continuous, years), maternal BMI (continuous, kg/m^2), parity (nulliparous, parous: 1, and parous: 2+), household income (continuous, USD/year), gestational tobacco smoke exposure (log₁₀-transformed serum cotinine concentrations), and child sex (male and female).

^b- Adjusted mean BMI is displayed for 1st tercile children with the average covariate pattern, otherwise difference in 2nd vs. 1st and 3rd vs. 1st tercile is shown.

^c- Ranges of 1st, 2nd, and 3rd PFOA concentration terciles were 0.5–4.2, 4.3–6.5, and 6.7–26 ng/mL, respectively. Ranges of 1st, 2nd, and 3rd PFOS concentration terciles were 0.4–10, 11–16, and 16–57 ng/mL, respectively. Ranges of 1st, 2nd, and 3rd PFNA concentration terciles were 0.1–0.8, 0.9–1.0, and 1.1–2.9 ng/mL, respectively. Ranges of 1st, 2nd, and 3rd PFHxS concentration terciles were 0.1–1.0, 1.1–2.0, and 2.1–33 ng/mL, respectively.