

Complete Genome Sequence of Bovine Polyomavirus Type 1 from Aborted Cattle, Isolated in Belgium in 2014

Steven Van Borm,^a Toon Rosseel,^{a*} Isabelle Behaeghel,^b Marc Saulmont,^c Laurent Delooz,^c Thierry Petitjean,^c Elisabeth Mathijs,^a Frank Vandebussche^a

Molecular Platform^a, Interactions and Surveillance^b, Veterinary and Agrochemical Research Center, Ukkel, Belgium; Regional Association for Animal Registration and Health (ARSIA) asbl, Ciney, Belgium^c

* Present address: Toon Rosseel, Centre for Medical Genetics, Ghent University Hospital, Ghent, Belgium.

S.V.B. and T.R. contributed equally to this work.

The complete and fully annotated genome sequence of a bovine polyomavirus type 1 (BPyV/BEL/1/2014) from aborted cattle was assembled from a metagenomics data set. The 4,697-bp circular dsDNA genome contains 6 protein-coding genes. Bovine polyomavirus is unlikely to be causally related to the abortion cases.

Received 30 November 2015 Accepted 15 January 2016 Published 3 March 2016

Citation Van Borm S, Rosseel T, Behaeghel I, Saulmont M, Delooz L, Petitjean T, Mathijs E, Vandebussche F. 2016. Complete genome sequence of bovine polyomavirus type 1 from aborted cattle, isolated in Belgium in 2014. *Genome Announc* 4(2):e01646-15. doi:10.1128/genomeA.01646-15.

Copyright © 2016 Van Borm et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Steven Van Borm, steven.vanborm@codac-cerva.be.

From July until November 2014, an increase in the number of bovine abortions showing atypical icteric appearance and splenomegaly was observed in Belgium (1). In addition to specific diagnostic efforts to diagnose pathogens known to be associated with bovine abortion, a viral metagenomics workflow was applied to samples (liver, spleen, placenta, abomasal fluid) from 5 aborted fetuses. Briefly, samples were homogenized in phosphate-buffered saline (10% wt/vol) and prepped with 0.4- μ M selective filtration and nuclease treatment, and DNA was extracted as previously described (2, 3). Sequencing libraries were prepared using the NexteraXT kit (Illumina) according to the manufacturer's instructions and quantified with the Kapa library quantification kit for Illumina platforms (Kapa Biosystems), and fragment length distribution was verified using the Bioanalyzer with the High Sensitivity DNA kit (Agilent Technologies). Sequencing was performed on a MiSeq sequencer using a MiSeq reagent kit version 3 (Illumina) with 2 \times 300-bp paired-end sequencing. Eighteen libraries were multiplexed using standard Illumina indexing primers.

The quality of the sequences was checked with the FastQC tool version 0.10.1 (<http://www.bioinformatics.babraham.ac.uk/projects/fastqc>). Stretches containing unidentified nucleotides ("N") were trimmed using Cutadapt version 1.3 (4) prior to quality-trimming using Sickle version 1.210 (Q score <30, length <50 bp) (5).

For the metagenomics analysis, rRNA reads and host reads were removed using the SortMeRNA tool version 1.7 (6) and mapping against the bovine reference genome (GenBank accession nos. AC_000158.1 to AC_000187.1, NC_006853.1) using BWA-MEM version 0.7.5a-r405 (7). Remaining reads were subjected to a megablast sequence alignment analysis against the NCBI nucleotide database using blast-2/2/27+ software (8), with an E value threshold of 0.001, followed by taxonomy assignment using MN version 4.70.4 (9, 10). Bovine polyomavirus type 1

(BPyV-1) sequences were present in all samples and could be confirmed in 5 out of 18 tested samples using real-time RT-PCR (11).

Members of the family *Polyomaviridae* are small, nonenveloped, double-stranded DNA viruses, which are widely distributed among vertebrates (12). They share a genome structure consisting of a 4.8- to 5.5-kbp circular double-stranded DNA. A BPyV-1 is unlikely to be causally involved in the abortion cases, as it is a known persistent contaminant of cattle that has been detected in bovine sera (13, 14), meat products (15, 16), and even waste and surface water (17–19).

A total of 50,230 quality-trimmed sequences from a highly positive liver sample were mapped against a reference genome (GenBank accession no. KM496323) using GS Mapper version 2.9 (Roche), resulting in a 4,697-nucleotide contig corresponding to the complete genome length. The protein-coding genes were predicted relative to reference sequence NC_001442 by GATU (20) and supplemented with manual annotation for spliced genes.

The complete genome of this bovine polyomavirus type 1 (BPyV/BEL/1/2014) is characterized as a double-stranded circular DNA comprising 4,697 bp, with an average GC content of 41.41%, and containing 6 protein-coding genes.

Nucleotide sequence accession number. The complete genome sequence of BPyV/BEL/1/2014 was deposited in DDBJ/EMBL/GenBank under the accession number [KU200259](https://www.ncbi.nlm.nih.gov/nuccore/KU200259).

ACKNOWLEDGMENTS

The expert technical assistance of Orkun Ozhelvacı is much appreciated. We acknowledge the collaborative effort of scientists from ARSIA asbl, DGZ vzw (Dierengezondheidszorg Vlaanderen), CODA-CERVA, and the Veterinary Faculties of the Universities of Liège, Gent, and Namur in investigating this syndrome.

This work was supported by an internal research grant of CODA-CERVA, European Union FP7 project RAPIDIA-FIELD (grant number FP7-289364), and Epi-SEQ, a transnational research project supported

under the second joint call for transnational research projects by EMIDA ERA-NET (FP7 project no. 219235).

FUNDING INFORMATION

This work was supported by an internal research grant of CODA-CERVA, European Union FP7 project RAPIDIA-FIELD (grant number FP7-289364), and Epi-SEQ, a transnational research project supported under the 2nd joint call for transnational research projects by EMIDA ERA-NET (FP7 project number 219235). The funders had no role in study design, data collection and interpretation, or the decision to submit the work for publication.

REFERENCES

1. Deloof L, Mori M, Petitjean T, Evrard J, Czaplicki G, Saegerman C. 2015. Congenital jaundice in bovine aborted foetuses: an emerging syndrome in southern Belgium. *Transbound Emerg Dis* 62:124–126. <http://dx.doi.org/10.1111/tbed.12326>.
2. Rosseel T, Pardon B, De Clercq K, Ozhelvaci O, Van Borm S. 2014. False-positive results in metagenomic virus discovery: a strong case for follow-up diagnosis. *Transbound Emerg Dis* 61:293–299. <http://dx.doi.org/10.1111/tbed.12251>.
3. Rosseel T, Ozhelvaci O, Freimanis G, Van Borm S. 2015. Evaluation of convenient pretreatment protocols for RNA virus metagenomics in serum and tissue samples. *J Virol Methods* 222:72–80. <http://dx.doi.org/10.1016/j.jviromet.2015.05.010>.
4. Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet J* 17:10–12. <http://dx.doi.org/10.14806/ej.17.1.200>.
5. Joshi NA, Fass J. 2011. Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ files. Version 1.33. <https://github.com/najoshi/sickle>.
6. Kopylova E, Noé L, Touzet H. 2012. SortMeRNA: fast and accurate filtering of ribosomal RNAs in metatranscriptomic data. *Bioinformatics* 28:3211–3217. <http://dx.doi.org/10.1093/bioinformatics/bts611>.
7. Li H. 2014. Toward better understanding of artifacts in variant calling from high-coverage samples. *Bioinformatics* 30:2843–2851. <http://dx.doi.org/10.1093/bioinformatics/btu356>.
8. Tao T. 2010. Standalone BLAST setup for Windows PC. In BLAST® help [Internet]. NCBI, Bethesda, MD. <http://www.ncbi.nlm.nih.gov/books/NBK52637>.
9. Huson DH, Auch AF, Qi J, Schuster SC. 2007. MegaN analysis of metagenomic data. *Genome Res* 17:377–386. <http://dx.doi.org/10.1101/gr.5969107>.
10. Huson DH, Mitra S. 2012. Introduction to the analysis of environmental sequences: metagenomics with MEGAN. *Methods Mol Biol* 856:415–429. http://dx.doi.org/10.1007/978-1-61779-585-5_17.
11. Hundesa A, Boffill-Mas S, Maluquer de Motes C, Rodriguez-Manzano J, Bach A, Casas M, Girones R. 2010. Development of a quantitative PCR assay for the quantitation of bovine polyomavirus as a microbial source-tracking tool. *J Virol Methods* 163:385–389. <http://dx.doi.org/10.1016/j.jviromet.2009.10.029>.
12. Pérez-Losada M, Christensen RG, McClellan DA, Adams BJ, Viscidi RP, Demma JC, Crandall KA. 2006. Comparing phylogenetic codivergence between polyomaviruses and their hosts. *J Virol* 80:5663–5669. <http://dx.doi.org/10.1128/JVI.00056-06>.
13. Van der Noordaa J, Sol CJ, Schuurman R. 1999. Bovine polyomavirus, a frequent contaminant of calf sera. *Dev Biol Stand* 99:45–47.
14. Wang J, Horner GW, O’Keefe JS. 2005. Detection and molecular characterisation of bovine polyomavirus in bovine sera in New Zealand. *NZ Vet J* 53:26–30.
15. Peretti A, FitzGerald PC, Bliskovsky V, Buck CB, Pastrana DV. 2015. Hamburger polyomaviruses. *J Gen Virol* 96:833–839. <http://dx.doi.org/10.1099/vir.0.000033>.
16. Zhang W, Li L, Deng X, Kapusinszky B, Delwart E. 2014. What is for dinner? Viral metagenomics of US store bought beef, pork, and chicken. *Virology* 468–470:303–310. <http://dx.doi.org/10.1016/j.virol.2014.08.025>.
17. Corsi SR, Borchardt MA, Spencer SK, Hughes PE, Baldwin AK. 2014. Human and bovine viruses in the Milwaukee River watershed: hydrologically relevant representation and relations with environmental variables. *Sci Total Environ* 490:849–860. <http://dx.doi.org/10.1016/j.scitotenv.2014.05.072>.
18. Lazic G, Grubac S, Lupulovic D, Bugarski D, Lazic S, Knezevic P, Petrovic T. 2015. Presence of human and animal viruses in surface waters in Vojvodina province of Serbia. *Food Environ Virol* 7:149–158. <http://dx.doi.org/10.1007/s12560-015-9187-3>.
19. Wong K, Xagorarakis I. 2011. Evaluating the prevalence and genetic diversity of adenovirus and polyomavirus in bovine waste for microbial source tracking. *Appl Microbiol Biotechnol* 90:1521–1526. <http://dx.doi.org/10.1007/s00253-011-3156-z>.
20. Tcherepanov V, Ehlers A, Upton C. 2006. Genome annotation transfer utility (GATU): rapid annotation of viral genomes using a closely related reference genome. *BMC Genomics* 7:150. <http://dx.doi.org/10.1186/1471-2164-7-150>.