

MEETING ABSTRACT

Open Access

Health outcomes, education, healthcare delivery and quality – 3048. From uterus to university: Recruitment and retention of a primary prevention birth cohort

Brenda Gerwing^{1*}, Rishma Chooniedass¹, Saiful Huq¹, Hao Huang², Anita Kozyrskyj³, Edmond Chan⁴, Clare Ramsey⁵, Moira Chan-Yeung⁶, Allan Becker¹

From 2nd WAO International Scientific Conference (WISC 2012)
Hyderabad, India. 6-9 December 2012

Background

It is important to identify predictors of retention in primary prevention studies as recruitment and retention are critical factors for a successful intervention study.

Methods

In 1994, the Canadian Asthma Primary Prevention Study (CAPPSS) was established. This high-risk birth cohort has 2 sites, Winnipeg and Vancouver, Canada. Expectant mothers were recruited during the third trimester. Enrollment criteria were a first degree relative with asthma or two first degree relatives with other allergic diseases. Participants were prenatally randomized into control and intervention groups. Intervention measures were introduced before birth and during baby's first year of life. Follow-up assessments by a Pediatric Allergist included skin prick testing (SPT) to common food and inhalants and pulmonary function testing.

Results

545 participants initially recruited. 266 randomized into control and 279 intervention. From recruitment to first year, 9.5% families (52) discontinued. At age 1, 493 infants were assessed; 52.3% males and 47.7% females, 49.1% control and 50.9% intervention. 76.8% high SES, 22.52% low SES. 9.7% maternal age ≤25 and 90.3%

maternal age >25. 17.6 % were diagnosed with asthma at 1 year. 22.1% with +SPT to food. 44.2% were 1stborn. Children were assessed at 2 (n=472, 95.7%) and 7 years (n=380, 77.1%). At 15 years, 326 (66.1%) participants returned; 55.8% males and 44.2% females (p=0.02), 44.5% control and 54.6% intervention (p=0.054). Maternal age >25 (OR=1.73, 95% CI 0.95-3.16, p=0.05), asthma diagnosis (OR=1.53, 95% CI 0.91-2.57, p=0.066), high SES (OR=1.37, 95% CI 0.88-2.11, p=0.1) and +SPT (OR=1.23, 95% CI 0.78-1.95, p=0.22) were all associated with higher rates of return. While 138 participants returned with no sibling(s) at enrollment (OR=0.81, 95% CI 0.56-1.18, p=0.16).

Conclusions

Participants with sibling(s) at birth had no significant difference in retention. Maternal age was the most likely predictor of participant drop out. Female participants, low SES, negative skin prick test to food and no asthma diagnosis at age 1 showed a trend towards drop out. When establishing future asthma and allergy cohorts, specific retention strategies should be considered for groups identified at risk for drop out, especially for younger mothers and female participants.

Author details

¹Pediatrics and Child Health, Section of Allergy, University of Manitoba, Canada. ²University of Manitoba, Winnipeg, MB, Canada. ³Pediatrics, University of Alberta, Edmonton, AB, Canada. ⁴Pediatrics, University of British Columbia, Vancouver, BC, Canada. ⁵Internal Medicine, University of Manitoba, Winnipeg, MB, Canada. ⁶Internal Medicine, University of British Columbia, Vancouver, BC, Canada.

¹Pediatrics and Child Health, Section of Allergy, University of Manitoba, Canada

Full list of author information is available at the end of the article

Published: 23 April 2013

doi:10.1186/1939-4551-6-S1-P218

Cite this article as: Gerwing et al.: Health outcomes, education, healthcare delivery and quality – 3048. From uterus to university: Recruitment and retention of a primary prevention birth cohort. *World Allergy Organization Journal* 2013 **6**(Suppl 1):P218.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

