

Resveratrol induces mitochondria-mediated, caspase-independent apoptosis in murine prostate cancer cells

Sanjay Kumar¹, Erdal Eroglu², James A. Stokes III¹, Karyn Scisum-Gunn¹, Sabita N. Saldanha¹, Udai P. Singh³, Upender Manne⁴, Selvarangan Ponnazhagan⁴, Manoj K. Mishra¹

¹Cancer Biology Research and Training, Department of Biological Sciences, Alabama State University, Montgomery, AL, USA

²Faculty of Engineering, Bioengineering Department, Celal Bayar University, Muradiye, Manisa, Turkey

³Department of Pathology, Microbiology and Immunology, University of South Carolina, Columbia, SC, USA

⁴Department of Pathology, School of Medicine, University of Alabama at Birmingham, Birmingham, AL, USA

Correspondence to: Manoj K. Mishra, email: mmishra@alasu.edu

Keywords: apoptosis, mitochondria, prostate cancer cells, resveratrol

Received: March 06, 2016

Accepted: January 10, 2017

Published: February 01, 2017

ABSTRACT

Found in the skins of red fruits, including grapes, resveratrol (RES) is a polyphenolic compound with cancer chemopreventive activity. Because of this activity, it has gained interest for scientific investigations. RES inhibits tumor growth and progression by targeting mitochondria-dependent or -independent pathways. However, further investigations are needed to explore the underlying mechanisms.

The present study is focused on examining the role of RES-induced, mitochondria-mediated, caspase-independent apoptosis of prostate cancer cells, namely transgenic adenocarcinoma of mouse prostate (TRAMP) cells. These cells were exposed to RES for various times, and cell killing, cell morphology, mitochondrial membrane potential ($\Delta\psi_m$), expression of Bax and Bcl2 proteins, the role of caspase-3, and DNA fragmentation were analyzed.

TRAMP cells exposed to RES showed decreased cell viability, altered cell morphology, and disrupted $\Delta\psi_m$, which led to aberrant expression of Bax and Bcl2 proteins. Furthermore, since the caspase-3 inhibitor, z-VAD-fmk (benzyloxycarbonyl-valine-alanine-aspartic acid-fluoromethyl ketone), had no appreciable impact on RES-induced cell killing, the killing was evidently caspase-independent. In addition, RES treatment of TRAMP-C1, TRAMP-C2, and TRAMP-C3 cells caused an appreciable breakage of genomic DNA into low-molecular-weight fragments.

These findings show that, in inhibition of proliferation of TRAMP cells, RES induces mitochondria-mediated, caspase-independent apoptosis. Therefore, RES may be utilized as a therapeutic agent to control the proliferation and growth of cancer cells.

INTRODUCTION

In Western populations, prostate cancer is the second leading cause of death (after heart disease) in men older than 65 years of age [1–6]. It arises through the change of pre-neoplastic lesions into adenocarcinomas, and thereafter progresses to metastatic disease [6–9]. Recent advances have found genetic alterations that enhance the probability of prostate cancer development [10, 11]. To limit the growth of prostate cancer, high doses

of chemotherapeutic drugs and high-frequency radiation have been used, but the limited efficacy and side effects of these treatments raise a concern [12, 13]. Therefore, it is desirable to find anti-cancer agents that are non-toxic and effective in inducing apoptosis in cancer cells. Previous studies have demonstrated that treatment with anti-androgens is beneficial in the early stages of prostate cancer development, suggesting that their growth may be dependent on androgens [14–17]. In contrast, androgen-independent prostate cancer cells do not undergo apoptosis

[14–17]. In view of these results, advanced therapies that target the proliferation of both androgen-dependent and -independent prostate cancer cells are needed.

Compounds naturally occurring and present in the human diet are generally nontoxic, and some have beneficial effects on human health. Among these dietary components, resveratrol (RES), a polyphenol found in the skin of red fruits, exhibits anti-cancer, anti-proliferative, anti-inflammatory, and anti-oxidative effects [18, 19]. The anti-cancer properties of RES are facilitated through various changes in apoptotic signaling, metabolic pathways, and other signaling pathways that regulate apoptosis, cell cycle progression, inflammation, proliferation, metastasis, and angiogenesis [18, 20]. Furthermore, cells exposed to RES show inhibition of PI3K/AKT signaling, which stimulates apoptosis [21–29]. RES induces the death receptor (TRAIL/DR4 and TRAIL-R2/DR5)-mediated apoptosis [7, 30–39], and it is involved in mitochondrial-mediated apoptosis [40–45]. Mitochondria are the primary provider of ATP in most mammalian cells, they regulate both necrotic and apoptotic cell death pathways [46]. Thus, apoptosis is considered to be the most likely mechanism adopted by cells after activation of death signals. Apoptosis may be triggered intrinsically or extrinsically, depending on the type of apoptotic signals [40, 41, 43–45].

In human lung adenocarcinoma cells, RES activates the intrinsic apoptotic pathway by inducing release of apoptosis-inducing factor (AIF) [47]. Intrinsic signals of apoptosis function mainly through mitochondria [48]. In healthy cells, the outer mitochondrial membrane expresses the B-cell lymphoma-2 (Bcl-2) family of proteins, which controls the release of pro-apoptotic factors from the inner-membrane space in mitochondria [49–52]. In response to internal damage to the cells, a Bcl-2 associated protein, Bax, migrates to the mitochondrial membrane and inhibits the action of Bcl-2, causing damage to the mitochondrial membrane that in turn releases cytochrome-c [49–52]. Cytochrome-c binds with the apoptotic protease activator factor-1 (Apaf-1) and forms a multimeric protein structure called the “apoptosome.” The apoptosome activates caspase-9, which triggers the activation of caspase-3 and caspase-7 [53–59]. Their activation initiates proteolytic activity that leads to cell death [53–59]. The extrinsic pathway of apoptosis, however, is triggered by external signals that stimulate death receptors, such as ligands Fas-L and TNF- α (tumor necrosis factor- α), which activate caspase-8 [60]. This activated molecule initiates a cascade of caspase activity, which facilitates cell death [60].

RES shows anti-cancer, anti-proliferative, anti-inflammatory, and anti-oxidative properties, which are involved in the mitochondrial pathway of apoptosis [18, 20, 40–44, 61, 62]. However, how RES-induced, mitochondria-mediated, caspase-independent apoptosis operates in controlling the progression of tumor cells is not clear.

In the present effort, we examined the effects of RES on mitochondria-mediated, caspase-independent apoptosis in transgenic adenocarcinoma of mouse prostate (TRAMP-C1, TRAMP-C2, and TRAMP-C3) cells. TRAMP cells exposed to RES showed, in a time- and dose-dependent manner, increased cell killing and altered cell morphology. Furthermore, RES treatment resulted in disrupted mitochondrial membrane potential ($\Delta\psi_m$), which triggered disproportionate expression of Bax and Bcl2 proteins. In addition, RES treatment did not induce marked fragmentation of DNA into low-molecular-weight segments. As determined by exposure of cells to the caspase-3 inhibitor, z-VAD-fmk, caspase-3 was not involved in RES-mediated cell killing. Thus, these findings indicate that RES induces mitochondria-mediated, caspase-independent apoptosis and delays proliferation of prostate cancer cells. Therefore, RES may be an agent for treatment of prostate cancer.

RESULTS

RES kills tumor cells

To test the effect of RES on TRAMP cells, a cell-killing assay was performed. First, we determined the optimal time and concentration of RES needed to kill TRAMP cells. Annexin V-FITC⁺ and double positive (FITC⁺, PI⁺) cells showed sign of early and late apoptosis respectively; however, PI⁺ cells considered as dead cells as shown in representative Figure 1A. We established that 16 hours was the optimal time for maximum killing (Figure 1B). These experiments were repeated 5 times independently in triplicates. Data are represented as mean values \pm SEM (Standard Error of the Mean) in Figure 1B. We also conducted experiments at 24 hours and 48 hours but did not find any significant difference in RES-mediated cell killing (data not shown). Cells incubated with 50 μ M or 100 μ M of RES showed a concentration-dependent increase in the percent of cells killed (Figure 1B). Further analysis revealed that RES (100 μ M) treatment resulted in a significantly greater (* P <0.001) killing of TRAMP-C3 cells (43 \pm 5%) as compared to TRAMP-C1 (21 \pm 5%) and TRAMP-C2 (6 \pm 5%) cells (Figure 1B). In addition, TRAMP cells were incubated with RES in the presence or absence of Nec-1, a necroptosis blocker, to confirm whether RES-mediate apoptosis or necroptosis. We found that RES exhibited a similar pattern of cell killing in the presence or absence of Nec-1 (Supplementary Figure 1A; TRAMP-C1, Supplementary Figure 1B; TRAMP-C2, and Supplementary Figure 1C; TRAMP-C3).

RES treatment alters cell morphology

To evaluate the impact of RES on cell morphology, phase contrast microscopy was conducted. Cells were treated with 100 μ M of RES for 16h exhibited altered

cell morphology in a concentration-dependent manner (Figure 1C). Furthermore, cells exposed to 100 μM of RES showed more prominent morphological alterations in TRAMP as compared to cells treated with 50 μM of RES (Figure 1C). Additionally, TRAMP-C3 cells were more sensitive to 100 μM of RES as compared to TRAMP-C1 and TRAMP-C2 cells (Figure 1C). TRAMP-C3 cells showed more oval shapes as compared to TRAMP-C1 and TRAMP-C2 cells, suggesting loss of adherence and loss of cell-to-cell contact.

RES induces mitochondrial membrane potential

To examine the effect of RES on mitochondria, the $\Delta\psi\text{m}$ was measured using fluorescence microscopy and flow cytometry. TRAMP cells treated with RES (50 or 100 μM) showed disrupted $\Delta\psi\text{m}$ as compared to appropriate control (Figure 2A). Observed under a fluorescence microscope, DePsipher-stained TRAMP cells exhibited a distinct fluorescence color: red, green, or an overlap of green and red that results in orange/yellow. Cells with red fluorescence were considered to be healthy and normal; cells with green fluorescence were considered to have disrupted $\Delta\psi\text{m}$, indicating apoptosis. Cells with orange/yellow fluorescence were considered to have collapsed mitochondria. Most of the treated TRAMP cells showed green and orange/yellowish color, indicating that these cells

had disrupted $\Delta\psi\text{m}$ (Figure 2A); however, TRAMP-C3 cells showed significantly disrupted $\Delta\psi\text{m}$ as compared to TRAMP-C1 and TRAMP-C2 cells (Figure 2A).

Further, the $\Delta\psi\text{m}$ was validated by flow cytometric analysis of TRAMP cells treated with 100 μM RES as demonstrated in representative Figure 2B. In this figure, three cell populations were evident: (i) TRAMP cells showing only green fluorescence (FL1: 488/530 nm) corresponding to those with disrupted (low) $\Delta\psi\text{m}$ following apoptosis as compared to control cells. (ii) Cells with different intensities of green and red or yellow/orange were also considered to have intermediate disrupted $\Delta\psi\text{m}$. (iii) Cells emitting red fluorescence (FL2: 488/585nm) were considered to demonstrate high $\Delta\psi\text{m}$. Further analysis revealed that, with exposure to 100 μM RES, the percentages of TRAMP-C3 cells showing green fluorescence, that is, with disrupted $\Delta\psi\text{m}$, were higher relative to TRAMP-C1 and TRAMP-C2 cells (Figure 2C). This experiment was performed 5 times independently in triplicates; the sum of all experimental data (\pm SEM) is shown in the histogram Figure 2C (* p <0.05 and ** p <0.01).

RES modulates the expression of Bax and Bcl2

Western blots were performed to investigate the effect of RES on the expression of Bax and Bcl2




Figure 1: RES kills TRAMP cells in a dose- and time-dependent manner. TRAMP cells were treated with RES (50 μM or 100 μM), and cell killing and cell morphology were examined. **A.** Representative figure of gating strategies to study percent cell killing after RES treatment using a flow cytometer. **B.** Mean average values with \pm SEM of cell death at 0, 2, 4, 8, 12, and 16 hours. **C.** Morphological changes in cells due to RES treatment (* indicates p <0.05 and ** indicates p <0.01).

proteins. It was found that RES treatment modulated the expression of Bax and Bcl2 proteins in TRAMP-C1, TRAMP-C2 and TRAMP-C3 cells as compared to the control (Figure 3A). Further, densitometric analysis revealed that treatment with 50 μ M or 100 μ M of RES resulted in significantly high expression of Bax in TRAMP-C1 (* p <0.02), TRAMP-C2 (* p <0.001) and TRAMP-C3 (* p <0.001) cells when compared to the respective control (Figure 3C). In contrast to Bax, Bcl2 expression repressed significantly in TRAMP-C2 (* p <0.03) and TRAMP-C3 (* p <0.02) cells after 100 μ M of RES treatment (Figure 3B). In addition, TRAMP-C1 (# p >0.05) cells did not show a significant difference in the expression of Bcl2 protein in comparison to the control (Figure 3B).

RES induces caspase-independent cell killing

To evaluate the role of caspase-3 in RES-mediated cell killing, caspase-3 activation was blocked with an inhibitor (z-VAD-fmk). RES treatment with and without z-VAD-fmk induced RES-mediated cell killing (Figure 4A). TRAMP-C3 cells exposed to 100 μ M of RES with or without z-VAD-fmk showed significantly (** p <0.001) higher percentage of cell death as compared to TRAMP-C1 (* p <0.05) and TRAMP-C2 (* p <0.05) cells (Figure 4A). However, there was no significant difference in TRAMP cells treated with either RES or with RES plus z-VAD-fmk (Figure 4A). There were corresponding results when cells were analyzed morphologically under a phase contrast microscope (Figure 4B: resultant morphology of TRAMP cell after 100 μ M of RES+z-VAD-fmk treatment). In addition,




Figure 2: RES disrupts $\Delta\psi$ m. Cells were treated with 100 μ M of RES for 16 hours, and $\Delta\psi$ m was examined by using DePsipher dye. **A.** Fluorescence microscopy of TRAMP cells. **B.** Representative gating of cells. (i) cells with disrupted (low) $\Delta\psi$ m are indicated as DePsipher-monomers (34.52%), (ii) cells with intermediate $\Delta\psi$ m (4.26%), and (iii) cell with high $\Delta\psi$ m indicated as DePsipher-aggregates (11.89%). **C.** Percent of cells with disrupted $\Delta\psi$ m. Data represented as mean values \pm SEM (* indicates p <0.05 and ** indicates p <0.01).

treatment with Nec-1 (an inhibitor of necroptosis) did not change RES-mediated cell killing in the presence or in the absence of z-VAD-fmk (Supplementary Figure 2). Thus, in TRAMP-C1, TRAMP-C2, and TRAMP-C3 cells, the caspase-3 inhibitor, z-VAD-fmk, and Nec-1, had negligible effects on RES-mediated apoptosis (Supplementary Figure 2A; TRAMP-C1, Supplementary Figure 2B; TRAMP-C2, and Supplementary Figure 2C; TRAMP-C3 cells).

RES modulates the γ -H2A.X expression

To test the effect of 50 and 100 μ M of RES treatment on the expression of γ -H2A.X in TRAMP cells. The expression of γ -H2A.X was examined using western blot analysis which demonstrated significantly ($*p < 0.05$) higher expression in TRAMP-C2 and TRAMP-C3 cells when compared to the control (Figure 5A and 5B). However, in TRAMP-C1 cells, γ -H2A.X expression was not significant ($\#p > 0.05$) as compared to the control (Figure 5A and 5B). These findings suggest that RES

treatment sensitizes DNA damage which further leads to apoptosis of TRAMP cells.

DISCUSSION

Resistance to anti-cancer therapies is facilitated through an array of mechanisms that vary across tumor types [63, 64]. In the present study, we demonstrated that RES modulates mitochondria-mediated, caspase-independent apoptosis in murine prostate cancer cells. These results reveal that dietary compounds such as RES may play a critical role in inducing mitochondria-dependent apoptotic pathway(s) in murine prostate cancer cells (Figure 6).

Dietary compounds can be used to target mitochondria-mediated, caspase-independent apoptosis. This pathway is characterized by changes in $\Delta\psi_m$ and by maintenance of an optimal ratio of Bcl2/Bax [65]. A high $\Delta\psi_m$ and a high Bcl2/Bax ratio are believed to promote cell proliferation and enhanced cell survival,




Figure 3: RES modulates the expression of Bax and Bcl2. Results indicated altered expression of Bax and Bcl2 in TRAMP cells after Res treatment. **A.** Representative western blot of Bax, Bcl-2 and β -actin for the respective doses of RES in TRAMP-C1, TRAMP-C2 and TRAMP-C3 cells. **B.** Expression of Bcl-2 relative to the control. **C.** Expression of Bax relative to the control. Results are representative of three independent experiments ($*$ indicates $p < 0.05$ and $\#$ indicates $p > 0.05$).

thereby contributing to cancer progression. Dietary compounds such as RES inhibit cell growth in several types of human cancers, including prostate cancers [8, 66–70]. Consistent with results of previous studies, our data demonstrate that RES treatment results in enhanced cell killing in a time and dose-dependent manner. As cell viability decreases, altered cell morphology, a characteristic of apoptosis, increases [71]. RES induces cellular morphological changes similar to those caused by other anti-cancer drugs [40–45, 61, 62, 72]. To confirm our preliminary results relating to viability, we determined the effect of RES on the expression of Bcl2 and Bax proteins, $\Delta\psi_m$, caspase-3 activity, and DNA fragmentation. TRAMP-C3 and TRAMP-C2 cells exhibited significant difference in Bax expression as compared to control at 50 μM and 100 μM of RES; however, TRAMP-C1 showed significant difference at 100 μM of RES, not at 50 μM when compared to the control (Figure 3C). In contrast to Bax, Bcl2 expression in TRAMP-C1, TRAMP-C2 and TRAMP-C3 was

observed altered as compared to control at both 50 μM and 100 μM of RES (Figure 3B). Additionally, Bcl2 was found significantly repressed at 50 μM and 100 μM of RES in TRAMP-C2 and TRAMP-C3 when compared to control (Figure 3B). Furthermore, RES treatment also caused disrupted $\Delta\psi_m$, which is associated with apoptosis [73]. Most TRAMP-C3 cells showed decreased $\Delta\psi_m$ relative to TRAMP-C1 and TRAMP-C2 cells. Thus, RES treatment induced killing of TRAMP cells in a caspase-3-independent manner, for the caspase-3 inhibitor (z-VAD-fmk), failed to prevent RES-mediated cell killing. Therefore, apoptosis of TRAMP cells induced by RES apparently acts through the mitochondrial mediated, caspase-independent pathway (Figure 6).

Changes in $\Delta\psi_m$ are evident in mitochondria-mediated, caspase-independent apoptosis [74, 75]. As shown in the present effort, RES treatment decreases $\Delta\psi_m$ in TRAMP-C3 cells compared to TRAMP-C1 and TRAMP-C2 cells. However, RES induces apoptosis of colon cancer cells independently of the tumor suppressor




Figure 4: RES causes caspase-independent cell killing. Cells were treated with 100 μM of RES with or without 10 μM of z-VAD-fmk (a broad-spectrum caspase-3 inhibitor) and incubated for 16 hours. Thereafter, annexin V-PI staining was accomplished. **A.** The percent of cell killing by RES in the presence or absence of 10 μM of z-VAD-fmk. **B.** Morphological alterations in TRAMP cells caused by 100 μM of RES treatment in the presence of z-VAD-fmk. Data indicated mean values of \pm SEM (* indicates $p < 0.05$ and ** indicates $p < 0.001$).

p53 via epithelial differentiation and mitochondrial membrane collapse [76, 77]. The present data correspond with these findings, which suggest that RES treatment induces a collapse of $\Delta\psi_m$ [76, 77].

Proteins of the Bcl2 family, particularly Bcl2 and Bax, are involved in mitochondria-mediated apoptotic pathways [65]. After treatment of HCT-116 colon carcinoma cells with RES, Bax is involved in alteration of




Figure 5: RES treatment induced DNA fragmentation in TRAMP cells. TRAMP cells were incubated with 50 μ M and 100 μ M of RES to examine the expression of γ -H2A.X in TRAMP cells. The resulting western blots of γ -H2A.X showed that the expression of γ -H2A.X was found significantly higher in TRAMP-C2 and TRAMP-C3 cells when compared to the control. Furthermore, TRAMP-C1 cells, did not show a significant difference in γ -H2A.X expression when compared to the control (* indicates p<0.05 and # indicates p>0.05).




Figure 6: RES induces apoptosis in TRAMP cells. A schematic representation of the mechanism of RES-mediated, caspase-independent apoptosis in TRAMP cells.

mitochondrial membrane permeability [78–80]. However, by up-regulation of Bcl2 and inhibition of p53 and Bax, RES reverses cadmium chloride-induced testicular damage and subfertility [79]. In bladder cancer cells, RES also induces apoptosis by down-regulating the expression of Bcl2 proteins [81, 82]. However, the current results show that RES modulates the expression of Bax and Bcl2 in TRAMP-C1, TRAMP-C2, and TRAMP-C3 cells compared to the control cells; however, in TRAMP-C1 cells Bax at 50 μ M of RES and Bcl2 at 50 μ M and 100 μ M of RES were found insignificant. These findings corroborated with other reports [65, 81, 82].

RES decreased cell viability and $\Delta\psi_m$, and modulated the expression of Bax and Bcl2 proteins in TRAMP-C1, TRAMP-C2, and TRAMP-C3 cells. In an investigation of the involvement of caspase-3 in RES-mediated cell killing and DNA fragmentation, we found that RES treatment with or without the caspase-3 inhibitor, z-VAD-fmk, resulted in similar cell killing. This result was supported by morphological examination of the cells after DePsipher staining. The mechanism of action, however, is not yet defined. RES induces apoptosis by depolarizing the mitochondrial membranes in a caspase-independent manner [73] [48, 83, 84]. Nevertheless, in U937 cells, overexpression of Bcl2 attenuates RES-mediated apoptosis by blocking caspase-3 activation [53]. Moreover, RES induces caspase-dependent and -independent apoptosis in various cancer cells [85–87]. In colon cancer cells, RES induces caspase-2 activation that subsequently triggers Bax-Bak-dependent and -independent cell death [79]. In human lung adenocarcinoma cells, RES stimulates mitochondria-mediated and caspase-dependent cell death [88]. Conversely, in primary mouse fibroblasts, RES exhibits a cytoprotective effect by acting against caspase-3 [89]. The present results show that, for TRAMP cells, the caspase-3 inhibitor, z-VAD-fmk, and Nec-1, which blocks necroptosis, had a negligible effect on RES-mediated cell killing. Thus, in these cells, RES induces caspase-independent apoptosis.

The present results show that RES treatment to TRAMP cells caused significant cleavage of genomic DNA, which was accomplished by the expression of γ -H2A.X, an evolutionary conserved variant of histone H2A, has been identified as one of the key histones to undergo various post-translational modification in response to double stranded DNA breaks [90, 91]. DNA damage caused by radiation, UV light, or anti-cancer agents results in phosphorylation of Histone γ -H2A.X at ser-139 by PI3K-like kinases, including ATM, ATR, and DNA-PK [92–94]. The DNA damage response during DNA fragmentation is required for DNA-damage response proteins including DNA-PK that phosphorylates γ -H2A.X [95, 96]. Phosphorylation of γ -H2A.X at Tyr142 inhibits the recruitment of DNA repair proteins and promotes binding of pro-apoptotic factors such as JNK1 [97, 98]. Thus, γ -H2A.X expression was significantly higher in

TRAMP-C2 and TRAMP-C3 cells after 50 and 100 μ M of RES treatment as compared to the control. However, in TRAMP-C1 cells, γ -H2A.X expression was not significant as compared to the control. These findings are consistent with the previous results, which show RES-mediated caspase-independent cell killing. Furthermore, previous studies have demonstrated that RES induces apoptosis and DNA fragmentation in several types of cancer cells [99–102]. These properties of RES suggest that it could be used as a therapeutic agent to treat prostate cancer.

Our results demonstrate that, for TRAMP-C1, TRAMP-C2, and TRAMP-C3 cells, RES increases cell killing in a dose- and time-dependent manner, induces morphological alterations, and triggers apoptosis. In these cells, RES causes a disrupted $\Delta\psi_m$ that leads to modulated expression of Bax and Bcl2 proteins. Additionally, caspase-3 is not involved in RES-mediated cell killing, showing that, in TRAMP cells, RES induces caspase-independent apoptosis. In these cells, RES treatment contributed to DNA fragmentation which enhanced γ -H2A.x expression in treated TRAMP-C2 and TRAMP-C3 cells, but not in TRAMP-C1 when compared to the control, indicating the sign of DNA damage after RES treatment [92]. Therefore, RES may be a promising dietary compound for the treatment of prostate cancer. However, further investigations are necessary to uncover the underlying mechanisms.

MATERIALS AND METHODS

Cell lines and culture conditions

TRAMP-C1, TRAMP-C2, and TRAMP-C3 cells were obtained from American Type Culture Collection (www.ATCC.org) and were maintained at 37^o C under 5% CO₂ in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% fetal bovine serum (FBS) (v/v), bovine insulin (0.005 mg/ml), dehydroisoandrosterone (10 nM), and antibiotics/antimycotics (1%).

Cell killing assay

To assess percentages of cell death, 70-80% confluent cells were harvested by trypsinization, counted, and seeded (10⁵) in 24-well plates in 1 ml of culture medium. Cells were treated with 50 μ M or 100 μ M of RES and were analyzed by flow cytometry at 0, 2, 4, 8, 12, and 16 hours. They were stained with annexin V-propidium iodide (PI) as directed by manufacturer. Briefly, into each tube, annexin V (5 μ l of 600 μ g/ml) and PI (5 μ l of 30 μ g/ml) were added, and tubes were then incubated for 15 min at 4^oC. Cells were washed with Dulbecco's phosphate buffered saline (DPBS) and centrifuged. For flow cytometric analysis, the cells were suspended in 200-500 μ l of annexin V binding buffer. Data were acquired by use of a 13-color flow cytometer (Novocyte,

Acea Biosciences, San Diego, CA). PI⁺, Annexin V⁺, and Annexin V⁺ PI⁺ cells were counted as dead populations; PI/Annexin V FITC⁻ cells were counted as live cells. All experiments were performed in triplicate.

Assessment of cell morphology

Cells were exposed to 50 μ M or 100 μ M of RES for 16 hours. Media were removed, and after media removal, cells were washed with DPBS and then suspended in 50 μ l of DPBS. To assess morphological changes, cells were observed under a phase-contrast microscope (Life Tech, Grand Island, NY).

Assessment of mitochondrial membrane potential

To examine $\Delta\psi_m$, cells in DMEM supplemented with 10% FBS were exposed to RES (50 μ M or 100 μ M) for 16 hours. After incubation, cells were harvested, washed, stained with the mitochondria-specific dye, DePsipher (Trevigen, Gaithersburg, MD) and flow analyzed as suggested by the manufacturer. DePsipher, a cationic dye (5,5',6,6'-tetrachloro-1,1',3,3'-tetraethylbenzimidazolylcarbocyanine iodide) stains both healthy cells and cells with disrupted $\Delta\psi_m$. The dye enters into healthy mitochondria and, in its multimeric form, fluoresces red. However, in apoptotic cells, the dye remains in the cytoplasm and fluoresces green while in its monomeric form. Thus, cells with disrupted mitochondria can be differentiated from healthy cells. Cells were observed under a fluorescent microscope for morphology (Nikon ECLIPSE Ti, Melville, NY).

Western blotting

Total protein was extracted from TRAMP cells (10⁶) treated with RES (50 μ M or 100 μ M) by use of 2x radioimmunoprecipitation assay buffer (RIPA). The concentrations of total proteins in lysates were estimated according to Bradford et al. [57], using bovine serum albumin as the standard. Estimated concentration of total proteins (30-40 μ g/well) was run on 12% SDS-PAGE. Protein complexes were then transferred to nitrocellulose membranes, which were blocked in 5% skimmed milk and then incubated overnight with mouse anti-Bax (1:500) or anti-Bcl2 (1:100) monoclonal antibodies (Trevigen, Gaithersburg, MD and Thermo Scientific, NY, respectively). After repeated washing, the membranes were treated with a goat anti-mouse secondary antibody (1:1000) for 1 hour at room temperature. Proteins on the membranes were detected using an ECL-liquid substrate system (BioRad, Hercules CA). As an internal control, β -actin antibody (Grand Island, NY) was used to measure β -actin.

Next, in a separate experiment, estimated concentration of total proteins (20-40 μ g/well) were

electrophoresed on 10% SDS-polyacrylamide gels. Protein complexes were transferred on nitrocellulose membranes (cat#162-0112: BioRad, CA, USA) and incubated with rabbit anti- γ -H2A.X polyclonal antibody (cat#2595: Cell signaling technology, MA, USA). Membranes were washed and incubated with goat anti-rabbit secondary antibody (cat#31460: ThermoScientific, NY, USA). Protein blots were visualized using super signal west femto ECL western blotting detection system (cat#34095: ThermoScientific, NY, USA), equal amount of proteins loading were tested by reprobing with anti-b-actin antibody (cat#3700S: Cell signaling technology, MA, USA).

Assessment of caspase-independent cell death

To determine the role of caspases in RES-mediated cell death, cells were exposed to RES (100 μ M) with or without 10 μ M of z-VAD-fmk (a broad-spectrum caspase inhibitor) (Thermo Fisher Scientific, Grand Island, NY) for 16 hours. After incubation, caspase-independent cell death was determined. To accomplish this, cells were stained with annexin V (5 μ l of 600 μ g/ml) and PI (5 μ l of 30 μ g/ml) for 15 min at 4°C. Cells were washed with DPBS and suspended in 200-500 μ l of annexin V buffer for flow cytometric analysis. Data were acquired by use of a 13-color flow cytometer (Novocyte, Acea Biosciences, San Diego, CA).

Statistical analysis

Statistical significance of data was determined using Student's *t* test to determine the *p* value. For comparison of differences among the groups, single factor or multifactor one-way analysis of variance (ANOVA) followed by post hoc Bonferroni and Tukey test was used. Data were considered statistically significant at value *p*<0.05.

ACKNOWLEDGMENTS

We thank Dr. Donald Hill for his critical review of the manuscript.

CONFLICTS OF INTEREST

There is no conflict of interest among the authors. The authors alone are responsible for the content and writing of the manuscript.

GRANT SUPPORT

The authors have been partially supported by National Institutes of Health grants P20CA192976 (MKM) and P20CA192973 (UM); US Department of Defense grants W911NF-12-1-0073 (MKM) and W911NF-14-1-0064 (MKM); and National Science Foundation grant 1154214 (MKM).

REFERENCES

1. Bieri U, Moch H, Dehler S, Korol D and Rohrmann S. Changes in autopsy rates among cancer patients and their impact on cancer statistics from a public health point of view: a longitudinal study from 1980 to 2010 with data from Cancer Registry Zurich. *Virchows Arch.* 2015; 466:637-643.
2. Chen W. Cancer statistics: updated cancer burden in China. *Chin J Cancer Res.* 2015; 27:1.
3. Jung KW, Won YJ, Kong HJ, Oh CM, Cho H, Lee DH and Lee KH. Cancer statistics in Korea: incidence, mortality, survival, and prevalence in 2012. *Cancer Res Treat.* 2015; 47:127-141.
4. Siegel RL, Miller KD and Jemal A. Cancer statistics, 2015. *CA Cancer J Clin.* 2015; 65:5-29.
5. Torre LA, Bray F, Siegel RL, Ferlay J, Lortet-Tieulent J and Jemal A. Global cancer statistics, 2012. *CA Cancer J Clin.* 2015; 65:87-108.
6. DeSantis CE, Lin CC, Mariotto AB, Siegel RL, Stein KD, Kramer JL, Alteri R, Robbins AS and Jemal A. Cancer treatment and survivorship statistics, 2014. *CA Cancer J Clin.* 2014; 64:252-271.
7. Ganapathy S, Chen Q, Singh KP, Shankar S and Srivastava RK. Resveratrol enhances antitumor activity of TRAIL in prostate cancer xenografts through activation of FOXO transcription factor. *PloS one.* 2010; 5:e15627.
8. Harper CE, Patel BB, Wang J, Arabshahi A, Eltoum IA and Lamartiniere CA. Resveratrol suppresses prostate cancer progression in transgenic mice. *Carcinogenesis.* 2007; 28:1946-1953.
9. Li J, Chong T, Wang Z, Chen H, Li H, Cao J, Zhang P and Li H. A novel anticancer effect of resveratrol: reversal of epithelial-mesenchymal transition in prostate cancer cells. *Mol Med Rep.* 2014; 10:1717-1724.
10. Dimitriadis E, Kalogeropoulos T, Velaeti S, Sotiriou S, Vassiliou E, Fasoulis L, Klapsas V, Synesiou M, Apostolaki A, Trangas T and Pandis N. Study of genetic and epigenetic alterations in urine samples as diagnostic markers for prostate cancer. *Anticancer Res.* 2013; 33:191-197.
11. Ozen M and Pathak S. Genetic alterations in human prostate cancer: a review of current literature. *Anticancer Res.* 2000; 20:1905-1912.
12. Prostate cancer. Part B: Imaging techniques, radiotherapy, chemotherapy, and management issues. Proceedings of the Second International Symposium on Prostate Cancer. Paris, France, June 16-18, 1986. *Prog Clin Biol Res.* 1987; 243B:1-545.
13. Xie H, Li C, Dang Q, Chang LS and Li L. Infiltrating mast cells increase prostate cancer chemotherapy and radiotherapy resistances via modulation of p38/p53/p21 and ATM signals. *Oncotarget.* 2016; 7:1341-53. doi: 10.18632/oncotarget.6372.
14. Halin S, Hammarsten P, Wikstrom P and Bergh A. Androgen-insensitive prostate cancer cells transiently respond to castration treatment when growing in an androgen-dependent prostate environment. *The Prostate.* 2007; 67:370-377.
15. Kyprianou N. Apoptosis: therapeutic significance in the treatment of androgen-dependent and androgen-independent prostate cancer. *World J Urol.* 1994; 12:299-303.
16. Li W. Treatment of prostate cancer cells with adenoviral vector-mediated antisense RNA using androgen-dependent and androgen-independent promoters. *Med Oncol.* 2010; 27:519-530.
17. Sirotiak FM, She Y, Lee F, Chen J and Scher HI. Studies with CWR22 xenografts in nude mice suggest that ZD1839 may have a role in the treatment of both androgen-dependent and androgen-independent human prostate cancer. *Clin Cancer Res.* 2002; 8:3870-3876.
18. Baur JA, Pearson KJ, Price NL, Jamieson HA, Lerin C, Kalra A, Prabhu VV, Allard JS, Lopez-Lluch G, Lewis K, Pistell PJ, Poosala S, Becker KG, et al. Resveratrol improves health and survival of mice on a high-calorie diet. *Nature.* 2006; 444:337-342.
19. Baur JA and Sinclair DA. Therapeutic potential of resveratrol: the *in vivo* evidence. *Nat Rev Drug Discov.* 2006; 5:493-506.
20. Delmas D, Solary E and Latruffe N. Resveratrol, a phytochemical inducer of multiple cell death pathways: apoptosis, autophagy and mitotic catastrophe. *Curr Med Chem.* 18:1100-1121.
21. Chang L, Graham PH, Ni J, Hao J, Bucci J, Cozzi PJ and Li Y. Targeting PI3K/Akt/mTOR signaling pathway in the treatment of prostate cancer radioresistance. *Crit Rev Oncol Hematol.* 2015.
22. Choudhury Y, Yang Z, Ahmad I, Nixon C, Salt IP and Leung HY. AMP-activated protein kinase (AMPK) as a potential therapeutic target independent of PI3K/Akt signaling in prostate cancer. *Oncoscience.* 2014; 1:446-456. doi: 10.18632/oncoscience.49.
23. Han I, Yun M, Kim EO, Kim B, Jung MH and Kim SH. Umbilical cord tissue-derived mesenchymal stem cells induce apoptosis in PC-3 prostate cancer cells through activation of JNK and downregulation of PI3K/AKT signaling. *Stem Cell Res Ther.* 2014; 5:54.
24. Hatashita M, Taniguchi M, Baba K, Koshiba K, Sato T, Jujo Y, Suzuki R and Hayashi S. Sinodiellide A exerts thermosensitizing effects and induces apoptosis and G2/M cell cycle arrest in DU145 human prostate cancer cells via the Ras/Raf/MAPK and PI3K/Akt signaling pathways. *Int J Mol Med.* 2014; 33:406-414.
25. Kim SM, Park JH, Kim KD, Nam D, Shim BS, Kim SH, Ahn KS, Choi SH and Ahn KS. Brassinin induces apoptosis in PC-3 human prostate cancer cells through the suppression of PI3K/Akt/mTOR/S6K1 signaling cascades. *Phytotherapy research : PTR.* 2014; 28:423-431.

26. Kumar D, Shankar S and Srivastava RK. Rottlerin induces autophagy and apoptosis in prostate cancer stem cells via PI3K/Akt/mTOR signaling pathway. *Cancer letters*. 2014; 343:179-189.
27. Lee SH, Johnson D, Luong R and Sun Z. Crosstalk between androgen and PI3K/AKT signaling pathways in prostate cancer cells. *The Journal of biological chemistry*. 2015; 290:2759-2768.
28. Liang F, Yue J, Wang J, Zhang L, Fan R, Zhang H and Zhang Q. GPCR48/LGR4 promotes tumorigenesis of prostate cancer via PI3K/Akt signaling pathway. *Med Oncol*. 2015; 32:49.
29. Blagosklonny MV. Inhibition of S6K by resveratrol: in search of the purpose. *Aging (Albany NY)*. 2009; 1:511-514. doi: 10.18632/aging.100059.
30. Delmas D, Rebe C, Micheau O, Athias A, Gambert P, Grazide S, Laurent G, Latruffe N and Solary E. Redistribution of CD95, DR4 and DR5 in rafts accounts for the synergistic toxicity of resveratrol and death receptor ligands in colon carcinoma cells. *Oncogene*. 2004; 23:8979-8986.
31. Horndasch M and Culig Z. SOCS-3 antagonizes proapoptotic effects of TRAIL and resveratrol in prostate cancer cells. *The Prostate*. 2011; 71:1357-1366.
32. Ivanov VN, Partridge MA, Johnson GE, Huang SX, Zhou H and Hei TK. Resveratrol sensitizes melanomas to TRAIL through modulation of antiapoptotic gene expression. *Experimental cell research*. 2008; 314:1163-1176.
33. Shankar S, Chen Q, Siddiqui I, Sarva K and Srivastava RK. Sensitization of TRAIL-resistant LNCaP cells by resveratrol (3, 4', 5 tri-hydroxystilbene): molecular mechanisms and therapeutic potential. *J Mol Signal*. 2007; 2:7.
34. Shankar S, Siddiqui I and Srivastava RK. Molecular mechanisms of resveratrol (3,4,5-trihydroxy-trans-stilbene) and its interaction with TNF-related apoptosis inducing ligand (TRAIL) in androgen-insensitive prostate cancer cells. *Molecular and cellular biochemistry*. 2007; 304:273-285.
35. Tameda M, Sugimoto K, Shiraki K, Inagaki Y, Ogura S, Kasai C, Yoneda M, Okamoto R, Yamamoto N, Takei Y, Ito M and Nobori T. Resveratrol sensitizes HepG2 cells to TRAIL-induced apoptosis. *Anti-cancer drugs*. 2014; 25:1028-1034.
36. Toume K, Habu T, Arai MA, Koyano T, Kowithayakorn T and Ishibashi M. Correction to Prenylated Flavonoids and Resveratrol Derivatives Isolated from *Artocarpus communis* with the Ability to Overcome TRAIL Resistance. *J Nat Prod*. 2015; 78:968.
37. Toume K, Habu T, Arai MA, Koyano T, Kowithayakorn T and Ishibashi M. Prenylated flavonoids and resveratrol derivatives isolated from *Artocarpus communis* with the ability to overcome TRAIL resistance. *J Nat Prod*. 2015; 78:103-110.
38. Fulda S and Debatin KM. Resveratrol-mediated sensitisation to TRAIL-induced apoptosis depends on death receptor and mitochondrial signalling. *Eur J Cancer*. 2005; 41:786-798.
39. Halicka HD, Zhao H, Li J, Lee YS, Hsieh TC, Wu JM and Darzynkiewicz Z. Potential anti-aging agents suppress the level of constitutive mTOR- and DNA damage- signaling. *Aging (Albany NY)*. 2012; 4:952-965. doi: 10.18632/aging.100521.
40. Bai T, Dong DS and Pei L. Resveratrol mitigates isoflurane-induced neuroapoptosis by inhibiting the activation of the Akt-regulated mitochondrial apoptotic signaling pathway. *Int J Mol Med*. 2013; 32:819-826.
41. Brito PM, Simoes NF, Almeida LM and Dinis TC. Resveratrol disrupts peroxynitrite-triggered mitochondrial apoptotic pathway: a role for Bcl-2. *Apoptosis : an international journal on programmed cell death*. 2008; 13:1043-1053.
42. Danz ED, Skramsted J, Henry N, Bennett JA and Keller RS. Resveratrol prevents doxorubicin cardiotoxicity through mitochondrial stabilization and the Sirt1 pathway. *Free radical biology & medicine*. 2009; 46:1589-1597.
43. Sun W, Wang W, Kim J, Keng P, Yang S, Zhang H, Liu C, Okunieff P and Zhang L. Anti-cancer effect of resveratrol is associated with induction of apoptosis via a mitochondrial pathway alignment. *Adv Exp Med Biol*. 2008; 614:179-186.
44. Tinhofer I, Bernhard D, Senfter M, Anether G, Loeffler M, Kroemer G, Kofler R, Csordas A and Greil R. Resveratrol, a tumor-suppressive compound from grapes, induces apoptosis via a novel mitochondrial pathway controlled by Bcl-2. *FASEB journal*. 2001; 15:1613-1615.
45. Zhou X, Chen M, Zeng X, Yang J, Deng H, Yi L and Mi MT. Resveratrol regulates mitochondrial reactive oxygen species homeostasis through Sirt3 signaling pathway in human vascular endothelial cells. *Cell death & disease*. 2014; 5:e1576.
46. Rizvi F, Heimann T, Herrnreiter A and O'Brien WJ. Mitochondrial dysfunction links ceramide activated HRK expression and cell death. *PLoS One*. 2011; 6:e18137.
47. Zhang W, Wang X and Chen T. Resveratrol induces apoptosis via a Bak-mediated intrinsic pathway in human lung adenocarcinoma cells. *Cellular signalling*. 2012; 24:1037-1046.
48. Dorrie J, Gerauer H, Wachter Y and Zunino SJ. Resveratrol induces extensive apoptosis by depolarizing mitochondrial membranes and activating caspase-9 in acute lymphoblastic leukemia cells. *Cancer research*. 2001; 61:4731-4739.
49. Bhatt K, Feng L, Pabla N, Liu K, Smith S and Dong Z. Effects of targeted Bcl-2 expression in mitochondria or endoplasmic reticulum on renal tubular cell apoptosis. *Am J Physiol Renal Physiol*. 2008; 294:F499-507.
50. Soane L, Siegel ZT, Schuh RA and Fiskum G. Postnatal developmental regulation of Bcl-2 family proteins in brain mitochondria. *J Neurosci Res*. 2008; 86:1267-1276.

51. Metrailler-Ruchonnet I, Pagano A, Carnesecchi S, Ody C, Donati Y and Barazzone Argiroffo C. Bcl-2 protects against hyperoxia-induced apoptosis through inhibition of the mitochondria-dependent pathway. *Free Radic Biol Med.* 2007; 42:1062-1074.
52. de Graaf AO, van den Heuvel LP, Dijkman HB, de Abreu RA, Birkenkamp KU, de Witte T, van der Reijden BA, Smeitink JA and Jansen JH. Bcl-2 prevents loss of mitochondria in CCCP-induced apoptosis. *Exp Cell Res.* 2004; 299:533-540.
53. Park JW, Choi YJ, Suh SI, Baek WK, Suh MH, Jin IN, Min DS, Woo JH, Chang JS, Passaniti A, Lee YH and Kwon TK. Bcl-2 overexpression attenuates resveratrol-induced apoptosis in U937 cells by inhibition of caspase-3 activity. *Carcinogenesis.* 2001; 22:1633-1639.
54. Hu Q, Wu D, Chen W, Yan Z and Shi Y. Proteolytic processing of the caspase-9 zymogen is required for apoptosome-mediated activation of caspase-9. *The Journal of biological chemistry.* 2013; 288:15142-15147.
55. Hu Q, Wu D, Chen W, Yan Z, Yan C, He T, Liang Q and Shi Y. Molecular determinants of caspase-9 activation by the Apaf-1 apoptosome. *Proceedings of the National Academy of Sciences of the United States of America.* 2014; 111:16254-16261.
56. Marsden VS, O'Connor L, O'Reilly LA, Silke J, Metcalf D, Ekert PG, Huang DC, Cecconi F, Kuida K, Tomaselli KJ, Roy S, Nicholson DW, Vaux DL, et al. Apoptosis initiated by Bcl-2-regulated caspase activation independently of the cytochrome c/Apaf-1/caspase-9 apoptosome. *Nature.* 2002; 419:634-637.
57. Saikumar P, Mikhailova M and Pandeswara SL. Regulation of caspase-9 activity by differential binding to the apoptosome complex. *Front Biosci.* 2007; 12:3343-3354.
58. Shi Y. Apoptosome: the cellular engine for the activation of caspase-9. *Structure.* 2002; 10:285-288.
59. Wurstle ML, Laussmann MA and Rehm M. The central role of initiator caspase-9 in apoptosis signal transduction and the regulation of its activation and activity on the apoptosome. *Experimental cell research.* 2012; 318:1213-1220.
60. Byun HS, Song JK, Kim YR, Piao L, Won M, Park KA, Choi BL, Lee H, Hong JH, Park J, Seok JH, Lee YJ, Kang SW, et al. Caspase-8 has an essential role in resveratrol-induced apoptosis of rheumatoid fibroblast-like synoviocytes. *Rheumatology (Oxford).* 2008; 47:301-308.
61. Gossiau A, Chen M, Ho CT and Chen KY. A methoxy derivative of resveratrol analogue selectively induced activation of the mitochondrial apoptotic pathway in transformed fibroblasts. *British journal of cancer.* 2005; 92:513-521.
62. Sha H, Ma Q, Jha RK, Xu F, Wang L, Wang Z, Zhao Y and Fan F. Resveratrol ameliorates hepatic injury via the mitochondrial pathway in rats with severe acute pancreatitis. *Eur J Pharmacol.* 2008; 601:136-142.
63. Silberstein JL, Taylor MN and Antonarakis ES. Novel Insights into Molecular Indicators of Response and Resistance to Modern Androgen-Axis Therapies in Prostate Cancer. *Curr Urol Rep.* 2016; 17:29.
64. Thelen P, Gschwend J, Wolff JM and Miller K. [Mechanisms of Resistance in Antihormone Therapies of Advanced Prostate Cancer]. *Aktuelle Urol.* 2016; 47:79-85.
65. Benadiba M, Dos Santos RR, Silva Dde O and Colquhoun A. Inhibition of C6 rat glioma proliferation by [Ru2Cl(lbp)4] depends on changes in p21, p27, Bax/Bcl2 ratio and mitochondrial membrane potential. *Journal of inorganic biochemistry.* 2010; 104:928-935.
66. Bhardwaj A, Sethi G, Vadhan-Raj S, Bueso-Ramos C, Takada Y, Gaur U, Nair AS, Shishodia S and Aggarwal BB. Resveratrol inhibits proliferation, induces apoptosis, and overcomes chemoresistance through down-regulation of STAT3 and nuclear factor-kappaB-regulated antiapoptotic and cell survival gene products in human multiple myeloma cells. *Blood.* 2007; 109:2293-2302.
67. Cai Y, Zhao L, Qin Y, Zhang M and He Y. Resveratrol inhibits proliferation and induces apoptosis of nasopharyngeal carcinoma cell line C666-1 through AMPK activation. *Die Pharmazie.* 2015; 70:399-403.
68. Ding XZ and Adrian TE. Resveratrol inhibits proliferation and induces apoptosis in human pancreatic cancer cells. *Pancreas.* 2002; 25:e71-76.
69. Fang Y, DeMarco VG and Nicholl MB. Resveratrol enhances radiation sensitivity in prostate cancer by inhibiting cell proliferation and promoting cell senescence and apoptosis. *Cancer Sci.* 2012; 103:1090-1098.
70. Garcia-Zepeda SP, Garcia-Villa E, Diaz-Chavez J, Hernandez-Pando R and Gariglio P. Resveratrol induces cell death in cervical cancer cells through apoptosis and autophagy. *European journal of cancer prevention.* 2013; 22:577-584.
71. Dai W, Wang F, Lu J, Xia Y, He L, Chen K, Li J, Li S, Liu T, Zheng Y, Wang J, Lu W, Zhou Y, et al. By reducing hexokinase 2, resveratrol induces apoptosis in HCC cells addicted to aerobic glycolysis and inhibits tumor growth in mice. *Oncotarget.* 2015; 6:13703-13717. doi: 10.18632/oncotarget.3800.
72. Kim HJ, Yang KM, Park YS, Choi YJ, Yun JH, Son CH, Suh HS, Jeong MH and Jo WS. The novel resveratrol analogue HS-1793 induces apoptosis via the mitochondrial pathway in murine breast cancer cells. *Int J Oncol.* 2012; 41:1628-1634.
73. Zhang J, Ma K, Qi T, Wei X, Zhang Q, Li G and Chiu JF. P62 regulates resveratrol-mediated Fas/Cav-1 complex formation and transition from autophagy to apoptosis. *Oncotarget.* 2015; 6:789-801. doi: 10.18632/oncotarget.2733.
74. Ho CH, Hsu JL, Liu SP, Hsu LC, Chang WL, Chao CC and Guh JH. Repurposing of phentolamine as a potential anticancer agent against human castration-resistant prostate cancer: A central role on microtubule stabilization and mitochondrial apoptosis pathway. *The Prostate.* 2015; 75:1454-1466.

75. Dussmann H, Rehm M, Kogel D and Prehn JH. Outer mitochondrial membrane permeabilization during apoptosis triggers caspase-independent mitochondrial and caspase-dependent plasma membrane potential depolarization: a single-cell analysis. *J Cell Sci.* 2003; 116:525-536.
76. Mahyar-Roemer M, Katsen A, Mestres P and Roemer K. Resveratrol induces colon tumor cell apoptosis independently of p53 and precede by epithelial differentiation, mitochondrial proliferation and membrane potential collapse. *Int J Cancer.* 2001; 94:615-622.
77. Ma XD, Yan F, Ma AD and Wang HJ. [Resveratrol induces HepG2 cell apoptosis by depolarizing mitochondrial membrane]. *Nan fang yi ke da xue xue bao = Journal of Southern Medical University.* 2006; 26:406-408, 413.
78. Crek C, Mutlu Altundag E, Karademir B, Kocturk S, Kartal Ozer N and Taga Y. The effect of resveratrol on signal transduction pathways and the role of pro-apoptotic Bax protein on apoptosis in HCT- 116 colon carcinoma cell lines. *Free Radic Biol Med.* 2014; 75:S27.
79. Mohan J, Gandhi AA, Bhavya BC, Rashmi R, Karunagaran D, Indu R and Santhoshkumar TR. Caspase-2 triggers Bax-Bak-dependent and -independent cell death in colon cancer cells treated with resveratrol. *J Biol Chem.* 2006; 281:17599-17611.
80. Gogada R, Prabhu V, Amadori M, Scott R, Hashmi S and Chandra D. Resveratrol induces p53-independent, X-linked inhibitor of apoptosis protein (XIAP)-mediated Bax protein oligomerization on mitochondria to initiate cytochrome c release and caspase activation. *J Biol Chem.* 2011; 286:28749-28760.
81. Zhou C, Ding J and Wu Y. Resveratrol induces apoptosis of bladder cancer cells via miR21 regulation of the Akt/Bcl2 signaling pathway. *Mol Med Rep.* 2014; 9:1467-1473.
82. Li YJ and Xu HJ. [Relationship between apoptotic effect of Resveratrol on KG-1 cells and expression of bcl-2/bax]. *Zhongguo Shi Yan Xue Ye Xue Za Zhi.* 2008; 16:1026-1029.
83. Shimizu T, Nakazato T, Xian MJ, Sagawa M, Ikeda Y and Kizaki M. Resveratrol induces apoptosis of human malignant B cells by activation of caspase-3 and p38 MAP kinase pathways. *Biochemical pharmacology.* 2006; 71:742-750.
84. Li Y, Liu J, Liu X, Xing K, Wang Y, Li F and Yao L. Resveratrol-induced cell inhibition of growth and apoptosis in MCF7 human breast cancer cells are associated with modulation of phosphorylated Akt and caspase-9. *Appl Biochem Biotechnol.* 2006; 135:181-192.
85. Pozo-Guisado E, Merino JM, Mulero-Navarro S, Lorenzo-Benayas MJ, Centeno F, Alvarez-Barrientos A and Fernandez-Salguero PM. Resveratrol-induced apoptosis in MCF-7 human breast cancer cells involves a caspase-independent mechanism with downregulation of Bcl-2 and NF-kappaB. *International journal of cancer.* 2005; 115:74-84.
86. Liu B, Zhou Z, Zhou W, Liu J, Zhang Q, Xia J, Liu J, Chen N, Li M and Zhu R. Resveratrol inhibits proliferation in human colorectal carcinoma cells by inducing G1/Sphase cell cycle arrest and apoptosis through caspase/cyclinCDK pathways. *Mol Med Rep.* 2014; 10:1697-1702.
87. Singh N, Nigam M, Ranjan V, Sharma R, Balapure AK and Rath SK. Caspase mediated enhanced apoptotic action of cyclophosphamide- and resveratrol-treated MCF-7 cells. *J Pharmacol Sci.* 2009; 109:473-485.
88. Zhang W, Wang X and Chen T. Resveratrol induces mitochondria-mediated AIF and to a lesser extent caspase-9-dependent apoptosis in human lung adenocarcinoma ASTC-a-1 cells. *Mol Cell Biochem.* 2011; 354:29-37.
89. Ulakesai Z, Bagamery F, Vincze I, Szoko E and Tabi T. Protective effect of resveratrol against caspase 3 activation in primary mouse fibroblasts. *Croat Med J.* 2015; 56:78-84.
90. Johansson P, Fasth A, Ek T and Hammarsten O. Validation of a flow cytometry-based detection of gamma-H2AX, to measure DNA damage for clinical applications. *Cytometry B Clin Cytom.* 2016.
91. Sedelnikova OA, Pilch DR, Redon C and Bonner WM. Histone H2AX in DNA damage and repair. *Cancer Biol Ther.* 2003; 2:233-235.
92. Rogakou EP, Pilch DR, Orr AH, Ivanova VS and Bonner WM. DNA double-stranded breaks induce histone H2AX phosphorylation on serine 139. *J Biol Chem.* 1998; 273:5858-5868.
93. Burma S, Chen BP, Murphy M, Kurimasa A and Chen DJ. ATM phosphorylates histone H2AX in response to DNA double-strand breaks. *J Biol Chem.* 2001; 276:42462-42467.
94. Walters DK, Wu X, Tschumper RC, Arendt BK, Huddleston PM, Henderson KJ, Dispenzieri A and Jelinek DF. Evidence for ongoing DNA damage in multiple myeloma cells as revealed by constitutive phosphorylation of H2AX. *Leukemia.* 2011; 25:1344-1353.
95. Yuan J, Adamski R and Chen J. Focus on histone variant H2AX: to be or not to be. *FEBS Lett.* 2010; 584:3717-3724.
96. Mukherjee B, Kessinger C, Kobayashi J, Chen BP, Chen DJ, Chatterjee A and Burma S. DNA-PK phosphorylates histone H2AX during apoptotic DNA fragmentation in mammalian cells. *DNA Repair (Amst).* 2006; 5:575-590.
97. Cook PJ, Ju BG, Telese F, Wang X, Glass CK and Rosenfeld MG. Tyrosine dephosphorylation of H2AX modulates apoptosis and survival decisions. *Nature.* 2009; 458:591-596.
98. Ferreira RF, Souza DR and Souza AS. Factors that Induce DNA Damage Involving Histone H2AX Phosphorylation. *Radiology.* 2015; 277:307-308.
99. Demoulin B, Hermant M, Castrogiovanni C, Staudt C and Dumont P. Resveratrol induces DNA damage in colon cancer cells by poisoning topoisomerase II and activates the ATM kinase to trigger p53-dependent apoptosis. *Toxicol In Vitro.* 2015; 29:1156-1165.

100. Sengottuvelan M, Deeptha K and Nalini N. Resveratrol ameliorates DNA damage, prooxidant and antioxidant imbalance in 1,2-dimethylhydrazine induced rat colon carcinogenesis. *Chem Biol Interact.* 2009; 181:193-201.
101. Seve M, Chimienti F, Devergnas S, Aouffen M, Douki T, Chantegrel J, Cadet J and Favier A. Resveratrol enhances UVA-induced DNA damage in HaCaT human keratinocytes. *Med Chem.* 2005; 1:629-633.
102. Tyagi A, Gu M, Takahata T, Frederick B, Agarwal C, Siriwardana S, Agarwal R and Sclafani RA. Resveratrol selectively induces DNA Damage, independent of Smad4 expression, in its efficacy against human head and neck squamous cell carcinoma. *Clin Cancer Res.* 2011; 17:5402-5411.