











## RESEARCH ARTICLE

**REVISED** **A tool for assessing the climate change mitigation and health impacts of environmental policies: the Cities Rapid Assessment Framework for Transformation (CRAFT) [version 2; peer review: 3 approved]**

Phil Symonds<sup>1</sup>, James Milner<sup>2</sup>, Nahid Mohajeri <sup>1</sup>, Juliette Aplin<sup>3</sup>, Joanna Hale <sup>4</sup>, Simon J Lloyd <sup>5</sup>, Henry Fremont <sup>6</sup>, Sam Younkin <sup>6</sup>, Clive Shrubsole <sup>1</sup>, Lawrie Robertson<sup>3</sup>, Jonathon Taylor <sup>1,7</sup>, Nici Zimmermann <sup>1</sup>, Paul Wilkinson<sup>2</sup>, Mike Davies<sup>1</sup>

<sup>1</sup>UCL Institute for Environmental Design and Engineering, London, WC1H 0NN, UK

<sup>2</sup>Centre on Climate Change and Planetary Health & Department of Public Health, Environments and Society, London School of Hygiene and Tropical Medicine, London, UK

<sup>3</sup>Buro Happold Consulting, London, W1T 1PD, UK

<sup>4</sup>Centre for Behaviour Change, University College London, London, UK

<sup>5</sup>Climate and Health Program (CLIMA), Barcelona Institute for Global Health (ISGlobal), Barcelona, Spain

<sup>6</sup>Global Health Institute, University of Wisconsin, Madison, USA

<sup>7</sup>Department of Civil Engineering, Tampere University, Tampere, Finland

**v2** **First published:** 12 Nov 2020, 5:269  
<https://doi.org/10.12688/wellcomeopenres.16345.1>  
**Latest published:** 18 May 2021, 5:269  
<https://doi.org/10.12688/wellcomeopenres.16345.2>

### Abstract

**Background:** A growing number of cities, including Greater London, have set ambitious targets, including detailed policies and implementation plans, to reach global goals on sustainability, health, and climate change. Here we present a tool for a rapid assessment of the magnitude of impact of specific policy initiatives to reach these targets. The decision-support tool simultaneously quantifies the environmental and health impacts of specified selected policies.




**Methods:** The 'Cities Rapid Assessment Framework for Transformation (CRAFT)' tool was applied to Greater London. CRAFT quantifies the effects of ten environmental policies on changes in (1) greenhouse gas (GHG) emissions, (2) exposures to environmental hazards, (3) travel-related physical activity, and (4) mortality (the number of attributable deaths avoided in one typical year). Publicly available data and epidemiological evidence were used to make rapid quantitative estimates of these effects based on proportional reductions in GHG emissions and environmental exposures from current baseline levels and to compute the mortality impacts.

**Results:** The CRAFT tool estimates that, of roughly 50,000 annual

### Open Peer Review

**Reviewer Status** 

	Invited Reviewers		
	1	2	3
<b>version 2</b>			
(revision)			
18 May 2021	report	report	report
			
<b>version 1</b>			
12 Nov 2020			
	report	report	

1. **Patrick L. Kinney** , Boston University School of Public Health, Boston, USA
2. **George D. Thurston** , New York University School of Medicine, New York City, USA
3. **Rebecca K. Saari** , University of Waterloo,

deaths in Greater London, the modelled hazards (PM<sub>2.5</sub> (from indoor and outdoor sources), outdoor NO<sub>2</sub>, indoor radon, cold, overheating) and low travel-related physical activity are responsible for approximately 10,000 premature environment-related deaths. Implementing the selected policies could reduce the annual mortality number by about 20% (~1,900 deaths) by 2050. The majority of these deaths (1,700) may be avoided through increased uptake in active travel. Thus, out of ten environmental policies, the 'active travel' policy provides the greatest health benefit. Also, implementing the ten policies results in a GHG reduction of around 90%.

**Conclusions:** The CRAFT tool quantifies the effects of city policies on reducing GHG emissions, decreasing environmental health hazards, and improving public health. The tool has potential value for policy makers through providing quantitative estimates of health impacts to support and prioritise policy options.

### Keywords

Health impact assessment, Rapid assessment tool, Mortality, Greenhouse gas emissions, Environmental exposures, City-scale policy assessment

Waterloo, Canada

Any reports and responses or comments on the article can be found at the end of the article.

**Corresponding author:** Nahid Mohajeri ([nahid.mohajeri.09@ucl.ac.uk](mailto:nahid.mohajeri.09@ucl.ac.uk))

**Author roles:** **Symonds P:** Conceptualization, Data Curation, Formal Analysis, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Milner J:** Conceptualization, Data Curation, Formal Analysis, Methodology, Writing – Original Draft Preparation, Writing – Review & Editing; **Mohajeri N:** Visualization, Writing – Original Draft Preparation, Writing – Review & Editing; **Aplin J:** Conceptualization, Writing – Review & Editing; **Hale J:** Conceptualization, Writing – Review & Editing; **J Lloyd S:** Data Curation, Formal Analysis, Methodology; **Fremont H:** Data Curation, Formal Analysis; **Younkin S:** Data Curation, Formal Analysis; **Shrubsole C:** Writing – Review & Editing; **Robertson L:** Conceptualization, Writing – Review & Editing; **Taylor J:** Data Curation, Formal Analysis, Writing – Review & Editing; **Zimmermann N:** Conceptualization, Writing – Review & Editing; **Wilkinson P:** Conceptualization, Funding Acquisition, Methodology, Writing – Review & Editing; **Davies M:** Conceptualization, Funding Acquisition, Methodology, Writing – Review & Editing

**Competing interests:** No competing interests were disclosed.

**Grant information:** This work is part of the Complex Urban Systems for Sustainability and Health (CUSSH) project, which is supported by Wellcome (project reference 209387).

*The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

**Copyright:** © 2021 Symonds P *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**How to cite this article:** Symonds P, Milner J, Mohajeri N *et al.* **A tool for assessing the climate change mitigation and health impacts of environmental policies: the Cities Rapid Assessment Framework for Transformation (CRAFT) [version 2; peer review: 3 approved]** Wellcome Open Research 2021, 5:269 <https://doi.org/10.12688/wellcomeopenres.16345.2>

**First published:** 12 Nov 2020, 5:269 <https://doi.org/10.12688/wellcomeopenres.16345.1>

**REVISED Amendments from Version 1**

The main changes in the revised version of the manuscript are as follows:

- We clarify the health estimation results related to the green space (Reviewer 1).
- Regarding the confidence intervals, we clarify that the CRAFT tool was intended to provide a broad understanding of what scale of impact specific policies might be expected to achieve and with what level of change. Detailed uncertainty analysis is beyond the scope and current capabilities of the CRAFT tool. Instead, complex dynamic models have the capability to assess the uncertainties in much greater detail (Reviewer 2).
- We justify and clarify in details the reasons for using NO<sub>2</sub> coefficient 1.023 per 10 µg/m<sup>3</sup> in this study which is taken from the 2018 COMEAP report. We also modify Figure 5 and Figure 6 and add a label to the NO<sub>2</sub> in the figure caption to make a clear distinction between NO<sub>2</sub> and PM<sub>2.5</sub> results. We make it clear that the health impacts of NO<sub>2</sub> should not be added to the PM<sub>2.5</sub> impacts (Reviewer 2).
- We have made textual improvements and clarifications throughout the text.

**Any further responses from the reviewers can be found at the end of the article**

## Introduction

With rapidly increasing urbanization, it is estimated that by 2050 nearly 70% of the world's population will be living in cities (United Nations, 2018). Cities are complex systems (Batty, 2008) and have long been known to be engines of innovation and creation of wealth. They are also key centres of pollution and diseases (Bettencourt *et al.*, 2007). While cities provide great potential for improving our lives and well-being, there exists considerable variation in the levels of environmental exposures including (i) air pollution, (ii) noise, (iii) temperature, (iv) green space, and (v) physical activity within and between cities. This is partly due to cities having different urban-planning practices and different infrastructure systems. There is rapidly growing knowledge as to the effects of environmental exposure on human health, and this knowledge is increasingly being translated into public policy (Caplin *et al.*, 2019).

With regard to environmental exposures, it is known that (i) air pollution, (ii) noise and (iii) temperature all have adverse health effects including increased premature mortality (Nieuwenhuijsen *et al.*, 2017a; Nieuwenhuijsen *et al.*, 2017b). By contrast, (iv) green spaces have predominantly positive health effects (Gascon *et al.*, 2016a; Gascon *et al.*, 2016b; Hartig *et al.*, 2014), but also some negative impacts such as contributing to urban sprawl and the spread of vector-borne diseases (Cucca, 2012; Hartig *et al.*, 2014; Löhmus & Balbus, 2015). As for (v) physical activity, it has many health benefits (Warburton & Bredin, 2017). In 2008, around 31% of adults above the age of 15 (28% of men and 34% of women) worldwide had insufficient physical activity. Each year globally about 3.2 million people die because of too little physical activity

(WHO, 2014). In Barcelona, 20% of premature mortality is related to urban and transport planning-related exposures, which include air pollution, noise, temperature, available green space, and physical activity levels (Mueller *et al.*, 2017). Many premature deaths could be prevented if recommendations for physical activity, exposure to air pollution, noise, and heat, as well as access to green space were followed.

Health impact assessment (HIA) has been recognized as one of the main tools to integrate evidence into decision-making processes, to include improved public health as a part of all policies, and to promote public health in multiple sectors (Davenport *et al.*, 2006; IFC, 2009; Mindell *et al.*, 2004; NHS, 2002; NRC, 2011; Shafiea *et al.*, 2013; Ståhl *et al.*, 2006). The EU-funded projects INTARESE and HEIMTSA represent rare examples of integrated environmental health impact assessments (Briggs, 2008; Hänninen *et al.*, 2014; Nieuwenhuijsen *et al.*, 2017a; Prüss-Ustün *et al.*, 2003). A combination of quantitative and participatory (citizen and stakeholder involvement) HIA tools has been used to allow policy makers to explore more accurately the positive and negative health impacts of current and future policy scenarios. However, HIAs that have been mainly conducted so far have been at the national or regional level (Briggs *et al.*, 2016; Forouzanfar *et al.*, 2017). This indicates the importance of developing such tools for assessing health-related policies at the city scale.

Action by cities will be crucial to the fulfilment of global goals on sustainability and health, including those related to climate change. A growing number of cities have set ambitious targets for environmental action and are developing detailed policies and implementation plans to reach them. Many of these policies are expected to have substantial benefits for health and arguments about health benefits are often invoked to strengthen the case for their adoption. However, our own experience of working with city authorities suggests that, at the early stages of policy formulation, decision-makers would benefit greatly if a tool could be provided for a rapid assessment of the magnitude of impact of specific policy initiatives regarding climate change mitigation and health. We hypothesized that the provision of evidence on such joint impacts would be particularly useful to the decision process and so aimed to design a tool that would provide broad estimates of these impacts. The result was an Excel-based modelling framework known as the *Cities Rapid Assessment Framework for Transformation* (CRAFT), which we describe in this paper.

The intended features of this tool were that it should be able to provide rapid and useful, approximate quantification of both CO<sub>2</sub>e reduction and mortality impacts of multiple policies so that 'order-of-magnitude' comparisons could be made at an early stage in the decision process to help guide initial discussions of policy choices. It was not intended as a tool for detailed simulation of the effects of policies. Such detailed modelling would need to be undertaken subsequently using other methods once initial choices were narrowed. The CRAFT model was intended to provide a broad understanding


of what scale of impact specific policies might be expected to achieve and with what level of change.

In this paper, we report the development and methods of the CRAFT tool (or model), illustrating its application to selected environment and health policies for Greater London, UK, in relation to transport, housing, energy, greenspace and waste. We discuss its potential strengths and weaknesses and its role as a decision-support tool and outline the process for the future evaluation of the utility of the model to policy makers and the public.

**Methods**

The development of the CRAFT tool was undertaken as part of the research project *Complex Urban Systems for Sustainability and Health* (CUSSH), supported by the Wellcome Trust. Figure 1 provides a flow diagram of the CRAFT tool.

In its current version, the tool provides estimates of the effects of specified policy choices with regard to (i) reduction in greenhouse gas (GHG) emissions in terms of tonnes of CO<sub>2</sub> equivalent avoided per year and (ii) the associated change in mortality quantified as the number of attributable deaths avoided in one year. To do this, the tool uses a common principle of fractional attribution. The precise details of how this principle is applied vary somewhat from policy to policy but the general principle entails estimating the fraction of total GHG emissions that are attributable to a particular sector (transport, housing, etc.) and then to the relevant target of action within each sector. The overall impact on GHG emissions for a specific policy is then computed by multiplying relevant fractions (e.g. the fraction of emissions attributable to the target sector × the fraction of the target that will be changed as a result of the policy × the fractional reduction in emissions per unit of the target consequent to the policy action). This is



**Figure 1. Flow diagram of the CRAFT tool.** The grey colour indicates the first step of the assessment using the CRAFT tool namely, policy identification. The orange colour indicates the second step of the assessment namely, the environmental impacts. This includes GHG emissions, air pollution, and indoor air quality, as well as travel related physical inactivity. The blue colour shows the final step of the assessment, that is, the health impacts. The arrows between stakeholder participation and the assessment steps represents engagement (e.g. presenting policy choices or showing model results) and receiving feedback from stakeholders.

more clearly understood by the example shown in [Figure 2](#). Throughout we assumed that the linear scaling implied by this approach is a reasonable approximation to real change even though in reality there may be some non-linearities. Initial estimates of the proportion of the city's emissions by sector, of specific sources within each sector and of the effect of the technology or behaviour change relevant to that source, were all taken from published sources, using the most relevant and up-to-date data available.

A similar process was applied for the computation of mortality impacts. The specific impacts we considered included those related to changes in (i) outdoor air pollution (particulate matter  $<2.5 \mu\text{m}$  ( $\text{PM}_{2.5}$ ), and nitrogen dioxide ( $\text{NO}_2$ )), (ii) the indoor environment (indoor  $\text{PM}_{2.5}$ , radon, cold, and over-heating), and (iii) travel-related physical activity. For each selected policy, we estimated what proportion of the population might be affected by the policy and what level of change in exposure or behaviour (physical activity) of an individual within the affected population as a result of the policy. For each form of exposure or health behaviour (physical activity), the policy-modifiable attributable mortality was computed using published epidemiological exposure-response relationships (relative risks) applied to current levels of exposure (ambient air pollution, winter indoor cold, levels of physical activity, etc.) and compared with exposure as modified by the specific policy. As with GHG emissions, the calculations generally reduced to a set of fractions to estimate the change to current exposure because of the policy, from which mortality calculations were computed by applying relative risks to the exposure change in the affected proportion of the population.


In the current implementation of the CRAFT tool, we do not include calculations of morbidity impacts which, for most outcomes, are less securely estimable than for mortality and often less important if judged in terms of monetized equivalent of quality-adjusted life-years. Morbidity impacts will be added

in future versions of the tool, however. For simplicity and transparency, for both GHG changes and attributable mortality, we provide calculations of the impact in one year relative to current levels of emission/exposure and assuming no time lag. In reality, there are bound to be time delays in implementation and in the evolution of impacts, sometimes substantial. However, characterizing year-by-year changes would require appreciable additional complexity and assumptions in the modelling process, which would detract from the simplicity and transparency of generating single-year estimates.

### Example application

To illustrate the use of the tool, we now describe its application to the assessment of selected environmental policies for Greater London drawn from key strategy documents published by the Mayor's Office. Greater London, a partner city of the CUSSH project, has ambitious goals to reduce GHG emissions and to adapt to the impacts of climate change through a wide range of cross-sector actions ([Belesova et al., 2018](#)). It is also a city with a strong focus on sustainability and health equity. We chose to focus on key policy objectives outlined in the 2018 Environment Strategy for Greater London ([GLA, 2018a](#)) and other documents, including the Mayor's Transport Strategy ([GLA, 2018b](#)), the Greater London Housing Strategy ([GLA, 2018c](#)), the GLA's Zero Carbon London: A 1.5°C Compatible Climate Action Plan ([GLA, 2018d](#)), and the Zero Carbon Pathways (ZCP) Tool ([GLA, 2018e](#)). We selected ten policies to model from among approximately 100 policies outlined in these documents, choosing policies that entailed relatively broad actions and which were primarily motivated by greenhouse gas emissions reduction ([Table 1](#)).

In this section, we describe the data sources and the key assumptions that are made to estimate the changes in GHG emissions and mortality impact of changes to environmental exposures and physical activity associated with the policy actions.



**Figure 2.** Simplified calculation steps used in CRAFT to estimate changes in GHG emissions.


**GHG emissions**

Our calculation for Greater London’s GHG emissions were based on Scope 1 & 2 emissions as defined by the Greenhouse Gas Protocol (GLA, 2016a), namely all direct GHG emissions produced within the city’s boundary (Scope 1) and indirect GHG emissions from consumption of imported electricity, heat or steam (Scope 2). We did not include Scope 3 emissions, i.e. those associated with manufactured and

purchased goods. It is important to note, however, that at 115 MtCO<sub>2</sub>e, Greater London’s Scope 3 emissions are estimated to account for more than three times Greater London’s direct emissions of around 34 MtCO<sub>2</sub>e. Estimates of the sector contributions to Greater London’s Scope 1 & 2 GHG emissions were derived from *The London Energy and Greenhouse Gas Inventory (LEGGI)* (GLA, 2016a) and are summarized in Figure 3.

**Table 1. Policy ID and a short description of the selected 10 policies.**

No.	Policy ID	Policy description
1	E-transport system	London’s entire transport system to be zero emission by 2050 (source: London Environment Strategy)
2	Active travel	8/10 trips made on foot, by cycle or by public transport (from 6/10 today) by 2041 (source: Mayor’s Transport Strategy)
3	Buildings upgraded	Up to 50% of buildings upgraded by 2025, 100% upgraded by 2050 (source: Zero Carbon London/ZCP Tool)
4	Heat pumps	Up to two million heat pumps installed across London by 2050 (source: Zero Carbon London/ZCP Tool)
5	Heat networks	Tenfold increase in heat networks by 2025, connecting up to 650,000 homes to waste and environmental heat sources by 2050 (source: Zero Carbon London/ZCP Tool)
6	PV installations	Up to 100,000 photovoltaic installations across London by 2025, increasing to 25% of all viable buildings by 2050 (source: Zero Carbon London/ZCP Tool)
7	Grid decarbonisation	Grid decarbonisation in line with UK carbon budgets. High penetration of renewables and nuclear, doubling capacity by 2030 (source: Zero Carbon London/ZCP Tool)
8	Green gas supply	Green gas in national supply increasing significantly from 2030, contributing 13% of gas supply by 2050 (source: Zero Carbon London/ZCP Tool)
9	Greenspace	Increase London’s green area from 47% to >50% by 2050 (source: London Environment Strategy)
10	Zero waste city	London will be a zero waste city. By 2026 no biodegradable or recyclable waste will be sent to landfill, and by 2030 65 per cent of London’s municipal waste will be recycled. (source: London Environment Strategy)



**Figure 3. Greater London GHG emissions by sector in % include both direct (Scope 1) and indirect (Scope 2) GHG emissions [reproduced with permission from LEGGI; GLA, 2016a].**

The GHG emissions associated with each of the ten policies was estimated using the principle of ‘fractional attribution’ outlined above, i.e. by the product of the fractions reflecting the proportion of overall GHG emissions attributable to the sector, the proportion of those sectoral emissions targeted by the policy and the proportional reduction in emissions from those targeted. The data sources and key assumptions used in these calculations are summarized in [Table 2](#).

### Mortality impact of change to environmental exposures

The method used to estimate changes in mortality related to environmental exposures and health-related behaviours as a result of each policy is broadly similar to that used for the GHG emissions. Namely, we calculate the proportional contributions from the relevant sector, the sources targeted by the policy within the sector, and then the relative change achieved among the targeted sources by the specific action (i.e. by the policy-related intervention). Baseline mortality rates for the Greater London population were based on data from the Office for National Statistics for the year 2017.


For outdoor air pollution, we assumed that 25% of ambient (background) PM<sub>2.5</sub> and 80% of the NO<sub>2</sub> concentrations in Greater London are due to emissions within the GLA ([Howard, 2015](#)). The changes in PM<sub>2.5</sub> and NO<sub>2</sub> concentrations from policies were then assumed to occur in proportion to the change in local source emissions for these pollutants. The

proportion of emissions produced within GLA by sector and transport type were obtained from the Greater London Atmospheric Emissions Inventory (LAEI) ([GLA, 2016b](#)) ([Figure 4](#)). Our primary calculations of outdoor air pollution-related mortality were based on changes to PM<sub>2.5</sub> alone, using an exposure-response coefficient derived from ([Pope et al., 2002](#)). However, because of increasing evidence of independent effects on mortality of NO<sub>2</sub>, we also carried out parallel calculations for change in NO<sub>2</sub> concentrations using exposure-response functions derived from ([COMEAP, 2018](#)). The impacts based on these calculations are not independent and should not be summed. They were included as alternatives to show the different estimates derived from these methods, especially for transport-related policies, which have much greater impact on NO<sub>2</sub> emissions and concentrations than on those for PM<sub>2.5</sub>.

For changes to the indoor environment, we used methods of health impact calculation outlined in [Hamilton et al. \(2015\)](#), for cold, heat ([Taylor et al., 2018](#)), indoor PM<sub>2.5</sub> and radon. A key uncertainty with these calculations was the change in ventilation characteristics of the dwelling consequent to home energy-efficiency interventions. Depending on the assumption about changes in ventilation following retrofit, the net impacts on mortality can be negative or positive. Additional data sources along with key assumptions used in the estimations are listed in [Table 3](#).

**Table 2. Data source and the assumptions used for GHG emissions calculations.**

NO.	Policy ID	Sources	Assumptions
1	E-transport system	( <a href="#">BEIS, 2018</a> )	Assume entire transport system is electrified. Use the BEIS Green book emissions factor is used which suggest that GHG emissions from the grid will reduce by ~95% from 2010 levels by 2050.
2	Active travel	London Travel Demand Survey ( <a href="#">TFL, 2018</a> ); ( <a href="#">EEA, 2015</a> )	The current mode share from the TFL's Travel Demand Survey is used. These mode shares are converted to typical distances travelled and then multiplied by the CO <sub>2</sub> per passenger kms from <a href="#">EEA (2015)</a> .
3	Buildings upgraded	<a href="#">Hamilton et al., 2015</a>	Predictions for changes in energy savings due to energy efficiency retrofit are used
4	Heat pumps	( <a href="#">BEIS, 2018</a> )	Assume electrification of heat and decarbonisation of the grid in line with BEIS Green book. Also assume a coefficient of performance (COP) of 3 for heat pumps.
5	Heat networks	<a href="#">DECC, 2009</a>	Take assumptions on CO <sub>2</sub> e savings from heat networks compared to conventional heating systems from a report commissioned by DECC
6	PV installations	( <a href="#">BEIS, 2018</a> ); ( <a href="#">GLA, 2018f</a> )	The ratio of energy demand that could be met by solar was taken from the GLA's Solar action plan. BEIS Green book emission factors were again applied.
7	Grid decarbonisation	( <a href="#">BEIS, 2018</a> )	Total electricity demand GHG emissions (LEGGI) and BEIS Green book emission factors were used.
8	Green gas supply	NA	No additional data requirements.
9	Greenspace	<a href="#">GLA, 2015</a>	Carbon Sequestration from the urban forest and energy savings from buildings reported in Valuing London's Urban Forest ( <a href="#">GLA, 2015</a> ) were used.
10	Zero waste city	<a href="#">GLA, 2017</a>	The reduction in CO <sub>2</sub> e from waste management used was taken from 'Greenhouse Gas Emissions Performance Standard for London's Local Authority Collected Waste – 2017 Update' ( <a href="#">GLA, 2017</a> ).



**Figure 4.** Greater London outdoor particulate matter, PM<sub>2.5</sub> emissions by sector [reproduced with permission from LAEI (GLA, 2016b)].

**Table 3.** Data source and the assumptions used for environmental exposure calculations.

NO.	Policy ID	Sources	Assumptions
1	E-transport system	NA	No additional data requirements.
2	Active travel	London Travel Demand Survey (TFL, 2018), (BEIS, 2012)	As with GHG emissions, the London Travel Demand Survey is used to establish the current proportion of passenger kms travelled by various modes. Transport emission factors for various pollutants per km were obtained from the National Atmospheric Emissions Inventory (BEIS, 2012).
3	Building upgraded	Hamilton <i>et al.</i> , 2015; Taylor <i>et al.</i> , 2018	Predictions for changes in indoor environmental exposures due to home energy efficiency retrofit were taken from Hamilton <i>et al.</i> , 2015. This includes indoor and outdoor PM <sub>2.5</sub> , Radon, mould and standard indoor temperature. Predicted changes in overheating were based on work by Taylor <i>et al.</i> (2018).
4	Heat pumps	NA	No additional data requirements. Heat pumps assumed to completely eliminate direct emissions of pollutants from homes.
5	Heat networks	NA	No additional data requirements. Heat networks assumed to eliminate direct emissions of pollutants.
6	PV installation	DEFRA, 2019	Emissions due to power generation were determined from DEFRA (2019). This was multiplied by the reduction in demand due to the adding solar PV.
7	Grid decarbonisation	DEFRA, 2019	Emissions due to power generation were determined from DEFRA (2019). It is assumed grid decarbonisation would eliminate the majority of these emissions.
8	Green gas supply	NA	No data currently available.
9	Greenspace	GLA, 2015	Proportion of pollution removed by the urban forest was obtained from Valuing London's Urban Forest (GLA, 2015).
10	Zero waste city	NA	No data currently available.

For transport-related physical activity, we assessed changes in transport-related metabolic equivalent of the task (MET) hours per week. The MET for a specific task is defined as the ratio of the rate of energy consumption by that person while performing that activity compared with a reference. By convention the reference is taken as 3.5 mL of oxygen per

kilogramme body mass per minute, which is approximately the energy expended when sitting at rest (Jetté *et al.*, 1990). Data from the London Travel Demand Survey (TFL, 2018) was used to estimate the current mean level of travel-related physical activity in Greater London (7.65 MET-hrs/week). The mean level of leisure related physical activity (not including travel



activity) was estimated (11.25 MET-hrs/week) using data from [Arem et al. \(2015\)](#). Changes to all-cause mortality risk were then based on estimates of change to the sum of these two sources of physical activity, applying a piecewise linear exposure-response function for all-cause mortality, constructed with data from [Arem et al. \(2015\)](#).

For this version of CRAFT the mortality impact associated with changes in green space was based only on calculations of the estimated effect of green space on ambient air pollution – specifically the proportion of pollution removed by the urban forest obtained from the 2015 report *Valuing London's Urban Forest* (GLA, 2015). This omits any effects associated with, for example, protection against heat risk in summer, increased levels of physical activity, improved mental well-being and other effects that depend on the form of green space change and for which the methods of quantification remains uncertain.

### Policy interactions

Where possible, the CRAFT tool attempts to account for the interactions between policies. We consider two general types of interaction. First, dependencies, where the benefits achieved by a given policy are contingent on the degree of implementation of at least one other policy. Second, double-counting, in which a given impact is (potentially) achieved independently by more than one policy. An example is the effect of the interaction between 'E-transport system' (policy 1) and 'Active travel' (policy 2), on GHG emissions which is mediated by a third policy namely, 'Grid decarbonisation' (policy 7). The first step is to determine which policy will be implemented first chronologically. In this case, policy 2 has a target of the year 2041, while policy 1 has a target of the year 2050. Therefore, reduced demand in non-active travel types due to modal shift is subtracted from policy 1. Policy 1 is also contingent on a third policy, 'Grid decarbonisation' being fully achieved. If grid decarbonisation is not achieved, the GHG reducing benefits of zero emission transport decrease because it is assumed to be electric powered. As indicated above, the possibility and the nature of any interactions depends on the assumptions embedded in the initial calculation of the impacts of a given policy.

### Results

The results for the ten policies assessed by the CRAFT tool are shown in [Figure 5](#) (central estimates of percentage changes in GHGs emissions and environmental exposures) and [Figure 6](#) (estimated impact on health due to the 10 key policy actions in Greater London). *Extended data*, Supplementary File 1 ([Symonds et al., 2020a](#)) presents the detailed calculation steps for each policy, along with assumptions used for current and minimum exposures and the exposure-mortality coefficients and population mortality statistics used. Overall, fully implementing all ten selected policies could reduce the environmental disease burden in Greater London by about 1,900 deaths in one year, or around 20% of current mortality. This total is based on calculations in which the impact of outdoor air pollution is based on changes to PM<sub>2.5</sub> only (to avoid potential double counting due to outdoor PM<sub>2.5</sub> and NO<sub>2</sub>). In terms of

reduction to CO<sub>2</sub>e emissions, the largest contribution would occur from grid decarbonization, followed by the 'zero emission' transport system, use of heat pumps and buildings energy efficiency upgrades ([Figure 5](#)).


Most policies were estimated to result in a reduction of harmful exposures. The transport related policies ('E-transport system' and 'Active travel') have the greatest potential for reduction of both CO<sub>2</sub>e emissions and mortality. By far the largest contributor to reducing mortality was estimated to be active travel, with a population impact for the increase in active travel greater than all other changes in exposure combined. However, if the calculation of mortality following changes to outdoor air pollution were based on NO<sub>2</sub> rather than PM<sub>2.5</sub>, these impacts would be only a little lower than those for physical activity ([Figure 6](#)).

The results relating to housing energy efficiency interventions ('Buildings upgraded') are mixed, partly because of the uncertainties over changes to ventilation characteristics. The 'buildings upgraded' policy has the potential to lead to worse indoor air quality in homes in Greater London if energy efficiency is not supported by provision of adequate compensatory ventilation. In the assumptions made for this analysis, the reduction in mortality from lower outdoor PM<sub>2.5</sub> are more than offset by home energy efficiency-related increases in indoor PM<sub>2.5</sub> derived from indoor sources. Radon-related mortality was also estimated to increase because of the reduced ventilation to homes. Overall, heat related mortality was also estimated to increase because of the nature of the interventions. However, the reduction in deaths due to cold was estimated to be appreciably larger than the increase in heat-related mortality. Energy efficiency installations can be net beneficial for health if ventilation is protected and steps taken to address overheating. Installation of two million heat pumps for homes ('Heat-pump' policy) would reduce Greater London's GHG emissions substantially (contingent on grid decarbonisation) but provide relatively modest benefits for health by reducing ambient air pollution. The 'Heat networks' and 'Green gas supply' policies would have more limited GHG reduction and health benefits.

Our estimates of the impact of increasing greenspace in Greater London from 47% to 50% (the 'Greenspace' policy) would have only relatively minor impact on GHG emissions and mortality, given the limited pathways to health included in the calculations. Making Greater London a zero-waste city would reduce GHG emissions by about 1–2%. Overall, the ten policies reduce outdoor PM<sub>2.5</sub> concentrations in Greater London by around 12% (excluding the effects of policy 3 on indoor exposures), which would reduce annual PM<sub>2.5</sub>-attributable premature mortality by around 300 deaths. *Extended data*, Supplementary File 2 ([Symonds et al., 2020b](#)) shows the results for each policy separately, ignoring any potential interactions.

### Discussion

The CRAFT tool is designed to be a decision-support tool that provides approximate, initial comparison of policy choices




**Figure 5.** (a) Central estimates of percentage changes in GHGs emissions (i.e. CO<sub>2</sub>e), air pollution (outdoor PM<sub>2.5</sub>, outdoor NO<sub>2</sub>), indoor environmental quality (indoor radon, indoor PM<sub>2.5</sub> (indoor sources), indoor PM<sub>2.5</sub> (outdoor sources)), and transport (travel-related physical activity) due to the 10 policy actions in London (policy IDs are shown in colour). Left-hand side represent decrease (values shown in minus) and right-hand side represent increase in % changes due to policy. (b) Central estimates of absolute degree changes of ‘Overheating’ and ‘Cold’ due to one policy action in London. \*Please see Figure 6.

in terms of changes to GHG emissions and mortality. Its results are based on approximate methods of calculation and are not intended to be a substitute for more detailed modelling that may be needed to examine specific policy choices in depth. However, its relatively approximate results do allow a quick comparison of multiple policy choices in terms of CO<sub>2</sub>e and mortality impacts, and thus have a potential role in the early phase of policy development. The results for our first analysis of selected environmental policies for Greater London show that there are large differences between major policy choices and that in some cases (as for housing) also important uncertainties about the health impacts depending on the detail of the policy implementation.

How valuable the analysis is to decision-makers remains to be determined in future evaluations. However, the key strengths of the CRAFT tool appear to be assessing the health impacts of city policies rapidly at an early stage to support the decision-making process and to help prioritise different actions.

Furthermore, the tool allows exploration of how policies can be modified so as to provide opportunities for increasing physical activities and mitigation of environmental hazards. The tool, however, has a number of limitations, specifically as regards validation and uncertainty assessment. Uncertainties arise at every stage in the analysis from the initial issue framing, through data collection, assumptions, and modelling to the interpretation and reporting of the results. In a rapid assessment such as this, detailed quantification of uncertainties is not possible. Complex dynamic models have the capability to assess the uncertainties in much greater detail.

Using the CRAFT tool, we estimate the percentage changes in GHGs emissions, air pollution (outdoor PM<sub>2.5</sub>, outdoor NO<sub>2</sub>), indoor environmental quality (indoor radon, indoor PM<sub>2.5</sub> (indoor source), indoor PM<sub>2.5</sub> (outdoor source)), and transport behaviours (travel-related physical activity) that may be achieved by ten key environmental policy actions in Greater London. Of these policies, the highest reduction in GHG



**Figure 6. Estimated impact on health due to the 10 key policy actions in London (the policy IDs are shown in colour).** \* Notice that the impacts for NO<sub>2</sub> has not been added to the PM<sub>2.5</sub> and thus to the total. We show the NO<sub>2</sub> results alongside those for PM<sub>2.5</sub> for completeness and to indicate the uncertainties involved. On the right-hand side is the decrease in number of deaths (avoided deaths) and on the left-hand side the increase in number of deaths due to 10 key policy actions.

emissions is 34% due to the policy action ‘Grid decarbonisation’. The lowest reduction, less than 1%, is due to the policy ‘Greenspace’. Implementing all ten policies results in a GHG reduction of around 90%, which is broadly in line with that projected in Greater London’s zero carbon plan (GLA, 2018d). It should be noted that these proportional reduction in emissions do not account for GHG emissions related to the manufacturing and transport of imported goods or due to projected population rises.

Regarding air pollution, the highest reduction in PM<sub>2.5</sub> is 5% and in NO<sub>2</sub> is 42% both due to the policy action ‘E-transport system’. Increasing the proportion of trips made by foot, cycle or public transport (‘Active travel’) reduces CO<sub>2</sub> emissions by 5%, PM<sub>2.5</sub> by 2%, and NO<sub>2</sub> by 9% and increases the level of transport related physical activity by 33%. This would have considerable benefit for public health, with about 1,700 annual premature deaths avoided, primarily through increased levels of transport-related physical activity.

In this study, health impact calculations have been performed for mortality only and therefore represent underestimates of the full positive impacts on health and quality of life that may be achieved. In some cases, impacts on morbidity may add appreciably to those on mortality; for example, recent evidence has suggested an effect of NO<sub>2</sub> on asthma incidence in children (Achakulwisut *et al.*, 2019; Bergmann *et al.*, 2020). There may also be additional environmental hazards and pathways to health which we have not identified or been able

to quantify. Nonetheless, we believe that the estimates we have generated indicate the main health impacts of different policies and broadly their relative scales.

Our method estimates the theoretical number of premature deaths per year attributable to each exposure at present levels and under an alternative scenario in which the policy has been implemented in full. However, the maximum health benefit achieved may be delayed in time due to a number of factors. This includes the rate of policy implementation and uptake (for example, how quickly the policy reduces the exposure or changes behaviours) and lag times between reductions in exposure and reductions in health risk. It is therefore not appropriate to add or multiply the annual estimates to derive estimates over longer time periods. These issues can be investigated in greater depth using more sophisticated modelling tools (including data-driven modelling) being developed through the CUSSH project. For example, a microsimulation model, MicroEnv, developed recently aimed at quantifying the impact of environmental exposures on health, outputs temporal health impacts at a high spatial resolution (Symonds *et al.*, 2019a).

In the current implementation of the tool, we have estimated health effects resulting from exposure to NO<sub>2</sub> for transport policy actions only. The methods for NO<sub>2</sub> health impact calculation remain uncertain and somewhat controversial. NO<sub>2</sub> and PM<sub>2.5</sub> tend to be highly correlated and it is difficult to separate their effects. Even from mutually adjusted two-pollutant models, it is not clear how much of apparent NO<sub>2</sub> effect is

causally independent of PM: PM<sub>2.5</sub> coefficients probably capture some, though not all, of the NO<sub>2</sub> effects, and vice versa (COMEAP, 2018). The more established method of mortality calculation remains that based on PM<sub>2.5</sub>, and it is this that we present as the main analysis. The NO<sub>2</sub> results are shown for comparison only and cannot be added to those of PM<sub>2.5</sub>. The NO<sub>2</sub> results are shown to indicate an alternative method of computation in which NO<sub>2</sub> is being used as an indicator of NO<sub>2</sub> and the pool of other causal pollutants with which it is correlated. We show the NO<sub>2</sub> results alongside those for PM 2.5 for completeness and to indicate the uncertainties involved. NO<sub>2</sub>-based calculations tend to give contrasting results for transport interventions because of the relative importance of transport sector emissions to ambient NO<sub>2</sub> concentrations.

As can be seen, estimates of the health impact of transport-related interventions are generally much larger when based on NO<sub>2</sub> rather than PM<sub>2.5</sub>, which is important for consideration of actions aimed at the transport sector. It still remains that estimates based on PM<sub>2.5</sub> are more widely accepted among epidemiologists and are the ones that we suggest as the principal estimates of air pollution-related health impact.

The fact that our calculations of the impact of green space were based only on estimates of change in ambient air quality is a limitation and may mean appreciable under-estimation of the potential benefits of interventions that increase green space. The possible connections between green space and health are multiple and complex but depend on the type and quality of green space and where it is implemented (Houlden *et al.*, 2018; Markevych *et al.*, 2017). A recent review (Twohig-Bennett & Jones, 2018) suggests that mortality is lower in areas with greater green space, and there is evidence that inequalities in mortality risk are lower in greener areas (Mitchell & Popham, 2008). However, it is far from certain that these associations can be translated into mortality benefits by intervention, especially when the form of new green space is not specified. It is possible that the proposed changes in green space in Greater London (as defined in terms of a satellite view) will have only a marginal effect on meaningful exposures for mortality that are not accounted for through other pathways (e.g. travel behaviour). But there could be an appreciable effect that we do not represent in the tabulations we present in this study.

The calculations for the 'Buildings upgraded' policy (*Up to 50% of buildings upgraded by 2025, 100% upgraded by 2050*), see *Extended data*, Supplementary Files 1 and 2 (Symonds *et al.*, 2020a; Symonds *et al.*, 2020b), for full details, represent our central estimate of the effects of home energy efficiency upgrades on health. Recent evidence regarding the provision of ventilation following retrofitting suggests that, if current practice continues, energy efficiency upgrades will most likely reduce levels of home ventilation and increase exposures to pollutants generated inside the home, as well as increasing over-heating related deaths (Shrubsole, *et al.*, 2015; Symonds, *et al.*, 2019b). Although there will be reduced exposures to outdoor-generated air pollution and cold-related deaths, our

estimates suggest the net impact on public health will be negative. A more substantial health benefit may be achieved if home energy efficiency measures are installed alongside ventilation that complies with UK building regulations, which would result in a net positive effect on health overall. Properly implemented, such measures can also provide increased protection against heat.

In this study, we focus on the mortality burden associated with exposures related to ten environmental policies in Greater London. There are, however, several other exposures that can be explored and their health impacts quantified in future work. This includes, quantifying road traffic injuries/deaths resulting from transport mode, noise (outdoor and indoor activity noise), and effects on mortality and morbidity relating to access to green space. Changes in green space in particular are likely to have appreciable effects additional to those quantified in this report that were confined to air quality changes alone. There are however challenges in doing such quantification, which ideally, needs to take account of the specific forms of green space change and a range of different health effects.

## Conclusion

We have developed a new rapid assessment tool (CRAFT), *Extended data*, Supplementary File 3 (Symonds *et al.*, 2020c), to quantify the effects of city policies on reducing GHG emissions, decreasing environmental health hazards and improving public health. With relatively simple assumptions and methods, the model can show appreciable differences in expected impact of different policy options. The tool has potential value for policy makers through providing quantitative estimates of health impacts to support and prioritise policy options, especially for the early phase of policy development. The assessment of the tool's utility to decision-makers will be an important next step. That assessment should consider how the policies to be compared are selected, the form of evidence presentation, the stakeholder groups involved in the process and the processes of (iterative) engagements with them.

## Data availability

### Underlying data

All data underlying the results are available as part of the article and no additional source data are required.

### Extended data

Figshare: A tool for assessing the climate change mitigation and health impacts of environmental policies. <https://doi.org/10.6084/m9.figshare.13011533.v3> (Symonds *et al.*, 2020a).

Supplementary File 1. This file contains assumptions; calculation steps; central estimates of percentage changes in GHGs emissions, indoor and outdoor air pollutants, housing-related risks, and travel-related physical activity due to ten key policy actions in Greater London.

Figshare: A tool for assessing the climate change mitigation and health impacts of environmental policies. <https://doi.org/10.6084/m9.figshare.13011539.v4> (Symonds *et al.*, 2020b)

Supplementary File 2. This file contains estimated impact on health (i.e. theoretical premature number of deaths avoided in one year) and CO<sub>2</sub> emissions due to ten key policy actions in Greater London without interactions between policies.

Figshare: A tool for assessing the climate change mitigation and health impacts of environmental policies. <https://doi.org/10.6084/m9.figshare.13125407.v3> (Symonds *et al.*, 2020c).

Supplementary File 3. The file contains a 'Cities Rapid Assessment Framework for Transformation (CRAFT)' tool (Excel format) which quantifies the effects of environmental policies on

changes in (1) greenhouse gas (GHG) emissions, (2) exposures to environmental hazards, (3) travel-related physical activity, and (4) mortality (the number of attributable deaths avoided in one typical year).

Extended data are available under the terms of the [Creative Commons Attribution 4.0 International license](#) (CC-BY 4.0).

## Acknowledgments

We would like to thank all colleagues within the CUSSH consortium who have helped support this work.

## References

- Achakulwisut P, Brauer M, Hystad P, *et al.*: **Global, national, and urban burdens of paediatric asthma incidence attributable to ambient NO<sub>2</sub> pollution: estimates from global datasets.** *Lancet Planet Health.* 2019; **3**(4): e166–e178.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Arem H, Moore SC, Patel A, *et al.*: **Leisure time physical activity and mortality: a detailed pooled analysis of the dose-response relationship.** *JAMA Intern Med.* 2015; **175**(6): 959–67.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Batty M: **The size, scale, and shape of cities.** *Science.* 2008; **319**(5864): 769–771.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- BEIS: **Road Transport Emission Factors: 2010 NAEI.** London, UK: Department for Business, Energy & Industrial Strategy. 2012.  
[Reference Source](#)
- BEIS: **Green Book Supplementary Guidance: Valuation of Energy Use and Greenhouse Gas Emissions for Appraisal.** London, UK: Department for Business, Energy & Industrial Strategy. 2018.  
[Reference Source](#)
- Bergmann S, Li B, Pilot E, *et al.*: **Effect modification of the short term effects of air pollution on morbidity by season: A systematic review and meta-analysis.** *Sci Total Environ.* 2020; **716**: 136985.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Bettencourt LMA, Lobo J, Helbing D, *et al.*: **Growth, innovation, scaling, and the pace of life in cities.** *Proc Natl Acad Sci U S A.* 2007; **104**(17): 7301–7306.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Belesova K, Hale J, Aplin J, *et al.*: **Complex Urban Systems for Sustainability and Health: A structured approach to support the development and implementation of city policies for population and planetary health.** In: *Proceedings of WHO International Healthy Cities Conference - 2018.* WHO: Belfast, Ireland. 2018.  
[Reference Source](#)
- Briggs D, Mason K, Borman B: **Rapid assessment of environmental health impacts for policy support: the example of road transport in New Zealand.** *Int J Environ Res Public Health.* 2015; **13**(1): ijerph13010061.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Briggs DJ: **A framework for integrated environmental health impact assessment of systemic risks.** *Environ Health.* 2008; **7**: 61.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Caplin A, Ghandehari M, Lim C, *et al.*: **Advancing environmental exposure assessment science to benefit society.** *Nat Commun.* 2019; **10**(1): 1236.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- COMEAP (Committee on the Medical Effects of Air Pollutants): **Associations of long-term average concentrations of nitrogen dioxide with mortality.** London, UK: Report of the Committee On the Medical Effects of Air Pollutants. PHE publishing gateway No. 2018238. 2018.  
[Reference Source](#)
- Cucca R: **The unexpected consequences of sustainability: green cities between innovation and ecogentrification.** *Sociologica.* 2012; **6**(2): 1–21.  
[Publisher Full Text](#)
- Davenport C, Mathers J, Parry J: **Use of health impact assessment in incorporating health considerations in decision making.** *J Epidemiol Commun Health.* 2006; **60**(3): 196–201.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- DECC: **The Potential and Costs of District Heating Networks.** London, UK: Department for Energy and Climate Change. 2009.  
[Reference Source](#)
- DEFRA: **Emissions of Air Pollutants in the UK, 1970 to 2017.** London, UK: Department for Environment, Food and Rural Affairs. 2019.  
[Reference Source](#)
- European Environmental Agency (EEA): **Specific CO<sub>2</sub> emissions per passenger-km and per mode of transport in Europe.** 2015.  
[Reference Source](#)
- Forouzanfar MH, Alexander L, Anderson HR, *et al.*: **Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks in 188 countries, 1990–2015: a systematic analysis for the Global Burden of Disease study 2015.** *Lancet.* 2017; **388**(10053): 1659–1724.
- Gascon M, Rojas-Rueda D, Torrico S, *et al.*: **Urban policies and health in developing countries: the case of Maputo (Mozambique) and Cochabamba (Bolivia).** *Public Health Open J.* 2016b; **1**(2): 24–31.  
[Publisher Full Text](#)
- Gascon M, Triguero-Mas M, Martínez D, *et al.*: **Green space and mortality: a systematic review and meta-analysis.** *Environ Int.* 2016a; **2**(86): 60–67.
- GLA: **Valuing London's Urban Forest.** London, UK: Greater London Authority. 2015.  
[Reference Source](#)
- GLA: **The London Energy and Greenhouse Gas Inventory (LEGGI).** London, UK: Greater London Authority. 2016a.  
[Reference Source](#)
- GLA: **London Atmospheric Emissions Inventory (LAEI) 2013 Air Quality Focus Areas - December 2016 update.** London, UK: Greater London Authority. 2016b.  
[Reference Source](#)
- GLA: **Greenhouse Gas Emissions Performance Standard for London's Local Authority Collected Waste - 2017 Update.** London, UK: Greater London Authority. 2017.  
[Reference Source](#)
- GLA: **London Environment Strategy.** London, UK: Greater London Authority. 2018a.  
[Reference Source](#)
- GLA: **Mayor's Transport Strategy.** London, UK: Greater London Authority. 2018b.  
[Reference Source](#)
- GLA: **London Housing Strategy.** London, UK: Greater London Authority. 2018c.  
[Reference Source](#)
- GLA: **Zero Carbon London: A 1.5°C Compatible Plan.** London, UK: Greater London Authority. 2018d.  
[Reference Source](#)
- GLA: **London's Zero Carbon Pathways Tool.** (version 9). London, UK: Greater London Authority. 2018e.  
[Reference Source](#)
- GLA: **Solar Action Plan for London.** London, UK: Greater London Authority. 2018f.  
[Reference Source](#)
- Hamilton I, Milner J, Chalabi Z, *et al.*: **Health effects of home energy efficiency interventions in England: a modelling study.** *BMJ Open.* 2015; **5**(4): e007298.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

- Hartig T, Mitchell R, de Vries S, *et al.*: **Nature and health.** *Annu Rev Public Health.* 2014; **35**: 207–228.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Hänninen O, Knol AB, Jantunen M, *et al.*: **Environmental Burden of Disease in Europe: Assessing Nine Risk Factors in Six Countries.** *Environ Health Perspect.* 2014; **122**(5): 439–446.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Houlden V, Weich S, Porto de Albuquerque J, *et al.*: **The relationship between greenspace and the mental wellbeing of adults: A systematic review.** *PLoS One.* 2018; **13**(9): e0203000.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Howard R: **Up in the Air. How to Solve London's Air Quality Crisis: Part 1.** Report by Policy Exchange and Kings College London. 2015.  
[Reference Source](#)
- IFC (International Finance Corporation): **Introduction to Health Impact Assessment.** The World Bank, US. 2009.  
[Reference Source](#)
- Jetté M, Sidney K, Blümchen G: **Metabolic equivalents (METs) in exercise testing, exercise prescription, and evaluation of functional capacity.** *Clin Cardiol.* 1990; **13**(8): 555–65.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Löhmus M, Balbus J: **Making green infrastructure healthier infrastructure.** *Infect Ecol Epidemiol.* 2015; **5**(1): 30082.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Markevych I, Schoierer J, Hartig T, *et al.*: **Exploring pathways linking greenspace to health: Theoretical and methodological guidance.** *Environ Res.* 2017; **158**: 301–17.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mindell J, Sheridan L, Joffe M, *et al.*: **Health impact assessment as an agent of policy change: Improving the health impacts of the mayor of London's draft transport strategy.** *J Epidemiol Commun Health.* 2004; **58**(3): 169–174.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Mitchell R, Popham F: **Effect of exposure to natural environment on health inequalities: an observational population study.** *Lancet.* 2008; **372**(9650): 1655–60.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Mueller N, Rojas-Rueda D, Basagaña X, *et al.*: **Urban and transport planning related exposures and mortality: a health impact assessment for cities.** *Environ Health Perspect.* 2017; **125**(1): 89–96.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- National Research Council (US) Committee on Health Impact Assessment: **Improving Health in the United States: The Role of Health Impact Assessment.** Washington (DC): National Academies Press (US); 2011.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- NHS: **Health Impact Assessment: A Review of Reviews.** Health Development Agency, UK. 2002.  
[Reference Source](#)
- Nieuwenhuijsen MJ, Khreis H, Verlinghieri E, *et al.*: **Participatory quantitative health impact assessment of urban and transport planning in cities: A review and research needs.** *Environ Int.* 2017a; **103**: 61–72.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Nieuwenhuijsen MJ, Khreis H, Triguero-Mas M, *et al.*: **Fifty shades of green: pathway to healthy urban living.** *Epidemiology.* 2017b; **28**(1): 63–71.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Pope CA 3rd, Burnett RT, Thun MJ, *et al.*: **Lung cancer, cardiopulmonary mortality, and long-term exposure to fine particulate air pollution.** *JAMA.* 2002; **287**(9): 1132–41.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Prüss-Ustün A, Mathers C, Corvalán C, *et al.*: **Introduction and Methods: Assessing the Environmental Burden of Disease at National and Global Levels.** Environmental burden of disease series No. 1, WHO: Geneva, Switzerland. 2003.  
[Reference Source](#)
- Shafiea F, Omara D, Karuppannanb S: **Environmental health impact assessment and urban planning.** *Procedia Soc Behav Sci.* 2013; **85**: 82–91.  
[Publisher Full Text](#)
- Shrubsole C, Das P, Milner J, *et al.*: **A tale of two cities: Comparison of impacts on CO<sub>2</sub> emissions, the indoor environment and health of home energy efficiency strategies in London and Milton Keynes.** *Atmos Environ.* 2015; **120**: 100–108.  
[Publisher Full Text](#)
- Ståhl T, Wismar M, Ollila E, *et al.*: **Health in All Policies: Prospects and Potentials.** Ministry of Social Affairs and Health, Finland. 2006.  
[Reference Source](#)
- Symonds P, Milner J, Mohajeri N, *et al.*: **A tool for assessing the climate change mitigation and health impacts of environmental policies: the Cities Rapid Assessment Framework for Transformation (CRAFT).** *figshare.* Journal contribution. 2020a.  
<http://www.doi.org/10.6084/m9.figshare.13011533.v4>
- Symonds P, Milner J, Mohajeri N, *et al.*: **A tool for assessing the climate change mitigation and health impacts of environmental policies: the Cities Rapid Assessment Framework for Transformation (CRAFT).** *figshare.* Journal contribution. 2020b.  
<http://www.doi.org/10.6084/m9.figshare.13011539.v3>
- Symonds P, Milner J, Mohajeri N, *et al.*: **A tool for assessing the climate change mitigation and health impacts of environmental policies: the Cities Rapid Assessment Framework for Transformation (CRAFT).** *figshare.* Journal contribution. 2020c.  
<http://www.doi.org/10.6084/m9.figshare.13125407.v3>
- Symonds P, Hutchinson E, Ibbetson A, *et al.*: **MicroEnv: A microsimulation model for quantifying the impacts of environmental policies on population health and health inequalities.** *Sci Total Environ.* 2019a; **697**: 134105.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Symonds P, Rees D, Daraktchieva Z, *et al.*: **Home energy efficiency and radon: An observational study.** *Indoor Air.* 2019b; **29**(5): 854–864.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Taylor J, Wilkinson P, Picetti R, *et al.*: **Comparison of built environment adaptations to heat exposure and mortality during hot weather, West Midlands region, UK.** *Environ Int.* 2018; **111**: 287–94.  
[PubMed Abstract](#) | [Publisher Full Text](#)
- Twohig-Bennett C, Jones A: **The health benefits of the great outdoors: A systematic review and meta-analysis of greenspace exposure and health outcomes.** *Environ Res.* 2018; **166**: 628–37.  
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
- Transport for London (TfL): **London Travel Demand Survey, 2018/19.** 2018.  
[Reference Source](#)
- United Nations: **World Urbanization Prospects.** The 2018 Revision, Highlights. 2018.  
[Reference Source](#)
- WHO: **Global Strategy on Diet, Physical Activity and Health.** 2014; (accessed 20/04/2020).  
[Reference Source](#)
- Warburton DER, Bredin SSD: **Health benefits of physical activity: a systematic review of current systematic reviews.** *Curr Opin Cardiol.* 2017; **32**(5): 541–556.  
[PubMed Abstract](#) | [Publisher Full Text](#)

# Open Peer Review

Current Peer Review Status:   

---

## Version 2

Reviewer Report 28 July 2021

<https://doi.org/10.21956/wellcomeopenres.18616.r44920>

© 2021 Saari R. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

 **Rebecca K. Saari** 

Department of Civil and Environmental Engineering, University of Waterloo, Waterloo, Canada

The myriad health-related benefits of GHG reductions are known to be significant. However, they are seldom integrated into policy decisions or designs, especially at the early screening phase. To address this, decision support tools at the city scale are critically needed. I commend the authors on their response to this challenge. They present an accessible, credible tool that helps to integrate these benefits into early decision processes. The methods are appropriate for the order-of-magnitude screening they aim to provide.

Following a careful technical review of the tool, I find the approaches defensible and grounded in solid data and research. They are well documented in the supplemental materials. However, I did wish that some of the key assumptions were better described in the manuscript itself to aid understanding and interpretation of the results. I have highlighted a few key assumptions that can affect the sign of the impact, or the magnitude of the largest impacts. I would encourage the authors to consider minor edits based on my specific suggestions below.

### Comments on assumptions and details for specific methods:

#### Active Travel:

More detail in this paragraph "For transport-related physical activity, we assessed changes in transport-related metabolic equivalent of the task (MET) hours per week..." would be helpful. The authors cite the current METs. I recommend citing the METs under the policy so the magnitude of the change is clear, and it's evident that only travel-related MET increases (as expected). The current wording "based on ... changes in the sum of these two sources of physical activity" implies that leisure related physical activity is also affected by policy.

I am aware that this is a particular detail that will not affect the conclusions of this work, but, for the sake of reproducibility, I also recommend describing the piecewise linear RR developed. The authors mentioned using a piece-wise linear RR developed from Arem et al. (2015), but, unless I am missing something, appear not to have described why or how they did so. They cite an RR of 0.75 for a MET of 11.25, yet Arem et al (2015) provides a piece-wise figure of hazard ratios, citing 0.69 for an MET range of 7.5-15. The authors have apparently performed a linear interpolation

within the categories that goes beyond the original epidemiologic evidence, the results of which depend on how the categories are represented.

#### Building Upgraded:

I recommend, in the Methods text or Table 3, describing the particular assumptions used here earlier on, in addition to mentioning that, “Depending on the assumption about changes in ventilation following retrofit, the net impacts on mortality can be negative or positive.” Otherwise, it is only in the final paragraphs of the Discussion (or in a footnote of the supplemental material) that readers learn that this reflects the “Installer Discretion” Scenario from Hamilton et al. (2015). This work also expands on Hamilton et al (2015) by including changes in outdoor air quality, another detail that could be mentioned, e.g., in Table 3, to better clarify the relation of this work to its sources.

#### Zero Waste City:

As a carbon policy, the literature is clear that zero waste / circular economy policies are valuable because they reduce carbon-intensive production of goods to meet demand. Scope 1 and 2 emissions associated with city-scale waste treatment are minor compared to the embodied carbon associated with the consumption driving that waste. The authors allude to this when mentioning the primacy of London’s Scope 3 emissions. I recommend that they also explicitly discuss how excluding these emissions can undervalue zero waste policies. While the system boundary chosen in CRAFT is clear and defensible, it will dictate the relative attractiveness of this option to decision makers.

#### **Other Specific comments:**

“what level of change in exposure or behaviour (physical activity) of an individual within the affected population as a result of the policy.” This phrase seems to be missing a verb since “change” and “result” are both used here as nouns... suggest replacing “as a result of” with “would result from”

#### Figure 1:

Can you please, when introducing Figure 1, clarify whether this is the ideal or actual flow diagram for the tool as described in this manuscript? The Figure suggests that stakeholders were involved throughout, but conclusions indicate that stakeholder participation is a next step – an apparent contradiction.

I found this wording confusing: “The overall impact on GHG emissions for a specific policy is then computed by multiplying relevant fractions (e.g. the fraction of emissions attributable to the target sector × the fraction of the target that will be changed as a result of the policy × the fractional reduction in emissions per unit of the target consequent to the policy action).” There appears to be a mid-sentence change in referring to a “target sector” and the “target of action”, but “target consequent to the policy action” is ambiguous and could still refer to a target sector. Possible alternative: replace “the fractional reduction in emissions per unit of the target consequent to the policy action” with “fraction of emissions reduced per unit change in the target of action”.

#### Results

“Extended data, Supplementary File 1 (Symonds et al., 2020a) presents the detailed calculation steps for each policy, along with assumptions used for current and minimum exposures and the



exposure-mortality coefficients and population mortality statistics used.”

Please introduce this early in the Methods section, e.g., near, “The precise details of how this principle is applied vary somewhat from policy to policy but the general principle entails estimating the fraction of total GHG emissions that are attributable to a particular sector (transport, housing, etc.) and then to the relevant target of action within each sector.” Also, please check the link here. The version linked (v4) threw a “DOI Not Found” error. For the purposes of my review, I used 10.6084/m9.figshare.13011533.v3 instead

#### Conclusions:

“new rapid assessment tool (CRAFT), *Extended data*, Supplementary File 3 ([Symonds et al., 2020c](#)),” Please reference the supplemental link to the tool on first mention of CRAFT in the manuscript.

#### **Minor Typos/wording:**

##### Abstract:

“including detailed policies” recommend replace with “and accompanying detailed policies” to better distinguish the roles of policy targets and policy instruments  
Suggest replacing “magnitude of impact... to reach” with “magnitude of impact... toward reaching”

##### Results

In “on the detail of the policy implementation” replace “detail” with “details”

##### Discussion:

Add “the” before “apparent NO2 effect”

Punctuation is odd, here, in its separation of the subject “calculations” from the verb “represent”.  
“The calculations for the ‘Buildings upgraded’ policy (*Up to 50% of buildings upgraded by 2025, 100% upgraded by 2050*), see *Extended data*, Supplementary Files 1 and 2 ([Symonds et al., 2020a](#); [Symonds et al., 2020b](#)), for full details, represent...”

Recommend instead:

“The calculations for the ‘Buildings upgraded’ policy (*Up to 50% of buildings upgraded by 2025, 100% upgraded by 2050*; see *Extended data*, Supplementary Files 1 and 2 ([Symonds et al., 2020a](#); [Symonds et al., 2020b](#)) for full details) represent...”

##### CRAFT tool

In the tab “Indoor Air Quality” cell J6 appears mislabeled. It should read Radon not PM2.5.

Suggest changing wording in cell G6 to reference outdoor-source PM2.5 to better distinguish it from cell D6 and map with other supplemental material.

#### **Is the work clearly and accurately presented and does it cite the current literature?**

Yes

#### **Is the study design appropriate and is the work technically sound?**

Yes

#### **Are sufficient details of methods and analysis provided to allow replication by others?**

Yes

**If applicable, is the statistical analysis and its interpretation appropriate?**

Yes

**Are all the source data underlying the results available to ensure full reproducibility?**

Yes

**Are the conclusions drawn adequately supported by the results?**

Yes

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Integrated modelling of air pollution and associated health and economic impacts under climate change and climate policy.

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

Reviewer Report 14 July 2021

<https://doi.org/10.21956/wellcomeopenres.18616.r43953>

© 2021 Thurston G. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**George D. Thurston** 

Department of Environmental Medicine, New York University School of Medicine, New York City, NY, USA

Nothing to add.

**Is the work clearly and accurately presented and does it cite the current literature?**

Yes

**Is the study design appropriate and is the work technically sound?**

Yes

**Are sufficient details of methods and analysis provided to allow replication by others?**

Yes

**If applicable, is the statistical analysis and its interpretation appropriate?**

Yes

**Are all the source data underlying the results available to ensure full reproducibility?**

Yes

**Are the conclusions drawn adequately supported by the results?**

Yes

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** air pollution exposure assessment and epidemiology

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

Reviewer Report 19 May 2021

<https://doi.org/10.21956/wellcomeopenres.18616.r43952>

© 2021 Kinney P. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Patrick L. Kinney** 

Department of Environmental Health, Boston University School of Public Health, Boston, MA, USA

No further comments, I approve the article.

**Is the work clearly and accurately presented and does it cite the current literature?**

Yes

**Is the study design appropriate and is the work technically sound?**

Yes

**Are sufficient details of methods and analysis provided to allow replication by others?**

Yes

**If applicable, is the statistical analysis and its interpretation appropriate?**

Yes

**Are all the source data underlying the results available to ensure full reproducibility?**

Yes

**Are the conclusions drawn adequately supported by the results?**

Yes

**Competing Interests:** No competing interests were disclosed.

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

**Version 1**

Reviewer Report 02 March 2021

<https://doi.org/10.21956/wellcomeopenres.17972.r42550>

© 2021 Thurston G. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**George D. Thurston**

Department of Environmental Medicine, New York University School of Medicine, New York City, NY, USA

Nations are doing poorly at achieving the climate forcing emission reductions needed to avoid a climate calamity. Even if the world's countries follow through on present plans, their combined impacts would put us on a path to achieve only a 1 % reduction in global emissions by 2030, while climate scientists tell us that we need more than that each and every year. It is clear that more progress is needed, and action by individual cities are one pathway to jump start action, but motivation is needed.

In "A tool for assessing the climate change mitigation and health impacts of environmental policies: the Cities Rapid Assessment Framework for Transformation (CRAFT)" Symonds *et al* provide a much needed city-level tool for the documentation of the potential local health benefits of climate mitigation measures, and their monetary valuations. The methods incorporated in the framework, while straightforward, are useful and consistent with present cost-benefit approaches.

I do have some concerns with the specifics, however. The estimates provided lack confidence intervals, as acknowledged by the authors, but which would be important for decision-makers to consider. In addition, the NO<sub>2</sub> mortality estimates (though not used in the final estimates) are not as confident as the PM<sub>2.5</sub> estimate, as the causality of that relationship has weaker causal foundation. The original COMEAP Report that is referenced, it is stated that: "A majority view of the Committee was that approximately 25-55% of the unadjusted summary coefficient derived from the meta-analysis of single pollutant models (1.023 per 10 µg/m<sup>3</sup> NO<sub>2</sub>) could be regarded as causally related to NO<sub>2</sub>. This gives a range of between 1.006 per 10 µg/m<sup>3</sup> and 1.013 per 10 µg/m<sup>3</sup>", not the 1.023 per 10 µg/m<sup>3</sup> used in the analysis. A smaller NO<sub>2</sub> estimate would be more defensible, but I would recommend dropping the NO<sub>2</sub> estimates from the report.

**Is the work clearly and accurately presented and does it cite the current literature?**

Yes

**Is the study design appropriate and is the work technically sound?**

Yes

**Are sufficient details of methods and analysis provided to allow replication by others?**

Yes

**If applicable, is the statistical analysis and its interpretation appropriate?**

Partly

**Are all the source data underlying the results available to ensure full reproducibility?**

Yes

**Are the conclusions drawn adequately supported by the results?**

Yes

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** air pollution exposure assessment and epidemiology

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.**

Reviewer Report 23 February 2021

<https://doi.org/10.21956/wellcomeopenres.17972.r42483>

© 2021 Kinney P. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Patrick L. Kinney** 

Department of Environmental Health, Boston University School of Public Health, Boston, MA, USA

This paper documents the structure and function of a new tool that has been developed to assist cities in weighing the climate and health benefits of various interventions. It is meant to be a rapid assessment tool that can be readily applied by non-scientists. I find this very solid and well supported. I did have a slight concern about the treatment of greenspace benefits, for which only a small subset of health benefit pathways are quantified. The authors fully acknowledge this in the discussion. However, I wonder if it's better to leave greenspace out of the example rather than to leave the impression that the benefits are so low. Alternatively, could include a sensitivity analysis using the exposure-response function from a recent meta analysis, such as *Lancet Planet Health* 2019;3: 469–77<sup>1</sup>.

Minor comment: second paragraph of page 12. First sentence. The parentheses seem mis-placed.

### References

1. Rojas-Rueda D, Nieuwenhuijsen M, Gascon M, Perez-Leon D, et al.: Green spaces and mortality: a systematic review and meta-analysis of cohort studies. *The Lancet Planetary Health*. 2019; **3** (11): e469-e477 [Publisher Full Text](#)

**Is the work clearly and accurately presented and does it cite the current literature?**

Yes

**Is the study design appropriate and is the work technically sound?**

Yes

**Are sufficient details of methods and analysis provided to allow replication by others?**

Yes

**If applicable, is the statistical analysis and its interpretation appropriate?**

Not applicable

**Are all the source data underlying the results available to ensure full reproducibility?**

Yes

**Are the conclusions drawn adequately supported by the results?**

Yes

**Competing Interests:** No competing interests were disclosed.

**Reviewer Expertise:** Health effects of air pollution and climate change. Health benefits of carbon mitigation strategies.

**I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.**

---