

Effects of an Emergency-Based FASE Strategy on Treating Geriatric Patients with Femoral Neck Fracture: A Retrospective Propensity Score-Matched Study [Letter]

Hejia Wan , Shuning Wang , Lanxia Pan 

School of Nursing, Henan University of Chinese Medicine, Zhengzhou, People's Republic of China

Correspondence: Lanxia Pan, Email panlanx@163.com

Dear editor

We have read with great interest the recent publication by Zhou et al, which presents the effects of the Fast Access to Surgery in Emergency (FASE) strategy on the treatment of geriatric patients with femoral neck fractures.¹ The authors argue that the FASE strategy significantly optimizes the preoperative evaluation workflow, shortens time to surgery and hospital stay, and reduces perioperative blood loss. While this constitutes an important contribution to the field of orthogeriatric care, we have identified some aspects that warrant further consideration.

First, the authors claimed that the FASE strategy significantly reduced operative time and hospitalization. However, we note that the retrospective design and single-center implementation of the study may limit the generalizability of these findings, although this is noted in the limitations of the paper. We suggest that the authors implement more comparisons, such as for relevant data or findings in other papers,²⁻⁴ which would help the generalizability of this paper. Further, the potential impact of confounding variables, such as level of care⁵ and resource availability (for example long-term care insurance) across centers,⁶ was not fully explored and is an issue that needs to be addressed.

Secondly, in the analysis of postoperative complications and mortality rates, we discovered that there were no significant differences between the FASE group and the control group. This finding contrasts with the expected benefits of an accelerated surgical approach, as per the Enhanced Recovery After Surgery (ERAS) principles, which the FASE strategy is based upon. Further investigation into the reasons for this lack of significant difference is necessary to better understand the implications of the FASE strategy on patient outcomes.

In summary, we appreciate the importance of the study in highlighting the potential benefits of the FASE strategy in optimizing surgical workflows for geriatric femoral neck fracture patients. However, we hope that future research will consider the broader context of patient care, including the impact of socioeconomic factors and the variability in healthcare settings, to provide a more comprehensive understanding of the FASE strategy's effectiveness (Figure 1). More organismal mechanisms need to be further elucidated, in cells or tissues, psychologically, and so on. We look forward to the authors' further exploration of these issues. It is our hope that a more nuanced consideration of the complexities involved in geriatric orthopedic care will lead to even more robust research and ultimately contribute to improved patient outcomes. We are eager to see how these insights will be integrated into future studies, potentially leading to a more refined and effective implementation of the FASE strategy in the management of geriatric hip fractures.

Data Sharing Statement

The present study did not involve the generation or analysis of any datasets.




Figure I General schematic of the proposal.

Funding

No specific funding was obtained for this letter from public, commercial, or non-profit organizations.

Disclosure

The authors disclose no conflicts of Interest in this communication. Figure was drawn using the figdraw website and is hereby acknowledged. The license code for this image is YTAOWa5b55. This Figure is not subject to copyright dispute.

References

1. Zhou J, Shi L, Tian C, et al. Effects of an emergency-based FASE strategy on treating geriatric patients with femoral neck fracture: a retrospective propensity score-matched study. *Clin Interv Aging*. 2024;19:1867–1880. doi:10.2147/CIA.S485809
2. Jiang M, Liu S, Deng H, Liang X, Bo Z. The efficacy and safety of fast track surgery (FTS) in patients after Hip fracture surgery: a meta-analysis. *J Orthop Surg Res*. 2021;16(1):162. doi:10.1186/s13018-021-02277-w
3. Li H, Liu Y, Li Q, Fan J, Gan L, Wang Y. Effects of a fast track surgery nursing program in perioperative care of older patients with a Hip fracture. *Eur Geriatr Med*. 2020;11(4):519–525. doi:10.1007/s41999-020-00298-y
4. Haugan K, Johnsen LG, Basso T, Foss OA. Mortality and readmission following Hip fracture surgery: a retrospective study comparing conventional and fast-track care. *BMJ Open*. 2017;7(8):e015574. doi:10.1136/bmjopen-2016-015574
5. Geldsetzer P, Manne-Goehler J, Marcus ME, et al. The state of hypertension care in 44 low-income and middle-income countries: a cross-sectional study of nationally representative individual-level data from 1.1 million adults. *Lancet*. 2019;394(10199):652–662. doi:10.1016/S0140-6736(19)30955-9
6. Tamiya N, Noguchi H, Nishi A, et al. Population ageing and wellbeing: lessons from Japan's long-term care insurance policy. *Lancet*. 2011;378(9797):1183–1192. doi:10.1016/S0140-6736(11)61176-8

Dove Medical Press encourages responsible, free and frank academic debate. The content of the Clinical Interventions in Aging 'letters to the editor' section does not necessarily represent the views of Dove Medical Press, its officers, agents, employees, related entities or the Clinical Interventions in Aging editors. While all reasonable steps have been taken to confirm the content of each letter, Dove Medical Press accepts no liability in respect of the content of any letter, nor is it responsible for the content and accuracy of any letter to the editor.

Clinical Interventions in Aging

Dovepress

Publish your work in this journal

Clinical Interventions in Aging is an international, peer-reviewed journal focusing on evidence-based reports on the value or lack thereof of treatments intended to prevent or delay the onset of maladaptive correlates of aging in human beings. This journal is indexed on PubMed Central, MedLine, CAS, Scopus and the Elsevier Bibliographic databases. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/clinical-interventions-in-aging-journal>