

POSTER PRESENTATION

Open Access

Sequential treatment with ipilimumab and BRAF inhibitors in patients with metastatic melanoma: data from the Italian ipilimumab expanded access programme (EAP)

Paolo Antonio Ascierto^{1*}, E Simeone¹, V Chiarion Sileni², P Queirolo³, M Del Vecchio⁴, L Di Guardo⁴, M Guidoboni⁵, P Marchetti^{6,7}, GC Antonini Cappellini⁶, PF Ferrucci⁸, F Cognetti⁹, MG Bernengo¹⁰, M Guida¹¹, R Marconcini¹², M Mandalà¹³, C Cimminiello¹⁴, G Rinaldi¹⁵, M Aglietta¹⁶, L Calabrò¹⁷, M Maio¹⁷

From Society for Immunotherapy of Cancer 28th Annual Meeting
National Harbor, MD, USA. 8-10 November 2013

Background

Data to guide the order in which ipilimumab and vemurafenib are used in patients with advanced melanoma are limited. Here are reported outcomes from patients treated in the ipilimumab EAP who received both drugs.

Methods

Patients with pretreated, BRAFV600 mutation-positive advanced melanoma who had received BRAF inhibitor before or after ipilimumab were eligible for analysis.

Results

93 patients were eligible: 48 patients received a BRAF inhibitor after ipilimumab and 45 patients ipilimumab after a BRAF inhibitor. Median overall survival (OS) was 14.5 and 9.9 months for the two groups, respectively ($P=0.04$). Among patients who received a BRAF inhibitor first, 18 (40%) had rapid disease progression and were unable to complete ipilimumab treatment as for protocol (rapid progressors). For this group median OS from the cessation of treatment with a BRAF inhibitor was 1.2 months. 27 patients had slower disease progression and were able to complete all four doses of ipilimumab (slow progressors); median OS was significantly longer (12.7 months; $P<0.0001$). Younger age and the presence of brain metastasis were significantly associated with a poorer outcome ($P=0.02$).

Conclusions

This EAP data suggests that pretreated, BRAF-mutated patients who have rapid disease progression upon failing treatment with a BRAF inhibitor die in one month, so they may benefit from receiving ipilimumab as the first part of their sequential regimen, otherwise clinical benefit may be limited due to them not being able to receive the full induction treatment.

Authors' details

¹Istituto Nazionale Tumori Fondazione "G. Pascale", Naples, Italy. ²Institute of Oncology IOV-IRCCS, Padua, Italy. ³San Martino Hospital-National Institute for Cancer Research, Genoa, Italy. ⁴National Cancer Institute, Milan, Italy. ⁵Romagna National Cancer Institute, Meldola, Italy. ⁶Dermatopathologic Institute of the Immaculate, Rome, Italy. ⁷Sant'Andrea Hospital, University Sapienza, Rome, Italy. ⁸Istituto Europeo di Oncologia, Milan, Italy. ⁹Regina Elena National Cancer Institute, Rome, Italy. ¹⁰University Hospital St John the Baptist, Turin, Italy. ¹¹National Cancer Research Center "Giovanni Paolo II", Bari, Italy. ¹²Gathered Hospitals of Santa Chiara", Pisa, Italy. ¹³Papa Giovanni XXIII Hospital, Bergamo, Italy. ¹⁴San Raffaele Hospital, Milan, Italy. ¹⁵Paolo Giaccone Polyclinic University Hospital, Palermo, Italy. ¹⁶Piedmont Oncology Foundation, Candiolo, Italy. ¹⁷Hospital of Siena Istituto Toscano Tumori, Siena, Italy.

Published: 7 November 2013

doi:10.1186/2051-1426-1-S1-P69

Cite this article as: Ascierto et al.: Sequential treatment with ipilimumab and BRAF inhibitors in patients with metastatic melanoma: data from the Italian ipilimumab expanded access programme (EAP). *Journal for ImmunoTherapy of Cancer* 2013 **1**(Suppl 1):P69.

¹Istituto Nazionale Tumori Fondazione "G. Pascale", Naples, Italy
Full list of author information is available at the end of the article