



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

**Reply to letter to the editor: “A response to ‘Male balding is a major risk factor for severe COVID-19’”**



*To the Editor:* We appreciate the interest regarding “Male balding is a major risk factor for severe COVID-19”<sup>1</sup> by Thatiparthi et al.<sup>2</sup> These authors propose a potential causal relationship between COVID-19 and scalp balding. They suggest that balding could possibly allow the virus to transverse the cutaneous barrier through accelerated photo-aging, oxidative stress, loss of elasticity, and reduced moisturization. Although the possibility of COVID-19 percutaneous transmissibility is a novel perspective, further investigation is needed to determine if SARS-CoV-2 infects epidermal keratinocytes and whether SARS-CoV-2 skin inoculation leads to systemic disease.

Prevalent in health care workers, hand eczema represents an alternative skin disease for further inquiry into the relationship between COVID-19 and skin barrier disruption. Hand eczema erodes the epidermis thereby shortening the transmission route needed for SARS-CoV-2 binding to ACE2 receptors in the epidermal basal layer. Among tested United Kingdom Biobank participants, 1450 of 18,221 (8.0%) participants tested COVID-19 positive compared with 14 of 139 (10.1%) participants with physician-diagnosed hand eczema.

The common types of balding in men are male pattern balding and senescent balding. Although our study did not differentiate between these 2 types, it is possible that both balding types are associated with COVID-19, as both male sex and advanced age are strong COVID-19 risk factors. In an analysis covering 40% of all patients in England, COVID-19–related death was associated with male sex (hazard ratio [HR], [95% confidence interval], 1.59 [1.53–1.65]) and increasing age (60–69 years; HR, 2.40 [2.16–2.66]; 70–79 years, HR, 6.07 [5.51–6.69]; 80+ years, HR, 20.60 [18.70–22.68]).<sup>3</sup>

The association between COVID-19 and male sex may be related to the androgen-mediated

COVID-19 hypothesis,<sup>4</sup> in which testosterone upregulates transmembrane serine protease 2 (TMPRSS2) facilitating SARS-CoV-2 entry into host cells via ACE2. Poor outcomes of COVID-19 in the elderly are most likely related to the aging immune system’s inability to neutralize virus thereby triggering a pro-inflammatory state in COVID-positive patients. As balding also presents a manifestation of cellular senescence, we suggest that balding may also function as a highly visible biomarker for immunosenescence. Thus, the association between balding and severe COVID-19 may be considered one of the factors that clinicians can use to rapidly identify patients at higher risk.

*Abmed Yousaf, BA, and Michael S. Kolodney, MD, PhD*

*From the Department of Dermatology, West Virginia University, Morgantown, WV.*

*IRB approval status: Not applicable.*

*Reprints not available from the authors.*

*Correspondence to: Michael S. Kolodney, MD, PhD, WVU School of Medicine, Department of Dermatology, PO Box 9158, 1 Medical Center Drive, Morgantown, WV 26506*

*E-mail: [michael.kolodney@hsc.wvu.edu](mailto:michael.kolodney@hsc.wvu.edu)*

**REFERENCES**

1. Lee J, Yousaf A, Fang W, Kolodney MS. Male balding is a major risk factor for severe COVID-19. *J Am Acad Dermatol.* 2020;83:e353–e354.
2. Thatiparthi A, Liu J, Martin A, Wu JJ. A response to “Male balding is a major risk factor for severe COVID-19.” *J Am Acad Dermatol.* 2021;84(2):e87–e88.
3. Williamson EJ, Walker AJ, Bhaskaran K, et al. Factors associated with COVID-19-related death using OpenSAFELY. *Nature.* 2020; 584:430–436.
4. Stopsack KH, Mucci LA, Antonarakis ES, Nelson PS, Kantoff PW. TMPRSS2 and COVID-19: serendipity or opportunity for intervention? *Cancer Disc.* 2020;10:779–782.

<https://doi.org/10.1016/j.jaad.2020.10.052>