


Article

Endosymbiotic RNA virus inhibits *Leishmania*-induced caspase-11 activation



Renan V.H. de Carvalho, Djalma S. Lima-Júnior, Caroline V. de Oliveira, Dario S. Zamboni

dszamboni@fmrp.usp.br

HIGHLIGHTS

LRV partially blocks NLRP3 inflammasome activation by *L.g.* via Casp11

LRV interferes directly with caspase-11 activation by *L.g.*

Caspase-11 blockage by LRV increases *L.g.* replication and disease pathogenesis

TLR3 and ATG5 are required for LRV-mediated inhibition of Casp11

de Carvalho et al., iScience 24, 102004
January 22, 2021 © 2021 The Authors.
<https://doi.org/10.1016/j.isci.2020.102004>



Article

Endosymbiotic RNA virus inhibits *Leishmania*-induced caspase-11 activationRenan V.H. de Carvalho,^{1,2} Djalma S. Lima-Júnior,^{1,3} Caroline V. de Oliveira,¹ and Dario S. Zamboni^{1,4,*}

SUMMARY

New World species of the intracellular protozoan parasites of the *Leishmania* genus can cause mucocutaneous leishmaniasis. The presence of an endosymbiotic *Leishmania* RNA virus (LRV) in *Leishmania guyanensis* (*L.g.*) promotes disease exacerbation and the development of mucocutaneous disease. It was previously reported that LRV blocks the NLRP3 inflammasome, but additional mechanisms remain unclear. Here, we investigated whether LRV interferes with the inflammasome via caspase-11, which induces non-canonical NLRP3 activation and was reported to be activated by *Leishmania*. By using macrophages and mice, we found that LRV inhibits caspase-11 activation and IL-1 β release by *L.g.* in a TLR3- and ATG5-dependent manner. Moreover, LRV exacerbates disease in C57BL/6 mice but not in *Casp11*^{-/-}, *Nlrp3*^{-/-}, and 129 mice, a mouse strain that is naturally mutant for caspase-11. These results demonstrate that LRV interferes with caspase-11 activation by *Leishmania*, expanding our understanding about the mechanisms by which LRV promotes disease exacerbation.

INTRODUCTION

Leishmaniasis comprise a large group of inflammatory insect-borne diseases that may range from self-healing cutaneous lesions to mucocutaneous and visceral outcomes, affecting millions of people worldwide and causing thousands of deaths per year (Hartley et al., 2012; Pigott et al., 2014; Scott and Novais, 2016). Although the host immune response against the parasite has extensively been studied, only recently the specific mechanisms of innate immune recognition began to be unraveled (reviewed in de Carvalho and Zamboni, 2020). Recent studies have shown that caspase-11 is efficiently activated by *Leishmania* parasites to induce non-canonical NLRP3 activation, expanding the importance of caspase-11 to parasitic diseases (de Carvalho et al., 2019a). Furthermore, it was demonstrated that parasites harboring the *Leishmania* RNA virus (LRV) induce less inflammasome activation by a TLR3- and autophagy-dependent mechanism, suggesting that this pathway could be responsible for the debilitating mucocutaneous form of leishmaniasis observed in patients (de Carvalho et al., 2019b; Hartley et al., 2016; Ives et al., 2011; Rossi et al., 2017). However, whether LRV affects caspase-11 and the non-canonical activation of NLRP3 by *Leishmania* remains unknown.

Here, we use mouse bone-marrow-derived macrophages (BMDMs) and mouse models of leishmaniasis to investigate the impact of LRV in non-canonical NLRP3 activation by *Leishmania* parasites. By using the M4147 *L. guyanensis* strain, which harbors high levels of LRV (*L.g.*+), and an M4147-derived LRV negative clone (*L.g.*-) that spontaneously lost LRV (de Carvalho et al., 2019b), we found that LRV blocks caspase-11 activation by *L.g.* We further show that this process is TLR3/ATG5 dependent and leads to inhibition of the non-canonical pathway for NLRP3 activation, thus favoring parasite persistence and increasing disease susceptibility in mice. Collectively, these data advance toward the understanding of the mechanisms by which LRV promotes inhibition of inflammasome activation and disease exacerbation.

RESULTS

***Leishmania* RNA virus (LRV) interferes with inflammasome activation via caspase-11**

NLRP3 activation and overall inflammation is a hallmark of many inflammatory and infectious diseases including leishmaniasis, although the complete mechanisms of inflammasome regulation in response to

¹Departamento de Biologia Celular e Molecular e Bioagentes Patogênicos, Faculdade de Medicina de Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto Medical School, FMRP/USP. Av. Bandeirantes 3900, Ribeirão Preto, SP 14049-900, Brazil

²Present address: Laboratory of Lymphocyte Dynamics, The Rockefeller University, 1230 York Av, New York, NY 10065, USA

³Present address National Institute of Allergy and Infectious Diseases, NIH, Bethesda, MD 20892, USA

⁴Lead contact

*Correspondence: dszamboni@fmrp.usp.br
<https://doi.org/10.1016/j.isci.2020.102004>






Figure 1. *Leishmania* RNA virus (LRV) inhibits the non-canonical activation of NLRP3 by *L.g.*

(A) Western blot (WB) analysis showing pro-caspase-1 (45 kDa) and its cleaved form (20 kDa), generated by inflammasome activation, in cell-free supernatants from LPS-primed wild-type C57BL/6 and *Casp11*^{-/-} macrophages, that were infected with either stationary phase (SP) *L. guyanensis* without the virus (*L.g.-*) or with the virus (*L.g.+*), at MOI (multiplicity of infection) 10, and processed for WB.

(B–D) Flow cytometry analysis of active caspase-1 staining (using FAM-YVAD fluorescent dye) in BMDMs from wild-type C57BL/6, *Nlrp3*^{-/-}, and *Casp11*^{-/-} mice, 24 hr after infection with SP *L.g.-* and *L.g.+* at MOI 10. A representative contour plot (B), the percentage of active caspase-1+ cells (FAM-YVAD+) (C), and the integrated mean of fluorescence intensity (iMFI) (D) are shown.

(E) ELISA assay quantifying IL-1β production in cell-free supernatants from LPS-primed wild-type C57BL/6, *Nlrp3*^{-/-}, and *Casp11*^{-/-} BMDMs, 24 hr after infection with SP *L.g.-* and *L.g.+* at MOI 10.

(F) ELISA assay quantifying IL-1β production in cell-free supernatants from LPS-primed wild-type C57BL/6 and *Casp11*^{-/-} BMDMs, 24 hr after infection with SP *L.g.-* and *L.g.+* at MOI 10 or after 20 hr of infection with parasites and 4 hr of infection with *Legionella pneumophila* lacking flagellin (*flaA*⁻).

The results are shown as mean ± SD, and statistical analysis was performed by unpaired Student's *t* test (C–F). *, *p* < 0.05. SD, standard deviation. One representative of at least three independent experiments performed with technical replicates is shown.

infection by *Leishmania* parasites are still under intense investigation (reviewed in Broz and Dixit, 2016; de Carvalho and Zamboni, 2020; Zamboni and Sacks, 2019). In this context, LRV emerged as an important modulator of innate immunity during *Leishmania* infection since it was shown to trigger TLR3 activation and negatively regulate NLRP3 assembly by the parasite (de Carvalho et al., 2019b; Hartley et al., 2016; Ives et al., 2011; Rossi et al., 2017). Importantly, studies from our and other labs recently demonstrated that *Leishmania* is capable of triggering caspase-11 activation (Chaves et al., 2019; de Carvalho et al., 2019a), a process that was previously reported to occur via bacterial lipopolysaccharide (LPS) (Hagar et al., 2013; Kayagaki et al., 2013). Thus, we decided to investigate whether LRV could exert any impact on caspase-11 and non-canonical inflammasome activation by *L. guyanensis*. To address this question, we initially infected wild-type (WT), *Nlrp3*^{-/-}, and *Casp11*^{-/-} BMDMs with both *L.g.+* and *L.g.-* and measured caspase-1 activation and IL-1β production. Western blotting for cleaved caspase-1 (Figure 1A), flow cytometric analysis by a fluorescent dye that binds specifically to active caspase-1 (Figures 1B–1D), and quantification of IL-1β levels by ELISA (Figure 1E) revealed that LRV blocks inflammasome activation in a




Figure 2. LRV interferes with caspase-11 activation in macrophages infected by *L. guyanensis*

C57BL/6 BMDMs were primed with PAM(3)CSK(4) (300 ng mL^{-1}) (A) or TNF- α (10 ng mL^{-1}) (B) for 4 hr and then infected with stationary phase (SP) *L. guyanensis* without the virus (L.g.-) or with the virus (L.g.+) at MOI 10. Active caspase-11 (45 and 38 kDa) was pulled down, and Western blotting of active (pull-down) or total (input, cell lysates) caspase-11 was performed. β -actin was used as a loading control. One representative of two independent experiments is shown.

CASP11-dependent manner. We also performed the coinfection experiment using *Legionella pneumophila* lacking flagellin (*flaA*), a bacteria that is known to induce caspase-11 activation and non-canonical activation of NLRP3 (Case et al., 2013). Briefly, we infected LPS-primed BMDMs with L.g.- or L.g.+ for 20 hr and then infected these cells with *flaA* for additional 4 hr. After these total 24 hr, cell-free supernatants from infected or non-infected (controls) were collected and IL-1 β production was assessed by ELISA. While we found that L.g.+ decreases IL-1 β production in *flaA*-infected C57BL/6 macrophages, this effect is completely abolished in *Casp11*^{-/-} BMDMs (Figure 1F). Collectively, these results suggest that *L. guyanensis* induces NLRP3 and caspase-1 activation via caspase-11, and the presence of LRV is capable of blocking non-canonical activation of the inflammasome.

LRV inhibits *Leishmania guyanensis*-induced caspase-11 activation

Next, we addressed the impact of LRV in caspase-11 activation by measuring caspase-11 directly, as previously described (Cunha et al., 2015). For this purpose, we primed macrophages with PAM(3)CSK(4), a TLR2 agonist, and incubated cells with biotin-VAD dye for 15 min prior to infection (more details in the Transparent methods section). This dye binds to all active forms of caspases, enabling its detection via incubation with streptavidin beads and subsequent antibody-mediated detection by Western blotting. By pulling down the active forms of caspase-11 upon infection, we found that different than L.g.-, L.g.+ fails to induce robust caspase-11 activation in PAM(3)CSK(4) (Figure 2A) or TNF- α -primed macrophages (Figure 2B). These data support the important role of LRV in blocking L.g.-induced caspase-11 activation and the non-canonical NLRP3 activation by L.g.

LRV exacerbates *L.g.* pathogenesis in mice and enhances parasite replication in BMDMs via caspase-11

Our results obtained using BMDMs demonstrate that LRV attenuates non-canonical inflammasome activation, suggesting that caspase-11 might play an important role for *L.g.* replication and pathogenesis *in vitro*




Figure 3. LRV increases *L. guyanensis* replication in macrophages via caspase-11 and NLRP3

C57BL/6, *Nlrp3*^{-/-}, *Tlr3*^{-/-}, and *Casp11*^{-/-} BMDMs were infected with *L. guyanensis* without the virus (L.g.-) or with the virus (L.g.+) at MOI 3. After 1 hr of infection, cells were washed and left in culture for 1 or 48 hr. Parasite killing was evaluated by Panotico Giemsa. The percentage of infected BMDMs at 1 (A) or 48 hr (C) and the average number of parasites at 1 (B) or 48 hr (D) after infection are shown. The results are shown as mean ± SD, and statistical analysis was performed by unpaired Student's t test. *, p < 0.05. SD, standard deviation. One representative of three independent experiments performed with technical replicates is shown.

and *in vivo*. In order to address these questions, we initially infected C57BL/6 WT, *Nlrp3*^{-/-}, and *Casp11*^{-/-} BMDMs with both parasites (L.g.- and L.g.+) and assessed parasite killing by Giemsa staining. At 1 hr after infection, no differences were observed in parasite internalization, as demonstrated by the percentage of infected cells (Figure 3A) and average number of parasites per infected cell (Figure 3B). By contrast, L.g.+ killing was significantly hampered compared to L.g.- after 48 hr of infection in WT BMDMs (Figures 3C and 3D). Interestingly, this phenomenon was not observed in NLRP3- and CASP11-deficient macrophages (and also not in *Tlr3*^{-/-} BMDMs), suggesting that LRV modulates parasite killing *in vitro* by interfering with the activation of the non-canonical inflammasome (Figures 3C and 3D).

These results encouraged us to investigate the participation of caspase-11 in LRV-mediated disease exacerbation *in vivo*. To tackle this question, we infected C57BL/6 WT, *Nlrp3*^{-/-}, and *Casp11*^{-/-} mice for four weeks and measured both ear thicknesses (weekly) and parasite titers. We found that L.g.+ induces increased ear thickness in WT mice compared to L.g.- (Figure 4A) in a NLRP3-dependent manner (Figure 4B), as reported previously (de Carvalho et al., 2019b). Interestingly, we also observed that this effect is also dependent on the presence of caspase-11 (Figure 4C). Likewise, parasite titers of L.g.+ infected WT mice are increased at four weeks after infection, but this effect is no longer observed in *Nlrp3*^{-/-} and *Casp11*^{-/-} infected mice, both in the infected ears (Figure 4D) and draining lymph nodes (Figure 4E). Taken together, these results support our findings that caspase-11 is a relevant LRV target upstream of the NLRP3 inflammasome for disease pathogenesis *in vivo*.

LRV does not interfere with inflammasome and disease pathogenesis in naturally caspase-11-deficient mice

The 129 mouse lineage has been used as a model to study infectious diseases, but differently from C57BL/6, this mouse strain carries polymorphisms that lead to a premature stop and inactive *casp11* gene (Kayagaki et al., 2011; Kenneth et al., 2012). Thus, we decided to investigate whether LRV-mediated inflammasome inhibition and disease progression would also be observed in this mouse strain. We initially infected C57BL/6 or 129 derived BMDMs with L.g.- and L.g.+ parasites and assessed IL-1β production by




Figure 4. LRV exacerbates *L. guyanensis* infection in vivo via caspase-11 and NLRP3

Wild-type C57BL/6 (A), *Nlrp3*^{-/-} (B), and *Casp11*^{-/-} mice (C) were infected with 10⁵ metacyclic promastigotes of *L. guyanensis* without the virus (L.g.-) or with the virus (L.g.+) and ear thicknesses were followed weekly, during 4 weeks. After that period, ears were removed and processed, and parasite titers were determined in the ear (D) and draining lymph nodes (E). The results are shown as mean ± SD, and statistical analysis was performed by unpaired Student's t test (D,E) or two-way ANOVA with Bonferroni's multiple comparison test (A-C). *, p < 0.05. SD, standard deviation. One representative of three independent experiments performed with biological replicates (n = 4–6 mice per group) is shown.

these cells. As expected, *L.g.*+ induced less cytokine secretion compared to *L.g.*- in C57BL/6-derived WT BMDMs, a process that was completely dependent on NLRP3, ASC, and caspase-1 (Figure 5A). However, in 129 derived WT BMDMs, *L.g.*+ and *L.g.*- induced similar levels of IL-1β (Figure 5B), differently from what were observed in C57BL/6-derived WT BMDMs. Remarkably, we observed a significant increase in the amount of IL-1β generated by 129 derived BMDMs upon infection with *L.g.*-, when compared to C57BL/6-derived macrophages, which we believe to be a direct result of genetic differences between these strains. For instance, the 129 mouse lineage has a NRAMP1 iron transporter that is functional and plays a role in antimicrobial immunity against intracellular pathogens, while C57BL/6 does not (Skamene et al., 1998). In addition to this, 129 and C57BL/6 express distinct functional alleles of the *Nlrp1* gene, which controls inflammasome activation and IL-1β production (Chavarría-Smith and Vance, 2015). Thus, these and possibly other genetic variations may explain different levels of IL-1β induced in response to *Leishmania* during infection in these two lineages of BMDMs. These data obtained *in vitro* further prompted us to evaluate the potential effects of LRV in *L.g.*-infected 129 mice. We found that *in vivo* infection of 129 mice with *L.g.*+ did not result in increased ear thickness (Figure 6A) compared to *L.g.*-, differently from what is observed in C57BL/6 mice (Figure 6B). Corroborating these data, we observed higher parasites titers in the ear (Figure 6C) and draining lymph nodes (Figure 6D) of *L.g.*+-infected C57BL/6 mice compared to *L.g.*-, but these differences were not observed in *L.g.*-infected 129 mice. Taken together, these results obtained using a mouse strain that is naturally deficient for caspase-11 provide additional support to our assertion that caspase-11 is required for LRV-induced inflammasome modulation.

LRV blocks *L.g.*-induced caspase-11 activation via TLR3 and ATG5

The results presented so far suggest that LRV interferes with the activation of caspase-11 and consequently with the non-canonical activation of the NLRP3 inflammasome in macrophages, a process that increases *L.g.* replication *in vitro* and disease pathogenesis *in vivo*. We previously demonstrated that viral dsRNA induces activation of TLR3, triggering autophagy and therefore blocking NLRP3 activation (de Carvalho et al., 2019b). We then decided to address the possible mechanisms by which LRV could potentially interfere with caspase-11 activation. For this purpose, we performed the pull-down experiment previously described




Figure 5. LRV does not impact NLRP3 inflammasome activation in macrophages from 129 mice

C57BL/6-derived (A) and wild-type (WT), *Nlrp3*^{-/-}, *Asc*^{-/-}, *Casp1*^{-/-} BMDMs, as well as 129 derived WT BMDMs (B), were primed for 4 hr with LPS (500 ng mL⁻¹) prior to infection with stationary phase *L. guyanensis* without the virus (*L.g.*-) or with the virus (*L.g.*+), at MOI 10. After 24 hr of infection, cell-free supernatants were collected, and the levels of IL-1 β were quantified by ELISA. The results are shown as mean \pm SD, and statistical analysis was performed by unpaired Student's *t* test. *, *p* < 0.05. SD, standard deviation. One representative of two independent experiments performed with technical replicates is shown.

utilizing *Tlr3*^{-/-} and *LysM*^{Cre/+} *Atg5*^{Fl/Fl} BMDMs, which lack TLR3 and ATG5 (a critical gene involved with autophagy induction), respectively. Interestingly, we found that caspase-11 activation was restored upon *L.g.*+ infection in BMDMs lacking either one of these molecules (Figures 7A and 7B), providing insights toward the mechanisms by which LRV modulates caspase-11. Although toll-like receptors (TLRs) are known to serve as a priming and therefore facilitate inflammasome activation (Swanson et al., 2019), these results corroborate previous findings suggesting that, during *Leishmania* infection, LRV-mediated TLR3 activation acts as a negative regulator of NLRP3, accounting to consolidate the concept that *Leishmania* orchestrates a differential innate immune signaling in macrophages. Although more studies are required to rule out the exact mechanisms by which LRV-induced TLR3 and ATG5 activation interferes with caspase-11 activation, this study advances in the understanding of NLRP3 modulation by LRV and reinforces the important participation of caspase-11 and the non-canonical inflammasome in the pathogenesis of leishmaniasis.

DISCUSSION

Activation of the NLRP3 inflammasome is a hallmark of leishmaniasis, and this platform is very important for the pathogenesis and the outcome of the disease (Zamboni and Sacks, 2019). We have previously demonstrated that the NLRP3 inflammasome is important to explain the previously reported effects of the LRV in disease exacerbation and induction of mucocutaneous leishmaniasis (de Carvalho et al., 2019b; Ives et al., 2011; Olivier and Zamboni, 2020). Here, we expand these findings indicating that caspase-11 is a target of LRV and required for the LRV-mediated inflammasome inhibition. Our data also highlight an important signaling axis involving caspase-11 activation to *Leishmania* pathogenesis and disease progression in infections caused by New World *Leishmania* species that harbors LRV. Although we show that this phenomenon is driven by TLR3 and ATG5, the precise mechanisms by which this pathway regulates caspase-11 activation were not evaluated in this study. Intriguingly, TLRs have been described as important receptors involved with the generation of first signal for inflammasome activation, often contributing to NLRP3-dependent cytokine production (Swanson et al., 2019). However, in the context of *Leishmania* infection, we show that LRV-driven TLR3 activation plays an opposing role in caspase-11 and NLRP3, contributing to inhibition, rather than assembly of the non-canonical inflammasome. We believe that this differential mechanism relies as a direct result of the parasite's manipulation of intracellular signaling pathways, such as NF- κ B-dependent inflammatory gene's transcription (Olivier and Zamboni, 2020; Zamboni and Sacks, 2019). However, the exact factors responsible for this different TLR3-mediated outcome remain to be addressed in future studies.

In this work, we combined a series of genetic tools (*in vitro* and *in vivo*) and different inflammasome readouts to identify caspase-11 as the target used by LRV to inhibit the inflammasome and promote




Figure 6. Disease progression upon *L.g.+* and *L.g.-* infection is similar in 129 mice

Wild-type 129 and C57BL/6 mice were infected with 10^5 metacyclic promastigotes of *L. guyanensis* without the virus (*L.g.-*) or with the virus (*L.g.+*) and ear thicknesses were followed weekly, during 4 weeks (A and B). After that period, ears and draining lymph nodes (LNs) were removed and processed, and parasite titers were determined in the ear (C) and LNs (D). The results are shown as mean \pm SD, and statistical analysis was performed by unpaired Student's *t* test (C and D) or two-way ANOVA with Bonferroni's multiple comparison test (A,B). *, $p < 0.05$. SD, standard deviation. One representative of three independent experiments performed with biological replicates ($n = 5$ mice per group) is shown.

parasite replication and disease exacerbation. So far, studies addressing the impact of LRV in the course of *Leishmania* infection employed genetically modified parasites (Castiglioni et al., 2017; Eren et al., 2016; Hartley et al., 2016; Ives et al., 2011) or clinical isolates (Kariyawasam et al., 2017) and they indeed reached important results. In this study, we used a previously obtained spontaneously derived *L.g.-* clone that lost the virus (de Carvalho et al., 2019b), offering us an excellent model to assess the LRV-mediated effects in the modulation of innate immunity and disease. Noteworthy, the differences observed in inflammasome activation by *L.g.+* and *L.g.-* parasites could be complemented by using extracellular vesicles containing LRV, as previously described (Atayde et al., 2019; de Carvalho et al., 2019b).

Although one could argue that *L.g.-* does not cause significant pathology (as indicated by an increase in skin swelling in 129 infected mice), *L.g.-* parasites are found in similar titers in both C57BL/6 and 129 mice, both in the ear and draining lymph node, suggesting that the parasite is resisting killing/multiplying even in the absence of substantial immunopathology in the infected site. Importantly, it is worth mentioning that the similar parasite titers found in 129 mice upon infection with either *L.g.-* or *L.g.+* do not necessarily represent a role for caspase-11 since other factors related to the background of this mouse could be responsible for this phenotype (examples: polymorphisms in genes involved with parasite replication or inflammatory reactions, natural resistance to infections, etc). A nice example of this is the *Slc11a11* gene, which codifies the NRAMP1 protein in myeloid cells. The 129 mouse strain produces a mature version of this protein and therefore is able to control intracellular pathogens such as *Salmonella enterica*, *Leishmania donovani*, and *Mycobacterium tuberculosis*. On the other hand, C57BL/6 mouse has a point mutation that prevents formation of a mature version of NRAMP1 and is therefore susceptible to these infections (Cellier et al., 1996; Skamene et al., 1998; Zhang et al., 2018). Nonetheless, our data in C57BL/6 mice




Figure 7. LRV interferes with caspase-11 activation by *L. guyanensis* via TLR3 and ATG5

C57BL/6 and *Tlr3*^{-/-} (A) as well as *LysM*^{Cre/+}*Atg5*^{FL/+} and *LysM*^{Cre/+}*Atg5*^{FL/FL} (B) BMDMs were primed with PAM(3)CSK(4) (300 ng mL⁻¹) for 4 hr and then infected with stationary phase (SP) *L. guyanensis* without the virus (*L.g.*⁻) or with the virus (*L.g.*⁺) at MOI 10. Active caspase-11 (45 and 38 kDA) was pulled down, and Western blotting of active (pull-down) or total (input, cell lysates) caspase-11 was performed. β -actin was used as a loading control. One representative of two independent experiments is shown.

unequivocally demonstrate that LRV utilizes caspase-11 and the inflammasome pathway to exacerbate infection in this mouse model of leishmaniasis.

Taken together, the data presented in this manuscript suggest that caspase-11 is an additional target for LRV, representing a potential molecule involved with increased disease pathology during infection with LRV + *Leishmania*. Understanding how to overcome the signaling axis triggered by the virus could represent a potential therapeutic target for the development of better therapies in patients suffering from severe and highly debilitating mucocutaneous leishmaniasis.

Limitations of the study

The current study demonstrates that *Leishmania* RNA virus promotes inhibition of the non-canonical inflammasome pathway induced upon *L. guyanensis* infection, by interfering with caspase-11 activation. Caspase-11 is known to operate upstream of NLRP3 (Hagar et al., 2013; Kayagaki et al., 2013), and it is activated in response to *Leishmania* infection (de Carvalho et al., 2019a). Although the findings represent an advance in our understanding of LRV-mediated effects on the NLRP3 inflammasome and intracellular signaling pathways in mammalian cells, we do not show the precise mechanisms by which LRV interferes with caspase-11 activation. This issue remains to be addressed and should be the subject of further studies.

Resource availability

Lead contact

Further information and requests for resources and reagents should be directed to our Lead Contact, Dario S. Zamboni (dszamboni@fmrp.usp.br).

Material availability

This study did not generate new materials or reagents.

Data and code availability

Data codes/sets were not generated or analyzed in this study. For questions regarding the raw data from the current study, please contact the corresponding author. All softwares used in this study are commercially available.

METHODS

All methods can be found in the accompanying [Transparent Methods supplemental file](#).

SUPPLEMENTAL INFORMATION

Supplemental Information can be found online at <https://doi.org/10.1016/j.isci.2020.102004>.

ACKNOWLEDGMENTS

We are thankful to Maira Nakamura for her technical support of this work. We would also like to thank Noburu Mizushima (University of Tokyo) for providing us the *Atg5^{Fl/Fl}* mice used in this study. This work was supported by grants from the Center for Research on Inflammatory Diseases (CRID), Pew, Training in Tropical Diseases/World Health Organization (TDR/WHO), INCTV/CNPq, and FAPESP (grants 2013/08216-2, 2018/14398-0 and 2019/11342-6). R.V.H.C. and D.S.L.J. were supported by fellowships from FAPESP, while D.S.Z. is a research fellow from CNPq. Currently, R.V.H.C. is a postdoctoral fellow from the Human Frontier Science Program Organization (HFSP – long-term Postdoctoral Fellowship, LT000892/2020-L).

AUTHOR CONTRIBUTIONS

R.V.H.C., D.S.L.J., and C.V.O. designed and performed experiments and generated tools for proper work development. R.V.H.C. and D.S.Z. analyzed the data, discussed hypotheses, generated figures, and wrote the article. All authors revised the manuscript.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: August 28, 2020

Revised: November 18, 2020

Accepted: December 23, 2020

Published: January 22, 2021

REFERENCES

- Atayde, V.D., da Silva Lira Filho, A., Chaparro, V., Zimmermann, A., Martel, C., Jaramillo, M., and Olivier, M. (2019). Exploitation of the *Leishmania* exosomal pathway by *Leishmania* RNA virus 1. *Nat. Microbiol.* 4, 714–723.
- Broz, P., and Dixit, V.M. (2016). Inflammasomes: mechanism of assembly, regulation and signalling. *Nat. Rev. Immunol.* 16, 407–420.
- Case, C.L., Kohler, L.J., Lima, J.B., Strowig, T., de Zoete, M.R., Flavell, R.A., Zamboni, D.S., and Roy, C.R. (2013). Caspase-11 stimulates rapid flagellin-independent pyroptosis in response to *Legionella pneumophila*. *Proc. Natl. Acad. Sci. U S A.* 110, 1851–1856.
- Castiglioni, P., Hartley, M.A., Rossi, M., Prevel, F., Desponds, C., Utzschneider, D.T., Eren, R.O., Zangger, H., Brunner, L., Collin, N., et al. (2017). Exacerbated leishmaniasis caused by a viral endosymbiont can be prevented by immunization with its viral capsid. *PLoS Negl. Trop. Dis.* 11, e0005240.
- Cellier, M., Belouchi, A., and Gros, P. (1996). Resistance to intracellular infections: comparative genomic analysis of *Nramp*. *Trends Genet.* 12, 201–204.
- Chavarría-Smith, J., and Vance, R.E. (2015). The NLRP1 inflammasomes. *Immunol. Rev.* 265, 22–34.
- Chaves, M.M., Sinflorio, D.A., Thorstenberg, M.L., Martins, M.D.A., Moreira-Souza, A.C.A., Rangel, T.P., Silva, C.L.M., Bellio, M., Canetti, C., and Coutinho-Silva, R. (2019). Non-canonical NLRP3 inflammasome activation and IL-1 β signaling are necessary to *L. amazonensis* control mediated by P2X7 receptor and leukotriene B4. *PLoS Pathog.* 15, e1007887.
- Cunha, L.D., Ribeiro, J.M., Fernandes, T.D., Massis, L.M., Khoo, C.A., Moffatt, J.H., Newton, H.J., Roy, C.R., and Zamboni, D.S. (2015). Inhibition of inflammasome activation by *Coxiella burnetii* type IV secretion system effector IcaA. *Nat. Commun.* 6, 10205.
- de Carvalho, R.V.H., Andrade, W.A., Lima-Junior, D.S., Dilucca, M., de Oliveira, C.V., Wang, K., Nogueira, P.M., Rugani, J.N., Soares, R.P., Beverley, S.M., et al. (2019a). *Leishmania* lipophosphoglycan triggers caspase-11 and the non-canonical activation of the NLRP3 inflammasome. *Cell Rep.* 26, 429–437 e425.
- de Carvalho, R.V.H., Lima-Junior, D.S., da Silva, M.V.G., Dilucca, M., Rodrigues, T.S., Horta, C.V., Silva, A.L.N., da Silva, P.F., Frantz, F.G., Lorenzon, L.B., et al. (2019b). *Leishmania* RNA virus exacerbates Leishmaniasis by subverting innate immunity via TLR3-mediated NLRP3 inflammasome inhibition. *Nat. Commun.* 10, 5273.
- de Carvalho, R.V.H., and Zamboni, D.S. (2020). Inflammasome activation in response to intracellular Protozoan parasites. *Trends Parasitol.* 36, 459–472.
- Eren, R.O., Reverte, M., Rossi, M., Hartley, M.A., Castiglioni, P., Prevel, F., Martin, R., Desponds, C., Lye, L.F., Drexler, S.K., et al. (2016). Mammalian innate immune response to a leishmania-resident RNA virus increases macrophage survival to promote parasite persistence. *Cell Host. Microbe* 20, 318–328.
- Hagar, J.A., Powell, D.A., Aachoui, Y., Ernst, R.K., and Miao, E.A. (2013). Cytoplasmic LPS activates

caspase-11: implications in TLR4-independent endotoxin shock. *Science* 341, 1250–1253.

Hartley, M.A., Bourreau, E., Rossi, M., Castiglioni, P., Eren, R.O., Prevel, F., Couppie, P., Hickerson, S.M., Launois, P., Beverley, S.M., et al. (2016). Leishmanavirus-dependent metastatic leishmaniasis is prevented by blocking IL-17a. *PLoS Pathog.* 12, e1005852.

Hartley, M.A., Ronet, C., and Fasel, N. (2012). Backseat drivers: the hidden influence of microbial viruses on disease. *Curr. Opin. Microbiol.* 15, 538–545.

Ives, A., Ronet, C., Prevel, F., Ruzzante, G., Fuertes-Marraco, S., Schutz, F., Zangger, H., Revaz-Breton, M., Lye, L.F., Hickerson, S.M., et al. (2011). Leishmania RNA virus controls the severity of mucocutaneous leishmaniasis. *Science* 331, 775–778.

Kariyawasam, R., Grewal, J., Lau, R., Pursell, A., Valencia, B.M., Llanos-Cuentas, A., and Boggild, A.K. (2017). Influence of Leishmania RNA virus 1 on proinflammatory biomarker expression in a human macrophage model of American tegumentary leishmaniasis. *J. Infect. Dis.* 216, 877–886.

Kayagaki, N., Warming, S., Lamkanfi, M., Vande Walle, L., Louie, S., Dong, J., Newton, K., Qu, Y.,

Liu, J., Heldens, S., et al. (2011). Non-canonical inflammasome activation targets caspase-11. *Nature* 479, 117–121.

Kayagaki, N., Wong, M.T., Stowe, I.B., Ramani, S.R., Gonzalez, L.C., Akashi-Takamura, S., Miyake, K., Zhang, J., Lee, W.P., Muszynski, A., et al. (2013). Noncanonical inflammasome activation by intracellular LPS independent of TLR4. *Science* 341, 1246–1249.

Kenneth, N.S., Younger, J.M., Hughes, E.D., Marcotte, D., Barker, P.A., Saunders, T.L., and Duckett, C.S. (2012). An inactivating caspase 11 passenger mutation originating from the 129 murine strain in mice targeted for c-IAP1. *Biochem. J.* 443, 355–359.

Olivier, M., and Zamboni, D.S. (2020). Leishmania Viannia guyanensis, LRV1 virus and extracellular vesicles: a dangerous trio influencing the faith of immune response during muco-cutaneous leishmaniasis. *Curr. Opin. Immunol.* 66, 108–113.

Pigott, D.M., Bhatt, S., Golding, N., Duda, K.A., Battle, K.E., Brady, O.J., Messina, J.P., Balard, Y., Bastien, P., Pratlong, F., et al. (2014). Global distribution maps of the leishmaniases. *Elife* 3, e02851.

Rossi, M., Castiglioni, P., Hartley, M.A., Eren, R.O., Prevel, F., Desponds, C., Utschneider, D.T.,

Zehn, D., Cusi, M.G., Kuhlmann, F.M., et al. (2017). Type I interferons induced by endogenous or exogenous viral infections promote metastasis and relapse of leishmaniasis. *Proc. Natl. Acad. Sci. U S A* 114, 4987–4992.

Scott, P., and Novais, F.O. (2016). Cutaneous leishmaniasis: immune responses in protection and pathogenesis. *Nat. Rev. Immunol.* 16, 581–592.

Skamene, E., Schurr, E., and Gros, P. (1998). Infection genomics: nramp1 as a major determinant of natural resistance to intracellular infections. *Annu. Rev. Med.* 49, 275–287.

Swanson, K.V., Deng, M., and Ting, J.P. (2019). The NLRP3 inflammasome: molecular activation and regulation to therapeutics. *Nat. Rev. Immunol.* 19, 477–489.

Zamboni, D.S., and Sacks, D.L. (2019). Inflammasomes and Leishmania: in good times or bad, in sickness or in health. *Curr. Opin. Microbiol.* 52, 70–76, <https://doi.org/10.1016/j.mib.2019.05.005>.

Zhang, J., Malo, D., Mott, R., Panthier, J.J., Montagutelli, X., and Jaubert, J. (2018). Identification of new loci involved in the host susceptibility to *Salmonella* Typhimurium in collaborative cross mice. *BMC Genomics* 19, 303.

iScience, Volume 24

Supplemental Information

**Endosymbiotic RNA virus
inhibits *Leishmania*-induced
caspase-11 activation**

Renan V.H. de Carvalho, Djalma S. Lima-Júnior, Caroline V. de Oliveira, and Dario S. Zamboni

1 **Supplemental information**

2

3 **Transparent Methods**

4

5 **Mice**

6 Six to eight weeks old female wild-type 129S1/SvImJ (JAX number 002448) and
7 C57BL/6, as well as knockout animals in C57BL/6 background *Nlrp3*^{-/-}, *Asc*^{-/-},
8 *Tlr3*^{-/-}, *Casp11*^{-/-}, *LysM*^{Cre/+}*Atg5*^{Fl/+} and *LysM*^{Cre/+}*Atg5*^{Fl/Fl} were used. Animals were
9 bred and maintained under specific pathogen-free conditions at the animal
10 facilities of the University of São Paulo, FMRP/USP. Experiments were approved
11 and conducted according to institutional guidelines for animal care (Protocol
12 number 014/2016).

13

14 **Parasites and infection**

15 The parasite strains used in this study were the wild-type strain of *L. guyanensis*
16 M4147 (MHOM/BR/75/M4147), harboring high levels of LRV1 (*L.g.*+) and its LRV
17 negative-derived clone (*L.g.*-), as previously described (de Carvalho et al., 2019).
18 Parasites were cultured at 25 °C in Schneider's Drosophila medium (Invitrogen,
19 Carlsbad, CA), pH 7.0, supplemented with 10% heat-inactivated fetal calf serum
20 (GIBCO BRL), 2 mM L-glutamine, and 2% bioppterin, pH 6.5. For in vivo
21 infections, mice were infected with 10⁵ metacyclic promastigotes (purified using
22 the gradient-density FICOLL method (Späth and Beverley, 2001) in 10 µL of
23 PBS, through an intradermal injection into the left ear. Ear thicknesses were
24 measured weekly with a dial gauge caliper and compared to the thickness of the
25 uninfected ear of each individual mouse. Parasite burdens were quantified by

26 limiting dilution assay, which consisted of preparing ear and lymph node's
27 homogenates from infected mice after 4 weeks of infection, plating undiluted
28 samples at the first lane of 96 well plates, and serial diluting each (1:5 for ear, 1:2
29 for lymph node) sample up to 12x in complete Schneider's medium, therefore
30 allowing parasite's growth. After 7 days in culture, each well in each dilution was
31 checked and parasites were quantified, as previously described (de Carvalho et
32 al., 2019). Coinfection experiments with *Legionella pneumophila* lacking flagellin
33 (*FlaA*-) were performed as previously described (de Carvalho et al., 2019).
34 Briefly, we infected LPS-primed BMDMs with *L.g.*- or *L.g.*+ for 20 hours, and then
35 infected these cells with *flaA* for additional 4 hours. After these total 24 hours,
36 cell-free supernatants from infected or non-infected (controls) were collected and
37 IL-1 β production was assessed by ELISA (R&D Systems), according to the
38 manufacturer's protocol.

39

40 **Bone marrow-derived macrophages (BMDM) and killing experiments**

41 Femurs and tibiae were isolated and flushed with PBS, and total bone marrow
42 cells were cultured in RPMI supplemented with 30% L929 cell-conditioned
43 medium and 20% FBS. After 7 days in culture, mature BMDMs were harvested
44 and infected with stationary phase or metacyclic promastigotes at MOI 3. 1 hour
45 after infection, wells were washed, and fresh media was added to the infected
46 cultures. The Leishmanicidal activity of the cells was determined at 48 hours after
47 infection by counting the *Giemsa*-stained cytopsin preparations under a light
48 microscope with a 40X objective. The infection rate was determined by scoring
49 the infected and uninfected cells (100 BMDMs) and the average number of
50 intracellular parasites per infected BMDM.

51

52 **Caspase-1 activation assays**

53 BMDMs were cultured and infected with stationary phase *L.g.+* and *L.g.-* (MOI
54 10). 24 hours after infection, macrophages were stained for 1 hour with FAM–
55 YVAD–fluoromethyl ketone (FAM–YVAD–FMK; Immunochemistry Technologies),
56 as recommended by the manufacturer’s instructions. The active CASP1 was then
57 measured by FACS. Both the percentage of active-caspase-1 macrophages and
58 the integrated mean of fluorescence intensity (iMFI, which represents the
59 average of caspase-1 activation per cell) were evaluated. Data were acquired on
60 a FACS ACCURI C6 flow cytometer (BD Biosciences) and analyzed with the
61 FlowJo software (Tree Star).

62

63

64 **Cytokine detection by ELISA**

65 BMDMs were primed for 4 hours with 500 ng/mL of LPS, a TLR4 agonist, and
66 then infected with *L.g.-* or *L.g.+* at MOI 10. After 24 hours of infection, cell-free
67 supernatants were collected and the levels of IL-1 β were measured by ELISA
68 (BD Biosciences), according to the manufacturer’s instructions.

69

70 **Pull-down of active caspase-11**

71 Macrophages were seeded in 6-well plates. Cells were plated in triplicates for
72 each condition, 5x10⁶ BMDMs/well. RPMI 10% FBS was used as media for the
73 entire assay. Cells were primed with either TNF- α (10 ng/mL) or PAM(3)CSK(4)
74 (300 ng/mL) for 4 hours, and 15 minutes prior to infection, medium was renewed
75 with 20 mM biotin-VAD-FMK (Enzo). Infection was performed with an additional 1

76 mL per well, totalizing 2 mL per well (not discarding the previous medium with b-
77 VAD). 24 hours after infection, medium was removed and 500 microliters of RIPA
78 4% protease inhibitor (PI) was added to each triplicate (ex: 200 microL in the first
79 replicate, and 150 microL on the other 2). Samples were centrifuged at 14000
80 G/15 min/4 degrees Celsius. As debris were sedimented, supernatant was
81 transferred to a new 1.5 mL eppendorf. Next, the total amount of proteins in each
82 sample was quantified by Bradford (Cepham Life Sciences). Samples were then
83 equalized to 1 microgram of protein and incubated in a rotator with streptavidin-
84 beads overnight (ON), at 4 degrees Celsius. Unincubated samples were used as
85 loading controls for caspase-11 and actin (total protein in cell lysates). After
86 incubating ON, the active caspases were bound to the streptavidin beads, and
87 500 microL of RIPA 4% PI to each sample was added. Next, samples were
88 centrifuged 11000G/2 min. We did that for a total of three times. Finally,
89 supernatants were discarded and we added 30 microL of sample buffer to each
90 sample, boiling them at 100 degrees Celsius for 5 minutes, what induces active
91 caspases dissociation from the beads. Supernatants were recovered in a new
92 eppendorf, and these samples containing active caspases were ready to be
93 runned in SDS-PAGE and analysis by western blotting. Rabbit anti-Casp11
94 antibody clone Ab180673 (Abcam) (1:1000), mouse anti- β -actin (C4) (Santa
95 cruz, sc-47778) and specific horseradish peroxidase-conjugated secondary
96 antibodies (1:3000; KPL) were diluted in blocking buffer for the incubations. The
97 ECL luminol reagent (GE Healthcare) was used for protein detection.

98

99 **Western blot for caspase-1**

100 A total of 10^7 BMDMs were seeded per well, primed with ultrapure LPS (500

101 ng/mL) for 4 h and left uninfected or infected for 48 hours. After this period, the
102 supernatants were collected and precipitated with 50% trichloroacetic acid (TCA)
103 and acetone. Cells were lysed in RIPA buffer containing protease inhibitor
104 cocktail (Roche). The lysates and supernatants were resuspended in Laemmli
105 buffer, boiled, resolved by SDS-PAGE and transferred (Semidry Transfer Cell,
106 Bio-Rad) to a nitrocellulose membrane (GE Healthcare). Rat anti-Casp1 p20
107 monoclonal antibody clone 4B4 (Genentech) (1:500) and specific anti-rat
108 horseradish peroxidase-conjugated antibodies (1:3000; KPL) were diluted in
109 blocking buffer for the incubations. The ECL luminol reagent (GE Healthcare)
110 was used for the protein detection.

111

112 **Statistical analysis**

113 Two-way analysis of variance (ANOVA) followed by Bonferroni post-test was
114 used for the comparison of multiple groups. The differences in the values
115 obtained for two different groups were determined using Student's *t* test. The
116 Prism 5.0 software (GraphPad, San Diego, CA) was used. A difference was
117 considered statistically significant when $P < 0.05$.

118

119 **Supplemental references**

120

121 de Carvalho, R.V.H., Lima-Junior, D.S., da Silva, M.V.G., Dilucca, M., Rodrigues,
122 T.S., Horta, C.V., Silva, A.L.N., da Silva, P.F., Frantz, F.G., Lorenzon, L.B., et al.
123 (2019b). Leishmania RNA virus exacerbates Leishmaniasis by subverting innate
124 immunity via TLR3-mediated NLRP3 inflammasome inhibition. *Nat Commun* 10,
125 5273.

126

127 Späth, G.F., Beverley, S.M. (2001). A Lipophosphoglycan-Independent Method for
128 Isolation of Infective Leishmania Metacyclic Promastigotes by Density Gradient
129 Centrifugation. *Exp Parasitol* 99, 97–103.

130