



# Cone-Beam CT-Guided Percutaneous Transthoracic Needle Lung Biopsy of Juxtaphrenic Lesions: Diagnostic Accuracy and Complications

Wonju Hong<sup>1</sup>, Soon Ho Yoon<sup>1, 2</sup>, Jin Mo Goo<sup>1, 2, 3</sup>, Chang Min Park<sup>1, 2, 3, 4</sup>

<sup>1</sup>Department of Radiology, Seoul National University Hospital, Seoul, Korea; <sup>2</sup>Institute of Radiation Medicine, Seoul National University Medical Research Center, Seoul, Korea; <sup>3</sup>Cancer Research Institute, Seoul National University, Seoul, Korea; <sup>4</sup>Institute of Medical and Biological Engineering, Medical Research Center, Seoul National University, Seoul, Korea

**Objective:** To investigate the diagnostic accuracy and complications of cone-beam CT-guided percutaneous transthoracic needle biopsy (PTNB) of juxtaphrenic lesions and identify the risk factors for diagnostic failure and complications.

**Materials and Methods:** In total, 336 PTNB procedures for lung lesions (mean size  $\pm$  standard deviation [SD],  $4.3 \pm 2.3$  cm) abutting the diaphragm in 326 patients (189 male and 137 female; mean age  $\pm$  SD,  $65.2 \pm 11.4$  years) performed between January 2010 and December 2014 were included. The accuracy, sensitivity, specificity, positive predictive value (PPV), and negative predictive value (NPV) of the PTNB procedures for the diagnosis of malignancy were measured based on the intention-to-diagnose principle. The risk factors for diagnostic failures and complications were evaluated using logistic regression analysis.

**Results:** The accuracy, sensitivity, specificity, PPV, and NPV were 92.7% (293/316), 91.3% (219/240), 91.4% (74/81), 96.9% (219/226), and 77.9% (74/95), respectively. There were 23 diagnostic failures (7.3%), and lesion sizes  $\leq 2$  cm ( $p = 0.045$ ) were the only significant risk factors for diagnostic failure. Complications occurred in 98 cases (29.2%), including 89 cases of pneumothorax (26.5%) and 7 cases of hemoptysis (2.1%). The multivariable analysis showed that old age ( $> 65$  years) ( $p = 0.002$ ), lesion size of  $\leq 2$  cm ( $p = 0.003$ ), emphysema ( $p = 0.006$ ), and distance from the pleura to the target lesion ( $> 2$  cm) ( $p = 0.010$ ) were significant risk factors for complications.

**Conclusion:** The diagnostic accuracy of cone-beam CT-guided PTNB of juxtaphrenic lesions for malignancy was fairly high, and the target lesion size was the only significant predictor of diagnostic failure. Complications of cone-beam CT-guided PTNB of juxtaphrenic lesions occurred at a reasonable rate.

**Keywords:** Lung; Biopsy; Image-guided biopsy; Cone-beam computed tomography

## INTRODUCTION

Percutaneous transthoracic needle biopsy (PTNB) is widely used as a diagnostic procedure to identify malignancy in

pulmonary lesions. The accuracy of PTNB for the diagnosis of malignancy has been reported to be as high as 90–99% [1-9]. Cone-beam CT-guided PTNB with virtual guidance has been established as a highly accurate and safe tissue sampling technique [1-4]. However, diagnostic failure and complications inevitably occur in at least a small proportion of PTNBs. The reported risk factors for diagnostic failure of PTNB include a small target lesion, a benign final diagnosis, and the use of fine-needle aspiration (FNA) as the diagnostic method [4-6]. A lower lobar location of the target lesion is also known to be associated with both diagnostic failure and the occurrence of pneumothorax [4,8-12]. However, the relationships between diagnostic failure or complications and the abutment of the diaphragm by the target lesion have not yet been established,

**Received:** October 10, 2020 **Revised:** December 10, 2020

**Accepted:** December 21, 2020

**Corresponding author:** Chang Min Park, MD, PhD, Department of Radiology, Institute of Radiation Medicine, Cancer Research Institute, Institute of Medical and Biological Engineering, Seoul National University, 101 Daehak-ro, Jongno-gu, Seoul 03080, Korea.

• E-mail: cmpark.morphius@gmail.com

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

although in practice, radiologists find the procedure more challenging when the target lesion is located adjacent to the diaphragm because it is strongly affected by respiratory motion. There is a risk of abdominal solid organ injury, and needle trajectory planning is limited. Some lesions are not sufficiently visible on fluoroscopy [12,13]. Patel et al. [12] investigated the impact of target lesion proximity to the diaphragm on biopsy yield and pneumothorax risk, but in their study, the mean distance of target lesions from the diaphragm was 8.3 cm, which was far from a juxtaphrenic location. To the best of our knowledge, the diagnostic accuracy and complication rate of PTNB for juxtaphrenic lesions have not been evaluated in prior studies.

Thus, the purpose of our study was to investigate the diagnostic accuracy and complications of cone-beam CT-guided PTNB of juxtaphrenic lesions and identify the risk factors for diagnostic failure and complications.


## MATERIALS AND METHODS

### Study Design and Population

This retrospective study was approved by the Institutional

Review Board of Seoul National University Hospital, and the requirement for written informed consent was waived (IRB No. 2006-065-1131).

We enrolled 3558 consecutive cone-beam CT-guided PTNBs that were performed at a single tertiary referral hospital between January 2010 and December 2014. We included 348 PTNBs that satisfied the selection criterion of the target lesion abutting the diaphragm on a sagittal scan of preprocedural cone-beam CT (Fig. 1). Twelve PTNBs that underwent repeat biopsies of the same target lesion within a week were excluded; post-procedural pneumothorax or pulmonary hemorrhage may affect the results or complications of repeat biopsies. Finally, 336 PTNBs in 326 patients were included in the study. With respect to the diagnostic performance analysis, 20 PTNBs that did not fulfill the predefined criteria for determining the final diagnosis were excluded. The patient cohort of our study has also been included in previous multi-institutional studies that reported the malignancy risk demonstrated by non-diagnostic results from PTNBs and the diagnostic accuracy and complications of PTNBs [1,14,15]. However, no analysis has been conducted on PTNBs with a focus on



**Fig. 1.** Cone-beam CT-guided PTNB of a lung nodule in the juxtaphrenic location in a 39-year-old male with hepatocellular carcinoma.

**A, B.** Diagnostic CT image shows a 1.3-cm nodule (arrows) in the left lower lobe abutting the diaphragm on axial (**A**) and coronal (**B**) images. **C.** On the sagittal scan of preprocedural cone-beam CT, the shortest distance between the target lesion and diaphragm is 0 cm. **D.** After confirming the location of the needle tip on cone-beam CT, PTNB was performed. The nodule was confirmed as poorly differentiated carcinoma on biopsy. The nodule increased in size on follow-up CT (not shown). PTNB = percutaneous transthoracic needle lung biopsy

juxtaphrenic lesions yet.

### Cone-Beam CT-Guided PTNB Procedures

Cone-beam CT-guided PTNBs were primarily performed by thoracic radiologists who had performed at least 100 PTNBs, but residents and fellows in training were involved under the supervision of experienced thoracic radiologists (with 15 years and 7 years of experience in image-guided PTNB). The PTNBs were performed using two different cone-beam CT systems (Axiom Artis dTA/VB30 flat-panel detector with a 2048 x 1538 element, Siemens; Allura Xper FD20 flat-panel detector with a 2480 x 1920 element, Philips Healthcare) with virtual guidance software (iGuide, Siemens Medical Solutions; XperGuide, Philips Healthcare). Preprocedural cone-beam CT and the insertion and extraction of the biopsy needle were performed when patients held their breath at the end of expiration.

### Data Collection

One radiologist (5 years of clinical experience) and a trained radiology technician (1 year of research experience in chest CT) retrospectively reviewed preprocedural CT images of the PTNBs using a picture archiving and communication system station. On the sagittal scans of preprocedural cone-beam CT, the shortest distance between the target lesion and diaphragm was measured. PTNBs with a distance of 0 cm were included in the study.

Variables related to patients, target lesions, and biopsy procedures were collected from our PTNB data registry. The patient variables included age, sex, and presence of pulmonary emphysema on CT images. The target characteristics were lesion size (long-axis diameter on a diagnostic CT image), location, and distance from the pleura to the target. The biopsy procedural information consisted of patient position, biopsy method (FNA or core needle biopsy [CNB]), and the number of tissue samples. Complications, including pneumothorax, pneumothorax requiring drainage catheter insertion, and hemoptysis, were recorded after reviewing the procedure records, medical charts, and follow-up images.

Based on the pathologic reports for the biopsy specimen, the cone-beam CT-guided PTNB results were categorized as positive, negative, or non-evaluable due to insufficient specimens, and the final diagnosis was blinded. Malignancy, atypical cells suggestive of malignancy, atypical cells suspicious for malignancy, and atypical cells of indeterminate malignancy were considered positive, based

on previous research [15]. Specific benign diagnoses, such as hamartoma and tuberculosis, and nonspecific benign pathologic results, including granulomatous inflammation, abscess, or organizing pneumonia, were considered negative. The results were considered non-evaluable when the cone-beam CT-guided PTNB reports indicated that the specimens were insufficient or inadequate for diagnosis due to cellular paucity.

### Reference Standards

For each lesion biopsied, the final diagnosis was classified as malignant or benign, and the reference standard for each lesion was established in one of the following four ways [4,8]. First, the final diagnosis was based on a surgical pathologic report if the lesion was surgically resected. Second, if the PTNB revealed a malignant or specific benign diagnosis and the target lesion was not surgically resected, the final diagnosis was based on a pathologic analysis of the PTNBs. Atypical cell results were not used to determine the final diagnosis. Third, the lesion was considered benign if its size decreased by 20% or more or remained stable for at least 2 years without treatment. Fourth, the lesion was identified as malignant if its clinical behavior showed obvious malignant processes. Lesions that did not fulfill the above criteria were classified as having incomplete reference standards and excluded from the diagnostic performance analysis.

### Statistical Analysis

The accuracy, sensitivity, specificity, positive predictive value (PPV), and negative predictive value (NPV) for the diagnosis of malignancy were calculated with 95% confidence intervals (CIs) on a per-lesion basis. Based on the intention-to-diagnose principle, non-evaluable results due to insufficient specimens were considered to be false negatives when calculating sensitivity and false positives when calculating specificity [1,16].

The cone-beam CT-guided PTNB results were categorized into two groups: the diagnostic success group and the diagnostic failure group. The diagnostic success group consisted of true-positive and true-negative results. The diagnostic failure group consisted of false-positive, false-negative, and non-evaluable results due to insufficient specimens. To determine the risk factors for diagnostic failure, the statistical significance of the differences between the diagnostic success and failure groups was analyzed using the Pearson chi-squared test or Fisher's

exact test, as appropriate. Variables with  $p < 0.010$  during the univariable analysis were used as input variables for multivariable logistic regression analysis to identify independent factors. In the multivariable logistic regression analysis, a backward stepwise conditional method was used.

The rates of overall complications, pneumothorax, pneumothorax requiring drainage catheter insertion, hemoptysis, and other rare complications were calculated. The risk factors for overall complications and pneumothorax were investigated as aforementioned.

All statistical analyses were performed using IBM SPSS Statistics for Windows, version 24.0 (IBM Corp.). A  $p$  value of less than 0.05 was considered to indicate statistical significance.

## RESULTS

### Patient Demographics and Lesions

In total, 326 patients underwent 336 cone-beam CT-guided PTNBs, and their mean age was  $65.2 \pm 11.4$  years

(age range, 30–86 years). After excluding 20 cases with incomplete reference standards, the diagnostic performance of 316 cone-beam CT-guided PTNBs in 306 patients with a mean age of  $65.5 \pm 11.2$  years (age range, 33–86 years) was analyzed. The characteristics of the patients, lesions, and procedures are shown in Table 1.

### Diagnostic Accuracy

Of the 316 cone-beam CT-guided PTNBs, 221 (69.9%) showed positive results, 90 (28.5%) showed negative results, and 5 (1.6%) were non-evaluable due to insufficient specimens. According to the final diagnosis, 235 (74.4%) lesions were confirmed as malignant, and 81 (25.6%) lesions were confirmed as benign. The final diagnosis of malignancy was based on surgical pathology ( $n = 59$ ), specific malignant biopsy results ( $n = 168$ ), and clinical malignant behavior ( $n = 8$ ). Benignity was confirmed based on surgical pathology ( $n = 9$ ), specific benign biopsy results ( $n = 8$ ), and a significant decrease or stability in size for at least 2 years ( $n = 64$ ).

**Table 1. Characteristics of Patients, Lesions, and Procedures**

|                                       | Analysis of Diagnostic Performance | Analysis of Complications |
|---------------------------------------|------------------------------------|---------------------------|
| Total no. of procedures               | 316                                | 336                       |
| Total no. of patients                 | 306                                | 326                       |
| Sex                                   |                                    |                           |
| Male                                  | 174 (56.9)                         | 189 (58.0)                |
| Female                                | 132 (43.1)                         | 137 (42.0)                |
| Age, year                             |                                    |                           |
| Mean $\pm$ SD                         | $65.5 \pm 11.2$                    | $65.2 \pm 11.4$           |
| Range                                 | 33–86                              | 30–86                     |
| Position                              |                                    |                           |
| Supine                                | 42 (13.3)                          | 45 (13.4)                 |
| Prone                                 | 274 (86.7)                         | 291 (86.6)                |
| Biopsy needle                         |                                    |                           |
| FNA only                              | 9 (2.8)                            | 11 (3.3)                  |
| CNB or combined                       | 307 (97.2)                         | 325 (96.7)                |
| No. of tissue samples                 |                                    |                           |
| $\leq 3$                              | 250 (79.1)                         | 265 (78.9)                |
| $\geq 4$                              | 66 (20.9)                          | 71 (21.1)                 |
| Lesion size, cm                       |                                    |                           |
| Mean $\pm$ SD                         | $4.3 \pm 2.4$                      | $4.3 \pm 2.3$             |
| Range                                 | 1.0–13.9                           | 1.0–13.9                  |
| Presence of pulmonary emphysema on CT |                                    |                           |
| Yes                                   | 55 (17.4)                          | 57 (17.0)                 |
| No                                    | 261 (82.6)                         | 279 (83.0)                |
| Distance from pleura to target, cm    |                                    |                           |
| Mean $\pm$ SD                         | $1.9 \pm 1.9$                      | $1.9 \pm 1.9$             |
| Range                                 | 0–9.3                              | 0–9.3                     |

Data are presented as n (%) unless specified otherwise. CNB = core needle biopsy, FNA = fine-needle aspiration, SD = standard deviation

The overall accuracy, sensitivity, specificity, PPV, and NPV for the diagnosis of malignancy were 92.7% (293 of 316; 95% CI, 89.3–95.3%), 91.3% (219 of 240; 95% CI, 86.9–94.5%), 91.4% (74 of 81; 95% CI, 83.0–96.5%), 96.9% (219 of 226; 95% CI, 93.9–98.5%), and 77.9% (74 of 95; 95% CI, 70.0–84.2%), respectively (Fig. 2). The diagnostic categories stratified by lesion size are presented in Table 2.


### Diagnostic Failure and Risk Factors

The diagnostic failure group included 7.3% cone-beam CT-guided PTNBs (95% CI, 4.4–10.1%; 23 PTNBs). Table 3 summarizes the results of the univariable analysis comparing the diagnostic success and failure groups. There was a significant difference in lesion size ( $\leq 2$  cm vs.  $> 2$  cm) ( $p = 0.037$ ). Logistic regression analysis revealed that

a lesion size of  $\leq 2$  cm was significantly associated with diagnostic failure (odds ratio [OR], 2.78; 95% CI, 1.02–7.55;  $p = 0.045$ ). The other variables did not significantly differ in the two groups.

### Cone-Beam CT-Guided PTNB-Related Complications

Cone-beam CT-guided PTNB-related complications occurred after 29.2% (95% CI, 24.1–33.9%; 98 PTNBs) of the 336 PTNBs. Pneumothorax occurred after 26.5% of the PTNBs (95% CI, 21.7–31.3%; 89 PTNBs), and drainage catheter insertion was, subsequently, required for 2.1% (95% CI, 0.6–3.9%; 7 PTNBs). Hemoptysis occurred after 2.1% of the PTNBs (95% CI, 0.6–3.9%; 7 PTNBs). Other rare complications included hemothorax after 0.9% (95% CI, 0.0–2.1%; 3 PTNBs), all of which resolved without



**Fig. 2.** Cone-beam CT-guided PTNB in a 55-year-old male with a solitary pulmonary nodule in the juxtaphrenic location. **A, B.** Diagnostic CT image shows a 2.2-cm nodule (arrows) in the right lower lobe abutting the diaphragm on axial (**A**) and coronal (**B**) images. **C.** On the sagittal scan of a preprocedural cone-beam CT, the shortest distance between the target lesion and the diaphragm is 0 cm. **D.** After confirming the location of the needle tip on cone-beam CT, PTNB was performed. The nodule was confirmed as chondroid hamartoma on biopsy. PTNB = percutaneous transthoracic needle lung biopsy

**Table 2.** Diagnostic Categories for Malignancy according to Lesion Size

| Diagnostic Categories | Lesion Size (cm)    |                   |                  |                  | Overall (n = 316) |
|-----------------------|---------------------|-------------------|------------------|------------------|-------------------|
|                       | $\leq 2.0$ (n = 39) | 2.1–4.0 (n = 143) | 4.1–6.0 (n = 70) | $> 6.0$ (n = 64) |                   |
| True-positive         | 26 (66.7)           | 91 (63.6)         | 53 (75.7)        | 49 (76.6)        | 219 (69.3)        |
| True-negative         | 7 (17.9)            | 41 (28.7)         | 15 (21.4)        | 11 (17.2)        | 74 (23.4)         |
| False-positive        | 1 (2.6)             | 0 (0.0)           | 1 (1.4)          | 0 (0.0)          | 2 (0.6)           |
| False-negative        | 1 (2.6)             | 10 (7.0)          | 1 (1.4)          | 4 (6.3)          | 16 (5.1)          |
| Non-evaluable         | 4 (10.3)            | 1 (0.7)           | 0 (0.0)          | 0 (0.0)          | 5 (1.6)           |

Data are presented as n (%).

**Table 3. Results of Univariable Analysis of Risk Factors for Diagnostic Failure**

| Variable                              | Diagnostic Success (n = 293) | Diagnostic Failure (n = 23) | P     |
|---------------------------------------|------------------------------|-----------------------------|-------|
| Sex                                   |                              |                             | 0.653 |
| Male                                  | 167 (57.0)                   | 12 (52.2)                   |       |
| Female                                | 126 (43.0)                   | 11 (47.8)                   |       |
| Age, year                             |                              |                             | 0.530 |
| ≤ 65                                  | 133 (45.4)                   | 12 (52.2)                   |       |
| > 65                                  | 160 (54.6)                   | 11 (47.8)                   |       |
| Position                              |                              |                             | 0.971 |
| Supine                                | 39 (13.3)                    | 3 (13.0)                    |       |
| Prone                                 | 254 (86.7)                   | 20 (87.0)                   |       |
| Biopsy needle                         |                              |                             | 0.133 |
| FNA only                              | 7 (2.4)                      | 2 (8.7)                     |       |
| CNB or combined                       | 286 (97.6)                   | 21 (91.3)                   |       |
| No. of tissue samples                 |                              |                             | 0.917 |
| ≤ 3                                   | 232 (79.2)                   | 18 (78.3)                   |       |
| ≥ 4                                   | 61 (20.8)                    | 5 (21.7)                    |       |
| Target size                           |                              |                             | 0.037 |
| ≤ 2 cm                                | 33 (11.3)                    | 6 (26.1)                    |       |
| > 2 cm                                | 260 (88.7)                   | 17 (73.9)                   |       |
| Emphysema                             |                              |                             | 0.392 |
| Yes                                   | 53 (18.1)                    | 2 (8.7)                     |       |
| No                                    | 240 (81.9)                   | 21 (91.3)                   |       |
| Pneumothorax                          |                              |                             | 0.664 |
| Yes                                   | 77 (26.3)                    | 7 (30.4)                    |       |
| No                                    | 216 (73.7)                   | 16 (69.6)                   |       |
| Hemoptysis                            |                              |                             | 1.000 |
| Yes                                   | 6 (2.0)                      | 0 (0)                       |       |
| No                                    | 287 (98.0)                   | 23 (100)                    |       |
| Distance from pleura to target lesion |                              |                             | 0.520 |
| ≤ 2 cm                                | 173 (59.0)                   | 12 (52.2)                   |       |
| > 2 cm                                | 120 (41.0)                   | 11 (47.8)                   |       |
| Final diagnosis                       |                              |                             | 0.584 |
| Benign                                | 74 (25.3)                    | 7 (30.4)                    |       |
| Malignant                             | 219 (74.7)                   | 16 (69.6)                   |       |

Data are presented as n (%). CNB = core needle biopsy, FNA = fine-needle aspiration

additional procedures, and vasovagal syncope after 0.3% (95% CI, 0.0–0.9%; 1 PTNB). There were no cases of abdominal solid organ injuries, air embolism, or mortality.

Regarding the risk factors for complications, there were significant differences in patient age (≤ 65 years vs. > 65 years) ( $p = 0.002$ ), lesion size (≤ 2 cm vs. > 2 cm) ( $p = 0.014$ ), presence of emphysema ( $p = 0.001$ ), and distance from the pleura to the target lesion (≤ 2 cm vs. > 2 cm) ( $p = 0.014$ ). Multivariable logistic regression analysis revealed that an age of > 65 years (OR, 2.36; 95% CI, 1.38–4.03;  $p = 0.002$ ), lesion size of ≤ 2 cm (OR, 2.93; 95% CI, 1.44–5.96;  $p = 0.003$ ), presence of emphysema (OR, 2.39; 95% CI, 1.29–4.44;  $p = 0.006$ ), and lesion distance of > 2

cm from the pleura (OR, 1.92; 95% CI, 1.16–3.18;  $p = 0.010$ ) were associated with the occurrence of complications. Table 4 summarizes the results of the multivariable analyses for the risk factors of complications.

With respect to pneumothorax, patient age (≤ 65 years vs. > 65 years) ( $p = 0.001$ ), presence of emphysema ( $p = 0.001$ ), and the distance from the pleura to the target lesion (≤ 2 cm vs. > 2 cm) ( $p = 0.007$ ) significantly differed in the PTNBs with pneumothorax and those without pneumothorax. The multivariable logistic regression analysis showed that patient age of > 65 years (OR, 2.78; 95% CI, 1.59–4.88;  $p = 0.001$ ), lesion size of ≤ 2 cm (OR, 2.50; 95% CI, 1.20–5.24;  $p = 0.015$ ), presence of emphysema (OR,

**Table 4. Results of Univariable and Multivariable Logistic Regression Analysis of Risk Factors for Complications**

| Variable                              | No Complication<br>(n = 238) | Complication<br>(n = 98) | Multivariable Analysis |       |
|---------------------------------------|------------------------------|--------------------------|------------------------|-------|
|                                       |                              |                          | AOR (95% CI)           | P     |
| Sex                                   |                              |                          |                        | 0.188 |
| Male                                  | 132 (55.5)                   | 62 (63.3)                |                        |       |
| Female                                | 106 (44.5)                   | 36 (36.7)                |                        |       |
| Age, year                             |                              |                          |                        | 0.002 |
| ≤ 65                                  | 125 (52.5)                   | 33 (33.7)                | Reference              |       |
| > 65                                  | 113 (47.5)                   | 65 (66.3)                | 2.360 (1.382–4.030)    | 0.002 |
| Position                              |                              |                          |                        | 0.271 |
| Supine                                | 35 (14.7)                    | 10 (10.2)                |                        |       |
| Prone                                 | 203 (85.3)                   | 88 (89.8)                |                        |       |
| Biopsy needle                         |                              |                          |                        | 0.736 |
| FNA only                              | 7 (2.9)                      | 4 (4.1)                  |                        |       |
| CNB or combined                       | 231 (97.1)                   | 94 (95.9)                |                        |       |
| No. of tissue samples                 |                              |                          |                        | 0.932 |
| ≤ 3                                   | 188 (79.0)                   | 77 (78.6)                |                        |       |
| ≥ 4                                   | 50 (21.0)                    | 21 (21.4)                |                        |       |
| Target size                           |                              |                          |                        | 0.014 |
| ≤ 2 cm                                | 23 (9.7)                     | 19 (19.4)                | 2.930 (1.439–5.964)    | 0.003 |
| > 2 cm                                | 215 (90.3)                   | 79 (80.6)                | Reference              |       |
| Emphysema                             |                              |                          |                        | 0.001 |
| Yes                                   | 30 (12.6)                    | 27 (27.6)                | 2.391 (1.288–4.437)    | 0.006 |
| No                                    | 208 (87.4)                   | 71 (72.4)                | Reference              |       |
| Distance from pleura to target lesion |                              |                          |                        | 0.014 |
| ≤ 2 cm                                | 151 (63.4)                   | 48 (49.0)                | Reference              | 0.010 |
| > 2 cm                                | 87 (36.6)                    | 50 (51.0)                | 1.924 (1.163–3.181)    |       |

Data are presented as n (%). AOR = adjusted odds ratio, CI = confidence interval, CNB = core needle biopsy, FNA = fine-needle aspiration

2.51; 95% CI, 1.34–4.71;  $p = 0.004$ ), and lesion distance of > 2 cm from the pleura (OR, 2.16; 95% CI, 1.28–3.63;  $p = 0.004$ ) were associated with pneumothorax. The risk factors for pneumothorax are summarized in Table 5.

## DISCUSSION

In this study, we evaluated the diagnostic accuracy of cone-beam CT-guided PTNB for juxtaphrenic lung lesions based on the intention-to-diagnose principle and analyzed the cone-beam CT-guided PTNB-related complications. The accuracy for the diagnosis of malignancy was 92.7%, and target size was the only significant predictor of diagnostic failure. Complications occurred after 29.2% of cone-beam CT-guided PTNBs of juxtaphrenic lesions, with pneumothorax occurring in 26.5%.

The sensitivity and specificity of the cone-beam CT-guided PTNBs were 91.3% and 91.4%, respectively, and diagnostic failure occurred after 7.3%. Although the technique we used in our study is limited to cone-beam CT with virtual

guidance, the results of our study are comparable to those of prior studies that analyzed the overall diagnostic accuracy of PTNB [1-4,6-9,17,18]. In a previous multicenter study, Lee et al. [1] reported that the sensitivity, specificity, and accuracy of PTNB with various types of imaging guidance, including cone-beam CT, CT, and fluoroscopy, were 92.5%, 86.5%, and 91.1%, respectively. The diagnostic accuracy of PTNB under cone-beam CT guidance has been reported to be equivalent or slightly higher in previous studies [1-4,17,18]. Considering this, the diagnostic accuracy of this study is reasonable for clinical situations where a biopsy is required to differentiate malignant tumors from benign lesions. The specificity recorded in our study was lower than that of previous studies, which ranged from 98–100% [2-9,18]. This discrepancy can be explained by the fact that non-evaluable results were considered as false positives in our study according to the intention-to-diagnose principle, whereas in previous studies, non-evaluable results were excluded or regarded as negative biopsy results [1-9,18].

A target lesion size of ≤ 2 cm was the only independent

**Table 5. Results of Univariable and Multivariable Logistic Regression Analysis of Risk Factors for Pneumothorax**

| Variable                              | No Pneumothorax<br>(n = 247) | Pneumothorax<br>(n = 89) | Univariable Analysis | Multivariable Analysis |
|---------------------------------------|------------------------------|--------------------------|----------------------|------------------------|
|                                       |                              |                          | P                    | AOR (95% CI) P         |
| Sex                                   |                              |                          | 0.098                | 0.704                  |
| Male                                  | 136 (55.1)                   | 58 (65.2)                |                      | 1.116 (0.633–1.967)    |
| Female                                | 111 (44.9)                   | 31 (34.8)                |                      | Reference              |
| Age, year                             |                              |                          | 0.001                | 0.001                  |
| ≤ 65                                  | 131 (53.0)                   | 27 (30.3)                |                      | Reference              |
| > 65                                  | 116 (47.0)                   | 62 (69.7)                |                      | 2.782 (1.586–4.880)    |
| Position                              |                              |                          | 0.289                |                        |
| Supine                                | 36 (14.6)                    | 9 (10.1)                 |                      |                        |
| Prone                                 | 211 (85.4)                   | 80 (89.9)                |                      |                        |
| Biopsy needle                         |                              |                          | 0.490                |                        |
| FNA only                              | 7 (2.8)                      | 4 (4.5)                  |                      |                        |
| CNB or combined                       | 240 (97.2)                   | 85 (95.5)                |                      |                        |
| No. of tissue samples                 |                              |                          | 0.718                |                        |
| ≤ 3                                   | 196 (79.4)                   | 69 (77.5)                |                      |                        |
| ≥ 4                                   | 51 (20.6)                    | 20 (22.5)                |                      |                        |
| Target size                           |                              |                          | 0.068                | 0.015                  |
| ≤ 2 cm                                | 26 (10.5)                    | 16 (18.0)                |                      | 2.504 (1.197–5.235)    |
| > 2 cm                                | 221 (89.5)                   | 73 (82.0)                |                      | Reference              |
| Emphysema                             |                              |                          | 0.001                | 0.004                  |
| Yes                                   | 31 (12.6)                    | 26 (29.2)                |                      | 2.510 (1.339–4.706)    |
| No                                    | 216 (87.4)                   | 63 (70.8)                |                      | Reference              |
| Distance from pleura to target lesion |                              |                          | 0.007                | 0.004                  |
| ≤ 2 cm                                | 157 (63.6)                   | 42 (47.2)                |                      | Reference              |
| > 2 cm                                | 90 (36.4)                    | 47 (52.8)                |                      | 2.157 (1.280–3.634)    |

Data are presented as n (%). AOR = adjusted odds ratio, CI = confidence interval, CNB = core needle biopsy, FNA = fine-needle aspiration

risk factor for diagnostic failure in our study. Although cone-beam CT-guided PTNB is known to be an accurate diagnostic method for even small lung nodules [2,3], adequate and safe tissue acquisition for small juxtaphrenic lesions requires high proficiency and expertise. Avoiding passing the needle through the diaphragm is technically challenging, particularly when the lesion is small because this location is strongly influenced by respiratory motion. Small juxtaphrenic lesions of ≤ 2 cm are more difficult for PTNB and have a higher diagnostic failure rate than larger lesions, but radiologists can expect successful diagnosis from a cone-beam CT-guided PTNB for larger lesions within the juxtaphrenic location. Other variables, including the final diagnosis and biopsy method (FNA or CNB), were not associated with diagnostic failure in this study, unlike in previous reports [1,5,8,9]. However, only 2.8% (9 of 316) of PTNBs were performed with FNA in our study, and more cases of FNA would be necessary for a confirmatory evaluation.

The complication rate of cone-beam CT-guided PTNB of

juxtaphrenic lesions was 29.2% in this study. Pneumothorax occurred after 26.5% of PTNBs, which is within the previously reported range of 16.2–38.4% [2,4,11,12,14,18–20]. Significant risk factors for pneumothorax were old age, presence of emphysema, small target size, and deep location, all of which are consistent with prior studies [4,9,10,14,21–23]. Regarding the association between a lower lobar location and the occurrence of pneumothorax, several studies have reported that a lower lobar location was related to the increased incidence of pneumothorax [4,9–11,17]. As our study evaluated only juxtaphrenic lesions, we did not investigate whether pneumothorax was correlated with a lobar location or abutment to the diaphragm. However, the incidence of pneumothorax in juxtaphrenic lesions was within the acceptable range in this study. Moreover, only 2.1% of cases required drainage catheter insertion, which is lower than the previously reported incidence, ranging from 2.4% to 15% [4,9,10,14,21]. We suggest that a juxtaphrenic location is unlikely to have a major influence on the occurrence of



pneumothorax.

In this study, the hemoptysis rate was 2.1%, which was within the previously reported range of 2.0–6.9% but in the lower range [4,9,10,14,18,22]. This can be explained by the fact that juxtaphrenic lesions are located in the periphery of the lung and have a lower chance of bronchovascular injuries along the needle course [24]. Furthermore, as no major complications, such as liver or spleen injury, air embolism, or mortality were noted, cone-beam CT-guided PTNB for juxtaphrenic lesions is a relatively safe procedure.

Our study had several limitations. First, this study had a retrospective design with unknown biases. Second, preprocedural cone-beam CT was performed at the end of expiration. However, diagnostic chest CT images are mostly performed when patients hold their breath after inspiration. The location of juxtaphrenic lesions may change according to respiration, and lesions located slightly apart from the diaphragm on diagnostic CT may have been included in this study. Nevertheless, the procedure was carried out during expiration, and this issue likely did not have a significant impact on the results. Third, our study did not compare the diagnostic accuracy and complications of juxtaphrenic lesions with those of lesions located farther from the diaphragm. Further studies are necessary to confirm the impact of lesions abutting the diaphragm. Lastly, the surgeons in our institution had already had experience and were familiar with cone-beam CT-guided PTNB. It is not certain that our results can be reproducible where surgeons were skilled with different methods such as CT or CT fluoroscopy and were not familiar with cone-beam CT guidance.

In conclusion, the accuracy of cone-beam CT-guided PTNB of juxtaphrenic lesions for the diagnosis of malignancy was fairly high, and the target size was the only significant predictor of diagnostic failure. Complications of cone-beam CT-guided PTNB of juxtaphrenic lesions occurred at a reasonable rate.

#### Conflicts of Interest

The authors have no potential conflicts of interest to disclose.

#### Author Contributions

Conceptualization: Chang Min Park. Data curation: Wonju Hong, Soon Ho Yoon. Formal analysis: Wonju Hong. Investigation: Wonju Hong, Chang Min Park. Methodology: Chang Min Park. Project administration: Chang Min Park. Supervision: Chang Min Park, Jin Mo Goo. Writing—original

draft: Wonju Hong. Writing—review & editing: Soon Ho Yoon, Jin Mo Goo, Chang Min Park.

#### ORCID iDs

Wonju Hong

<https://orcid.org/0000-0001-6891-5483>

Soon Ho Yoon

<https://orcid.org/0000-0002-3700-0165>

Jin Mo Goo

<https://orcid.org/0000-0003-1791-7942>

Chang Min Park

<https://orcid.org/0000-0003-1884-3738>

#### REFERENCES

1. Lee KH, Lim KY, Suh YJ, Hur J, Han DH, Kang MJ, et al. Diagnostic accuracy of percutaneous transthoracic needle lung biopsies: a multicenter study. *Korean J Radiol* 2019;20:1300-1310
2. Choi MJ, Kim Y, Hong YS, Shim SS, Lim SM, Lee JK. Transthoracic needle biopsy using a C-arm cone-beam CT system: diagnostic accuracy and safety. *Br J Radiol* 2012;85:e182-e187
3. Choo JY, Park CM, Lee NK, Lee SM, Lee HJ, Goo JM. Percutaneous transthoracic needle biopsy of small ( $\leq 1$  cm) lung nodules under C-arm cone-beam CT virtual navigation guidance. *Eur Radiol* 2013;23:712-719
4. Lee SM, Park CM, Lee KH, Bahn YE, Kim JI, Goo JM. C-arm cone-beam CT-guided percutaneous transthoracic needle biopsy of lung nodules: clinical experience in 1108 patients. *Radiology* 2014;271:291-300
5. Choi SH, Chae EJ, Kim JE, Kim EY, Oh SY, Hwang HJ, et al. Percutaneous CT-guided aspiration and core biopsy of pulmonary nodules smaller than 1 cm: analysis of outcomes of 305 procedures from a tertiary referral center. *AJR Am J Roentgenol* 2013;201:964-970
6. Yeow KM, Tsay PK, Cheung YC, Lui KW, Pan KT, Chou AS. Factors affecting diagnostic accuracy of CT-guided coaxial cutting needle lung biopsy: retrospective analysis of 631 procedures. *J Vasc Interv Radiol* 2003;14:581-588
7. Priola AM, Priola SM, Cataldi A, Errico L, Di Franco M, Campisi P, et al. Accuracy of CT-guided transthoracic needle biopsy of lung lesions: factors affecting diagnostic yield. *Radiol Med* 2007;112:1142-1159
8. Hiraki T, Mimura H, Gobara H, Iguchi T, Fujiwara H, Sakurai J, et al. CT fluoroscopy-guided biopsy of 1,000 pulmonary lesions performed with 20-gauge coaxial cutting needles: diagnostic yield and risk factors for diagnostic failure. *Chest* 2009;136:1612-1617
9. Takeshita J, Masago K, Kato R, Hata A, Kaji R, Fujita S, et al. CT-guided fine-needle aspiration and core needle biopsies of pulmonary lesions: a single-center experience with 750

- biopsies in Japan. *AJR Am J Roentgenol* 2015;204:29-34
10. Hiraki T, Mimura H, Gobara H, Shibamoto K, Inoue D, Matsui Y, et al. Incidence of and risk factors for pneumothorax and chest tube placement after CT fluoroscopy-guided percutaneous lung biopsy: retrospective analysis of the procedures conducted over a 9-year period. *AJR Am J Roentgenol* 2010;194:809-814
  11. Huang MD, Weng HH, Hsu SL, Hsu LS, Lin WM, Chen CW, et al. Accuracy and complications of CT-guided pulmonary core biopsy in small nodules: a single-center experience. *Cancer Imaging* 2019;19:51
  12. Patel MV, Ahmed O, Jilani D, Zangan S. Computed tomography-guided percutaneous lung biopsy: impact of lesion proximity to diaphragm on biopsy yield and pneumothorax rate. *J Thorac Imaging* 2014;29:344-349
  13. Ikezoe J, Morimoto S, Arisawa J, Takashima S, Kozuka T, Nakahara K. Percutaneous biopsy of thoracic lesions: value of sonography for needle guidance. *AJR Am J Roentgenol* 1990;154:1181-1185
  14. Yoon SH, Park CM, Lee KH, Lim KY, Suh YJ, Im DJ, et al. Analysis of complications of percutaneous transthoracic needle biopsy using CT-guidance modalities in a multicenter cohort of 10568 biopsies. *Korean J Radiol* 2019;20:323-331
  15. Lee KH, Lim KY, Suh YJ, Hur J, Han DH, Kang MJ, et al. Nondiagnostic percutaneous transthoracic needle biopsy of lung lesions: a multicenter study of malignancy risk. *Radiology* 2019;290:814-823
  16. Schuetz GM, Schlattmann P, Dewey M. Use of 3x2 tables with an intention to diagnose approach to assess clinical performance of diagnostic tests: meta-analytical evaluation of coronary CT angiography studies. *BMJ* 2012;345:e6717
  17. Fior D, Vacirca F, Leni D, Pagni F, Ippolito D, Riva L, et al. Virtual guidance of percutaneous transthoracic needle biopsy with C-arm cone-beam CT: diagnostic accuracy, risk factors and effective radiation dose. *Cardiovasc Intervent Radiol* 2019;42:712-719
  18. Jin KN, Park CM, Goo JM, Lee HJ, Lee Y, Kim JI, et al. Initial experience of percutaneous transthoracic needle biopsy of lung nodules using C-arm cone-beam CT systems. *Eur Radiol* 2010;20:2108-2115
  19. Ohno Y, Hatabu H, Takenaka D, Higashino T, Watanabe H, Ohbayashi C, et al. CT-guided transthoracic needle aspiration biopsy of small (< or = 20 mm) solitary pulmonary nodules. *AJR Am J Roentgenol* 2003;180:1665-1669
  20. Geraghty PR, Kee ST, McFarlane G, Razavi MK, Sze DY, Dake MD. CT-guided transthoracic needle aspiration biopsy of pulmonary nodules: needle size and pneumothorax rate. *Radiology* 2003;229:475-481
  21. Manhire A, Charig M, Clelland C, Gleeson F, Miller R, Moss H, et al. Guidelines for radiologically guided lung biopsy. *Thorax* 2003;58:920-936
  22. Yeow KM, Su IH, Pan KT, Tsay PK, Lui KW, Cheung YC, et al. Risk factors of pneumothorax and bleeding: multivariate analysis of 660 CT-guided coaxial cutting needle lung biopsies. *Chest* 2004;126:748-754
  23. Anderson JM, Murchison J, Patel D. CT-guided lung biopsy: factors influencing diagnostic yield and complication rate. *Clin Radiol* 2003;58:791-797
  24. Kim H, Kwon D, Yoon SH, Kim H, Park CM, Goo JM, et al. Bronchovascular injury associated with clinically significant hemoptysis after CT-guided core biopsy of the lung: radiologic and histopathologic analysis. *PLoS One* 2018;13:e0204064