

Commentary

Importation of Ticks on Companion Animals and the Risk of Spread of Tick-Borne Diseases to Non-Endemic Regions in Europe

Alicja Buczek * and Weronika Buczek

Chair and Department of Biology and Parasitology, Faculty of Health Sciences, Medical University of Lublin, 20-080 Lublin, Poland; wera1301@gmail.com

* Correspondence: alicja.buczek@umlub.pl; Tel.: +48-814-486-060

Simple Summary: The risk of transmission of pathogen-infected ticks by dogs and cats transported by humans has increased substantially in recent decades due to the rise in tourist and economic migration rates. Therefore, we highlight the role of companion animals, mainly dogs, travelling with their owners or importation of these animals in the transmission of ticks and tick-borne diseases to non-endemic areas. The brown dog tick *Rhipicephalus sanguineus*, which is a vector of numerous pathogens causing diseases in animals and humans, is imported most frequently to many European countries from endemic areas in the Mediterranean region or from other parts of the world. Additionally, alien tick species with high epizootic and epidemiological importance can be imported while attached to dog skin to Europe from other continents. Companion animals play an even greater role in the spread of autochthonous tick species and transmission of pathogens to other animals and humans. Before travelling to endemic areas of tick-borne diseases, tourists should be acquainted with prophylaxis methods to protect themselves and their companion animals against tick attacks.



Citation: Buczek, A.; Buczek, W. Importation of Ticks on Companion Animals and the Risk of Spread of Tick-Borne Diseases to Non-Endemic Regions in Europe. *Animals* **2021**, *11*, 6. <https://dx.doi.org/10.3390/ani11010006>

Received: 17 November 2020

Accepted: 16 December 2020

Published: 22 December 2020

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2020 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Increased human mobility elevates the risk of exposure of companion animals travelling with their owners or imported from other regions to tick attacks. In this study, we highlight the potential role of dogs and cats taken for tourist trips or imported animals in the spread of ticks and tick-borne pathogens. The *Rhipicephalus sanguineus* tick, which is a vector of numerous pathogens causing diseases in animals and humans, is imported most frequently from endemic areas to many European countries. Additionally, alien tick species with high epizootic and epidemiological importance can be imported on dogs from other continents. Companion animals play an even greater role in the spread of autochthonous tick species and transmission of tick pathogens to other animals and humans. Although the veterinary and medical effects of the parasitism of ticks carried by companion animals travelling with owners or imported animals are poorly assessed, these animals seem to play a role in the rapid spread of tick-borne diseases. Development of strategies for protection of the health of companion animals in different geographic regions should take into account the potential emergence of unknown animal tick-borne diseases that can be transmitted by imported ticks.

Keywords: ticks; companion animals; importation of ticks; ticks on travelling dogs; transmission of tick-borne pathogens

1. Introduction

Companion animals, such as dogs (*Canis lupus familiaris*) and cats (*Felis catus*), can be hosts for various tick species from the family Ixodidae [1–9]. In Europe, these animals are parasitized by tick species from the genera *Dermacentor*, *Haemaphysalis*, *Hyalomma*, *Rhipicephalus*, and *Ixodes* [10,11]. In the northern, western, and central parts of the continent, companion animals are most often attacked by the sheep tick *Ixodes ricinus*, the hedgehog tick *Ix. hexagonus*, the dog tick *Ix. canisuga*, and the meadow tick *Dermacentor reticulatus*. The most frequent attacks in the north-eastern part of Europe are ascribed to the taiga tick

Ix. persulcatus, whereas the brown dog tick *Rhipicephalus sanguineus* and *Rh. turanicus* have been reported to infest animals primarily in the southern part of Europe. Other tick species (e.g., the red sheep tick *Haemaphysalis punctata*, the rodent tick *Ha. concinna*, the ornate sheep tick *De. marginatus*, and the southern rodent tick *Ix. acuminatus*) infest dogs and cats less frequently [12–19]. The Mediterranean ticks *Hyalomma marginatum marginatum* and *Hy. lusitanicum* [16,20,21] and the rodent tick *Ix. trianguliceps* [16,22] have been found to attack dogs sporadically. In the group of ticks that parasitize dogs and cats, there are two species (i.e., *Rh. sanguineus* and *Ix. canisuga*) associated with the human environment, in which their entire development cycle can be completed. A majority of ticks parasitizing companion animals can attack humans (e.g., [23–25]).

Companion animals travelling with their owners over long distances on the same continent or between continents are exposed to an increased risk of infestations by various species of ticks with considerable vector competence and infection with pathogens transmitted by these ticks. Most often, ticks are imported on dogs or cats in the parasitic stage. However, unattached ticks can also be found in animal fur or transport containers, and later reach the skin of other animals or their keepers.

The spread of ixodid ticks by companion animals and infection of these hosts with tick-borne pathogens during tick feeding is facilitated by specific biological and physiological features distinguishing these arthropods from other bloodsucking ectoparasites.

Ticks are attached to host skin for several days or even weeks, depending on the species and developmental stage [26–28], and their feeding dynamics may be influenced by various factors, including the intensity of invasion [29], the presence of ticks from the same or different species co-feeding on the same host [30], the sexual structure of foraging specimens [31], and climate conditions [32].

During blood ingestion, infected ticks can introduce factors of tick-borne diseases contained in tick saliva (e.g., viruses, bacteria, and protozoa) into the host. Canine babesiosis (*Babesia* spp.), borreliosis (*Borrelia burgdorferi* s.l.), ehrlichiosis (*Ehrlichia* spp.), rickettsiosis (*Rickettsia* spp.), and anaplasmosis (*Anaplasma* spp.) [19,33–39] are the most common tick-borne diseases in dogs and/or cats.

Skin lesions (granuloma, alopecia) and allergic reactions [16,40–47] as well as tick paralysis caused by the neurotoxic components of tick saliva [48–53] are the direct effects of tick parasitism in companion animals. In dogs, tick bites may induce production of specific immunoglobulin E antibodies to oligosaccharide galactose- α -1, 3-galactose (anti- α -Gal antibodies), which can be a source of red meat allergy [54].

Incidents of importation of ticks to new areas by animals transported by humans have been repeatedly described in the world literature [55–60]. A majority of reports have described importation of ticks by exotic reptiles transported from endemic areas to other regions of the world [55,61–67]. However, the scale of this phenomenon and the potential threat to animal and human health posed by imported ticks and pathogens in the new environment have rarely been assessed.

In this study, we highlight the role of companion animals, mainly dogs, travelling with their owners or importation of these animals in the transmission of ticks and tick-borne diseases to non-endemic areas. We also emphasize the role of dogs and cats in the spread of autochthonous tick and pathogen species.

The risk of transmission of pathogen-infected ticks by dogs and cats transported by humans has increased substantially in recent decades due to the rise in tourist and economic migration rates. Most agents of tick-borne diseases (e.g., *Borrelia* spirochetes [13,68–71], *Rickettsia* [69,70,72–74], *Anaplasma phagocytophilum* [68,71,73], and tick-borne encephalitis virus (TBEV) [75,76]) are pathogenic to not only companion animals but also humans [25,73,77]. Companion animals and humans can also be infected with bacteria, protozoa, and helminths transmitted by other vectors (e.g., fleas, mosquitoes, and phlebotomine sand flies; [78–80]). Therefore, clinical and laboratory methods for diagnostics of diseases transmitted by ticks and other bloodsucking arthropods, which facilitate

identification of the pathogenic factor and effective treatment, arouse increasing interest (e.g., [25,81–87]).

Due to the increase in the incidence of animal and human tick-borne diseases recorded worldwide, there is a need to trace the potential routes of transmission thereof to areas where they did not occur or rarely appeared previously. Determination of these routes will prompt effective measures for limiting the harmful results of tick parasitism.

2. Alien and Autochthonous Ticks and Tick-Borne Pathogens Carried by Travelling and Imported Companion Animals

Cases of transport of ticks and associated pathogens by companion animals have been reported in various countries, but only some countries have monitored this process. In Europe, the role of dogs in the transmission of autochthonous and exotic tick species has been assessed, specifically in Germany [88–95], the UK [17,20,96,97], Benelux countries [98,99], Sweden [100], Switzerland [101], Hungary [15,102], Cyprus [103], Greece [104], Italy [105], Spain [106], and Portugal [107]. However, there are very few reports on the role of transported cats in the transmission of ticks to other habitats [108–110], which is probably related to the fact that cats accompany their travelling owners less frequently than dogs. Additionally, cats usually stay indoors when they accompany their owners and are thus not exposed to contact with ticks present in the environment.

Tick infestations and transmission of pathogens in dogs and cats are supported by the presence of tick species with high epidemiological importance in urban and suburban areas that are often visited by owners of these animals. An example is the occurrence of *Rh. sanguineus*, *Rh. turanicus*, *Rh. bursa*, *Ix. ricinus*, *De. marginatus*, and *Ha. punctata* in the parks of Rome and other Italian cities [111–113] and the *Rh. sanguineus* group, *Ha. parva*, *Hy. marginatum*, *Rh. bursa*, *Ix. ricinus*, *De. marginatus*, *Hy. anatolicum*, *Hy. detritum*, *Hy. excavatum*, and *Ha. punctata* in Ankara (Turkey) [114]. The likelihood of infection of companion animals with pathogens in these areas is associated with the level of the prevalence of these microorganisms in ticks, which is higher than in natural habitats [115–118].

Dogs are usually infested by tick adult stages [15,16], less often by nymphs [15,16,106], and sporadically by larvae [16,20,22,46,119]. Dog's head (especially the eye area, ears, and muzzle), legs, and neck are the most commonly attacked body areas [16,120,121]. The frequency of tick attachment to the skin is associated with the breed of dogs and their temperament [120]. In cats, the greatest number of ticks attach to the head (ears, chin, face, and muzzle) and the neck [121].

In the world, *Rh. sanguineus* sensu lato is the most commonly imported dog tick (e.g., [20,101,109,122–124]). This species is distributed in large areas in warm climate regions in both hemispheres, where it can be found in kennels, buildings, and gardens [4,125].

Rh. sanguineus is a common tick species in the Mediterranean region, from where specimens are most often imported on dogs to countries in the central and northern part of the continent, including the UK [109,126–129], Germany [88,92,130–135], Poland [136], the Czech Republic [137,138], the Netherlands [98,139], Denmark [140–143], Belgium [108,144], and Switzerland [145]. Less frequently, *Rh. sanguineus* ticks are transferred to European countries from other endemic areas of this species, like the Dominican Republic [88] and Sudan [140]. As shown in literature, the number of these cases clearly increased after the economic and political transformations in Europe at the end of the 20th century. This is associated with the growing interest in tourism and business trips to remote locations in Europe, Americas, Africa, and Asia, where high prevalence of ticks from the *Rh. sanguineus* group in dogs from urban and rural environments is observed. For instance, *Rh. sanguineus* sensu lato (s.l.) has been found to attack 63.6% of autochthonous dogs in Italy [146], 53% in Spain [106], 51.32% in Northern Algeria [147], 59.6% on the Mexico-USA border [148], 80% in north-central Nigeria [149], 80% in north-eastern Thailand [150], and as many as 98.33% in Punjab, Pakistan [151]. Similarly, in the endemic areas of the *Rh. sanguineus* group (*Rh. sanguineus* sensu stricto, and *Rh. pusillus*) in southern Italy, these species most often infest cats [152].

Travelling and imported dogs have been shown to be infested by the largest number of adults and nymphs but only single larvae of *Rh. sanguineus* and other tick species [20]. An unusual case was the transfer of approximately 50 or more mobile engorged and unfed *Rh. sanguineus* larvae infesting a five-old Jack Russel, which was noticed 18 h after the dog had returned from Greece [153].

The importation of *Rh. sanguineus* s.l. ticks on animals to non-endemic areas has considerable importance for public health, as they are vectors and/or potential reservoirs of numerous pathogens. These include *Ehrlichia canis*, *Anaplasma platys*, and *A. phagocytophilum*, protozoa *Babesia vogeli*, *B. canis*, *Babesia vulpes*, and *Hepatozoon canis*, zoonotic *Rickettsia* species (*Rickettsia conorii* complex and *R. massiliae*), and the filarioid nematode *Cercopithifilaria bainae* [154–160]. Besides *Rh. sanguineus*, other alien tick species were occasionally transported by dogs to various European countries (Tables 1 and 2).

Table 1. Examples of alien ticks imported on dogs and cats into non-endemic areas in Europe.

Tick Species	Areas of Tick Infestations of Dogs or Cats	Place of Import of Ticks	Source of Information Authors, [Ref.]
<i>Ixodes holocyclus</i>	Australia	United Kingdom (UK)	Adamantos et al. [49]
<i>Ixodes pacificus</i>	USA	UK	Jameson et al. [109]
<i>Amblyomma americanum</i>	USA	UK	Jameson et al. [109]
<i>Dermacentor variabilis</i>	USA	UK	Abdullah et al. [17]; Jameson et al. [109]
<i>Dermacentor albipictus</i>	USA	UK	Jameson et al. [109]
<i>Haemaphysalis leachi</i>	South Africa, Tanzania, Zambia	UK	Dutto and Selmi [16]
	Rhodesia (Zimbabwe), South Africa, Tanzania, Zambia	UK	Jameson et al. [109]
<i>Haemaphysalis elliptica</i>	South Africa	UK	Hansford et al. [123]
<i>Hyalomma lusitanicum</i>	Portugal	UK	Hansford et al. [20]
<i>Rhipicephalus sanguineus</i>	Sudan	Denmark	Winding and Haarløv [140]
	Zambia	Denmark	Haarløv [141]
	No data	Denmark	Winding et al. [142]
	No data	Denmark	Willeberg [143]
	Spain, Dominican Republic, Italy, France, Greece	Germany	Dongus et al. [88]
	No data	Belgium	Claerebouts et al. [108]
	No data	UK	Hansford et al. [123]
	Cyprus, Spain	UK	Hansford et al. [123]
	Africa, Australia, Canada, Iran, Malta, Philipines, Singapore, West Indies, Saudi Arabia, South Africa, Mideterranean region (Spain, Cyprus, Tunisia), USA	UK	Jameson et al. [109]
	Greece	UK (London)	Wright et al. [121]
<i>Haemaphysalis leachi</i> *	Africa	UK	Jameson et al. [109]

* ticks imported on cats.

Table 2. Examples of autochthonous tick species carried by imported and travelling dogs in Europe

Tick Species	Areas of Tick Infestations of Dogs	Place of Import of Ticks	Source of information Authors, [Ref.]
<i>Ixodes ricinus</i>	Belgium, The Netherlands, Germany, Finland	United Kingdom (UK) UK	Jameson et al. [109] Hansford et al. [123]
<i>Ixodes hexagonus</i>	Germany France	UK UK	Jameson et al. [109] Hansford et al. [123]
<i>Ixodes canisuga</i>	France	UK	Hansford et al. [123]
<i>Ixodes ventalloi</i>	France (Montpellier)	Northwest Italy	Dutton and Selmi [16]
<i>Dermacentor reticulatus</i>	France, Bulgaria	UK	Hansford et al. [123]

Some of these tick species can live in new habitats with favourable conditions. Engorged females lay eggs, and juvenile stages (larvae and nymphs) transform into subsequent developmental stages. Hungry active stages of these ticks can attack hosts present in the surroundings [88,161]. The introduction of some species (*Rh. sanguineus*, *Hy. marginatum*, *Hy. suspense*) to non-endemic areas is possible due to the climate and environmental changes occurring worldwide (e.g., [145,162,163]).

An interesting case of introduction of *Rh. sanguineus* to Poland and mass reproduction of this tick in an apartment in Warsaw was described by Szymański [136]. These ticks from a dog travelling with an Italian family infested the dog of the Polish family when they spent holidays together in the Mazury region. After return to Warsaw, an engorged *Rh. sanguineus* female detached from the host skin and laid eggs, from which larvae developed. *Rh. sanguineus* ticks can complete their full life cycle indoors. Cases of household *Rh. sanguineus* infestations have been reported from other countries as well, including Germany [88], England [96,97,164], and France [165].

In Europe, dogs and cats most frequently transfer autochthonous species of ticks, which are vectors of numerous animal tick-borne diseases. For instance, a group of 2373 ticks removed from dogs and cats in Belgium was dominated by the castor bean tick *Ix. ricinus* (76.4%) and the European hedgehog tick *Ix. hexagonus* (22.6%); single specimens of the meadow tick *De. reticulatus* (0.8%) were found as well. *Rh. sanguineus* specimens, which accounted for 0.3% of the total number, were found on dogs after return from endemic areas of this tick species [108].

In the UK in 2005–2016, *Rh. sanguineus* (46.2%) and *I. ricinus* (33.8%) ticks were the most common species of ticks found on animals with a history of travel mainly to southern Europe, the USA, and the United Arab Emirates [20]. In Spain, *Rh. sanguineus* s.l. (53%) was the most abundant autochthonous species in a group of 1628 collected adult ticks. Other ticks accounted for a lower percentage: *De. reticulatus* (9%), *Ix. ricinus* (9%), and *Ix. hexagonus* (4%). In a group of 660 dogs sampled in 26 veterinary clinics, as many as 507 dogs (76.8%) were infested by at least one adult tick [86]. In Greece, the majority of ticks infesting 150 dogs (48.4%) were *Rh. sanguineus* s.l. (70.1%), whereas other species were less numerous (i.e., *Ha. parva* (14.7%), *Rh. turanicus* (11.4%), and *Ha. concinna* (2.4%)). A total of 11.1% of 344 sampled ticks were vectors of at least one microorganism (5.5%: *Cercopithifilaria bainae*, 2.9%: *Hepatozoon canis*, 1.7%: *Rickettsia hoogstraalii*, 1.2%: *Hepatozoon felis*, 0.6%: *Rickettsia massiliae*, 0.6%: *Theileria ovis*, 0.3%: *Anaplasma platys*, and 0.3%: *Coxiella* like-endosymbiont) [166].

Rh. sanguineus transferred on travelling and imported dogs into non-endemic areas may pose a health risk by transmission of pathogens (e.g., *Ehrlichia canis*) to native dogs [167,168]. Canine hepatosporosis caused by the apicomplexan protozoan *Hepatozoon canis* was diagnosed in three dogs imported to the United Kingdom from Mediterranean

countries [169]. The infection by pathogens acquired by animals in the travel destination area is often confirmed by serological tests only after returning home [170]. Dogs are a reservoir of infectious agents of human diseases, like Mediterranean spotted fever caused by *Rickettsia conorii* [171,172].

Various microorganisms were detected in *Rh. sanguineus* ticks carried into new areas on dogs, including *Ri. conorii* [173,174], *Ri. massiliae*/Bar29, and *Coxiella* sp. [175]. Contact with dogs or cats infested by pathogen-infected *Rh. sanguineus* ticks increases the likelihood of the spread of tick-borne pathogens, like spotted fever group (SFG) rickettsiae. Symptoms of spotted fever group (SFG) rickettsiosis were confirmed in members of a family from south France living in a house where 22 *Rh. sanguineus* ticks were collected from the floor behind furniture. Of these ticks, 20 were infected with *Rickettsia*; more specifically, nine and seven specimens were infected by *Ri. conorii* subsp. *caspia* and *Ri. massiliae*, respectively, whereas both rickettsiae were identified in four ticks [165]. As shown by Edouard et al. [176], some patients in France were diagnosed with rickettsiosis caused by *Ri. sibirica mongolitimonae*, another representative of the spotted fever group. *Ri. massiliae* and *Ri. sibirica mongolitimonae* were detected in *Rh. sanguineus* and *Rh. pusillus* ticks sampled from their dogs and cats, respectively. *Rh. pusillus* ticks infected by *Ri. sibirica mongolitimonae* were also collected from cats' litter boxes and beds. There are reports of cases of human infection with *Ri. conorii* contracted during removal of ticks from dogs [177].

An especially high risk of development of tick-borne and other arthropod-borne diseases in companion animals has been observed in attractive tourist regions characterized by a high prevalence of pathogens transmitted by ticks and other arthropods (e.g., [79,178–185]). For instance, it was reported that the prevalence of *Ri. conorii* in dogs in southern Italy ranged from 72% to 73.60%. The prevalence of *An. phagocytophilum* was in the range from 32.80% to 38%, *B. canis* from 5.17% do 70%, and *Eh. canis* from 21.70% to 46%. The prevalence of *Co. burnetii* was estimated at 31.50% [181,186].

In the north of Spain, antibodies against these pathogens were detected in 11.33% of tested dogs, with prevalence of 1.26% for *Anaplasma* spp., 0.9% for *Eh. canis*, and 0.72% for *Bo. burgdorferi* transmitted by ticks and 8.99% for *Le. infantum* and 0.18% for *Di. immitis* transmitted by sand flies and culicid mosquitoes, respectively [187].

3. Tick Preventive Behaviors and Practices of Dog and Cat Owners

The effects of tick parasitism on companion animals and their owners can be limited by application of various methods for protection against tick attacks recommended by the Centers for Disease Control and Prevention [188] and the European Scientific Counsel Companion Animal Parasites [189].

The risk of human and animal contact with ticks can be minimized by avoidance of habitats with high numbers of ticks at the peak of their seasonal and diurnal activity. Another important element of tick prophylaxis is the preventive behavior of animal owners outdoors (e.g., limitation of contact with plants, avoidance of sitting on grass and under shrubs, wearing protective clothes to prevent ticks from attachment to the skin, inspection of clothes or skin to find ticks), and removal of the feeding tick from the skin as soon as possible after attachment. Tick attacks can be limited with effective repellents (e.g., [190]). As shown in various studies, inspection of the body on return home, wearing protective clothes, and the use of repellents are the most preferred methods used by humans for personal protection against tick attachment to the skin [191–197].

Owners of dogs and cats are advised to inspect the hair and skin of their animals, especially the head, ears, shoulders, and upper leg areas, when the animals stay outdoors during the tick activity period. They should also use various acaricidal and/or repellent substances contained in numerous formulations available on the market (e.g., spot-on, pour-on, baths, and insecticide-impregnated collars).

Orally administered parasiticides against ticks are also effective in protecting dogs against ticks and tick-borne pathogens (e.g., [198–202]). A single oral dose reduced the

survival rate of over 98% of ticks present on the host 48 h after treatment and effectively protected against new infestations for several weeks [198,199].

Specimens attached to host's skin are removed with various methods described in various publications (e.g., [203,204]).

4. Conclusions

The most important role in rapid transmission of ticks and associated pathogens over long distances is ascribed to seasonally migrating birds (e.g., [205–210]). Nevertheless, it seems that animals accompanying their owners during foreign and domestic journeys or animals imported from other geographic regions may contribute considerably as well. The problem has to be emphasised to draw attention of owners of dogs and cats as well as veterinary staff, due to the dramatic increase in the prevalence of tick-borne diseases in these animals (e.g., [19,79,211–217]).

Before travelling to endemic areas of tick-borne diseases, tourists should be acquainted with prophylaxis methods to protect themselves and their companion animals against tick attacks. Information about tick species that infest dogs and cats most frequently, periods of the highest tick questing activity, and threats of zoonotic tick-borne diseases posed to animal health is indispensable for owners of companion animals.

Author Contributions: Conceptualization, A.B.; writing and reviewing of the original draft and supervision, A.B.; writing of the original draft and editing, W.B. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Acknowledgments: In this section you can acknowledge any support given which is not covered by the author contribution or funding sections. This may include administrative and technical support, or donations in kind (e.g., materials used for experiments).

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Hoogstraal, H. A preliminary, annotated list of ticks (Ixodoidea) of the Anglo-Egyptian Sudan. *J. Parasitol.* **1954**, *40*, 304–310. [[CrossRef](#)]
2. Arthur, D.R. *Ticks and Disease*; Pergamon Press: Oxford, UK, 1962; p. 445.
3. Nijhof, A.M.; Bodaan, C.; Postigo, M.; Nieuwenhuijs, H.; Opsteegh, M.; Fransses, L.; Jebbink, F.; Jongejan, F. Ticks and associated pathogens collected from domestic animals in the Netherlands. *Vector Borne Zoonotic Dis.* **2007**, *7*, 585–595. [[CrossRef](#)] [[PubMed](#)]
4. Kolonin, G.V. Fauna of Ixodid Ticks of the World (Acari, Ixodidae), Moscow. 2009. Available online: <https://www.kolonin.org/3.html> (accessed on 24 August 2009).
5. Tsai, Y.-L.; Shyu, C.-L.; Yao, C.-T.; Lin, J.A. The ixodid ticks collected from dogs and other animals in Taiwan and Kinmen Island. *Int. J. Acarol.* **2012**, *38*, 110–115. [[CrossRef](#)]
6. Barker, S.C.; Walker, A.R. Ticks of Australia. The species that infest domestic animals and humans. *Zootaxa* **2014**, *3816*, 1–144. [[CrossRef](#)]
7. Portugaliza, H.P.; Bagot, M.A. Different species of lice (Phthiraptera), fleas (Siphonaptera) and ticks (Ixodida) collected from livestock, poultry, reptile and companion animal in Leyte Island, Philippines. *LRRD* **2015**, *27*, 1–10.
8. Durden, L.A.; Beckmen, K.B.; Gerlach, R.F. New records of ticks (Acari: Ixodidae) from dogs, cats, humans, and some wild vertebrates in Alaska: Invasion potential. *J. Med. Entomol.* **2016**, *53*, 1391–1395. [[CrossRef](#)]
9. Greay, T.L.; Oskam, C.L.; Gofton, A.W.; Rees, R.L.; Ryan, U.M.; Irwin, P.J. A survey of ticks (Acari: Ixodidae) of companion animals in Australia. *Parasites Vectors* **2016**, *9*, 207. [[CrossRef](#)]
10. Estrada-Peña, A.; Bouattour, A.; Camicas, J.L.; Walker, A.R. *Ticks of Domestic Animals in the Mediterranean Region. A Guide of Identification of Species*; University of Zaragoza Press: Zaragoza, Spain, 2004; p. 131.
11. Nowak-Chmura, M. *Fauna Kleszczy (Ixodida) Europy Środkowej*; Wyd. Naukowe Uniwersytetu Pedagogicznego: Kraków, Poland, 2013; p. 290.
12. Encinas Grandes, A. Ticks of the province of Salamanca (Central/NW Spain). Prevalence and parasitization intensity in dogs and domestic ungulates. *Ann. Parasitol. Hum. Comp.* **1986**, *61*, 95–107. [[CrossRef](#)]
13. Beichel, E. Tick infestation patterns and prevalence of *Borrelia burgdorferi* in ticks collected at a veterinary clinic in Germany. *Vet. Parasitol.* **1996**, *65*, 147–155. [[CrossRef](#)]
14. Ogden, N.H.; Cripps, P.; Davison, C.C.; Owen, G.; Parry, J.M.; Timms, B.J.; Forbes, A.B. The ixodid tick species attaching to domestic dogs and cats in Great Britain and Ireland. *Med. Vet. Entomol.* **2000**, *14*, 332–338. [[CrossRef](#)]

15. Földvari, G.; Farkas, R. Ixodid tick species attaching to dogs in Hungary. *Vet. Parasitol.* **2005**, *129*, 125–131. [CrossRef] [PubMed]
16. Dutto, M.; Selmi, M. Report on ticks (Acaria: Ixodidae) collected on dogs in Piedmont: The potential transmission of tick-borne pathogens to man. *Parassitologia* **2009**, *51*, 113–117.
17. Abdullah, S.; Helps, C.; Tasker, S.; Newbury, N.; Wall, R. Ticks infesting domestic dogs in the UK: A large-scale surveillance programme. *Parasites Vectors* **2016**, *9*, 391. [CrossRef] [PubMed]
18. Medlock, J.M.; Hansford, K.; Vaux, A.G.C.; Cull, B.; Pietzsch, M.E.; Gillingham, E.L.; Johnson, N.; Phipps, L.P. Has the red sheep tick, *Haemaphysalis punctata*, recently expanded its range in England?: *Haemaphysalis punctata* in the U.K. *Med. Vet. Entomol.* **2018**, *32*. [CrossRef]
19. Rudek, M.; Buczek, W.; Buczek, A.M.; Błaszkiewicz, P.S.; Kulina, D.; Ciura, D.; Buczek, A. The state of research on the infestation of companion animals by ticks (Ixodida) and their epizootiological role in Poland. In *Parasitic and Allergic Arthropods*; Part 1; Buczek, A., Błaszkak, C., Eds.; Koliber: Lublin, Poland, 2019; pp. 73–89.
20. Hansford, K.M.; Medlock, J.M.; Atkinson, B.; Santos-Silva, M.M. Importation of a *Hyalomma lusitanicum* tick into the UK on a dog. *Vet. Rec.* **2016**, *179*, 415. [CrossRef]
21. Hansford, K.M.; Pietzsch, M.E.; Cull, B.; Gillingham, E.L.; Medlock, J.M. Potential risk posed by the importation of ticks into the UK on animals: Records from the Tick Surveillance Scheme. *Vet. Rec.* **2018**, *182*, 107. [CrossRef]
22. Manilla, G. *Acaria, Ixodida (Fauna d’Italia)*; Edizioni Cal-Derini: Bologna, Italy, 1998.
23. Bartosik, K.; Sitarz, M.; Szymańska, J.; Buczek, A. Tick bites on humans in the agricultural and recreational areas in south-eastern Poland. *Ann. Agric. Environ. Med.* **2011**, *18*, 151–157.
24. Földvári, G.; Rigó, K.; Lakos, A. Transmission of Rickettsia slovaca and Rickettsia raoultii by male *Dermacentor marginatus* and *Dermacentor reticulatus* ticks to humans. *Diagn. Microbiol. Infect. Dis.* **2013**, *76*, 387–389. [CrossRef]
25. Blanda, V.; Torina, A.; La Russa, F.; D’Agostino, R.; Randazzo, K.; Scimeca, S.; Giudice, E.; Caracappa, S.; Cascio, A.; de la Fuente, J. A retrospective study of the characterization of *Rickettsia* species in ticks collected from humans. *Ticks Tick Borne Dis.* **2017**, *8*, 610–614. [CrossRef]
26. Balashov, Y.S. Blood sucking ticks (Ixodidae)-vectors of diseases of man and animals. *Misc. Publ. Entomol. Soc. Am.* **1972**, *8*, 161–376.
27. Sonenshine, D.E.; Roe, M. *Biology of Ticks*; Oxford University Press: New York, NY, USA, 2014; Volume 1, p. 560.
28. Levin, M.L.; Schumacher, L.B.M. Manual for maintenance of multi-host ixodid ticks in the laboratory. *Exp. Appl. Acarol.* **2016**, *70*, 343–367. [CrossRef] [PubMed]
29. Bartosik, K.; Buczek, A. The impact of intensity of invasion of *Ixodes ricinus* and *Dermacentor reticulatus* on the course of the parasitic phase. *Ann. Agric. Environ. Med.* **2012**, *19*, 651–655. [PubMed]
30. Buczek, A.; Bartosik, K.; Zajac, Z.; Stanko, M. Host-feeding behaviour of *Dermacentor reticulatus* and *Dermacentor marginatus* in mono-specific and inter-specific infestations. *Parasites Vectors* **2015**, *8*, 470. [CrossRef] [PubMed]
31. Bartosik, K.; Buczek, A.; Buczek, W.; Buczek, A.M.; Kulina, D.; Koman-Iżko, A. Host feeding behaviour of *Dermacentor reticulatus* males in relation to the transmission of pathogens. *Ann. Agric. Environ. Med.* **2019**, *26*, 227–230. [CrossRef]
32. Karbowiak, G.; Izdebska, J.N.; Czaplińska, U.; Wita, I. Cases of survival of the winter by Ixodidae ticks on the hosts in the Białowieża primeval forest. In *Arthropods and Hosts*; Buczek, A., Błaszkak, C., Eds.; Koliber: Lublin, Poland, 2003; pp. 77–82.
33. Zygnier, W.; Wiśniewski, M. Tick-transmitted diseases which may threaten health of dogs in Poland. *Wiad. Parazytol.* **2006**, *52*, 85–92. (In Polish)
34. Otranto, D.; Capelli, G.; Genchi, C. Changing distribution patterns of canine vector borne diseases in Italy: Leishmaniosis vs. dirofilariosis. *Parasites Vectors* **2009**, *2* (Suppl. 1), S2. [CrossRef]
35. Pérez Vera, C.; Kapiainen, S.; Junnikkala, S.; Aaltonen, K.; Spillmann, T.; Vapalahtiet, O. Survey of selected tick-borne diseases in dogs in Finland. *Parasites Vectors* **2014**, *7*, 285. [CrossRef]
36. Pantchev, N.; Pluta, S.; Huiszinga, E.; Nather, S.; Scheufelen, M.; Vrhovec, M.G.; Schweinitz, A.; Hampel, H.; Straubinger, R.K. Tick-Borne diseases (Borreliosis, Anaplasmosis, Babesiosis) in German and Austrian dogs: Status quo and review of distribution, transmission, clinical findings, diagnostics and prophylaxis. *Parasitol. Res.* **2015**, *114*, 19–54. [CrossRef]
37. Swainsbury, C.; Bengtson, G.; Hill, P. Babesiosis in dogs. *Vet. Rec.* **2016**, *178*, 172. [CrossRef]
38. Livanova, N.N.; Fomenko, N.V.; Akimov, I.A.; Ivanov, M.J.; Tikunova, N.V.; Armstrong, R.; Konyaev, S.V. Dog survey in Russian veterinary hospitals: Tick identification and molecular detection of tick-borne pathogens. *Parasites Vectors* **2018**, *1*, 591. [CrossRef]
39. Skotarczak, B. The role of companion animals in the environmental circulation of tick-borne bacterial pathogens. *Ann. Agric. Environ. Med.* **2018**, *25*, 473–480. [CrossRef] [PubMed]
40. Theis, J.H.; Budwiser, P.D. *Rhipicephalus sanguineus*: Sequential histopathology at the host-arthropode interface. *Exp. Parasitol.* **1974**, *36*, 77–105. [CrossRef]
41. Bechara, G.H.; Szabó, M.P.J.; Mukai, L.S.; Rosa, P.C.S. Immunisation of dogs, hamsters and guinea pigs against *Rhipicephalus sanguineus* using crude unfed adult tick extracts. *Vet. Parasitol.* **1994**, *52*, 79–90. [CrossRef]
42. Szabó, M.P.J.; Bechara, G.H. An insight into the histopathology caused by the tick *Rhipicephalus sanguineus* (Acarina: Ixodidae) in the skin of previously infested, vaccinated or tick-bite naive dogs, guinea pigs and hamsters. *Braz. J. Vet. Res. Anim. Sci.* **1995**, *32*, 37–42. [CrossRef]

43. Szabó, M.P.J.; Mukai, L.S.; Rosa, P.C.S.; Bechara, G.H. Differences in the acquired resistance of dogs, hamsters, and guinea pigs to repeated infestations with adult ticks *Rhipicephalus sanguineus* (Acari: Ixodidae). *Braz. J. Vet. Res. Anim. Sci.* **1995**, *32*, 43–50. [[CrossRef](#)]
44. Szabó, M.P.J.; Bechara, G.H. Sequential histopathology at the *Rhipicephalus sanguineus* tick feeding site on dogs and guinea pigs. *Exp. Appl. Acarol.* **1999**, *23*, 915–928. [[CrossRef](#)]
45. Pappalardo, B.L.; Brown, T.; Gookin, J.L.; Morrill, C.L.; Breitschwerdt, E.B. Granulomatous disease associated with *Bartonella* infections in two dogs. *J. Vet. Med.* **2000**, *14*, 37–42.
46. Kumsa, B.E.; Mekonnen, S. Ixodid ticks, fleas and lice infesting dogs and cats in Hawassa, southern Ethiopia. *Onderstepoort J. Vet. Res.* **2011**, *78*, E326. [[CrossRef](#)]
47. Christensson, D.; Zakrisson, G. Ticks, *Ixodes ricinus* in the sub-cutaneous tissues of a dog and foxes. *Svensk Vet.* **2010**, *62*, 25–27. (In Swedish)
48. Atwell, R.B.; Campbell, F.E.; Evans, E.A. Prospective survey of tick paralysis in dogs. *Aust. Vet. J.* **2001**, *79*, 412–418. [[CrossRef](#)]
49. Adamantos, S.; Boag, A.; Church, D. Australian tick paralysis in a dog imported into the UK. *Aust. Vet. J.* **2005**, *83*, 352–353. [[CrossRef](#)] [[PubMed](#)]
50. Otranto, D.; Dantas-Torres, F.; Tarallo, V.D.; do Nascimento Ramos, R.A.; Stanneck, D.; Baneth, G.; de Caprariis, D. Apparent tick paralysis by *Rhipicephalus sanguineus* (Acari: Ixodidae) in dogs. *Vet. Parasitol.* **2012**, *188*, 325–329. [[CrossRef](#)] [[PubMed](#)]
51. Eppleston, K.R.; Kelman, M.; Ward, M.P. Distribution, seasonality and risk factors for tick paralysis in Australian dogs and cats. *Vet. Parasitol.* **2013**, *196*, 460–468. [[CrossRef](#)] [[PubMed](#)]
52. Leister, E.; Morton, J.; Atwell, R.; Webster, R. Clinical presentations, treatments and risk factors for mortality in cats with tick paralysis caused by *Ixodes holocyclus*: 2077 cases (2008–2016). *J. Feline Med. Surg.* **2018**, *20*, 465–478. [[CrossRef](#)]
53. Padula, A.M.; Leister, E.M.; Webster, R.A. Tick paralysis in dogs and cats in Australia: Treatment and prevention deliverables from 100 years of research. *Vet. J.* **2020**, *98*, 53–59. [[CrossRef](#)]
54. Hodžić, A.; Mateos-Hernández, L.; Leschnik, M.; Alberdi, P.; Rego, R.O.M.; Contreras, M.; Villar, M.; de la Fuente, J.; Cabezas-Cruz, A.; Duscher, G.G. Tick bites induce anti- α -gal antibodies in dogs. *Vaccines* **2019**, *7*, 114. [[CrossRef](#)]
55. Becklund, W.W. Ticks of veterinary significance found on imports in the United States. *J. Parasitol.* **1968**, *54*, 622–628. [[CrossRef](#)]
56. Keirans, J.E.; Durden, L.A. Invasion: Exotic ticks (Acari: Argasidae, Ixodidae) imported into the United States. A review and new records. *J. Med. Entomol.* **2001**, *38*, 850–861. [[CrossRef](#)]
57. Liebisch, G.; Liebisch, A.; Paufler, S. Importation of a New World tick, *Dermacentor albipictus* (Acari: Ixodidae), with a horse from the USA into Germany. In *Trends in Acarology, Proceedings of the 12th International Congress, Heraklion, Greece, 21–25 September 2016*; Sabelis, M., Bruun, J., Eds.; Springer: Dordrecht, The Netherlands, 2010.
58. Navajas, M.; Migeon, A.; Estrada-Peña, A.; Mailleux, A.C.; Servigne, P.; Petanović, R.; Roques, A. Mites and ticks (Acari). Chapter 7.4. Arthropod invasions in Europe. *BioRisk* **2010**, *4*, 149–192. [[CrossRef](#)]
59. Schvartz, G.; Epp, T.; Burgess, H.J.; Chilton, N.B.; Armstrong, J.S.; Lohmann, K.L. Passive surveillance for ticks on horses in Saskatchewan. *Can. Vet. J.* **2015**, *56*, 486–489.
60. Chitimia-Dobler, L.; Schaper, S.; Rieß, R.; Bitterwolf, K.; Frangoulidis, D.; Bestehorn, M.; Springer, A.; Oehme, R.; Drehmann, M.; Lindau, A.; et al. Imported *Hyalomma* ticks in Germany in 2018. *Parasites Vectors* **2019**, *12*, 134. [[CrossRef](#)] [[PubMed](#)]
61. Simmons, L.A.; Burridge, M.J. Introduction of the exotic ticks *Amblyomma humerale* Koch and *Amblyomma geoemydae* (Cantor) (Acari: Ixodidae) into the United States on imported reptiles. *Int. J. Acarol.* **2000**, *26*, 239–242. [[CrossRef](#)]
62. Simmons, L.A.; Burridge, M.J. Introduction of the exotic tick *Amblyomma chabaudi* rageau (Acari: Ixodidae) into Florida on imported tortoises. *Fla. Entomol.* **2002**, *85*, 288–289. [[CrossRef](#)]
63. Burridge, M.J.; Simmons, L.A. Exotic ticks introduced into the United States on imported reptiles from 1962 to 2001 and their potential roles in international dissemination of diseases. *Vet. Parasitol.* **2003**, *113*, 289–320. [[CrossRef](#)]
64. Nowak-Chmura, M. Teratological changes in tick morphology in ticks feeding on exotic reptiles. *J. Nat. Hist.* **2012**, *46*, 911–921. [[CrossRef](#)]
65. Nowak-Chmura, M. A biological/medical review of alien tick species (Acari: Ixodida) accidentally transferred to Poland. *Ann. Parasitol.* **2014**, *60*, 49–59. [[PubMed](#)]
66. Paștiu, A.I.; Matei, I.A.; Mihalca, A.D.; D’Amico, G.; Dumitache, M.O.; Kalmár, Z.; Sándor, A.; Lefkaditis, M.; Gherman, C.M.; Cozma, V. Zoonotic pathogens associated with *Hyalomma aegyptium* in endangered tortoise: Evidence for host-switching behavior in ticks? *Parasites Vectors* **2012**, *5*, 301–306. [[CrossRef](#)]
67. Mihalca, A.D. Ticks imported to Europe with exotic reptiles. *Vet. Parasitol.* **2015**, *213*, 67–71. [[CrossRef](#)]
68. Zygnier, W.; Górska, P.; Wędrychowicz, H. Detection of the DNA of *Borrelia afzelii*, *Anaplasma phagocytophilum* and *Babesia canis* in blood samples from dogs in Warsaw. *Vet. Rec.* **2009**, *164*, 465–467. [[CrossRef](#)]
69. Schreiber, C.; Krücken, J.; Beck, S.; Maaz, D.; Pachnicke, S.; Krieger, K.; Gross, M.; Kohn, B.; von Samson-Himmelstjerna, G. Pathogens in ticks collected from dogs in Berlin/Brandenburg, Germany. *Parasites Vectors* **2014**, *7*, 535. [[CrossRef](#)]
70. Morganti, G.; Gavaudan, S.; Canonico, C.; Ravagnan, S.; Olivieri, E.; Diaferia, M.; Marenzoni, M.L.; Antognoni, M.T.; Capelli, G.; Silaghi, C.; et al. Molecular survey on *Rickettsia* spp., *Anaplasma phagocytophilum*, *Borrelia burgdorferi* sensu lato, and *Babesia* spp. in *Ixodes ricinus* ticks infesting dogs in Central Italy. *Vector Borne Zoonotic Dis.* **2017**, *17*, 743–748. [[CrossRef](#)] [[PubMed](#)]
71. Namina, A.; Capligina, V.; Seleznova, M.; Krumins, R.; Aleinikova, D.; Kivrane, A.; Akopjana, S.; Lazovska, M.; Berzina, I.; Ranka, R. Tick-Borne pathogens in ticks collected from dogs, Latvia, 2011–2016. *BMC Vet. Res.* **2019**, *15*, 398. [[CrossRef](#)] [[PubMed](#)]

72. Mannelli, A.; Mandola, M.L.; Pedri, P.; Tripoli, M.; Nebbia, P. Associations between dogs that were serologically positive for *Rickettsia conorii* relative to the residences of two human cases of Mediterranean spotted fever in Piemonte (Italy). *Prev. Vet. Med.* **2006**, *60*, 13–26. [[CrossRef](#)]
73. Nicholson, W.L.; Allen, K.E.; McQuiston, J.H.; Breitschwerdt, E.B.; Little, S.E. The increasing recognition of rickettsial pathogens in dogs and people. *Trends Parasitol.* **2010**, *26*, 205–212. [[CrossRef](#)]
74. Król, N.; Obiegala, A.; Pfeffer, M.; Lonic, E.; Kiewra, D. Detection of selected pathogens in ticks collected from cats and dogs in the Wrocław Agglomeration, South-West Poland. *Parasites Vectors* **2016**, *9*, 351. [[CrossRef](#)]
75. Leschnik, M.W.; Kirtz, G.C.; Thalhammer, J.G. Tick-Borne encephalitis (TBE) in dogs. *Int. J. Med. Microbiol.* **2002**, *291* (Suppl. 33), 66–69. [[CrossRef](#)]
76. Lindhe, K.E.; Meldgaard, D.S.; Jensen, P.M.; Houser, G.A.; Berendt, M. Prevalence of tick-borne encephalitis virus antibodies in dogs from Denmark. *Acta Vet. Scand.* **2009**, *51*, 56. [[CrossRef](#)]
77. Nelder, M.P.; Russell, C.B.; Sheehan, N.J.; Sander, B.; Moore, S.; Li, Y.; Johnson, S.; Patel, S.N.; Sider, D. Human pathogens associated with the blacklegged tick *Ixodes scapularis*: A systematic review. *Parasites Vectors* **2016**, *9*, 265. [[CrossRef](#)]
78. Tabar, M.D.; Roura, X.; Francino, O.; Altet, L.; Ruiz de Gopegui, R. Detection of *Leishmania infantum* by real-time PCR in a canine blood bank. *J. Small Anim. Pract.* **2008**, *49*, 325–328. [[CrossRef](#)]
79. Otranto, D.; Dantas-Torres, F. Canine and feline vector-borne diseases in Italy: Current situation and perspectives. *Parasites Vectors* **2010**, *3*, 2. [[CrossRef](#)]
80. Adem, P.V. Emerging and re-emerging rickettsial infections. *Semin. Diagn. Pathol.* **2019**, *36*, 146–151. [[CrossRef](#)] [[PubMed](#)]
81. Giudice, E.; Domina, F.; Britti, D.; Di Pietro, S.; Pugliese, A. Clinical findings associated with *Borrelia burgdorferi* infection in the dog. *Vet. Res. Commun.* **2003**, *27* (Suppl. 1), 767–770. [[CrossRef](#)] [[PubMed](#)]
82. Giudice, E.; Giannetto, C.; Torina, A.; GIANESELLA, M. *Anaplasma phagocytophilum* intragranulocytic morulae in aborting sheep: A herd case in Sicily. *Transbound. Emerg. Dis.* **2011**, *58*, 263–267. [[CrossRef](#)]
83. Maia, C.; Campino, L. Methods for diagnosis of canine leishmaniasis and immune response to infection. *Vet. Parasitol.* **2008**, *158*, 274–287. [[CrossRef](#)]
84. Giulieri, S.; Jaton, K.; Cometta, A.; Trellu, L.T.; Greub, G. Development of a duplex real-time PCR for the detection of *Rickettsia* spp. and typhus group rickettsia in clinical samples. *FEMS Immunol. Med. Microbiol.* **2012**, *64*, 92–97. [[CrossRef](#)]
85. Adaszek, Ł.; Banach, T.; Bartnicki, M.; Winiarczyk, D.; Łyp, P.; Winiarczyk, S. Application the mass spectrometry MALDI-TOF technique for detection of *Babesia canis canis* infection in dogs. *Parasitol. Res.* **2014**, *113*, 4293–4295. [[CrossRef](#)]
86. Blanda, V.; D’Agostino, R.; Giudice, E.; Randazzo, K.; La Russa, F.; Villari, S.; Vullo, S.; Torina, A. New real-time PCRs to differentiate *Rickettsia* spp. and *Rickettsia conorii*. *Molecules* **2020**, *25*, 4431. [[CrossRef](#)]
87. Di Pietro, S.; Crinò, C.; Falcone, A.; Crupi, R.; Francaviglia, F.; Vitale, F.; Giudice, E. Parasitemia and its daily variation in canine leishmaniasis. *Parasitol. Res.* **2020**, *119*, 3541–3548. [[CrossRef](#)]
88. Dongus, H.; Zahler, M.; Gothe, R. Die braune Hundezecke, *Rhipicephalus sanguineus* (Ixodidae), in Deutschland: Eine epidemiologische Studie und Bekämpfungsmassnahmen [The brown dog tick, *Rhipicephalus sanguineus* (Ixodidae), in Germany: An epidemiologic study and control measures]. *Berl. Münch. Tierärztl. Wochenschr.* **1996**, *109*, 245–248. (In German)
89. Glaser, B.; Gothe, R. Imported arthropod-borne parasites and parasitic arthropods in dogs: Spectrum of species and epidemiological analysis of the cases diagnosed in 1995 and 1996. *Tierärztl. Prax.* **1998**, *26*, 40–46.
90. Glaser, B.; Gothe, R. Tourism and import of dogs: An inquiry in Germany on the extent as well as on the spectrum and preference of countries concerning stay abroad and origin, respectively. *Tierärztl. Prax.* **1998**, *26*, 197–202.
91. Hirsch, M.; Pantchev, N. Occurrence of the travel diseases leishmaniosis, ehrlichiosis, babesiosis and dirofilariosis in dogs living in Germany. *Kleintierpraxis* **2008**, *53*, 154–165. (In German)
92. Menn, B.; Lorentz, S.; Naucke, T.J. Imported and travelling dogs as carriers of canine vector-borne patogens in Germany. *Parasites Vectors* **2010**, *3*, 34. [[CrossRef](#)] [[PubMed](#)]
93. Naucke, T.; Lorentz, S.; Menn, B.; Mencke, N. Prevalence of canine vector-borne pathogens in imported and travelling dogs in Germany and studies on the prevalence of CVBD. *Tierärztl. Umsch.* **2011**, *66*, 311–317.
94. Hamel, D.; Silaghi, C.; Pfister, K. Arthropod-Borne infections in travelled dogs in Europe. *Parasite* **2013**, *20*, 9. [[CrossRef](#)]
95. Schafer, I.; Weingart, C.; Kohn, B. Infection with *Anaplasma phagocytophilum* in a cat in Germany. *Kleintierpraxis* **2019**, *64*, 205–215. (In German)
96. Best, J.M.; Butt, K.M.; Rohrbach, J.A. Occurrence of *Rhipicephalus sanguineus* in London. *Vet. Rec.* **1969**, *85*, 633. [[CrossRef](#)]
97. Fox, M.T.; Sykes, T.J. Establishment of the tropical dog tick, *Rh. sanguineus*, in a house in London. *Vet. Rec.* **1985**, *116*, 661–662. [[CrossRef](#)]
98. Garben, A.F.; Bosman, B.T.; van Bronswijk, J.E. [The brown dog tick *Rhipicephalus sanguineus* Latreille 1806 in the Netherlands, an analysis of imported cases, including their veterinary and medical significance]. *Tijdschr. Diergeneesk.* **1980**, *105*, 192–203. (In Dutch)
99. Jongejan, F.; de Jong, S.; Voskuilen, T.; van den Heuvel, L.; Bouman, R.; Heesen, H.; Carlijn Ijzermans, C.; Berger, L. “Tekenscanner”: A novel smartphone application for companion animal owners and veterinarians to engage in tick and tick-borne pathogen surveillance in the Netherlands. *Parasites Vectors* **2019**, *12*, 116. [[CrossRef](#)]
100. Jaenson, T.G.; Jaenson, D.G.; Eisen, L.; Petersson, E.; Lindgren, E. Changes in the geographical distribution and abundance of the tick *Ixodes ricinus* during the past 30 years in Sweden. *Parasites Vectors* **2012**, *5*, 8. [[CrossRef](#)] [[PubMed](#)]

101. Eichenberger, R.M.; Peter, D.; Alexander, M. Ticks on dogs and cats: A pet owner-based survey in a rural town in northeastern Switzerland. *Ticks Tick Borne Dis.* **2015**, *6*, 267–271. [CrossRef] [PubMed]
102. Földvári, G.; Márialigeti, M.; Solymosi, N.; Lukács, Z.; Majoros, G.; Kósa, J.P.; Farkas, R. hard ticks infesting dogs in Hungary and their infection with *Babesia* and *Borrelia* species. *Parasitol. Res.* **2007**, *101*, 25–34. [CrossRef]
103. Tsatsaris, A.; Chochlakis, D.; Papadopoulos, B.; Petsa, A.; Georgalis, L.; Angelakis, E.; Ioannou, I.; Tselentis, Y.; Psaroulaki, A. Species composition, distribution, ecological preference and host association of ticks in Cyprus. *Exp. Appl. Acarol.* **2016**, *70*, 523–542. [CrossRef]
104. Papazahariadou, M.G.; Saridomichelakis, M.N.; Koutinas, A.F.; Papadopoulos, E.G.; Leontides, L. Tick infestation of dogs in Thessaloniki, northern Greece. *Med. Vet. Entomol.* **2003**, *17*, 110–113. [CrossRef]
105. Zanet, S.; Battisti, E.; Pepe, P.; Ciucà, L.; Colombo, L.; Trisciuoglio, A.; Ferroglio, E.; Cringoli, G.; Rinaldi, L.; Maurelli, M.P. Tick-borne pathogens in Ixodidae ticks collected from privately-owned dogs in Italy: A country-wide molecular survey. *BMC Vet. Res.* **2020**, *16*, 46. [CrossRef]
106. Estrada-Peña, A.; Roura, X.; Sainz, A.; Miró, G.; Solano-Gallego, L. Species of ticks and carried pathogens in owned dogs in Spain: Results of a one-year national survey. *Ticks Tick Borne Dis.* **2017**, *8*, 443–452. [CrossRef]
107. Dantas-Torres, F.; Maia, C.; Latrofa, M.S.; Annoscia, G.; Cardoso, L.; Otranto, D. Genetic characterization of *Rhipicephalus sanguineus* (sensu lato) ticks from dogs in Portugal. *Parasites Vectors* **2017**, *10*, 133. [CrossRef]
108. Claerebout, E.; Losson, B.; Cochez, C.; Casaert, S.; Dalemans, A.C.; De Cat, A.; Madder, M.; Saegerman, C.; Heyman, P.; Lempereur, L. Ticks and associated pathogens collected from dogs and cats in Belgium. *Parasites Vectors* **2013**, *6*, 183. [CrossRef]
109. Jameson, L.J.; Phipps, L.P.; Medlock, J.M. Surveillance for exotic ticks on companion animals in the UK. *Vet. Rec.* **2010**, *166*, 202–203. [CrossRef]
110. Tulloch, J.S.P.; McGinley, L.; Sánchez-Vizcaíno, F.; Medlock, J.M.; Radford, A.D. The passive surveillance of ticks using companion animal electronic health records. *Epidemiol. Infect.* **2017**, *145*, 2020–2029. [CrossRef] [PubMed]
111. Corrain, R.; Drigo, M.; Fenati, M.; Menandro, M.L.; Mondin, A.; Pasotto, D.; Martini, M. Study on ticks and tick-borne zoonoses in public parks in Italy. *Zoonoses Public Health* **2012**, *59*, 468–476. [CrossRef] [PubMed]
112. Di Luca, M.; Toma, L.; Bianchi, R.; Quarchioni, E.; Marini, L.; Mancini, F.; Ciervo, A.; Khouri, C. Seasonal dynamics of tick species in an urban park of Rome. *Ticks Tick Borne Dis.* **2013**, *4*, 513–517. [CrossRef] [PubMed]
113. Scarpulla, M.; Barlozzari, G.; Salvato, L.; De Liberato, C.; Lorenzetti, R.; Macrì, G. *Rickettsia helvetica* in human-parasitizing and free-living *Ixodes ricinus* from urban and wild green areas in the metropolitan city of Rome, Italy. *Vector Borne Zoonotic Dis.* **2018**, *18*, 404–407. [CrossRef] [PubMed]
114. Beyhan, Y.E.; Mungan, M.; Babur, C. The species of ticks bites on human and their seasonal distribution in Ankara, Turkey. *Ank. Univ. Vet. Fak. Derg.* **2016**, *63*, 115–119.
115. Buczek, A.; Ciura, D.; Bartosik, K.; Zajac, Z.; Kulisz, J. Threat of attacks of *Ixodes ricinus* ticks (Ixodida: Ixodidae) and Lyme borreliosis within urban heat islands in south-western Poland. *Parasites Vectors* **2014**, *7*, 562. [CrossRef] [PubMed]
116. Kowalec, M.; Szewczyk, T.; Welc-Fałęciak, R.; Siński, E.; Karbowiak, G.; Bajer, A. Rickettsiales occurrence and co-occurrence in *Ixodes ricinus* Ticks in natural and urban areas. *Microb. Ecol.* **2019**, *77*, 890–904. [CrossRef]
117. Reis, C.; Cote, M.; Paul, R.E.; Bonnet, S. 2011. Questing ticks in suburban forest are infected by at least six tick-borne pathogens. *Vector Borne Zoonotic Dis.* **2011**, *11*, 907–916. [CrossRef]
118. Rizzoli, A.; Silaghi, C.; Obiegala, A.; Rudolf, I.; Hubálek, Z.; Földvári, G.; Eplantard, O.; Evayssier-Taussat, M.; Bonnet, S.I.; Špitálská, E.; et al. *Ixodes ricinus* and its transmitted pathogens in urban and peri-urban areas in Europe. *Front. Public Health* **2014**, *2*, 251. [CrossRef]
119. Saleh, M.N.; Sundstrom, K.D.; Duncan, K.T.; Ientile, M.M.; Jordy, J.; Ghosh, P.; Litt, S.E. Show us your ticks: A survey of ticks infesting dogs and cats across the USA. *Parasites Vectors* **2019**, *12*, 595. [CrossRef]
120. Lorusso, V.; Dantas-Torres, F.; Lia, R.P.; Tarallo, V.D.; Mencke, N.; Capelli, G.; Otranto, D. Seasonal dynamics of the brown dog tick, *Rhipicephalus sanguineus*, on a confined dog population in Italy. *Med. Vet. Entomol.* **2010**, *24*, 309–315. [CrossRef] [PubMed]
121. Wright, I.; Cull, B.; Gillingham, E.L.; Hansford, K.M.; Medlock, J. Be tick aware: When and where to check cats and dogs for ticks. *Vet. Rec.* **2018**, *182*, 514. [CrossRef] [PubMed]
122. McColl, H.P.; Tenquist, J.D. Accidental importation of the brown dog tick, *Rhipicephalus sanguineus*. *N. Z. Vet. J.* **1980**, *28*, 168–169.
123. Hansford, K.M.; Pietzsch, M.E.; Cull, B.; Medlock, J.M. Importation of *Rhipicephalus sanguineus* into the UK via dogs. *Vet. Rec.* **2014**, *175*, 385–386. [CrossRef]
124. Bates, P.; Rankin, M.; Shickle, L. Importation of the brown dog or kennel tick (*Rhipicephalus sanguineus*) into the UK. *Vet. Rec.* **2002**, *150*, 224.
125. European Centre for Disease Prevention and Control and European Food Safety Authority. Mosquito Maps. Stockholm: ECDC. 2018. Available online: <https://ecdc.europa.eu/en/disease-vectors/surveillance-and-disease-data> (accessed on 12 December 2018).
126. Hunter, A.; Walker, A.; Sumption, K.; Eelesten, M.; Scott, G.; Sewell, M.; Morrow, A.; Hopper, D. Health risks from increased movement of companion animals in Europe. *Vet. Rec.* **1998**, *142*, 23.
127. Hoyle, D.V.; Walker, A.R.; Craig, R.S.; Woolhouse, M.E.J. Survey of parasite infections not endemic to the United Kingdom in quarantined animals. *Vet. Rec.* **2001**, *149*, 457–458. [CrossRef]
128. Trees, A.J.; Ridge, A. Threat of imported diseases to UK dogs. *Vet. Rec.* **2016**, *178*, 347. [CrossRef]

129. Gillingham, E.L.; Cull, B.; Pietzsch, M.E.; Phipps, L.P.; Medlock, J.M.; Hansford, K. The unexpected holiday souvenir: The public health risk to UK travellers from ticks acquired overseas. *Int. J. Environ. Res. Public Health* **2020**, *17*, 7957. [[CrossRef](#)]
130. Gothe, R. On the occurrence of *Rhipicephalus sanguineus* (Latreille, 1806) in Germany. *Z. Tropenmed. Parasitol.* **1968**, *19*, 305–307. (In German)
131. Gothe, R. *Rhipicephalus sanguineus* (Ixodidae): Frequency of infestation and ehrlichial infections transmitted by this tick in dogs in Germany: An epidemiological study and consideration. *Wiener Tierärztl. Mon.* **1999**, *86*, 49–56. (In German)
132. Gothe, R.; Hamel, H.D. Epizootics due to *Rhipicephalus sanguineus* (Latreille 1806) in Germany. *Zent. Veterinärmed. B* **1973**, *20*, 245–249. (In German) [[CrossRef](#)]
133. Centurier, C.; Gothe, R.; Hoffmann, G.; Liebisch, A.; Schein, E. [The brown dog tick *Rhipicephalus sanguineus* (Latreille, 1806) in Germany, a problem in small animal practice]. *Berl. Münch. Tierärztl. Wochenschr.* **1979**, *92*, 472–477. [[PubMed](#)]
134. Röhrig, E.; Hamel, D.; Pfister, K. Retrospective evaluation of laboratory data on canine vector-borne infections from the years 2004–2008. *Berl. Münch. Tierärztl. Wochenschr.* **2011**, *124*, 411–418.
135. Schäfer, I.; Volkmann, M.; Beelitz, P.; Merle, R.; Müller, E.; Kohn, B. Retrospective evaluation of vector-borne infections in dogs imported from the Mediterranean region and southeastern Europe (2007–2015). *Parasites Vectors* **2019**, *12*, 30. [[CrossRef](#)]
136. Szymański, S. Przypadek masowego rozwoju kleszcza *Rhipicephalus sanguineus* (Latreille, 1806) w warszawskim mieszkaniu [Case of a mass development of tick *Rhipicephalus sanguineus* (Latreille, 1806) in a Warsaw residence]. *Wiad. Parazytol.* **1979**, *25*, 453–459. (In Polish)
137. Černý, V. First case of introduction of the tick *Rhipicephalus sanguineus* to Czechoslovakia. *Folia Parasitol. (Praha)* **1985**, *32*, 162.
138. Černý, V. Introduction of the tick *Rhipicephalus sanguineus* to Czechoslovakia: An additional case. *Folia Parasitol. (Praha)* **1989**, *36*, 184.
139. Van Bronswijk, J.E.M.H. A bibliography on ticks and tick-borne diseases in the countries of the Benelux (1567–1978). *Vet. Q.* **1980**, *2*, 220–233. [[CrossRef](#)]
140. Winding, O.; Haarløv, N. Accidental introduction into Denmark from Sudan of the kennel tick (*Rhipicephalus sanguineus* (Latr. 1806)). *Nord. Veterinaerm.* **1968**, *20*, 284–285.
141. Haarløv, N. Another case of introduction into Denmark from Africa of the kennel tick (*Rhipicephalus sanguineus* (Latr. 1806)). *Nord. Veterinaerm.* **1969**, *21*, 445–448.
142. Winding, O.; Willeberg, P.; Haarløv, N. *Rhipicephalus sanguineus*, an ectoparasite of dogs of current interest. *Nord. Veterinaerm.* **1970**, *22*, 48–58.
143. Willeberg, P. Forekomst af husflæt, *Rhipicephalus sanguineus*, i Danmark [Existence of the brown dog tick, *Rhipicephalus sanguineus*, in Denmark]. *Ugeskr. Laeger* **1970**, *132*, 1270–1271. (In Danish with Abstract in English) [[PubMed](#)]
144. Sibomana, G.; Geerts, S.; De Vries, T. Establishment of *Rhipicephalus sanguineus* (Latreille, 1806) inside houses in Belgium. *Ann. Soc. Belg. Méd. Trop.* **1986**, *66*, 79–81. [[PubMed](#)]
145. Péter, O.; Burgdorfer, W.; Aeschlimann, A.; Chatelanat, P. *Rickettsia conorii* isolated from *Rhipicephalus sanguineus* introduced into Switzerland on a pet dog. *Z. Parasitenkd.* **1984**, *70*, 265–270. [[CrossRef](#)]
146. Maurelli, M.P.; Pepe, P.; Colombo, L.; Armstrong, R.; Battisti, E.; Morgoglion, M.E.; Counturis, D.; Rinaldi, L.; Cringoli, G.; Ferroglio, E.; et al. A national survey of Ixodidae ticks on privately owned dogs in Italy. *Parasites Vectors* **2018**, *11*, 420. [[CrossRef](#)]
147. Kebbi, R.; Nait-Mouloud, M.; Hassissen, L.; Ayad, A. Seasonal activity of ticks infesting domestic dogs in Bejaia province, Northern Algeria. *Onderstepoort J. Vet. Res.* **2019**, *86*, e1–e6. [[CrossRef](#)]
148. Tinoco-Gracia, L.; Quiroz-Romero, H.; Quintero-Martínez, M.T.; Rentería-Evangelista, T.B.; González-Medina, Y.; Barreras-Serrano, A.; Hori-Oshima, S.; Moro, M.H.; Vinasco, J. Prevalence of *Rhipicephalus sanguineus* ticks on dogs in a region on the Mexico-USA border. *Vet. Rec.* **2009**, *164*, 59–61. [[CrossRef](#)]
149. Abah, O.O.I.; Audu, P.A. Prevalence of brown dog tick (*Rhipicephalus sanguineus*) infestation of dogs in Lokoja Metropolis, Kogi State, North-central Nigeria. *Niger. J. Parasitol.* **2013**, *34*, 91–98.
150. Nithikathkul, C.; Polseela, R.; Iamsaard, J.; Wongsawad, C.; Jittapalapong, S. A study of ectoparasites of *Canis lupus familiaris* in Mueang district, Khon Kaen, Thailand. *Southeast Asian J. Trop. Med. Public Health* **2005**, *36* (Suppl. 4), 149–151.
151. Ul-Hasan, M.; Abubakar, M.; Muhamad, G.; Khan, M.N.; Hussain, M. Prevalence of tick infestation (*Rhipicephalus sanguineus* and *Hyalomma anatomicum anatomicum* in dogs in Punjab, Pakistan. *Vet. Ital.* **2012**, *48*, 95–98. [[PubMed](#)]
152. Persichetti, M.F.; Solano-Gallego, L.; Serrano, L.; Altet, L.; Reale, S.; Masucci, M.; Pennisi, M.G. Detection of vector-borne pathogens in cats and their ectoparasites in southern Italy. *Parasites Vectors* **2016**, *9*, 247. [[CrossRef](#)] [[PubMed](#)]
153. Wright, I.; Wildgoose, W.H.; Wall, R.; McGarry, J. *Rhipicephalus sanguineus* in an imported dog. *Vet. Rec.* **2018**, *182*, 25–26.
154. Sparagano, O.A.; de Vos, A.P.; Paoletti, B.; Cammà, C.; de Santis, P.; Otranto, D.; Giangaspero, A. Molecular detection of *Anaplasma platys* in dogs using polymerase chain reaction and reverse line blot hybridization. *J. Vet. Diagn. Investig.* **2003**, *15*, 527–534. [[CrossRef](#)]
155. Alberti, A.; Addis, M.F.; Sparagano, O.; Zobba, R.; Chessa, B.; Cubeddu, T.; Parpaglia, M.L.; Ardu, M.; Pittau, M. *Anaplasma phagocytophylum*, Sardinia, Italy. *Emerg. Infect. Dis.* **2005**, *11*, 1322–1324. [[CrossRef](#)]
156. Uspensky, I. Brown dog tick or kennel tick, *Rhipicephalus sanguineus* (Latreille) (Acari: Ixodida: Ixodidae). In *Encyclopedia of Entomology*, 2nd ed.; Capinera, J.L., Ed.; Springer: Dordrecht, The Netherlands, 2008; Volume 1, pp. 577–580.
157. Cassini, R.; Zanutto, S.; di Regalbono, A.F.; Gabrielli, S.; Calderini, P.; Moretti, A.; Tampieri, M.P.; Pietrobelli, M. Canine piroplasmosis in Italy: Epidemiological aspects in vertebrate and invertebrate hosts. *Vet. Parasitol.* **2009**, *165*, 30–35. [[CrossRef](#)]

158. Dantas-Torres, F.; Otranto, D. Further thoughts on the taxonomy and vector role of *Rhipicephalus sanguineus* group ticks. *Vet. Parasitol.* **2015**, *208*, 9–13. [[CrossRef](#)]
159. Santos, M.A.B.; de Souza, I.B.; de Macedo, L.O.; Ramos, C.A.D.N.; Rego, A.G.D.O.; Alves, L.C.; Ramos, R.A.N.; De Carvalho, G.A. *Cercopithifilaria bainae* in *Rhipicephalus sanguineus* sensu lato ticks from dogs in Brazil. *Ticks Tick Borne Dis.* **2017**, *8*, 623–625. [[CrossRef](#)]
160. Soares, R.L.; Silva, A.O.; Coelho, M.L.; Echeverria, J.T.; Souza, M.L.; Babo-Terra, V.J.; Pasquatti, T.N.; Ramos, R.A.N.; Ramos, C.A.N. Molecular detection of *Cercopithifilaria bainae* and other tick-borne pathogens in *Rhipicephalus sanguineus* s.l. isolated from dogs in Midwest Brazil. *Braz. J. Vet. Parasitol.* **2020**, *29*, e018019. [[CrossRef](#)]
161. Gehrke, F.S.; Gazeta, G.S.; Souza, E.R.; Ribeiro, A.; Marrelli, M.T.; Schumaker, T.T.S. *Rickettsia rickettsii*, *Rickettsia felis* and *Rickettsia* sp. TwKM03 infecting *Rhipicephalus sanguineus* and *Ctenocephalides felis* collected from dogs in a Brazilian spotted fever focus in the State of Rio De Janeiro/Brazil. *Clin. Microbiol. Infect.* **2009**, *15* (Suppl. 2), 267–268. [[CrossRef](#)]
162. Macaigne, F.; Pérez-Eid, C. *Hyalomma scupense* Schulze, 1919 (Acarina, Ixodoidea) tique autochtone du sud-ouest de la France. *Ann. Parasitol. Hum. Comp.* **1993**, *68*, 199–200. [[CrossRef](#)]
163. Vial, L.; Stachurski, F.; Leblond, A.; Hubera, K.; Vourc'h, G.; René-Martellet, M.; Desjardins, I.; Balançae, G.; Grosboise, V.; Pradier, S.; et al. Strong evidence for the presence of the tick *Hyalomma marginatum* Koch, 1844 in southern continental France. *Ticks Tick Borne Dis.* **2016**, *7*, 1162–1167. [[CrossRef](#)] [[PubMed](#)]
164. Hansford, K.M.; Pietzsch, M.E.; Cull, B.; Medlock, J.M. Surveillance: Brown dog tick infestation of a home in England. *Vet. Rec.* **2015**, *176*, 129–130. [[CrossRef](#)] [[PubMed](#)]
165. Renvoisé, A.; Delaunay, P.; Blanchouin, E.; Cannavo, I.; Cua, C.; Socolovschi, C.; Parola, P.; Raoult, D. Urban family cluster of spotted fever rickettsiosis linked to *Rhipicephalus sanguineus* infected with *Rickettsia conorii* subsp. *caspia* and *Rickettsia massiliae*. *Ticks Tick Borne Dis.* **2012**, *3*, 389–392. [[CrossRef](#)]
166. Latrofa, M.S.; Angelou, A.; Giannelli, A.; Annoscia, G.; Ravagnan, S.; Dantas-Torres, F.; Capelli, G.; Halos, L.; Beugnet, F.; Papadopoulos, E.; et al. Ticks and associated pathogens in dogs from Greece. *Parasites Vectors* **2017**, *10*, 301. [[CrossRef](#)]
167. Jameson, L.J.; Medlock, J.M. Tick surveillance in Great Britain. *Vector Borne Zoonotic Dis.* **2011**, *11*, 403–412. [[CrossRef](#)]
168. Wright, I. Exotic disease threats from pet travel, importation and changing parasite distributions in Europe. *Vet. CPD* **2017**, *4*, 3.
169. Attipa, C.; Maguire, D.; Solano-Gallego, L.; Szladovits, B.; Barker, E.N.; Farr, A.; Baneth, G.; Tasker, S. *Hepatozoon canis* in three imported dogs: A new tickborne disease reaching the United Kingdom. *Vet. Rec.* **2018**, *183*, 716. [[CrossRef](#)]
170. Senneville, E.; Ajana, F.; Lecocq, P.; Chidiac, C.; Mouton, Y. *Rickettsia conorii* isolated from ticks introduced to northern France by a dog. *Lancet* **1991**, *337*, 676. [[CrossRef](#)]
171. Rovery, C.; Raoult, D. Mediterranean spotted fever. *Infect. Clin. N. Am.* **2008**, *22*, 515–530. [[CrossRef](#)]
172. Rovery, C.; Brouqui, P.; Raoult, D. Questions on Mediterranean spotted fever a century after its discovery. *Emerg. Infect. Dis.* **2008**, *14*, 1360–1367. [[CrossRef](#)] [[PubMed](#)]
173. Alieva, E.E.; Bondarenko, E.I.; Maliy, K.D.; Shvalov, A.N.; Verbenets, E.A.; Gafarova, M.T. The role of *Rhipicephalus sanguineus* ticks parasitizing dogs in the spread of tick-borne rickettsial pathogens in the city of Sevastopol. *New Microbes New Infect.* **2020**, *36*, 100704. [[CrossRef](#)] [[PubMed](#)]
174. Khrouf, F.; M'Ghirbi, Y.; Znazen, A.; Ben Jemaa, M.; Hammami, A.; Bouattour, A. Detection of *Rickettsia* in *Rhipicephalus sanguineus* ticks and *Ctenocephalides felis* fleas from southeastern Tunisia by reverse line blot assay. *J. Clin. Microbiol.* **2014**, *52*, 268–274. [[CrossRef](#)] [[PubMed](#)]
175. Bernasconi, M.V.; Casati, S.; Péter, O.; Piffaretti, J.-C. *Rhipicephalus* ticks infected with *Rickettsia* and *Coxiella* in Southern Switzerland (Canton Ticino). *Infect. Genet. Evol.* **2002**, *2*, 111–120. [[CrossRef](#)]
176. Edouard, S.; Parola, P.; Socolovschi, C.; Davoust, B.; La Scola, B.; Raoult, D. Clustered cases of *Rickettsia sibirica mongolimoniae* infection, France. *Emerg. Infect. Dis.* **2013**, *19*, 337–338. [[CrossRef](#)] [[PubMed](#)]
177. Oztoprak, N.; Celebi, G.; Aydemir, H.; Pişkin, N.; Bektaş, S.; Koca, R.; Kuloglu, F. Köpek kenesi ile temas sonrasında gelisen akdeniz benekli ateşi [Mediterranean spotted fever due to contact with dog-tick]. *Mikrobiyol. Bul.* **2008**, *42*, 701–706. (In Turkish with English Abstract)
178. Solano-Gallego, L.; Morell, P.; Arboix, M.; Alberola, J.; Ferrer, L. Prevalence of *Leishmania infantum* infection in dogs living in an area of canine leishmaniasis endemicity using PCR on several tissues and serology. *J. Clin. Microbiol.* **2001**, *39*, 560–563. [[CrossRef](#)]
179. Lachaud, L.; Chabbert, E.; Dubessay, P.; Dereure, J.; Lamothe, J.; Dedet, J.P.; Bastien, P. Value of two PCR methods for the diagnosis of canine visceral leishmaniasis and the detection of asymptomatic carriers. *Parasitology* **2002**, *125*, 197–207. [[CrossRef](#)]
180. Mircean, V.; Dumitrache, M.O.; Györke, A.; Pantchev, N.; Jodies, R.; Mihalca, A.D.; Cozma, V. Seroprevalence and geographic distribution of *Dirofilaria immitis* and tick-borne infections (*Anaplasma phagocytophilum*, *Borrelia burgdorferi* sensu lato, and *Ehrlichia canis*) in dogs from Romania. *Vector Borne Zoonotic Dis.* **2012**, *12*, 595–604. [[CrossRef](#)]
181. Pennisi, M.G.; Caprì, A.; Solano-Gallego, L.; Lombardo, G.; Torina, A.; Masucci, M. Prevalence of antibodies against *Rickettsia conorii*, *Babesia canis*, *Ehrlichia canis*, and *Anaplasma phagocytophilum* antigens in dogs from the Stretto di Messina area (Italy). *Ticks Tick Borne Dis.* **2012**, *3*, 315–318. [[CrossRef](#)]
182. Morchón, R.; Carretón, E.; González-Miguel, J. Heartworm disease (*Dirofilaria immitis*) and their vectors in Europe—new distribution trends. *Front. Physiol.* **2012**, *3*, 196. [[CrossRef](#)] [[PubMed](#)]
183. Farkas, R.; Gyurkovszky, M.; Lukács, Z.; Aladics, B.; Solymosi, N. Seroprevalence of some vector-borne infections of dogs in Hungary. *Vector Borne Zoonotic Dis.* **2014**, *14*, 256–260. [[CrossRef](#)] [[PubMed](#)]

184. Miró, G.; Montoya, A.; Roura, X.; Gálvez, R.; Sainz, A. Seropositivity rates for agents of canine vector-borne diseases in Spain: A multicentre study. *Parasites Vectors* **2013**, *6*, 117. [CrossRef] [PubMed]
185. Morganti, G.; Veronesi, F.; Stefanetti, V.; Di Muccio, T.; Fiorentino, E.; Diaferia, M.; Santoro, A.; Passamonti, F.; Gramiccia, M. Emerging feline vector-borne pathogens in Italy. *Parasites Vectors* **2019**, *12*, 193. [CrossRef] [PubMed]
186. Torina, A.; Caracappa, S. Dog tick-borne diseases in Sicily. *Parassitologia* **2006**, *48*, 145–147. [PubMed]
187. Díaz-Regañón, D.; Roura, X.; Suárez, M.L.; León, M.; Sainz, Á. Serological evaluation of selected vector-borne pathogens in owned dogs from northern Spain based on a multicenter study using a commercial test. *Parasites Vectors* **2020**, *13*, 301. [CrossRef]
188. Pantchev, N.; Vrhovec, M.G.; Pluta, S.; Straubinger, R.K. Seropositivity of *Borrelia burgdorferi* in a cohort of symptomatic cats from Europe based on a C6-peptide assay with discussion of implications in disease aetiology. *Berl. Münch. Tierarztl. Wochenschr.* **2016**, *129*, 333–339.
189. Izzard, L.; Cox, E.; Stenos, J.; Waterston, M.; Fenwick, S.; Graves, S. Serological prevalence study of exposure of cats and dogs in Launceston, Tasmania, Australia to spotted fever group rickettsiae. *Aust. J. Vet.* **2010**, *88*, 29–31. [CrossRef]
190. Montenegro, V.M.; Bonilla, M.C.; Kaminsky, D.; Romero-Zúñiga, J.J.; Siebert, S.; Krämer, F. Serological detection of antibodies to *Anaplasma* spp., *Borrelia burgdorferi* sensu lato and *Ehrlichia canis* and of *Dirofilaria immitis* antigen in dogs from Costa Rica. *Vet. Parasitol.* **2017**, *236*, 97–107. [CrossRef]
191. CDC. Centers for Disease Control, Prevention. Preventing Tick-Bites. 2017. Available online: <https://www.cdc.gov/lyme/prevention/index.html> (accessed on 1 June 2017).
192. European Scientific Counsel Companion Animal Parasites. ESCCAP Guideline 5: Control of Vector-Borne Diseases in Dogs and Cats, 3rd ed. 2019. Available online: <https://www.esccap.org> (accessed on 30 January 2020).
193. Bissinger, B.W.; Roe, R.M. Tick repellents: Past, present, and future. *Pestic. Biochem. Physiol.* **2010**, *96*, 63–79. [CrossRef]
194. Phillips, C.B.; Liang, M.H.; Sangha, O.; Wright, E.A.; Fossel, A.H.; Lew, R.A.; Fossel, K.K.; Shadick, N.A. Lyme disease and preventive behaviors in residents of Nantucket Island, Massachusetts. *Am. J. Prev. Med.* **2001**, *20*, 219–224. [CrossRef]
195. Bartosik, K.; Kubrak, T.; Olszewski, T.; Jung, M.; Buczek, A. Prevention of tick bites and protection against tick-borne diseases in south-eastern Poland. *Ann. Agric. Environ. Med.* **2008**, *15*, 181–185. [PubMed]
196. Butler, A.D.; Sedghi, T.; Petrini, J.R.; Ahmadi, R. Tick-Borne disease preventive practices and perceptions in an endemic area. *Ticks Tick Borne Dis.* **2016**, *7*, 331–337. [CrossRef] [PubMed]
197. Eisen, L.; Dolan, M.C. Evidence for personal protective measures to reduce human contact with blacklegged ticks and for environmentally based control methods to suppress host-seeking blacklegged ticks and reduce infection with lyme disease spirochetes in tick vectors and rodent reservoirs. *J. Med. Entomol.* **2016**, *53*, 1063–1092.
198. Nejedzlebová, H.; Kiewra, D.; Žákovská, A.; Ovesná, P. Students' attitudes to tick risks. *Ann. Agric. Environ. Med.* **2016**, *23*, 437–441. [CrossRef]
199. Pańczuk, A.; Tokarska-Rodak, M.; Mikul'áková, W.; Kendrová, L.; Magurová, D. Exposure to ticks and undertaking Lyme borreliosis prevention activities among students from Poland and Slovakia. *Ann. Agric. Environ. Med.* **2019**, *26*, 217–221. [CrossRef]
200. Buczek, A.; Pilch, J.; Buczek, W. Tick preventive behaviors and practices adopted by medical students from Poland, Germany, and Thailand in relation to socio-demographic conditions and their knowledge of ticks and tick-borne diseases. *Insects* **2020**, *863*. [CrossRef]
201. Dumont, P.; Blair, J.; Fourie, J.J.; Chester, T.S.; Larsen, D.L. Evaluation of the efficacy of afoxolaner against two European dog tick species: *Dermacentor reticulatus* and *Ixodes ricinus*. *Vet. Parasitol.* **2014**, *201*, 216–219. [CrossRef]
202. Kunkle, B.; Daly, S.; Dumont, P.; Drag, M.; Larsen, D. Assessment of the efficacy of orally administered afoxolaner against *Rhipicephalus sanguineus* sensu lato. *Vet. Parasitol.* **2014**, *201*, 226–228. [CrossRef]
203. Six, R.H.; Young, D.R.; Holzmer, S.J.; Mahabir, S.P. Comparative speed of kill of sarolaner (Simparica) and afoxolaner (NexGard) against induced infestations of *Rhipicephalus sanguineus* s.l. on dogs. *Parasites Vectors* **2016**, *9*, 91. [CrossRef]
204. Cavalleri, D.; Murphy, M.; Gorbea, R.L.; Seewald, W.; Drake, J.; Nanchen, S. Laboratory evaluations of the immediate and sustained effectiveness of lotilaner (Credelio™) against three common species of ticks affecting dogs in Europe. *Parasites Vectors* **2017**, *10*, 527. [CrossRef] [PubMed]
205. Becskei, C.; Fias, D.; Mahabir, S.P.; Farkas, R. Efficacy of a novel oral chewable tablet containing sarolaner, moxidectin and pyrantel (Simparica Trio™) against natural flea and tick infestations on dogs presented as veterinary patients in Europe. *Parasites Vectors* **2020**, *13*, 72. [CrossRef] [PubMed]
206. Coleman, N.; Coleman, S. Methods of tick removal: A systematic review of the literature. *AMJ* **2017**, *10*, 53–62. [CrossRef]
207. Buczek, A.M.; Buczek, W. Can freezing feeding ticks during removal from host skin be an effective method in prevention of tick-borne diseases? In *Parasitic and Allergenic Arthropods*; Buczek, A., Błaszkak, C., Eds.; Koliber: Lublin, Poland, 2020; pp. 37–49.
208. Hoogstraal, H. *African Ixodoidea. I. Ticks of the Sudan (with Special Reference to Equatoria Province and with Preliminary Reviews of the Genera Boophilus, Magaropus, and Hyalomma)*; U.S. Naval Medical Research Report NM 005 050.29.07; U.S. Navy: Philadelphia, PA, USA, 1956.
209. Hasle, G. Transport of ixodid ticks and tick-borne pathogens by migratory birds. *Front. Cell. Infect. Microbiol.* **2013**, *3*, 48. [CrossRef]
210. Movila, A.; Alekseev, A.N.; Dubinin, H.V.; Toderas, I. Detection of tick-borne pathogens in ticks from migratory birds in the Baltic region of Russia. *Med. Vet. Entomol.* **2013**, *27*, 113–117. [CrossRef]

211. Capek, M.; Literak, I.; Kocanova, E.; Sychra, O.; Najar, T.; Trnka, A.; Kverek, P. Ticks of the *Hyalomma marginatum* complex transported by migratory birds into Central Europe. *Ticks Tick Borne Dis.* **2014**, *5*, 489–493. [[CrossRef](#)]
212. Cohen, E.B.; Auckland, L.D.; Marra, P.P.; Hamer, S.A. Avian migrants facilitate invasions of neotropical ticks and tick-borne pathogens into the United States. *Appl. Environ. Microbiol.* **2015**, *81*, 8366–8378. [[CrossRef](#)]
213. Buczek, A.M.; Buczek, W.; Buczek, A.; Bartosik, K. The potential role of migratory birds in the rapid spread of ticks and tick-borne pathogens in the changing climatic and environmental conditions in Europe. *Int. J. Environ. Res. Public Health* **2020**, *17*, 2117. [[CrossRef](#)]
214. Beugnet, F.; Marié, J.-L. Emerging arthropod-borne diseases of companion animals in Europe. *Vet. Parasitol.* **2009**, *163*, 298–305. [[CrossRef](#)]
215. Mencke, N. Future challenges for parasitology: Vector control and ‘One health’ in Europe: The veterinary medicinal view on CVBDs such as tick borreliosis, rickettsiosis and canine leishmaniosis. *Vet. Parasitol.* **2013**, *195*, 256–271. [[CrossRef](#)]
216. Maia, C.; Ramos, C.; Coimbra, M.; Bastos, F.; Martins, Â.; Pinto, P.; Nunes, M.; Vieira, M.L.; Cardoso, L.; Campino, L. Bacterial and protozoal agents of feline vector-borne diseases in domestic and stray cats from southern Portugal. *Parasites Vectors* **2014**, *7*, 115. [[CrossRef](#)] [[PubMed](#)]
217. Willeberg, P.; Kjaersgaard, P. Babesiose hos en hund i Danmark [A case of canine babesiosis in Denmark]. *Nord. Veterinaermed.* **1973**, *25*, 308–312. (In Danish with Abstract in English).