



Complete Genome Sequence of a Virulent *Leptospira interrogans* Serovar Copenhageni Strain, Assembled with a Combination of Nanopore and Illumina Reads

Alejandro Llanes,^a Dhani Prakoso,^b Carlos Mario Restrepo,^a Sreekumari Rajeev^b

^aCentro de Biología Celular y Molecular de Enfermedades, Instituto de Investigaciones Científicas y Servicios de Alta Tecnología, Panama City, Panama

^bCollege of Veterinary Medicine, University of Florida, Gainesville, Florida, USA

ABSTRACT Here, we present the complete genome sequence of a highly virulent *Leptospira interrogans* serovar Copenhageni strain isolated from a dog with severe leptospirosis. In this work, a gapless genome draft was assembled with a combination of Nanopore and Illumina data of relatively low coverage.

Leptospira species are spirochete bacteria that colonize proximal renal tubules of mammals, and infections may cause fatal disease in humans and animals. In this study, we assembled the genome of *Leptospira interrogans* strain SK1 and compared it with those of *L. interrogans* serovar Copenhageni (strain Fiocruz L1-130) (1) and *L. interrogans* serovar Lai (strain 56601) (2). The SK1 strain was isolated from the blood of a canine patient that succumbed to severe clinical leptospirosis (3). The strain was isolated through routine *Leptospira* culture protocols using Ellinghausen-McCullough-Johnson-Harris medium, as described previously (1).

Genomic DNA from strain SK1 was isolated with the MasterPure complete DNA and RNA purification kit (Epicentre, USA). Sequencing libraries for the MiSeq and MinION platforms were prepared with the Nextera XT DNA library preparation kit (Illumina, USA) and the ligation sequencing kit (SQK-LSK109; Oxford Nanopore Technologies, UK), respectively, following the manufacturers' instructions. Quality control was performed with FastQC (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc>). In the case of Nanopore reads, we used Guppy v.3.2.4 for base calling and Porechop v.0.2.4 (<https://github.com/rwick/Porechop>) for adapter trimming. *De novo* assembly was performed with Canu v.2.0 (4) using only the Nanopore reads. The Canu assembly was initially polished with Medaka v.0.11.4 (<https://nanoporetech.github.io/medaka>) using probabilistic model r941_min_fast_g303. Pilon v.1.23 (5) was used for hybrid polishing with the Illumina reads. To estimate the number of sequencing errors after each polishing step, Illumina reads were aligned to the Nanopore assembly with BWA v.0.7.12 (6). BCFTools v.1.9 (7) was then used to call for single-nucleotide variants and insertion/deletions (indels), which were considered to be errors if their mapping quality score was above 20. RATT was used to transfer the genes annotated in the Fiocruz L1-130 genome to the SK1 assembly. Prokka v.1.14.3 (8) was used to complement the RATT results. Default parameters were used for all software except where otherwise noted.

Illumina and Nanopore sequencing generated 1.1 million 300-bp paired-end reads and 142,300 reads with a median length of 8 kb, respectively. *De novo* assembly of Nanopore reads resulted in 2 contigs, consistent with the 2 *Leptospira* chromosomes. The number of suspected errors decreased from 14,672 (0.3%) in the original Canu assembly to 92 (0.002%) after polishing with Pilon. Of these, only 6 errors seemed to affect putative coding sequences and were corrected manually.

The SK1 genome is very similar to that of strain Fiocruz L1-130 in terms of synteny

Citation Llanes A, Prakoso D, Restrepo CM, Rajeev S. 2020. Complete genome sequence of a virulent *Leptospira interrogans* serovar Copenhageni strain, assembled with a combination of Nanopore and Illumina reads. *Microbiol Resour Announc* 9:e00200-20. <https://doi.org/10.1128/MRA.00200-20>.

Editor Irene L. G. Newton, Indiana University, Bloomington

Copyright © 2020 Llanes et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Sreekumari Rajeev, sree63rajeev@gmail.com.

Received 25 February 2020

Accepted 7 April 2020

Published 23 April 2020




FIG 1 Comparison of the genomes of *L. interrogans* serovar Copenhageni strains SK1 and Fiocruz L1-130. Pink bands represent regions of shared sequence similarity. Coverage of Nanopore and Illumina reads is plotted along the SK1 genome, averaged over a 500-bp sliding window. Chr, chromosome.

(Fig. 1) and global metrics, including a total size of 4,630,180 bp, a G+C content of 35%, and 3,731 protein-coding genes. However, at least 11 pseudogenes from Fiocruz L1-130 appear to have functional orthologs in the genomes of strains SK1 and 56601, despite the latter belonging to *L. interrogans* serovar Lai. Conversely, genes *folA* and *folC*, encoding dihydrofolate reductase and dihydrofolate synthase, respectively, appear to have become pseudogenes in SK1, which suggests impaired folate metabolism in this strain.

Data availability. The genome sequence of the SK1 strain and associated data were deposited under BioProject number [PRJNA600490](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA600490), GenBank accession numbers [CP048830](https://www.ncbi.nlm.nih.gov/nuccore/CP048830) (chromosome I) and [CP048831](https://www.ncbi.nlm.nih.gov/nuccore/CP048831) (chromosome II), and SRA accession numbers [SRX7657977](https://www.ncbi.nlm.nih.gov/sra/SRX7657977) (Illumina reads) and [SRX7730232](https://www.ncbi.nlm.nih.gov/sra/SRX7730232) (MinION reads).

ACKNOWLEDGMENTS

This research was funded by internal grant support from the University of Florida College of Veterinary Medicine (Gainesville, FL) and intramural grants from the Ross University School of Veterinary Medicine Center for One Health and Tropical Medicine (Saint Kitts, West Indies).

REFERENCES

- Nascimento A, Ko AI, Martins EAL, Monteiro-Vitorello CB, Ho PL, Haake DA, Verjovski-Almeida S, Hartskeerl RA, Marques MV, Oliveira MC, Menck CFM, Leite LCC, Carrer H, Coutinho LL, Degraeve WM, Dellagostin OA, El-Dorry H, Ferro S, Ferro MIT, Furlan LR, Gamberini M, Giglioti EA, Góes-Neto A, Goldman GH, Goldman MHS, Harakava R, Jerônimo SMB, Junqueira-de-Azevedo ILM, Kimura ET, Kuramae EE, Lemos EGM, Lemos MVF, Marino CL, Nunes LR, De Oliveira RC, Pereira GG, Reis MS, Schriefer A, Siqueira WJ, Sommer P, Tsai SM, Simpson AJG, Ferro JA, Camargo LEA, Kitajima JP, Setubal JC, Van Sluys MA. 2004. Comparative genomics of two *Leptospira interrogans* serovars reveals novel insights into physiology and pathogenesis. *J Bacteriol* 186:2164–2172. <https://doi.org/10.1128/jb.186.7.2164-2172.2004>.
- Ren SX, Fu G, Jiang XG, Zeng R, Miao YG, Xu H, Zhang YX, Xiong H, Lu G, Lu LF, Jiang HQ, Jia J, Tu Y-F, Jiang J-X, Gu W-Y, Zhang Y-Q, Cai Z, Sheng H-H, Yin H-F, Zhang Y, Zhu G-F, Wan M, Huang H-L, Qian Z, Wang S-Y, Ma W, Yao Z-J, Shen Y, Qiang B-Q, Xia Q-C, Guo X-K, Danchin A, Saint Girons I, Somerville RL, Wen Y-M, Shi M-H, Chen Z, Xu J-G, Zhao G-P. 2003. Unique physiological and pathogenic features of *Leptospira interrogans* revealed by whole-genome sequencing. *Nature* 422:888–893. <https://doi.org/10.1038/nature01597>.
- Larson CR, Dennis M, Nair RV, Llanes A, Peda A, Welcome S, Rajeev S. 2017. Isolation and characterization of *Leptospira interrogans* serovar Copenhageni from a dog from Saint Kitts. *JMM Case Rep* 4:e005120. <https://doi.org/10.1099/jmmcr.0.005120>.
- Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. 2017. Canu: scalable and accurate long-read assembly via adaptive *k*-mer weighting and repeat separation. *Genome Res* 27:722–736. <https://doi.org/10.1101/gr.215087.116>.
- Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q, Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated tool for comprehensive microbial variant detection and genome assembly improvement. *PLoS One* 9:e112963. <https://doi.org/10.1371/journal.pone.0112963>.
- Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows-Wheeler transform. *Bioinformatics* 26:589–595. <https://doi.org/10.1093/bioinformatics/btp698>.
- Li H. 2011. A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. *Bioinformatics* 27:2987–2993. <https://doi.org/10.1093/bioinformatics/btr509>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.