

## Concise Review: Advances in Generating Hepatocytes from Pluripotent Stem Cells for Translational Medicine

DAGMARA SZKOLNICKA, DAVID C. HAY

**Key Words.** Pluripotent stem cells • Hepatocyte-like cells • Cell-based modeling • Technology scale-up • Translational medicine

Medical Research Council Centre for Regenerative Medicine, University of Edinburgh, Edinburgh, Scotland, United Kingdom

Correspondence: David C. Hay, Ph.D., MRC Centre for Regenerative Medicine, University of Edinburgh, 5 Little France Drive, Edinburgh EH16 4UU, Scotland, United Kingdom. Telephone: +44(0)1316519500; e-mail: davehay@talktalk.net

Received December 22, 2015; accepted for publication March 10, 2016; first published online in *STEM CELLS EXPRESS* March 26, 2016.

© AlphaMed Press  
1066-5099/2016/\$30.00/0

[http://dx.doi.org/  
10.1002/stem.2368](http://dx.doi.org/10.1002/stem.2368)

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

### ABSTRACT

The liver is one of the major organs in the human body. Severe or prolonged exposure of the liver to different factors may cause life-threatening disease, which necessitates donor organ transplantation. While orthotopic liver transplantation can be used to effectively treat liver failure, it is an invasive procedure, which is severely limited by organ donation. Therefore, alternative sources of liver support have been proposed and studied. This includes the use of pluripotent stem cell-derived hepatocytes as a renewable source of cells for therapy. In addition to cell-based therapies, in vitro engineered liver tissue provides powerful models for human drug discovery and disease modeling. This review focuses on the generation of hepatocyte-like cells from pluripotent stem cells and their application in translational medicine. *STEM CELLS* 2016;34:1421–1426

### SIGNIFICANCE STATEMENT

There has been tremendous progress in the development of efficient and defined hepatocyte differentiation from pluripotent stem cells, yet instability of hepatocyte cell phenotype still exists. We have shown that this is not specific to stem cell-derived hepatocytes, but also observed in gold standard primary hepatocytes cultured in two or three dimensions [48]. Therefore, we need to stabilize cell phenotype, so that somatic cell technology can be dependably scaled for application. Key to this will be the building of supportive liver niches in vitro. This review focuses on the new advances in the generation of hepatocyte-like cells and their application in translational medicine.

### THE LIVER

The liver is a multifunctional and highly regenerative organ, playing an important role in human physiology [1]. While resilient, the liver is susceptible to tissue damage and, therefore, degenerative diseases. Significant morbidity, mortality, and economic burden are associated with human liver disease. Therefore, the development of new systems that improve the study and treatment of liver disease are essential.


The structure of the liver is essential to its multifunctional performance. In the context of disease, liver structure becomes gradually more distorted with the loss of the hepatocyte compartment and consequently organ function [2]. Hepatocytes are located in the parenchyma and comprise approximately 70–80% of the liver mass [3]. Their function is supported by the non-parenchymal cells, forming a functional unit termed the acinus [4]. Hepatocyte polarization

is essential for proper function. The basolateral surface of hepatocytes is directly connected with sinusoidal endothelial cells, which facilitate mass transport between the parenchyma and the blood stream. At the apical surface, tight junction formation between hepatocytes is required for canaliculus formation and bile acid transport [5].

Although hepatocytes are extremely stable in vivo, they rapidly lose their phenotype in vitro [6]. This has significant consequences for scientists and clinicians who wish to build models of human liver biology “in a dish” or develop pioneering treatments for human liver disease.

### CELL-BASED MODELS

Although human hepatocytes are scarce and inherently unstable in vitro, they have been successfully deployed to model human biology



**Figure 1.** Hepatocyte differentiation and expansion. Hepatocytes can be derived from human pluripotent stem cells (HLC) and via conversion of human somatic cells (hiHeps). Hepatocytes can also be expanded from primary human hepatocytes (PHH) using cell transformation techniques and small molecules. Abbreviations: hESC, human embryonic stem cell; hiPSC, human induced pluripotent stem cell; PHH: primary human hepatocytes.

and bridge patients until their liver recovers or a transplant becomes available [7]. To bypass the issues of scarcity and instability, several groups have immortalized human hepatocytes (Fig. 1). Unfortunately, the derivative cell lines exhibited both poor function and karyotypic instability, limiting their large-scale application (for a review see ref. 28).

Given the practical issues associated with primary material, researchers have turned to cancer-derived cell lines to perform *in vitro* studies. Hepatic cancer cell lines such as HepG2, Hep3B, HepaRG, or Fa2N-4 have been extensively used in drug metabolism studies. Although these cell lines have their uses, poor cell phenotype and their resilience to toxicological insult, limit their accuracy and extrapolation to human biology. To overcome the issues associated with cancer or immortalized cell lines, researchers have focused on improving hepatocyte cell expansion and phenotype using synthetic polymers [29] and small molecules [10, 24]. More recently, Levy *et al.* [30] have expanded primary human hepatocytes, for up to 40 population doublings, by ectopic expression of Human Papilloma Virus (HPV) E6 and E7 oncoproteins. Although the cells produced *in vitro* will be useful for *in vitro* modeling studies, their utility is limited, as they are not derived from self-renewing populations and are not appropriate for use in the clinic.

Human pluripotent stem cells (PSCs), human embryonic (hESCs), and induced pluripotent (hiPSCs) stem cells, offer a scalable alternative to primary and transformed cells [31]. hESCs are derived from the inner cell mass of blastocysts that are unsuitable for human implantation. The cells display two important attributes, self-renewal and pluripotency, promising an unlimited supply of human somatic cells *in vitro* [32, 33].

Induced pluripotent stem cells (iPSCs) were initially generated by the introduction of four transcription factors (Oct 3/4, Sox2, c-Myc, Klf4) using integrative retrovirus technology [11, 34]. This led to multiple genomic insertions and was a major concern for the field. Since those seminal studies there have been numerous attempts to generate insertion-free human iPSCs, using Sendai virus [35], adenovirus [36], episomal vectors [37], the piggyBac system [38, 39] or mRNAs [40]. Those methods have proved successful, with Sendai virus reprogramming system currently considered as the most efficient reprogramming system [41, 42].

PSCs have created new opportunities to model human biology and provide the prospect of personalized medicine. In recent years, significant progress has been made in differentiating PSCs into HLCs (Fig. 1). The use of defined factors and serum-free media has facilitated the development of efficient

procedures for HLC specification using either spontaneous or directed differentiation. Spontaneous differentiation involves formation of three-dimensional (3D) multicellular aggregates, termed embryoid bodies. In the presence of particular growth factors (e.g., Wnt, BMP, FGF, HGF, or OSM) these 3D structures differentiate into HLCs [15–17]. Although this method is reproducible, its spontaneous nature has several drawbacks, including low efficiency and mixed cell type generation, which limit technology scale-up. In contrast, directed differentiation is performed in two dimensions and has proved to be more efficient [18–23, 43–48]. Some of the achievements to date using HLCs include; the accurate prediction of human drug metabolism [49–51], the mechanistic analysis of drug-induced liver injury [52, 53], the use of noncoding RNAs to modulate drug overdose [54], the study of virus infection and replication [55–59] and the ability to model inherited monogenic metabolic disorders of the liver [60, 61].

Such findings demonstrate the importance of PSC based liver models in the development of platform technologies to study human biology. This will likely lead to the identification of new medicines or the re-purposing of existing medicines to treat human disease.

In addition to PSCs, somatic cells have been transdifferentiated to hepatocytes (iHeps). Like iPSCs, iHeps can be derived from defined genetic background. Importantly, the production of iHeps bypasses the need for pluripotency (Fig. 1). Initially, Huang et al. [62] and Sekiya & Suzuki [63] generated induced hepatocytes (iHeps) from mouse fibroblasts using viral transduction and expression of either GATA4, HNF1A, and FOXA3 and inactivation of p19<sup>Arf</sup> or HNF4A in combination with FOXA1, FOXA2, or FOXA3. Three years later Huang et al. [25] and Du et al. [26] generated functional hiHeps from human somatic cells using viral transduction and expression of either FOXA3, HNF1A, and HNF4A or overexpression of hepatic nuclear factors (HNF1A, HNF4A, HNF6) in combination with ATF5, PROX1, and CEBPA. The same year, Zhu et al. [27] used a panel of small molecules to initiate hepatocyte differentiation from incompletely reprogrammed human fibroblasts. While the small molecules studies were of interest, the incompletely reprogrammed nature of the cells plus the cocktail of small molecules will complicate technology transfer and scale-up. The issue of hiHep cell scale-up has recently been addressed in an elegant study by Shi et al. [64]. Excitingly, the scaled hiHep populations were deployed in an artificial liver device and corrected abnormal blood biochemistry following acute liver failure in pigs, offering a significant therapeutic potential for the future.

#### IN VIVO TRANSPLANTATION OF LIVER PROGENITORS AND HEPATOCYTE LIKE-CELLS

Currently, the only cure for advanced liver disease is donor organ transplant. Although highly successful, the lack of donors has forced scientists to look for the alternative sources of liver support. Hepatocyte transplantation has been used to successfully treat compromised liver function. However, routine access to good quality donor livers, as for organ transplant, remains a significant limitation. Therefore, the development of a scalable and renewable source of hepatocytes would be a game changing addition.

Although several studies have demonstrated successful transplantation of pluripotent-derived hepatic cells in rodents

[65, 66], the limited capacity of cell proliferation in vivo, poor engraftment, and immune rejection rates are major challenges to clinical application. To address this, Song et al. [67] have efficiently transplanted hiPSC-derived cells in immunocompetent mice by pre-engineering 3D cell coaggregates with stromal cells followed by hydrogel encapsulation. Nagamoto et al. [68] took a different approach, improving hepatocyte engraftment and animal survival by attaching the PSC-derived hepatocyte sheets onto the surface of the liver during acute liver failure. In addition, noncoding RNAs have also been used to improve cell-based therapies. In a study by Möbus et al. [69] miR-199a-5p inhibition in hESC-derived HLCs enhanced cell engraftment in the liver.

Transplantation of adult hepatic progenitors is another promising cell-based therapy. Recently, Lu et al. [70] transplanted hepatic progenitor cells (HPCs) from wild-type mice to adult mice livers where the hepatocyte compartment had been conditionally deleted. Wild-type HPCs successfully engrafted and expanded in vivo, restoring both hepatic and biliary compartments. In addition, Huch et al. [71] have demonstrated that murine Lgr5+ liver stem cells can be expanded as epithelial organoids using a Wnt agonist and subsequently differentiated into functional hepatocytes and bile ducts in vivo. Two years later, the same group reported the successful isolation, expansion, and differentiation of human bile duct-derived progenitor cells [72].

#### CULTURE DEFINITION AND TECHNOLOGY SCALE-UP

Human PSC biology has revolutionary potential for modern medicine. The majority of procedures, published to date rely on undefined and/or xenobiotic containing culture systems. The undefined components found in bovine serum, Matrigel<sup>TM</sup>, or from feeder cell layers, among others, elicit unknown biological effects and lead to phenotypic variability in vitro. This is a significant limitation, which hampers the scale-up and application of PSCs and their derivatives [73].

To overcome these issues, researchers and companies have focused on developing defined, xeno-free, and serum-free media formulations [74–81]. In a multicenter trial, Akopian et al. [73] examined eight different serum-free media formulations in five different laboratories and concluded that StemPro and mTeSR1 were the only formulations, which supported stem cell self-renewal for at least 10 passages.

In addition to stem cell self-renewal and the maintenance of pluripotency, defined culture is essential for cellular differentiation, scale-up, and biomedical application. To improve the definition of the differentiation procedures, serum-free processes have been developed [45] and used in combination with recombinant extracellular matrices [47]. To improve culture definition and further reduce differentiation costs, small molecules have also been used to hepatic differentiation [19, 20, 24, 27]. While promising, those small molecule studies relied on undefined culture components, demonstrating the need for further research and development in this space.

#### IMPROVING THE CURRENT STATE OF THE ART AND TECHNOLOGY SCALE-UP

Many studies have focused on improving hepatocyte physiology and biology “in the dish.” Studies by Miki et al. [82] demonstrated

that oxygenation as well as continuous supply of nutrients, using hollow fiber technology, promoted mature hepatocyte gene expression. Additional studies using defined media and synthetic tissue culture substrata, have shown promise in cell specification and the maintenance of hepatic phenotype from both research and GMP hESC lines [46, 47]. These provide GMP ready options for large-scale manufacture. In addition to culture definition, faithful markers of cell specification are required to ensure proper cellular differentiation and to control for quality. Currently, the most widely used markers for tracking endoderm induction from PSCs are C-KIT, CXCR4, and EPCAM [14, 83]. Although efficient, these markers are not endoderm specific. Therefore, new lineage markers are required to track endoderm and ultimately hepatic specification.

Recently, Holtzinger et al. [84] have identified two new endodermal markers, HDE1 and HDE2. HDE1 marked the component of the definitive endoderm population with high hepatic potential, whereas HDE2 tracked developing hepatic progenitors and hepatocytes.

Those markers are important additions to the field and allow for the formation of purer DE population from stem cells, improving cell identity and phenotype in culture. Along similar lines, Kido et al. [85] have shown that carboxypeptidase M is an efficient marker to isolate and culture hepatic progenitors from PSCs. In addition to cell tracking, the identification of gene signatures, which predict stable cell phenotype are also important. Recently, we demonstrated that the use of a defined polymer substrate, in conjunction with serum-free hepatic differentiation, revealed a unique gene signature (MMP13, CTNND2, and THBS2), which predicted stable hepatocyte performance from both research and GMP hESC lines [46]. In the future, gene signatures may serve as important criteria for large-scale manufacturing and product release.

In addition to directing and monitoring cell differentiation, it is important to provide a supportive niche, which transmits key stimuli to support somatic cell phenotype. While two dimensional hepatic differentiation systems are efficient and functional, they are not equivalent to freshly isolated human hepatocytes [47, 48]. To overcome these issues, and mimic 3D tissue architecture, efforts have focused on developing new in vitro platforms using natural and/or synthetic materials, fluid flow and bioprinting [29, 86–89]. Excitingly, hiPSC-based systems have also been shown to generate functional and implantable human liver tissue [90].

In the quest for large-scale automated tissue production, bioprinting is an attractive approach. Recently, Faulkner-Jones et al. [88] bioprinted hiPSC-derived hepatocytes in a 3D alginate matrix. The printed cells survived this process and expressed the hepatic markers hepatocyte nuclear factor 4 alpha, albumin, and zona occludin 1. While these studies are encouraging, future

experimentation should extensively characterise the performance of bioprinted tissue in vitro and in vivo.

Bioinformatics has also proved highly effective in studying cell specification and phenotype. We have recently performed a genome-wide study where PSC-derived hepatocytes were compared to freshly isolated and cultured primary human hepatocytes (PHHs) [48]. In these studies, we identified unfavorable gene regulatory networks present in PSC-derived hepatocytes. In addition, the expression of essential nuclear factors such as constitutive androstane receptor, pregnane X receptor and the farnesoid X receptor were much lower in stem cell-derived hepatocytes than in PHHs. Therefore, to further differentiate PSC-derived hepatocytes to mature populations, the modulation of factors responsible for appropriate and inappropriate gene expression are required. Most recently, we have made progress in this space removing Matrigel™ extracellular matrix from our differentiation system and replacing this with recombinant laminins. This has resulted in improved hepatocyte maturity, organization, and phenotype [47]. We believe that this offers the prospect that stem cell-derived hepatocytes can be fabricated from GMP grade hESC lines, under defined conditions and may be close to clinical application [47].

## CONCLUSION

There has been tremendous progress in the development of hepatocyte differentiation systems from PSCs. Yet, hepatocyte immaturity and instability still persist. We have shown that this is not specific to stem cell-derived hepatocytes, but also observed in adult hepatocytes cultured in vitro [48]. We believe that key to solving this issue is the provision of a supportive cell niche, which ensures faithful hepatic differentiation and long-term hepatocyte performance in vitro and in vivo.

## ACKNOWLEDGMENTS

Dr. Dagmara Szkolnicka was supported by a MRC PhD Studentship. Dr David Hay was supported by awards from the UK Regenerative Medicine Platform (MR/K026666/1 and MR/L022974/1).

## AUTHOR CONTRIBUTIONS

D.S.: manuscript writing. D.C.H.: manuscript writing, financial support, final approval of manuscript.

## DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

The authors indicate no potential conflicts of interest.

## REFERENCES

- Sadri A-R, Jeschke MG, Amini-Nik S. Advances in liver regeneration: Revisiting hepatic stem/progenitor cells and their origin. *Stem Cells Int* 2016;2016:7920897.
- Malhi H, Gores GJ. Cellular and molecular mechanisms of liver injury. *Gastroenterology* 2008;134:1641–1654.
- Blouin A, Bolender RP, Weibel ER. Distribution of organelles and membranes between hepatocytes and nonhepatocytes in the rat liver parenchyma. A serological study. *J Cell Biol* 1977;72:441–455.
- Kmiec Z. Cooperation of liver cells in health and disease. *Adv Anat Embryol Cell Biol* 2001;161:1–151.
- Wang L, Boyer JL. The maintenance and generation of membrane polarity in hepatocytes. *Hepatology* 2004;39:892–899.
- Mitry RR, Hughes RD, Dhawan A. Progress in human hepatocytes: Isolation, culture and cryopreservation. *Stem Cell Dev Biol* 2002;13:463–467.
- Strom SC, Fisher RA, Thompson MT et al. Hepatocyte transplantation as a bridge to orthotopic liver transplantation in terminal liver failure. *Transplantation* 1997;63:559–569.
- Eva R, De Craene B, De Kock J et al. Strategies for immortalization of primary hepatocytes. *J Hepatol* 2014;61:925–943.
- Wege H, Le HT, Chui MS et al. Telomerase reconstitution immortalizes human fetal hepatocytes without disrupting their differentiation potential. *Gastroenterology* 2003;124:432–444.

- 10 Shan J, Schwartz RE, Ross NT et al. Identification of small molecules for human hepatocyte expansion and iPS differentiation. *Nat Chem Biol* 2013;9:514–520.
- 11 Takahashi K, Tanabe K, Ohnuki M et al. Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell* 2007;131:1–12.
- 12 Aasen R, Raya A, Barrero MJ et al. Efficient and rapid generation of induced pluripotent stem cells from human keratinocytes. *Nat Biotechnol* 2008;26:1276–1284.
- 13 Yu J, Vodyanik MA, Smuga-Otto K et al. Induced pluripotent stem cell lines derived from human somatic cells. *Science* 2007;318:1917–1920.
- 14 Green MD, Chen A, Nostro MC et al. Generation of anterior foregut endoderm from human embryonic and induced pluripotent stem cells. *Nat Biotechnol* 2011;29:267–272.
- 15 Lavon N, Yanuka O, Benvenisty N. Differentiation and isolation of hepatic-like cells from human embryonic stem cells. *Differentiation* 2004;72:230–238.
- 16 Duan Y, Catan A, Meng Y. Differentiation and enrichment of hepatocyte-like cells from human embryonic stem cells in vitro and in vivo. *STEM CELLS* 2007;25:3058–3068.
- 17 Basma H, Soto-Gutierrez A, Yannam GR, et al. Differentiation and transplantation of human embryonic stem cell-derived hepatocytes. *Gastroenterology* 2009;136:990–999.
- 18 Cai J, Zhao Y, Liu Y et al. Directed differentiation of human embryonic stem cells into functional hepatic cells. *Hepatology* 2007;45:1229–1239.
- 19 Hay DC, Zhao D, Ross A et al. Direct differentiation of human embryonic stem cells to hepatocyte-like cells exhibiting functional activities. *Cloning Stem Cells* 2007;9:61–62.
- 20 Hay DC, Fletcher J, Payne C et al. Highly efficient differentiation of hESCs to functional hepatic endoderm requires Activin A and Wnt3a signalling. *Proc Natl Acad Sci USA* 2008;105:12301–12306.
- 21 Agarwal S, Holton KL, Lanza R. Efficient differentiation of functional hepatocytes from human embryonic stem cells. *STEM CELLS* 2008;26:1117–1127.
- 22 Sullivan GJ, Hay DC, Park IH et al. Generation of functional human hepatic endoderm from human induced pluripotent stem cells. *Hepatology* 2010;51:329–335.
- 23 Si-Tayeb K, Noto FK, Nagaoka M et al. Highly efficient generation of human hepatocyte-like cells from induced pluripotent stem cells. *Hepatology* 2010;51:297–305.
- 24 Siller R, Greenhough S, Naumovska E et al. Small-molecule-driven hepatocyte differentiation of human pluripotent stem cells. *Stem Cell Rep* 2015;4:939–952.
- 25 Huang P, Zhang L, Gao Y, et al. Direct reprogramming of human fibroblasts to functional and expandable hepatocytes. *Cell Stem Cell* 2014;14:370–384.
- 26 Du Y, Wang J, Jia J et al. Human hepatocytes with drug metabolic function induced from fibroblasts by lineage reprogramming. *Cell Stem Cell* 2014;14:394–403.
- 27 Zhu S, Rezvani M, Harbell J et al. Mouse liver repopulation with hepatocytes generated from human fibroblasts. *Nature* 2014;508:93–97.
- 28 Szkolnicka D, Zhou W, Lucendo-Villarin B et al. Pluripotent stem cell-derived hepatocytes: Potential and challenges in pharmacology. *Annu Rev Pharmacol Toxicol* 2013;53:147–159.
- 29 Hay DC, Pernagallo S, Diaz-Mocho J et al. Unbiased screening of polymer libraries to define novel substrates for functional hepatocytes with inducible drug metabolism. *Stem Cell Res* 2011;6:92–102.
- 30 Levy G, Bomze D, Heinz S et al. Long-term culture and expansion of primary human hepatocytes. *Nat Biotechnol* 2015;33:1264–1271.
- 31 Barbosa MS, Schlegel R. The E6 and E7 genes of HPV-18 are sufficient for inducing two-stage in vitro transformation of human keratinocytes. *Oncogene* 1989;4:1529–1532.
- 32 Thomson J, Itskovitz-Eldor J, Shapiro SS et al. Embryonic stem cell lines derived from human blastocysts. *Science* 1998;282:1145–1147.
- 33 Reubinoff BE, Pera MF, Fong CY et al. Embryonic stem cell lines from human blastocysts: Somatic differentiation in vitro. *Nat Biotechnol* 2000;18:399–404.
- 34 Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell* 2006;124:663–676.
- 35 Fusaki N, Ban H, Nishiyama A et al. Efficient induction of transgene-free human pluripotent stem cells using a vector based virus that does not integrate into the host genome. *Proc Jpn Acad Ser B Phys Biol Sci* 2009;85:348–362.
- 36 Zhou W, Freed CR. Adenoviral gene delivery can reprogram human fibroblasts to induced pluripotent stem cells. *STEM CELLS* 2009;27:2667–2674.
- 37 Yu J, Hu K, Smuga-Otto K et al. Human induced pluripotent stem cells free of vector and transgene sequences. *Science* 2009;324:797–801.
- 38 Kaji K, Norrby K, Paca A et al. Virus-free induction of pluripotency and subsequent excision of reprogramming factors. *Nature* 2008;458:771–775.
- 39 Woltjen K, Michael IP, Mohseni P et al. piggyBac transposition reprograms fibroblasts to induced pluripotent stem cells. *Nature* 2009;458:766–770.
- 40 Warren L, Manos PD, Anfeldt T et al. Highly efficient reprogramming to pluripotency and directed differentiation of human cells with synthetic modified mRNA. *Cell Stem Cell* 2010;7:618–630.
- 41 Lieu PT, Fontes A, Vemuri MC et al. Generation of induced pluripotent stem cells with CytoTune, a non-integrating Sendai virus. *Methods Mol Biol* 2013;997:45–56.
- 42 Beers J, Linask KL, Chen JA et al. A cost-effective and efficient reprogramming platform for large-scale production of integration-free human induced pluripotent stem cells in chemically defined culture. *Sci Rep* 2015;5:Article number: 11319.
- 43 Duan Y, Ma X, Zou W et al. Differentiation and characterization of metabolically functioning hepatocytes from human embryonic stem cells. *STEM CELLS* 2010;28:678–686.
- 44 Touboul T, Hannan NR, Corbinau S et al. Generation of functional hepatocytes from human embryonic stem cells under chemically defined conditions that recapitulate liver development. *Hepatology* 2010;51:1754–1765.
- 45 Szkolnicka D, Farnworth SL, Lucendo-Villarin S et al. Deriving functional hepatocytes from pluripotent stem cells. *Curr Protoc Stem Cell Biol* 2014;30:1G.5.1.
- 46 Lucendo-Villarin B, Cameron K, Szkolnicka D et al. Polymer supported directed differentiation reveals a unique gene signature predicting stable hepatocyte performance. *Adv Healthcare Mater* 2015;4:1820–1825.
- 47 Cameron K, Tan R, Schmidt-Heck W et al. Recombinant laminins drive the differentiation and self-organisation of hESC-derived hepatocytes. *Stem Cell Rep* 2015;5:1250–1262.
- 48 Godoy P, Schmidt-Heck W, Natarajan K et al. Gene networks and transcription factor motifs defining the differentiation of stem cell into hepatocyte-like cells. *J Hepatol* 2015;63:934–942.
- 49 Medine CN, Lucendo-Villarin B, Storck C et al. Developing high-fidelity hepatotoxicity models from pluripotent stem cells. *STEM CELLS TRANSL MED* 2013;2:505–509.
- 50 Holmgren G, Sjögren AK, Barragan I et al. Long-term chronic toxicity testing using human pluripotent stem cell-derived hepatocytes. *Drug Metab Dispos* 2014;42:1401–1406.
- 51 Takayama K, Morisaki Y, Kuno S et al. Prediction of interindividual differences in hepatic functions and drug sensitivity by using human iPS-derived hepatocytes. *Proc Natl Acad Sci USA* 2014;111:16772–16777.
- 52 Takayama K, Inamura M, Kawabata K et al. Efficient generation of functional hepatocytes from human embryonic stem cells and induced pluripotent stem cells by HNF4a transduction. *Mol Ther* 2012;20:127–137.
- 53 Szkolnicka D, Farnworth SL, Lucendo-Villarin B et al. Accurate prediction of drug-induced liver injury using stem cell-derived populations. *Stem Cells Transl Med* 2014;3:141–148.
- 54 Szkolnicka D, Lucendo-Villarin B, Moore J et al. Reducing hepatocyte injury and necrosis in response to paracetamol using non-coding RNAs. *Stem Cells Transl Med* (in press). DOI: 10.5966/sctm.2015-0117.
- 55 Shlomai A, de Jong YP, Rice CM. Virus associated malignancies: The role of viral hepatitis in hepatocellular carcinoma. *Sem Cancer Biol* 2014;26:78–88.
- 56 Zhou X, Sun P, Lucendo-Villarin B et al. Modulating innate immunity improves hepatitis C virus infection and replication in stem cell-derived hepatocytes. *Stem Cell Rep* 2014;3:204–214.
- 57 Roelandt P, Obeid S, Paeshuyse J et al. Human pluripotent stem cell-derived hepatocytes support complete replication of hepatitis C virus. *J Hepatol* 2012;57:246–251.
- 58 Schwartz RE, Trehan K, Andrus L et al. Modeling hepatitis C virus infection using human induced pluripotent stem cells. *Proc Natl Acad Sci USA* 2012;109:2544–2548.
- 59 Wu CY, Chen YJ, Ho HJ et al. Association between nucleoside analogues and risk of hepatitis B virus-related hepatocellular carcinoma recurrence following liver resection. *JAMA* 2012;308:1906–1914.

- 60 Rashid S, Corbineau S, Hannan N et al. Modelling inherited metabolic disorders of the liver using human induced pluripotent stem cells. *J Clin Invest* 2010;120:3127–3136.
- 61 Cayo MA, Cai J, DeLaForest A et al. JD induced pluripotent stem cell-derived hepatocytes faithfully recapitulate the pathophysiology of familial hypercholesterolemia. *Hepatology* 2012;56:2163–2171.
- 62 Huang P, He Z, Ji S et al. Induction of functional hepatocyte-like cells from mouse fibroblast by defined factors. *Nature* 2011;475:386–389.
- 63 Sekiya S, Suzuki A. Direct conversion of mouse fibroblasts to hepatocyte-like cells by defined factors. *Nature* 2011;475:390–393.
- 64 Shi XL, Gao Y, Yan Y et al. Improved survival of porcine acute liver failure by a bioartificial liver device implanted with induced human functional hepatocytes. *Cell Res* 2016;26:206–216.
- 65 Liu H, Kim Y, Sharkis S et al. In vivo liver regeneration potential of human induced pluripotent stem cells from diverse origins. *Sci Transl Med* 2011;3:82ra39.
- 66 Chen Y, Li Y, Wang X et al. Amelioration of hyperbilirubinemia in Gunn rats after transplantation of human induced pluripotent stem cell-derived hepatocytes. *Stem Cell Rep* 2015;5:1–9.
- 67 Song W, Lu YC, Frankel AS et al. Engraftment of human induced pluripotent stem cell-derived hepatocytes in immunocompetent mice via 3D co-aggregation and encapsulation. *Sci Rep* (in press). DOI: 10.1038/srep16884.
- 68 Nagamoto Y, Takayama K, Ohashi K et al. Enhancement of survival rate by human iPSC-derived hepatocyte sheet transplantation in acute liver failure mice. *J Hepatol* (in press). DOI: 10.1016/j.hep.2016.01.004.
- 69 Möbus S, Yang D, Yuan Q et al. MicroRNA-199a-5p inhibition enhances the liver repopulation ability of human embryonic stem cell-derived hepatic cells. *J Hepatol* 2014;62:101–110.
- 70 Lu WY, Bird TG, Boulter L et al. Hepatic progenitor cells of biliary origin with liver repopulation capacity. *Nat Cell Biol* 2015;17:971–983.
- 71 Huch M, Dorrell C, Boj SF et al. In vitro expansion of single Lgr5+ liver stem cells induced by Wnt-driven regeneration. *Nature* 2013;494:247–250.
- 72 Huch M, Gehart H, Van Boxtel R et al. Long-term culture of genome-stable bipotent stem cells from adult human liver. *Cell* 2015;160:299–312.
- 73 Akopian V, Andrews PW, Beil S et al. Comparison of defined culture systems for feeder cell free propagation of human embryonic stem cells. *In Vitro Cell Dev Biol Anim* 2010;46:247–258.
- 74 Wang G, Zhang H, Zhao Y et al. Noggin and bFGF cooperate to maintain the pluripotency of human embryonic stem cells in the absence of feeder layers. *Biochem Biophys Res Commun* 2005;330:934–942.
- 75 Beattie GM, Lopez AD, Bucay N et al. Activin A maintains pluripotency of human embryonic stem cells in the absence of feeder layers. *STEM CELLS* 2005;23:489–495.
- 76 James D, Levine AJ, Besser D et al. TGFbeta/activin/nodal signalling is necessary for the maintenance of pluripotency in human embryonic stem cells. *Development* 2005;132:1273–1282.
- 77 Vallier L, Alexander M, Pedersen R. Activin/Nodal and FGF pathways cooperate to maintain pluripotency of human embryonic stem cells. *J Cell Sci* 2005;118:4495–4509.
- 78 Xiao L, Yuan X, Sharkis SJ. Activin A maintains self-renewal and regulates fibroblast growth factor, Wnt, and bone morphogenetic protein pathways in human embryonic stem cells. *STEM CELLS* 2006;24:1476–1486.
- 79 Greber B, Lehrach H, Adjaye J. Fibroblast growth factor 2 modulates transforming growth factor beta signalling in mouse embryonic fibroblasts and human ESCs (hESCs) to support hESC self-renewal. *STEM CELLS* 2007;25:455–464.
- 80 Xu RH, Sampsel-Barron TL, Gu F et al. NANOG is a direct target of TGB/Activin mediated SMAD signalling in human ESCs. *Cell Stem Cell* 2008;3:196–206.
- 81 Chen G, Gulbranson DR, Hou Z et al. Chemically defined conditions for human iPSC cell derivation and culture. *Nat Methods* 2011;8:424–429.
- 82 Miki T, Ring A, Gerlach J. Hepatic differentiation of human embryonic stem cells is promoted by three-dimensional dynamic perfusion culture conditions. *Tissue Eng Part C Methods* 2011;17:557–568.
- 83 Cheng X, Ying L, Lu L et al. Self-renewing endodermal progenitors lines generated from human pluripotent stem cells. *Cell Stem Cell* 2012;10:371–384.
- 84 Holtzinger A, Streeter PR, Sarangi F et al. New markers for tracking endoderm induction and hepatocyte differentiation from human pluripotent stem cells. *Development* 2015;142:4253–4265.
- 85 Kido T, Kouji Y, Suzuki K et al. CPM is a useful cell surface marker to isolate expandable bi-potential liver progenitor cells derived from human iPSCs. *Stem Cell Rep* 2015;5:508–515.
- 86 Skardal A, Smith L, Bharadwaj S et al. Tissue specific synthetic ECM hydrogels for 3D in vitro maintenance of hepatocyte function. *Biomaterials* 2012;33:4565–4575.
- 87 Rashidi H, Alhaque S, Szkolnicka D et al. Fluid shear stress modulation of hepatocyte-like cell function. *Arch Toxicol* (in press). DOI: 10.1007/s00204-016-1689-8.
- 88 Faulkner-Jones A, Fyfe C, Cornelissen D-J et al. Bioprinting of human pluripotent stem cells and their directed differentiation into hepatocyte-like cells for the generation of mini-livers in 3D. *Biofabrication* 2015;7:044102.
- 89 Gieseck III RL, Hannan NRF, Bort R et al. Maturation of induced pluripotent stem cell derived hepatocytes by 3D culture. *PLoS One* 2014;9:e86372.
- 90 Takebe T, Sekine K, Enomura M et al. Vascularized and functional human liver and iPSC-derived organ bud transplant. *Nature* 2014;499:481–484.