

ORAL PRESENTATION

Open Access

New cytokinin derivatives – their discovery, development and use for micropropagation of endangered tree species

Karel Dolezal^{1*}, Jana Malá², Pavlína Máčková², Helena Cvrcková², Michal Karady¹, Ondrej Novák¹, Lucie Szucova³, Jaromir Mikulik¹, Lukas Spichal³, Miroslav Strnad³

From IUFRO Tree Biotechnology Conference 2011: From Genomes to Integration and Delivery
Arraial d'Ajuda, Bahia, Brazil. 26 June - 2 July 2011

Elms (*Ulmus* sp.) are highly valuable trees for their great strength, tightly twisted grain, and durability, as well as for their cold and salt tolerance[1]. Wych elm (*Ulmus glabra*, Huds.) is a native species in Europe. In the Czech Republic, it is spread especially in mountain regions. During the years 1970–80, the second epidemic of Dutch elm disease destroyed most elm populations in Europe. The elm trees, which have survived the disease, are capable to maintain a high degree of resistance against the repeated infection. Micropropagation represents promising alternative of their vegetative reproduction.

The wild service tree (*Sorbus torminalis* (L.) Crantz) is a slow-growing tree reaching maximum height of 20–25 m at around 80–100 years. According to the latest data, the species is also disappearing from forests of the Czech Republic and it is considered to be endangered. On the other hand, the wild service tree is rated as one of the most valuable hardwoods [2] with a great potential for wider use in forestry and forest ecology, and also for its importance in the timber industry.

The *in vitro* induction of organogenesis, rooting and acclimatization of above mentioned elm species as well as wild service tree were optimized and standardized. Widely used BAP is an important and affordable cytokinin routinely utilized for its effective stimulatory properties in micropropagation practice. On the other hand, BAP may negatively influence the growth, rooting and acclimatization processes in some crops [3]. It is known that hydroxylated aromatic cytokinins (topolins) are more resistant to cytokinin oxidase (CKX), more stable

and in some biological systems active at lower concentrations than the isoprenoid CKs. In addition, *meta*-topolin does not inhibit root formation which is a typical inconvenient effect of high concentrations of BAP. These properties might help to enhance the future productivity of plant tissue culture industry crops [3,4]. The development of other new CK derivatives exhibiting high morphogenetic activity might consequently be of a great practical importance in plant biotechnology. During our recent search for naturally occurring aromatic cytokinins (ARCKs) in plants, another new group of endogenous BAP derivatives – methoxytopolins – exhibiting high biological, especially anti-senescence (and surprisingly also anti-cancer) activity, was discovered. Based on these results, we synthesized several groups of their synthetic analogues that exhibited high activity in three different CK bioassays and showed the ability to activate cytokinin receptors and/or to inhibit CKX. Best compounds so far (6-(3-hydroxybenzylamino)purine (mT) and the 6-(3-methoxybenzylamino)purine-9-ribofuranoside (MeomTR)) were used for retardation of senescence during multiplication stage of micropropagation of above mentioned tree species as well as for rooting support. Subsequently, wide range of endogenous plant hormones (isoprenoid cytokinins, IAA, ethylene) were quantified (using UPLC-MS/MS and/or GC/FID, respectively) and compared in relation to cytokinin exogenously used.

The results about optimal endogenous plant hormone concentrations and their dependence on different exogenous cytokinin used in the cultivation media, may improve *in-vitro* micropropagation efficiency as well as quality of *ex vitro* acclimatized plants of above mentioned

* Correspondence: karel.dolezal@upol.cz

¹Laboratory of Growth Regulators, Palacký University and Institute of Experimental Botany AS CR, 783 71 Olomouc-Holice, The Czech Republic
Full list of author information is available at the end of the article

elm species, wild service tree and possibly also of other tree species.

This work was supported by the Ministry of Agriculture of the Czech Republic (Mze 0002070202, QI 92A247), Czech Ministry of Education (MSM 6198959216), ED0007/01/01 Centre of the Region Haná for Biotechnological and Agricultural Research.

Author details

¹Laboratory of Growth Regulators, Palacký University and Institute of Experimental Botany AS CR, 783 71 Olomouc-Holice, The Czech Republic.

²Institute of Forest and Game Management, Strnady 136, 252 02 Jíloviště, The Czech Republic. ³Centre of the Region Haná for Biotechnological and Agricultural Research, Faculty of Science, Palacký University, Šlechtitelů 11, CZ 783 71 Olomouc, Czech Republic.

Published: 13 September 2011

References

1. Heybroek HM, Elgersma DM, Scheffer RJ: **Dutch elm disease - an ecological accident.** *Outlook Agric* 1982, **11**:1-9.
2. Demesure B, Leguerroué B, Lucchi G, Prat D, Petit RJ: **Genetic variability of a scattered temperate forest tree *Sorbus torminalis* L-(Crantz).** *Ann. For. Sci* 2000, **57**:63-71.
3. Strnad M: **The aromatic cytokinins.** *Physiol. Plant* 1997, **101**:674-688.
4. Werbrouck SPO, Strnad M, Van Onckelen HA, Debergh PC: **Meta-topolin, an alternative to benzyladenine in tissue culture?** *Physiol. Plant* 1996, **98**:291-298.

doi:10.1186/1753-6561-5-S7-O46

Cite this article as: Dolezal *et al.*: New cytokinin derivatives – their discovery, development and use for micropropagation of endangered tree species. *BMC Proceedings* 2011 **5**(Suppl 7):O46.

Submit your next manuscript to BioMed Central
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

