



Translational relevance of rodent models of hypothalamic-pituitary-adrenal function and stressors in adolescence

Cheryl M. McCormick ^{a, b,*}, Matthew R. Green ^a, Jonathan J. Simone ^b

^a Department of Psychology, Brock University, 1812 Sir Isaac Brock Way, St. Catharines, ON, L2S 3A1, Canada

^b Department of Biological Sciences, Brock University, 1812 Sir Isaac Brock Way, St. Catharines, ON, L2S 3A1, Canada



ARTICLE INFO

Article history:

Received 1 June 2016

Received in revised form

18 August 2016

Accepted 20 August 2016

Available online 29 August 2016

Keywords:

Puberty

Rats

Corticosterone

Social stress

Gonadal

ABSTRACT

Elevations in glucocorticoids that result from environmental stressors can have programming effects on brain structure and function when the exposure occurs during sensitive periods that involve heightened neural development. In recent years, adolescence has gained increasing attention as another sensitive period of development, a period in which pubertal transitions may increase the vulnerability to stressors. There are similarities in physical and behavioural development between humans and rats, and rats have been used effectively as an animal model of adolescence and the unique plasticity of this period of ontogeny. This review focuses on benefits and challenges of rats as a model for translational research on hypothalamic-pituitary-adrenal (HPA) function and stressors in adolescence, highlighting important parallels and contrasts between adolescent rats and humans, and we review the main stress procedures that are used in investigating HPA stress responses and their consequences in adolescence in rats. We conclude that a greater focus on timing of puberty as a factor in research in adolescent rats may increase the translational relevance of the findings.

© 2016 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Contents

1. Introduction	31
2. Pubertal development in humans and rats and the definition of adolescence	32
3. Hypothalamic-pituitary-adrenal (HPA) function in adolescence in humans and rats	33
4. Choice of stress procedure for a rat model of adolescent stress	34
4.1. Administration of exogenous corticosterone	34
4.2. Predator stress	37
4.3. Restraint stress	37
4.4. Social stressors	37
4.4.1. Social defeat/Resident-intruder stress	37
4.4.2. Social isolation	38
4.4.3. Social instability	38
4.5. Chronic unpredictable stress/chronic variable stress	38
5. Conclusions	38
Acknowledgements	39
References	39

1. Introduction

* Corresponding author. Centre for Neuroscience, Department of Psychology, Brock University, 1812 Sir Isaac Brock Way, St. Catharines, ON, L2S 3A1, Canada.

E-mail address: cmccormick@brocku.ca (C.M. McCormick).

Glucocorticoid hormones (mostly cortisol in humans and corticosterone in rats), the release of which is under the control of

the hypothalamic-pituitary-adrenal (HPA) axis, initiate mechanisms that enable the individual to adapt to the immediate demands of the environment. For example, elevations in glucocorticoids that result from the activation of the HPA axis in response to stressors have immunosuppressive and anti-inflammatory effects, influence lipid and glucose metabolism, and inhibit reproductive functions. In the CNS, through actions on neurogenesis, synaptic and dendritic remodeling, neurotransmission, and learning and memory functions, glucocorticoids shape future brain function and behaviour. Programming effects of stress exposures on the developing fetus provide a means by which glucocorticoids experienced in early life have effects across the lifespan and possibly into future generations (Bale, 2014; Meaney et al., 2007). The preclinical literature has made great inroads into mechanisms involved in the maladaptive consequences of excessive glucocorticoid exposures in early life and in adulthood that result from repeated or chronic stressors, or a severe stress exposure. So, why the interest in the adolescent period, an interest that has grown exponentially in the last 15 years?

Adolescence is of great clinical importance. It is the time in which many mental health problems such as mood disorders emerge. Further, the risk of drug abuse and addiction is greater from drug exposures in adolescence than in adulthood (e.g., Blomeyer et al., 2013). The clinical problems of many adult psychiatric patients originate in adolescent experiences (Oldehinkel and Bouma, 2011). In adolescents, as in adults, atypical HPA function and/or a history of stress exposures, are precursors to clinical depression (Guerry and Hastings, 2011). Thus, the growing interest in the adolescent period is tied to understanding risks that might be associated specifically with this time of life, a time that had been relatively neglected by researchers (Grant et al., 2003) until recently. Further, the idea that developmental plasticity (defined as a lasting phenotypic change in response to cues received in the past, Fawcett and Frankenhuys, 2015) was greatest during critical periods of prenatal and neonatal life and diminished thereafter with age is confronted with evidence that there are a series of sensitive periods across the lifespan in many species. Adolescence, specifically, has been cast as a sensitive period for the development of social functions in species as diverse as zebra finches (Ruploh et al., 2014), guinea pigs (Sachser, 1992), rats (McCormick et al., 2015), and humans (Blakemore, 2012), among others. Challenges to the view that adolescence was a specialization unique to humans (e.g., Bogin and Smith, 1996) also promoted the possibility of animal models for understanding human adolescence.

The importance of animal models for understanding mental health is well documented (Stevens and Vaccarino, 2015). This review focuses on benefits and challenges of rats as a model for translational research on hypothalamic-pituitary-adrenal function and stressors in adolescence, highlighting important parallels and contrasts between adolescent rats and humans. We also review the main stress procedures that are used in investigating HPA function and stress in adolescence in rats, and factors that should be considered for rat models of adolescent stress for translational research.

2. Pubertal development in humans and rats and the definition of adolescence

Adolescence involves a transition between childhood and adulthood, and requires a reorganization of a physiology and a behaviour repertoire that is adapted to one ontogenetic period to enable adaptations for a new ontogenetic period during which reproductive function is attained. As such, one of the important hallmarks of the adolescent period is puberty. Although there is some independence between adolescent development and the

onset of puberty in that some maturational processes in the brain occur irrespective of a pubertal rise in gonadal function (Sisk and Foster, 2004), the importance of puberty to adolescence cannot be overstated. The World Health Organization defined adolescence in humans as people between the ages of 10 and 19 years, with the onset of puberty marking the transition from childhood to adolescence (http://www.who.int/maternal_child_adolescent/topics/adolescence/dev/en/).

Puberty begins at about 8–10 years of age in girls and about a year later in boys and involves a rise in kisspeptin signaling, which results in increased gonadotrophin releasing hormone release, the hypothalamic hormone of the hypothalamic-pituitary-gonadal pathway (Cortés et al., 2015). The increase in estrogenic function results in the development of breast buds in girls typically between the ages of 10 and 11 years, and menarche at about 12 years of age (Parent et al., 2003). Menarche is a relatively late manifestation of puberty that usually (but not always) is preceded by the first ovulation (Cortés et al., 2015). Further, the mean age of onset of menarche is later in underprivileged populations than in “well-off” populations (Parent et al., 2003). An increase in testicular volume is an early marker of pubertal onset in boys, and the mean age of the time of this increase is 11.5 years (Lee et al., 2010). The completion of spermatogenesis is the culmination of puberty in boys. Another feature of the pubertal process in both girls and boys is a peak in growth velocity. Growth in height ceases in girls 4–5 years after menarche at a median age of 17.3 years (Spear, 2002). The growth velocity is higher in boys and growth in height stops at a median age of 21.2 years (Spear, 2002). A 4–5 year age range in pubertal onset is considered normal variation in both sexes (Parent et al., 2003). The variation in onset of puberty in humans relative to mean life expectancy, however, is negligible compared with that in other mammals, and rats in particular (Bronson and Rissman, 1986).

One of the challenges of a preclinical model of adolescence is defining the ages that are comparable to humans. Adolescence in rats has been defined liberally as being from postnatal day (PND) 21–59 (Tirelli et al., 2003) and conservatively as being from PND 28 to 42 (Spear, 2000). There are similarities and differences in the pubertal process of rats and humans. For example, whereas the gonads are quiescent until puberty and spermatogenesis only begins at puberty in humans (Plant, 2015), in Wistar rats, spermatogenesis was found to begin at postnatal day 5 and completed at PND 43 (van Haaster and de Rooij, 1993). Comparable to the growth spurt in adolescent humans, rats have a steep increase in the length of the tibia in both males and females from about PND 25–60 and a less steep rise thereafter until reaching asymptote at about PND 175 (Horton et al., 2008). Growth rates ($\mu\text{m}/\text{day}$) in the length of the tibia are highest at about PND 45. Nevertheless, rats do not have the quiescent period in growth that is evident in humans prior to the pubertal growth spurt. Instead, in rats, skeletal growth is continuous and displays an exponential trajectory that decays at about PND 64 (Horton et al., 2008). Other markers of pubertal development than growth trajectories, however, are more commonly used in studies of adolescent rats.

Physical markers of puberty in rats that coincide with increased hypothalamic-pituitary-gonadal function are the onset of vaginal opening in females, which coincides with a surge in estradiol and the onset of ovulation (Castellano et al., 2011; Ojeda and Urbanski, 1994), and balanopreputial separation in males, which coincides with a rise in androgen concentrations and with sperm in the epididymis (reviewed in McCormick and Mathews, 2010). Regular estrous cycles typically are evident about a week after vaginal opening, and sperm production is optimal only several weeks after balanopreputial separation (Lohmiller and Sonya, 2006). Although the ages of PND 25–42 in rats has been suggested to be analogous

to the ages of 10–18 years in humans (Saalfeld and Spear, 2016; Spear, 2015), this time span would involve mostly a prepubertal period in females based on age at vaginal opening and in males based on age at balanopreputial separation, which is inconsistent with the post-pubertal definition of adolescence for humans. Nevertheless, the overwhelming majority of research into the adolescent period has been conducted in prepubertal male rats.

Korenbrot et al. (1977) reported that about 50% of Sprague Dawley male rats had attained balanopreputial separation by PND 40 or PND 43 (in two graphs of different samples), and that 50% had mature motile sperm by about 47 days of age. They found an increase of about 20% in circulating androgen (testosterone and dihydrotestosterone) from PND 30 to 40, after which there was a steep increase (more than 200%) from PND 40 to PND 60 (Korenbrot et al., 1977). A more recent study with Sprague Dawley rats has results consistent with those of Korenbrot et al. (1977): No Sprague Dawley female had a vaginal opening at PND 28, just over 20% had vaginal openings at PND 32 and 100% had vaginal openings at postnatal day 36 (Vetter-O'Hagen and Spear, 2012). For Sprague Dawley males, none had balanopreputial separation at PND 36, about 60% did at PND 40, and 100% did at PND 48 (Vetter-O'Hagen and Spear, 2012). There was a gradual increase in estradiol from PND 28 to PND 48, at which point there was no further increase, and the increase from PND 28 was significant only by PND 40. Detectable concentrations of testosterone were only obtained at PND 40 (approx. 0.25 ng), and rose to about 1.3 ng/mL at PND 48, and rose again to about 2.5 ng/mL in PND 75 rats (Vetter-O'Hagen and Spear, 2012). Others have reported estradiol to peak at PND 35 in females, but this may be in part because of the use of a different strain of rats (Wistar rats) (Zapatero-Caballero et al., 2004). Across different strains of males, however, the pattern of a slight rise to about PND 45, and a rise again to adult concentrations at about PND 60 is the typical result (Wistar rats, Pignatelli et al., 2006; Zapatero-Caballero et al., 2003).

There are strain differences in the age at which specific male sexual behaviours (e.g., genital grooming, intromission) are demonstrated, which are thought to reflect, in part, strain differences in pubertal development (Hernandez-Gonzalez, 2000). Long Evans rats are reported to attain reproductive milestones earlier than do Wistar rats (Hernandez-Gonzalez, 2000). A recent study reports the mean age of vaginal opening in Long Evans rats to be PND 34.9, with a range of 32–38, and the mean age of balanopreputial separation to be 44.9, with a range of 42–48 (Drzewiecki et al., 2016). We find a similar mean age for vaginal opening in Long Evans female rats from a different supplier (Charles River rather than Harlan), 34.4 days, although we found a wider range in days (30–40, unpublished observations). In addition, we found no effect of daily injection stress or of daily 1 h isolation/restraint stress beginning at PND 30 on mean day of vaginal opening (unpublished observations), although there is evidence in the literature of effects of chronic stressors on timing of puberty in both humans and rodents (Parent et al., 2015). We have not investigated age of balanopreputial separation systematically, although when we have examined our Long Evans males at PND 46, all had reached this milestone. A direct comparison of two strains of males and females found an earlier onset of puberty in Long Evans than in Wistar rats in both sexes (Keeley et al., 2015).

A potential caveat in using physical markers as indicators of pubertal status is that environmental factors can disrupt the linkage between the outward physical markers and inner physiology. For example, first estrous varied from a mean age of PND 36.5 to 44.9 depending on housing conditions (grouped or not, presence of male or not) without a similar effect on day of vaginal opening (mean of 35.6–36.1 across groups) (Vandenbergh, 1976). Thus reliance on external physical markers to determine pubertal status

may be problematical in some experimental designs.

These studies of pubertal development in rats highlight a number of considerations for their use as translational models. First, when using rats as a model for human adolescence, there should be greater focus on the time post-puberty. This point does not detract from the importance of the prepubertal period; there is much evidence that prepubertal rats differ from adult rats in the experience of and consequences of exposure to stressors (see reviews by Green and McCormick, 2013b; McCormick and Green, 2013). Nevertheless, there also is evidence that postpubertal rats differ from both prepubertal and adult rats: for example, in response to amphetamine (Mathews et al., 2011); in expression of tyrosine hydroxylase in the caudate nucleus and medial prefrontal cortex (Mathews et al., 2009); in HPA function and its regulation by testosterone release (Green et al., 2015, 2016); in testosterone's influence on sonic hedgehog signaling (Bond et al., 2010); in corticotropin releasing factor receptor expression in various brain regions (Lukkes et al., 2016); in performance on associative learning tasks and D1 receptors in the orbitofrontal cortex and in piriform cortex (Garske et al., 2013).

Another important example is in the investigation of differences between prepubertal and postpubertal rats in sensitivity to reward; an argument that has been made in favour of rats as an animal model for investigating adolescence is that both adolescent rats and adolescent humans differ from their adult counterparts in motivation and reward (higher reward sensitivity to both natural and drug rewards in the younger age groups than in the adult age groups) (Doremus-Fitzwater et al., 2010). Many such adolescent versus adult comparisons, however, have involved prepubertal and adult rats (e.g., Doremus-Fitzwater et al., 2012; Hammerslag and Gulley, 2014; Torres et al., 2008). In a study of reward value of palatable foods, the intake of sweetened condensed milk by male rats during a 15 min period of intermittent access increased to a peak at PND 50 after which there was a steep decline to PND 70 (Friemel et al., 2010). Postpubertal PND 50 rats displayed more lever presses in a progressive ratio test of motivational incentive than did pre-pubescent PND 40 and adult PND 70 rats, which did not differ (Friemel et al., 2010). Reward sensitivity thus was greatest during the restricted period of puberty than either before or after in rats, which suggests that the postpubertal rat may provide the best comparison for human adolescents. During adolescence, humans also show higher preferences for sweet tastes (Desor and Beauchamp, 1987) and have a higher intake of calories relative to body weight than in adulthood (Post and Kemper, 1993), which parallels the results found in rats.

In sum, the majority of research in adolescence in rats has involved primarily male rats in the prepubertal phase of adolescence. Nevertheless, postpubertal rats may provide the best model for translational research.

3. Hypothalamic-pituitary-adrenal (HPA) function in adolescence in humans and rats

In humans, there is a rise in baseline HPA function and reactivity after puberty in both sexes (Gunnar et al., 2009). Adolescent and adult men typically show greater HPA reactivity to many types of stressors, including social stressors, than do adolescent and adult women (Bouma et al., 2009; Lopez-Duran et al., 2015; Stephens et al., 2016). The greater susceptibility of adolescent girls than boys to stressors, however, may not involve sex differences in HPA function per se, but rather sex differences in terms of the types of stressors to which they are exposed and/or sex differences in whether, and the extent to which, an event is considered a stressor (Oldehinkel and Bouma, 2011). A recent study highlights how pubertal phase is critical to understanding the relationship between

HPA (dys)function and psychopathology; whereas the onset of a mood disorder was predicted by a hyporeactive HPA response to a laboratory stressor in girls in the earliest stages of puberty, a hyperreactive HPA response was predictive in later stages of pubertal development (Colich et al., 2015). Yet some studies that find pubertal status to be a relevant factor in stress reactivity in adolescence do not find sex to be a relevant factor (e.g., Hankin et al., 2015; Zhang et al., 2016). More direct studies of adolescents compared with adults are required to investigate whether there is an age-related change in the magnitude of the sex difference, or in the reversal of the direction of the sex difference, in people.

In rats, there have been few comparisons of adolescents and adults in glucocorticoid release in response to stressors, and the majority of these have involved prepubertal male adolescents. Prepubertal male rats tend to have higher and/or more prolonged release of corticosterone than do adults in response to 30 min of restraint stress (Bingham et al., 2011; Lui et al., 2012; Romeo et al., 2004a), intermittent footshock (Goldman et al., 1973), and ether inhalation (Vazquez and Akil, 1993). Prepubertal adolescents had lower corticosterone release than did adults after injection of nicotine (Cao et al., 2010; Cruz et al., 2008), paroxetine (Karanges et al., 2016), or lipopolysaccharide (Goble et al., 2011), and the two groups did not differ after administration of ethanol (Willey et al., 2012) or tetrahydrocannabinol (Schramm-Saptya et al., 2007). The extent and direction of age differences depends in part on the type of stressor, which suggests that the critical differences between adolescents and adults may be in neural regions upstream from the paraventricular nucleus rather than in HPA function specifically.

We have conducted several studies involving acute responses to stressors in postpubertal male rats. Neither prepubertal (PND 30) nor postpubertal male rats (PND 45) differed in corticosterone concentrations from those of adults (either PND 70 or 85) after 1 h of isolation in small, ventilated containers and at time points during recovery (Hodges et al., 2014; Hodges and McCormick, 2015). The adolescents, however, showed more prolonged activation in the paraventricular nucleus of the hypothalamus (as indicated by expression of immediate early genes) after the isolation than did adults (Hodges et al., 2014; Hodges and McCormick, 2015), which is consistent with evidence of more prolonged stress responding in adolescents than in adults. In postpubertal male adolescents (PND 45), we found lower corticosterone concentrations compared with adults after 15 min of confinement to an elevated platform, but faster recovery to baseline concentrations in the adults than in the adolescents (McCormick et al., 2008). We found greater corticosterone release in response to 15 min of forced swim (Mathews et al., 2008b; Waters and McCormick, 2011) and in response to 30 min of restraint in postpubertal male adolescents (PND 45–47) than in adults. Thus, stress responses differ from adults for both prepubertal and postpubertal males, although there are differences between the two adolescent groups (Green and McCormick, 2016).

In adolescent females, there are reports of greater corticosterone release in response to 30 min of restraint in prepubertal than in adult females (Romeo et al., 2004b; Viau et al., 2005), and a report of reduced corticosterone release at 15 and 30 min into a 90 min restraint session compared with adult females (Doremus-Fitzwater et al., 2009). Postpubertal females did not differ from adults in corticosterone release in response to 15 min of confinement to an open arm (McCormick et al., 2008) or forced swim (Mathews et al., 2008b). PND 45, however, is a longer time post-puberty for female rats than it is for male rats; female adolescents may have greater corticosterone release to stressors at an earlier time after puberty than do adults. In contrast to the direction of the sex difference in humans, female rats have greater corticosterone release than do males in response to a wide-range of

stressors (Goel et al., 2014). The sex differences in the HPA response to stress tend to increase after puberty in rats in keeping with the dampening effect of testosterone and the enhancing effect of estradiol on HPA function in rats (Handa and Weiser, 2014). Although results from a recent study indicate that testosterone may have the dampening effect on HPA responding in men that also is found in male rats (Stephens et al., 2016), the results from studies of sex hormones and HPA function in humans are largely inconsistent (Kajantie and Phillips, 2006). Nevertheless, there is evidence that the expression of human corticosteroid receptor genes is influenced by sex hormones, as has been found for rats (DeRijk and de Kloet, 2008).

In sum, the evidence of differences in HPA function across periods of ontogeny support that the risk associated with exposure to glucocorticoids will be specific to developmental stage. Further, the extensive, documented differences between adolescents and adults in neural structures that are substrates for glucocorticoid actions (Ahmed et al., 2015; Juraska and Willing, 2016; Shulman et al., 2016) indicate that even when faced with the same degree of glucocorticoid exposure, the consequences of the exposures will inevitably differ. Humans and rats, however, differ in the distribution of corticosteroid receptors across brain regions; there is relatively greater expression of glucocorticoid receptor expression in the neocortex in primates than in rodents (Pryce, 2008), and regions with highest receptor densities may be more susceptible to stressors. Glucocorticoid receptor polymorphisms are proving to be important factors in individual differences in HPA responses to stressors and the consequences thereof in humans, but comparable studies in rats are lacking. There is evidence, however, for stress-induced epigenetic modifications of the glucocorticoid receptor gene in both humans and rats (Li-Tempel et al., 2016; Radtke et al., 2015; Zhang et al., 2013).

In sum, puberty marks a change in HPA function in both humans and rats, which supports the use of rats as models in translational research on stress in adolescence.

4. Choice of stress procedure for a rat model of adolescent stress

Our review of chronic stress procedures in adolescent rats is not exhaustive, although we have tried to capture the main procedures used within the last 15 years. In addition, for studies involving rats of prepubertal ages, we have only included those that refer to that age as adolescence; our use of the search term adolescence may have omitted papers referring to the same age as juvenile. Table 1 describes the differences across labs in the procedures used and the ages, sex, and strain of rats involved.

4.1. Administration of exogenous corticosterone

Administering glucocorticoids either in drinking water, by subcutaneous injection, or through surgical implant of pellets rather than relying on stressor-induced elevations is an approach that has had much success, particularly as a preclinical model of depression (Sterner and Kalynchuk, 2010). Some of the advantages of exogenous administration of glucocorticoids is that it allows for better control of dosage, for example, by reducing the variability across individuals that stressors produce because of individual differences in the perception of stressors and by the reduction in corticosterone release that can occur to repeated exposures to a stressor (Sterner and Kalynchuk, 2010). There are a number of disadvantages. Direct comparisons of exogenous administration of corticosterone and of chronic stress show that the effects are sometimes different quantitatively (Conrad et al., 2007; Lussier et al., 2009), qualitatively (Conrad et al., 2004), or directionally

Table 1

Procedures used as repeated or chronic stressors in studies of adolescent rats.

Repeated/Chronic stress procedure			
Sex ^a & strain ^b	Ages ^c (Days)	Description of procedure	References
Exogenous Corticosterone			
♀/♂ SD	30–50	Drinking water (50 µg/mL for 14 days, then 25 and 12.5 µg/mL for 3 days each)	(Bertholomey et al., 2016)
♂ SD	30–50	Drinking water (50 µg/mL for 14 days, then 25 and 12.5 µg/mL for 3 days each)	(Torregrossa et al., 2012)
♀/♂ Wistar outbred	56–70	Drinking water (50 µg/mL)	(Hill et al., 2014)
♂ SD	27–33	Drinking water (200 µg/mL)	(Den et al., 2014)
♀/♂ SD	30–58	Drinking water (150 or 300 µg/mL)	(Kaplowitz et al., 2016)
♂ LE	30–45	Drinking water (400 µg/mL)	(Waters and McCormick, 2011)
♂ Lister	24–30	S.C. pellet (62.5 mg)	(Bush et al., 2003)
♂ Wistar outbred	-56–70	S.C. pellet (100 mg)	(Choy and van den Buuse, 2008)
♂ SD	28–42	S.C. pellet (50 or 200 mg)	(Lee et al., 2003)
♂ Wistar Han	28–42	Injection (7x, 1 per day, 5 mg/kg)	(Veenit et al., 2013)
♂ LE	30–45	Injection (16x, daily, 40 mg/kg)	(Waters and McCormick, 2011)
Restraint			
♂ SD	28–55	5 min (28 sessions)	(Suo et al., 2013)
♂ Wistar	~100 g age ??	10 min (6 sessions, 2 per day)	(Kusek et al., 2013)
♂ SD	29–37	20 min (7 sessions)	(Zhang and Rosenkranz, 2016)
♂ SD	32–40	20 min (7 sessions)	(Padival et al., 2015)
♀ Fischer	55–62	20 min (8 sessions)	(Panagiotakopoulos et al., 2015)
♂ Wistar Han	28–42	20 min (15 sessions)	(Hetzler and Rosenkranz, 2014)
♂ SD	35–44	30 min (10 sessions)	(Lee and Hill, 2013)
♀/♂ Wistar	26–32	1 h (7 sessions)	(Traslaviña et al., 2014)
♀/♂ Wistar	31–37	1 h (7 sessions)	(Dayi et al., 2015)
♂ Wistar	28–37	1 h (10 sessions)	(Duarte et al., 2015a; Duarte et al., 2015b)
♀/♂ SD	30–52	1 h (12 sessions)	(Barha et al., 2011)
♂ SD	29–33	1.5 h (5 sessions)	(Anderson et al., 2013)
♀/♂ SD	38–42	1.5 h (5 sessions)	(Varliinskaya et al., 2013)
♂ SD or Wistar	30–34	2 h (5 sessions)	(Fernández et al., 2016)
♀ SD	25–31	2 h (7 sessions)	(Lee and Noh, 2015)
♂ SD	28–34 or 42–48	2 h (7 sessions)	(Bingham et al., 2011)
♂ SD	42–48	3 h (7 sessions)	(Negrón-Oyarzo et al., 2014)
♂ SD	42–49	3 h (7 sessions)	(Negrón-Oyarzo et al., 2015)
♂ SD	26–46	6 h (21 sessions)	(Gillette et al., 2015)
Predation Stress			
♀/♂ LE	40–48	Cat odour (5 sessions)	(Wright et al., 2008)
♀/♂ LE	~38–46	Cat odour (5 sessions)	(Wright et al., 2012; Wright et al., 2013)
♂ Wistar albino	28–60	Cat fur (17 sessions)	(Kendig et al., 2011)
Social Stressors			
<i>Social Isolation</i>			
♂ Wistar	21–36	Housed singly continuously	(Cuenya et al., 2015)
♀ SD	21–42		(Lukkes et al., 2012)
♂ SD	21–42		(Lukkes et al., 2009a; Lukkes et al., 2009b; Lukkes et al., 2009c)
♂ Wistar	21–48		(Sonei et al., 2016)
♂ SD	~21–51		(Biggio et al., 2014)
♂ SD	22–28		(Granholm et al., 2015)
♂ SD	28–46		(Caruso et al., 2014)
♂ Wistar	28–48		(Cruz et al., 2016)
♂ Fischer	28–53		(Hori et al., 2014)
♂ LE	28–70		(Karkhanis et al., 2015)
♂ LE	28–70		(Skelly et al., 2015)
♂ LE	28–85		(Rau et al., 2015)
♀/♂ SD	30–50		(Hong et al., 2012; Weintraub et al., 2010)
♀ LE	30–70		(Butler et al., 2014)
<i>Social Defeat or Resident-Intruder</i>			
♂ Wistar or WTG	45–46	Defeated by resident (2x)	(Vidal et al., 2011b)
♂ WTG	45–46	Defeated by resident then placed behind wire mesh in resident's cage (2x)	(Coppens et al., 2011)
♂ LE	35–44	Defeated by resident (4x)	(Burke and Miczek, 2015)
♂ Wistar or WTG	45–58	Defeated by resident (5x)	(Vidal et al., 2011a)
♂ Wistar	45–57	Defeated by resident (2x), placed behind wire mesh in resident's cage (3x)	(Vidal et al., 2007)
♂ WTG	45–57	Defeated by resident (2x), placed behind wire mesh in resident's cage (3x)	(Coppens et al., 2014)
♂ Roman	45–57	Defeated by resident (2x), placed behind wire mesh in resident's cage (3x)	(Vidal et al., 2007)
♂ SD	42–55	Defeated by resident then placed behind wire mesh in resident's cage (5x)	(Zitnik et al., 2015)
♂ SD	35–39	Defeated by resident then placed behind wire mesh in resident's cage (5x)	(Burke et al., 2010; Burke et al., 2011; Novick et al., 2016; Watt et al., 2009; Watt et al., 2014)
♀ SD	28–32 or 42–46		(Snyder et al., 2015a)

(continued on next page)

Table 1 (continued)

Repeated/Chronic stress procedure			
Sex ^a & strain ^b	Ages ^c (Days)	Description of procedure	References
♂ SD	28–32 or 42–46	Defeated by resident then placed behind wire mesh in resident's cage (5x)	(Snyder et al., 2015b)
♂ SD	28–34 or 42–48	Defeated by resident then placed behind wire mesh in resident's cage (5x)	(Bingham et al., 2011)
♀ SD	~36–~45	Defeated by resident (7x)	(Ver Hoeve et al., 2013)
♂ and ♀ LE	45–54	Defeated by resident then placed behind wire mesh in resident's cage (7x)	(Furuta et al., 2015)
♂ Wistar (1) or SD (2)	28–34 (1) or 35–41 (2)	Defeated by resident then placed behind wire mesh in resident's cage (10x) (1) Defeated by resident (2x), placed behind wire mesh in resident's cage (1x) (2): Defeated by resident (10x)	(Buwalda et al., 2013)
♂ Wistar	28–50	Defeated by resident then placed behind wire mesh in resident's cage (23x)	(Bourke et al., 2014)
<i>Social Instability</i>			
♀ Wistar	30–38	1 h isolation then paired with new cage partner (9 sessions)	(Raftogianni et al., 2012)
♂ LE	30–45	1 h isolation then paired with new cage partner (16 sessions)	(Cumming et al., 2014; Green et al., 2013; Green and McCormick, 2013a; Hodges and McCormick, 2015; McCormick et al., 2013a; McCormick et al., 2012; Morrissey et al., 2011) (McCormick et al., 2013b; McCormick et al., 2010) (Mathews et al., 2008a; Mathews et al., 2008b; McCormick et al., 2008) (McCormick et al., 2004; McCormick et al., 2005) (Tsai et al., 2014)
♀ LE	30–45	1 h isolation then paired with new cage partner (16x)	
♀/♂ LE	30–45	1 h isolation then paired with new cage partner (16 sessions)	
♀/♂ LE	33–48	1 h isolation then paired with new cage partner (16 sessions)	
♂ SD	28–62	1 h isolation then paired with new cage partners (35 sessions)	
<i>Chronic Unpredictable Stress/Chronic Variable Stress</i>			
♀ SD	31–41	2 per day: soiled bedding, cage tilt, elevated platform, restraint, novel bedding, overnight isolation, water deprivation.	(Comeau et al., 2015)
♂ SD	30–70	6 per week: 2 physical (small cage, wet bedding, cage tilt); 2 social (isolation, crowding, foreign bedding); 2 predation (taxidermied bobcat nearby, cat fur, feline vocalizations).	(Chaby et al., 2015b; Chaby et al., 2015c)
♂ SD	30–78	6 per week: 2 physical (small cage, wet bedding, cage tilt); 2 social (isolation, crowding, foreign bedding); 2 predation (taxidermied bobcat, cat and fox urine, feline vocalizations).	(Chaby et al., 2015a)
♂ LE	30–70	6 per week: 3 physical (small cage, wet bedding, cage tilt), 3 social (isolation, crowding, foreign bedding)	(Chaby et al., 2013)
♀ SD	42–48	Each stressor once: forced swim (warm and cold water), isolation, food deprivation, water deprivation, overnight light, elevated platform (3x), foot shock (10x), crowding with constant light.	(Zaidan and Gaisler-Salomon, 2015)
♂ bLR SD	35–60	>9 per week with increasing frequency: exposure to damp bedding, white noise, lighting, food deprivation, water deprivation, cage tilt, stroboscopic light, predator odour.	(Rana et al., 2016)
♀/♂ Wistar Han	28–42	1-2 stressors on 7 of the days: novel box, TMT odour, bright light, elevated platform.	(Toledo-Rodriguez and Sandi, 2011)
♀/♂ Wistar Han	28–30	Daily exposure: TMT odour, elevated platform	(Toledo-Rodriguez and Sandi, 2007)
♂ SD	28–55	1 stressor per day; for social stress: isolation, novel environment, crowding, litter-shifting, subordination (resident-intruder); for physical stress: cold, ether, forced swim, restraint, loud noise	(Kabbaj et al., 2002)
♀/♂ Fischer	37–44	2-3 per day: restraint, exposure to cold (4° C), food deprivation, wet bedding, swim stress, crowding.	(Taylor et al., 2013)
♂ SD	33–35	One per day: forced swim, elevated platform (3x), foot shock (6x)	(Tsoory and Richter-Levin, 2006)
♂ SD	27–29	One per day: forced swim, elevated platform (3x), foot shock (6x)	(Saul et al., 2012)
♂ LE	26–35	2 per day: forced swim, tail pinch, cat fur, restraint.	(Wright et al., 2015)
♂ Wistar	28–37	2 per day: forced swim, restraint, lights on overnight, lights off during the day, humid sawdust, cold stress, food and water deprivation, isolation.	(Duarte et al., 2015b)
♂ SD	~38–42		(Comeau et al., 2014)

Table 1 (continued)

Repeated/Chronic stress procedure			
Sex ^a & strain ^b	Ages ^c (Days)	Description of procedure	References
♂ Wistar Han	28–42	2 per day: soiled bedding, cage tilt, elevated platform, restraint, novel cage, overnight isolation, tail pinching. On 7 of 15 days: open field, TMT, elevated platform.	(Veenit et al., 2014)
♂ Wistar	28–37	2 per day: restraint, wet bedding, cold exposure, lights off, lights on, food and water deprivation, isolation, forced swim.	(Cruz et al., 2012)
♂ SD	27–33	1 per day, every other day: restraint, elevated platform (2x), footshock (40x)	(Luo et al., 2014)
♂ SD	35–50	2 per day: restraint, rotation, forced swim, cage tilt, wet sawdust, crowding, cold, reverse light-cycle, food and water deprivation, tail pinch.	(Xu et al., 2016)
♀ SD	45–58	2 per day: agitation, cold, open field, hypoxia, restraint.	(Wulsin et al., 2016)
♂ LE ♂ SD	45–51 28–48	One per day: restraint, TMT, tail pinch 2 per day: cold, water deprivation, agitation, tilted cage, forced swim (cold), crowding, soiled bedding, light-cycle reversal, food deprivation, tail pinch.	(Handy et al., 2016) (Suo et al., 2013)
♀/♂ LE	22–33 or 35–46	6 out of 12 days: water immersion, elevated platform, or foot shock.	(Wilkin et al., 2012)
♂ SD	~42–62	One per day: forced swim, cage rotation, isolation, damp bedding, food and water deprivation, restraint, strobe light, cage tilt.	(Reich et al., 2013)
♀/♂ Wistar ♀/♂ Wistar	37–48 37–49	One per day: social defeat, restraint One per day: social defeat, restraint	(Bourke et al., 2013) (Bourke and Neigh, 2011; Harrell et al., 2015; Kelly et al., 2014; Pyter et al., 2013)
♀/♂ SD	37–49	One per day: social defeat, restraint	(Harrell et al., 2013)

Notes.

^a Grey shading is used when the experiments were in female rats or included female rats.^b LE = Long Evans; SD = Sprague Dawley; WTG = wild-type Groeningen; bred Low Responders Sprague Dawley.^c Grey shading is used to indicate when stress procedures were applied at peripubertal ages (includes ages > 35 for females and >42 for males). Bold font indicates that stress procedures were applied post-pubertally only (>35 for females and >42 for males).

(Nacher et al., 2004). Because of individual differences in intake, administration in the drinking water does not permit the control over dosage that injection does; age differences in intake also make the comparison of adolescent versus adult treatment difficult because of the resulting dosage differences (Waters and McCormick, 2011). A problem with the other modes of administration, however, is the greater susceptibility of adolescents than adults to stress of injection and of surgery (e.g., Keeley et al., 2015; O'Shea et al., 2004; Raap et al., 2000).

4.2. Predator stress

Predator stress is a naturalistic stressor that can be readily adapted for use in the laboratory. Olfaction is a critical sensory system for guiding behaviour in rats, and thus olfactory stimuli can be used without introducing prey animals to the lab. Scents as diverse as 2-propylthietane (in weasel anal secretions), 2,5-dihydro-2,4,5-trimethylthiazoline (in fox anal secretions), cat fur or ferret fur odour, fox urine, all result in increased HPA responding (Hegab and Wei, 2014), although the specific neural regions activated differed across odours (Takahashi, 2015). In addition, some strains of rats are more sensitive to some odours than are other strains (Staples, 2010). A direct comparison of adolescent and adults rats found greater neuroendocrine responses to cat odour in the younger group than in the older group (Wright et al., 2012), which is of translational relevance for exploring the greater susceptibility of adolescents than adults to stressors.

4.3. Restraint stress

Restraint stress involves confining the animal in a space,

typically a hemi-cylindrical ventilated plastic tube or similar device made of mesh. Restraint stress differs from immobilization stress; whereas immobilization stress involves preventing movement with devices attached to head and paws, restraint stress restricts movement because of the dimensions of the space but does not prevent movement. An advantage of the use of restraint stress is that it is one of the most commonly used means of investigating stress responses and their consequences, and thus there is a vast literature in adult rats available for reference. The effects of restraint stress are considered to be the result of distress of being confined rather than any lasting effect of any physical discomfort (Buynitsky and Mostofsky, 2009). In studies with adult rats, restraint stress has been applied in repeated sessions of 5 min to several hours, and up to 6 h per day for 21 days (e.g., Conrad et al., 2003). The studies that have investigated the lasting effects of the most severe exposures have found many of the effects to dissipate with time when the chronic stress was experienced in adulthood (e.g., Radley et al., 2005). When restraint was experienced for shorter durations in adolescence, effects on HPA function and neurogenesis were evident several weeks after the last exposure (Barha et al., 2011), which is consistent with the hypothesis that adolescents are more vulnerable than are adults to stressors.

4.4. Social stressors

4.4.1. Social defeat/Resident-intruder stress

There are many different models of social defeat stress, though the commonality across procedures involves subjugating the experimental animal (intruder) to attack from an aggressive male (resident) that is defending its territory (reviewed in Hammels et al., 2015). After a number of physical attack sessions, the

psychological stress can be maintained by keeping the intruder near the resident but separated by a barrier. This model has proven to be an excellent model for understanding stress-induced dysfunction in adult rats, but it is a model that involves a number of challenges. Finding an aggressive male to act as the aggressor can be challenging, and retired breeders are more likely to be effective residents than are randomly selected male rats. Females are not as territorial as are males, and high probabilities of attacking are found usually only when the female is lactating (Toth and Neumann, 2013); thus the procedure is more difficult to apply in females. Further, there is significant variation between residents and within residents from bout to bout, which may increase the variability in the results obtained (reviewed in MacKay, 2016). Further, residents are much less aggressive toward an adolescent than toward an adult intruder, and there are many qualitative differences between the interactions of resident-adolescent intruder pairs and interactions of resident-adult intruder pairs (Burke and Miczek, 2015). Both adolescent and adult rats that experience social defeat, however, display a number of long-lasting changes in behavioural and neural function, particularly if housed singly after the exposures (Buwalda et al., 2011).

4.4.2. Social isolation

Social isolation deprives the animal of all social contact by having it housed alone for extended periods (usually more than three weeks or more) and has been referred to as a form of “sociogenic brain damage” or social malnourishment (Montagu, 1977). Although adult rats show a number of deficits after long periods of social deprivation from social isolation housing, adolescents are more susceptible (Einon and Morgan, 1977; Panksepp and Beatty, 1980), and some remediation may be possible from a return to social housing (Hellemans et al., 2004; Hol et al., 1999). Social isolation housing has been referred to as a stressor; nevertheless, the procedure does not involve a steep rise and prolonged elevation in glucocorticoids that is characteristic of other stress exposures, and instead may involve dysfunction resulting from minimal stimulation (reviewed in Green and McCormick, 2013a). Thus, although this model highlights the importance of social contact particularly in the adolescent period and has been used as a model of more severe psychopathology (e.g., schizophrenia, Fone and Porekess, 2008), it may be less useful as a translational model for adolescent stress. In humans, variation in the quality of social relationships rather than the absence versus presence of social contact is likely of greater relevance for understanding risk and resilience for the majority of psychiatric conditions.

4.4.3. Social instability

Social instability stress involves changing the social housing conditions of rats. Although this procedure usually involves changing the membership of individuals that are housed together, sometimes it also has involved periods of isolation housing and periods of overcrowding (Herzog et al., 2009). The rotation of membership in mixed-sex colonies increases aggression and results in elevations in circulating corticosterone (Haller et al., 1999). When the social instability involves changes in pair-housed rats of the same sex, little aggression is observed in either adult or adolescent males and females and the elevation in corticosterone that arises from a change in cage partner is short-lived (Hodges and McCormick, 2015; McCormick et al., 2007). We have used change of cage partners after 1 h of isolation in small containers (akin to restraint stress) as our model of social instability stress. Whereas adults readily habituate to the repeated change of cage partner after the 1 h isolation, adolescents instead show potentiated corticosterone release to repeated change in cage partners despite some habituation to repeated 1 h isolation stress (Hodges and

McCormick, 2015). We have found long-lasting effects of our social instability procedure on cognitive and emotional behaviours and behavioural responses to drugs of abuse, effects that are not observed when the procedure is applied to adults (reviewed in McCormick, 2010; McCormick et al., 2015), which suggests that the procedure may capture adolescent-specific plasticity. Many of the differences in social behaviour evident in rats as adults after adolescent social instability stress (reviewed in McCormick et al., 2015) parallel those observed after depriving adolescents of social play, suggesting that the quality of social interactions in adolescence may be as important as their presence versus absence (Hol et al., 1999; van den Berg et al., 1999).

4.5. Chronic unpredictable stress/chronic variable stress

The use of a lengthy schedule involving the application of diverse physical and psychological stressors (e.g., electric shock, period of isolation, immobilization, cage tilt, wet bedding, water deprivation, etc.) one to several stressors experienced daily for 21 days and up to 3 months has been used as an effective model of depression in adult rats (reviewed in Qiao et al., 2016). The use of different stressors prevents the reduction in corticosterone release that can occur when the same stressor is administered repeatedly. Nevertheless, the impairments typically dissipate with time when applied to adult male rats whether the procedure involves repeated exposure to the same stressor (repeated restraint) or involves varied, and more severe stressors than restraint (Bian et al., 2012; Heine et al., 2004). In contrast, a lengthy period of exposures to chronic mild stressors from PND days 30 through 78 resulted in increased anxiety that continued to be evident six months after the exposures (Chaby et al., 2015a). Typically, a much shorter schedule of stressor exposures are used in adolescent rats than in adult rats. Nevertheless, direct comparisons of chronic variable stress exposures administered in adolescence versus administered in adulthood are lacking; such studies would help pinpoint the extent to which adolescents are uniquely vulnerable.

5. Conclusions

Although the wide variety in stress procedures, ages at which they are applied, and strain of rat makes comparison of findings across studies difficult, the evidence is consistent with the hypothesis that adolescence is a period of life in which the response to stressors and the consequences of stressors differ from those in other times of life. The definition of adolescence for rats, however, is not always consistent with that used for humans. As indicated in Table 1, the majority of stress procedures have been applied peripubertally, and usually the stressors are applied mostly in the prepubertal period, and, in several studies, only on prepubertal days of age. There are far fewer studies of female rats, and when females have been included, they typically were tested at the same age as males. Because of their earlier onset of puberty, the studies of females were more likely to include a lengthier postpubertal period than those in males. As we described earlier, there are marked differences between the prepubertal and postpubertal rat beyond gonadal status.

There is increasing awareness in human research that both the timing of puberty and the phase of puberty are important factors in adolescent psychological development, and that these factors should receive greater attention by researchers (Berenbaum et al., 2015). For example, a recent study found that exposure to trauma during puberty (defined as the two years after the onset of menarche) led to greater risk of developing an anxiety disorder than exposures at earlier or later times (Marshall, 2016). Thus, greater attention to the timing of puberty in research with rats may

increase the translational relevance of the research.

Acknowledgements

CMM is funded by a Natural Sciences and Engineering Research Council (NSERC) (288348) of Canada Discovery Grant and MRG and JJS each hold an Alexander Graham Bell Canada Graduate Scholarship (NSERC).

References

- Ahmed, S.P., Bittencourt-Hewitt, A., Sebastian, C.L., 2015. Neurocognitive bases of emotion regulation development in adolescence. *Dev. Cogn. Neurosci.* 15, 11–25.
- Anderson, R.I., Agoglia, A.E., Morales, M., Varlinskaya, E.I., Spear, L.P., 2013. Stress, kappa manipulations, and aversive effects of ethanol in adolescent and adult male rats. *Neuroscience* 249, 214–222.
- Bale, T., 2014. Lifetime stress experience: transgenerational epigenetics and germ cell programming. *Dialogues Clin. Neurosci.* 16, 297–305.
- Barha, C., Brummelte, S., Lieblich, S.E., Galea, L.A.M., 2011. Chronic restraint stress in adolescence differentially influences hypothalamic-pituitary-adrenal axis function and adult hippocampal neurogenesis in male and female rats. *Hippocampus* 21, 1216–1227.
- Berenbaum, S.A., Beltz, A.M., Corley, R., 2015. The importance of puberty for adolescent development: conceptualization and measurement. *Adv. Child. Dev. Behav.* 48, 53–92.
- Bertholomey, M.L., Nagarajan, V., Torregrossa, M.M., 2016. Sex differences in reinstatement of alcohol seeking in response to cues and yohimbine in rats with and without a history of adolescent corticosterone exposure. *Psychopharmacology* 233, 2277–2287.
- Bian, Y., Pan, Z., Hou, Z., Huang, C., Li, W., Zhao, B., 2012. Learning, memory, and glial cell changes following recovery from chronic unpredictable stress. *Brain Res. Bull.* 88, 471–476.
- Biggio, F., Pisù, M.G., Garau, A., Boero, G., Locci, V., Mostallino, M.C., Olla, P., Utzeli, C., Serra, M., 2014. Maternal separation attenuates the effect of adolescent social isolation on HPA axis responsiveness in adult rats. *Eur. Neuropsychopharmacol.* 24, 1152–1161.
- Bingham, B., McFadden, K., Zhang, X.Y., Bhatnagar, S., Beck, S., Valentino, R., 2011. Early adolescence as a critical window during which social stress distinctly alters behavior and brain norepinephrine activity. *Neuropsychopharmacology* 36, 896–909.
- Blakemore, S.J., 2012. Development of the social brain in adolescence. *J. R. Soc. Med.* 105, 111–116.
- Blomeyer, D., Friemel, C.M., Buchmann, A.F., Banaschewski, T., Laucht, M., Schneider, M., 2013. Impact of pubertal stage at first drink on adult drinking behavior. *Alcoh. Clin. Exp. Res.* 37, 1804–1811.
- Bogin, B.A., Smith, H.B., 1996. Evolution of the human life cycle. *Am. J. Hum. Biol.* 8, 703–716.
- Bond, C.W., Angeloni, N.L., Podlasek, C.A., 2010. Analysis of testosterone effects on sonic hedgehog signaling in juvenile, adolescent and adult Sprague Dawley rat penis. *J. Sex. Med.* 7, 1116–1125.
- Bouma, E.M., Riese, H., Ormel, J., Verhulst, F.C., Oldehinkel, A.J., 2009. Adolescents' cortisol responses to awakening and social stress; effects of gender, menstrual phase and oral contraceptives. The TRAILS study. *Psychoneuroendocrinology* 34, 884–893.
- Bourke, C.H., Glasper, E.R., Neigh, G.N., 2014. SSRI or CRF antagonism partially ameliorate depressive-like behavior after adolescent social defeat. *Behav. Brain Res.* 270, 295–299.
- Bourke, C.H., Neigh, G.N., 2011. Behavioral effects of chronic adolescent stress are sustained and sexually dimorphic. *Horm. Behav.* 60, 112–120.
- Bourke, C.H., Raees, M.Q., Malviya, S., Bradburn, C.A., Binder, E.B., Neigh, G.N., 2013. Glucocorticoid sensitizers Bag1 and Ppid are regulated by adolescent stress in a sex-dependent manner. *Psychoneuroendocrinology* 38, 84–93.
- Bronson, F.H., Rissman, E.F., 1986. The biology of puberty. *Biol. Rev.* 61, 157–195.
- Burke, A.R., Miczek, K.A., 2015. Escalation of cocaine self-administration in adulthood after social defeat of adolescent rats: role of social experience and adaptive coping behavior. *Psychopharmacology* 232, 3067–3079.
- Burke, A.R., Renner, K.J., Forster, G.L., Watt, M.J., 2010. Adolescent social defeat alters neural, endocrine and behavioral responses to amphetamine in adult male rats. *Brain Res.* 1352, 147–156.
- Burke, A.R., Watt, M.J., Forster, G.L., 2011. Adolescent social defeat increases adult amphetamine conditioned place preference and alters D2 dopamine receptor expression. *Neuroscience* 197, 269–279.
- Bush, V.L., Middlemiss, D.N., Marsden, C.A., Fone, K.C., 2003. Implantation of a slow release corticosterone pellet induces long-term alterations in serotonergic neurochemistry in the rat brain. *J. Neuroendocrinol.* 15, 607–613.
- Butler, T.R., Carter, E., Weiner, J.L., 2014. Adolescent social isolation does not lead to persistent increases in anxiety-like behavior or ethanol intake in female Long Evans rats. *Alcoh. Clin. Exp. Res.* 38, 2199–2207.
- Buwalda, B., Geerdink, M., Vidal, J., Koolhaas, J.M., 2011. Social behavior and social stress in adolescence: a focus on animal models. *Neurosci. Biobehav. Rev.* 32, 1713–1721.
- Buwalda, B., Stubbendorff, C., Zickert, N., Koolhaas, J.M., 2013. Adolescent social stress does not necessarily lead to a compromised adaptive capacity during adulthood: a study on the consequences of social stress in rats. *Neuroscience* 249, 258–270.
- Buynitsky, T., Mostofsky, D.I., 2009. Restraint stress in biobehavioral research: recent developments. *Neurosci. Biobehav. Rev.* 33, 1089–1098.
- Cao, J.R., Belluzzi, J.D., Loughlin, S.E., Dao, J.M., Chen, Y.L., Leslie, F.M., 2010. Locomotor and stress responses to nicotine differ in adolescent and adult rats. *Pharmacol. Biochem. Behav.* 96, 82–90.
- Caruso, M.J., McClintock, M.K., Cavigelli, S.A., 2014. Temperament moderates the influence of periadolescent social experience on behavior and adrenocortical activity in adult male rats. *Horm. Behav.* 66, 517–524.
- Castellano, J.M., Bentsen, A.H., Sánchez-Garrido, M.A., Ruiz-Pino, F., Romero, M., García-Galiano, D., Aguilar, E., Pinilla, L., Diéguez, C., Mikkelsen, J.D., Tenias-Sempere, M., 2011. Early metabolic programming of puberty onset: impact of changes in postnatal feeding and rearing conditions on the timing of puberty and development of the hypothalamic kisspeptin system. *Endocrinology* 152, 3396–3408.
- Chaby, L.E., Cavigelli, S.A., Hirrlinger, A.M., Caruso, M.J., Braithwaite, V.A., 2015a. Chronic unpredictable stress during adolescence causes long-term anxiety. *Behav. Brain Res.* 278, 492–495.
- Chaby, L.E., Cavigelli, S.A., Hirrlinger, A.M., Lim, J., Warg, K.M., Braithwaite, V.A., 2015b. Chronic stress during adolescence impairs and improves learning and memory in adulthood. *Front. Behav. Neurosci.* 9, 327.
- Chaby, L.E., Cavigelli, S.A., White, A., Wang, K., Brathwaite, V.A., 2013. Long-term changes in cognitive bias and coping response as a result of chronic unpredictable stress during adolescence. *Front. Hum. Neurosci.* 7, 1–10.
- Chaby, L.E., Sheriff, M.J., Hirrlinger, A.M., Braithwaite, V.A., 2015c. Does early stress prepare individuals for a stressful future? Stress during adolescence improves foraging under threat. *Anim. Behav.* 105, 37–45.
- Choy, K.H., van den Buuse, M., 2008. Attenuated disruption of prepulse inhibition by dopaminergic stimulation after maternal deprivation and adolescent corticosterone treatment in rats. *Eur. Neuropsychopharmacol.* 18, 1–13.
- Colich, N.L., Kircanski, K., Foland-Ross, L.C., Gotlib, I.H., 2015. HPA-axis reactivity interacts with stage of pubertal development to predict the onset of depression. *Psychoneuroendocrinology* 55, 94–101.
- Comeau, W.L., Lee, K., Anderson, K., Weinberg, J., 2015. Prenatal alcohol exposure and adolescent stress increase sensitivity to stress and gonadal hormone influences on cognition in adult female rats. *Physiol. Behav.* 148, 157–165.
- Comeau, W.L., Winstanley, C.A., Weinberg, J., 2014. Prenatal alcohol exposure and adolescent stress—unmasking persistent attentional deficits in rats. *Eur. J. Neurosci.* 40, 3078–3095.
- Conrad, C.D., Grote, K.A., Hobbs, R.J., Ferayorni, A., 2003. Sex differences in spatial and non spatial Y-maze performance after chronic stress. *Neurobiol. Learn. Mem.* 79, 32–40.
- Conrad, C.D., MacMillan, D.D., Tsekhmanov, S., Wright, R.L., Baran, S.E., Fuchs, R.A., 2004. Influence of chronic corticosterone and glucocorticoid receptor antagonism in the amygdala on fear conditioning. *Neurobiol. Learn. Mem.* 81, 185–199.
- Conrad, C.D., McLaughlin, K.J., Harman, J.S., Foltz, C., Wieczorek, L., Lightner, E., Wright, R.L., 2007. Chronic glucocorticoids increase hippocampal vulnerability to neurotoxicity under conditions that produce CA3 dendritic retraction but fail to impair spatial recognition memory. *J. Neurosci.* 27, 8278–8285.
- Coppens, C.M., Coolen, A., de Boer, S.F., Koolhaas, J.M., 2014. Adolescent social defeat disturbs adult aggression-related impulsivity in wild-type rats. *Behav. Process* 108, 181–196.
- Coppens, C.M., Siripornmongkolchai, T., Wibrand, K., Alme, M.N., Buwalda, B., de Boer, S.F., Koolhaas, J.M., Bramham, C.R., 2011. Social defeat during adolescence and adulthood differentially induce BDNF-regulated immediate early genes. *Front. Behav. Neurosci.* 5.
- Cortés, M.E., Carrera, B., Rioseco, H., del Río, J.P., Vigil, P., 2015. The role of kisspeptin in the onset of puberty and in the ovulatory mechanism: a mini-review. *J. Ped. Adolsc. Gynecol.* 28, 286–291.
- Cruz, F., DeLucia, R., Planeta, C., 2008. Effects of chronic stress on nicotine-induced locomotor activity and corticosterone release in adult and adolescent rats. *Addict. Biol.* 13, 63–69.
- Cruz, F.C., Duarte, J.O., Leão, R.M., Hummel, L.F., Planeta, C.S., Crestani, C.C., 2016. Adolescent vulnerability to cardiovascular consequences of chronic social stress: immediate and long-term effects of social isolation during adolescence. *Dev. Neurobiol.* 76, 34–46.
- Cruz, F.C., Marin, M.T., Leao, R.M., Planeta, C.S., 2012. Behavioral and neuroendocrine effects of the exposure to chronic restraint or variable stress in early adolescent rats. *Int. J. Dev. Neurosci.* 30, 19–23.
- Cuenya, L., Mustaca, A., Kamenetzky, G., 2015. Postweaning isolation affects responses to incentive contrast in adulthood. *Dev. Psychobiol.* 57, 177–188.
- Cumming, M.J., Thompson, M.A., McCormick, C.M., 2014. Adolescent social instability stress increases aggression in a food competition task in adult male Long-Evans rats. *Dev. Psychobiol.* 56, 1575–1588.
- Dayi, A., Cetin, F., Sisman, A.R., Aksu, I., Tas, A., Gönenç, S., Uysal, N., 2015. The effects of oxytocin on cognitive deficit caused by chronic restraint stress applied to adolescent rats and on hippocampal VEGF and BDNF levels. *Med. Sci. Monit.* 21, 69–75.
- Den, M.L., Altmann, S.R., Richardson, R., 2014. A comparison of the short-and long-term effects of corticosterone exposure on extinction in adolescence versus adulthood. *Behav. Neurosci.* 128, 722–735.

- DeRijk, R.H., de Kloet, E.R., 2008. Corticosteroid receptor polymorphisms: determinants of vulnerability and resilience. *Eur. J. Pharmacol.* 583, 303–311.
- Desor, J.A., Beauchamp, G.K., 1987. Longitudinal changes in sweet preferences in humans. *Physiol. Behav.* 39, 639–641.
- Doremus-Fitzwater, T.L., Barreto, M., Spear, L.P., 2012. Age-related differences in impulsivity among adolescent and adult Sprague-Dawley rats. *Behav. Neurosci.* 126, 735–741.
- Doremus-Fitzwater, T.L., Varlinskaya, E.I., Spear, L.P., 2009. Social and non-social anxiety in adolescent and adult rats after repeated restraint. *Physiol. Behav.* 97, 484–494.
- Doremus-Fitzwater, T.L., Varlinskaya, E.I., Spear, L.P., 2010. Motivational systems in adolescence: possible implications for age differences in substance abuse and other risk-taking behaviors. *Brain Cogn.* 72, 114–123.
- Drzewiecki, C.M., Willing, J., Juraska, J.M., 2016. Synaptic number changes in the medial prefrontal cortex across adolescence in male and female rats: a role for pubertal onset. *Synapse* 70, 361–368.
- Duarte, J.O., Cruz, F.C., Leao, R.M., Planeta, C.S., Crestani, C.C., 2015a. Stress vulnerability during adolescence: comparison of chronic stressors in adolescent and adult rats. *Psychosom. Med.* 77, 186–199.
- Duarte, J.O., Planeta, C.S., Crestani, C.C., 2015b. Immediate and long-term effects of psychological stress during adolescence in cardiovascular function: comparison of homotypic vs heterotypic stress regimens. *Int. J. Dev. Neurosci.* 40, 52–59.
- Einon, D.F., Morgan, M.J., 1977. A critical period for social isolation in the rat. *Dev. Psychobiol.* 10, 123–132.
- Fawcett, T.W., Frankenhuys, W.E., 2015. Adaptive explanations for sensitive windows in development. *Front. Zool.* 12 (S1), 1–14.
- Fernández, M.S., Fabio, M.C., Miranda-Morales, R.S., Virgolini, M.B., De Giovanni, L.N., Hansen, C., Wille-Bille, A., Nizhnikov, M.E., Spear, L.P., Pautassi, R.M., 2016. Age-related effects of chronic restraint stress on ethanol drinking, ethanol-induced sedation, and on basal and stress-induced anxiety response. *Alcohol* 51, 89–100.
- Fone, K.C., Porkness, M.V., 2008. Behavioural and neurochemical effects of post-weaning social isolation in rodents-relevance to developmental neuropsychiatric disorders. *Neurosci. Biobehav. Rev.* 32, 1087–1102.
- Friemel, C.M., Spanagel, R., Schneider, M., 2010. Reward sensitivity for a palatable food reward peaks during pubertal development in rats. *Front. Behav. Neurosci.* 4, 39.
- Furuta, M., Ninomiya-Baba, M., Chiba, S., Funabashi, T., Akema, T., Kunugi, H., 2015. Exposure to social defeat stress in adolescence improves the working memory and anxiety-like behavior of adult female rats with intrauterine growth restriction, independently of hippocampal neurogenesis. *Horm. Behav.* 70, 30–37.
- Garske, A.K., Lawyer, C.R., Peterson, B.M., Illig, K.R., 2013. Adolescent changes in dopamine D1 receptor expression in orbitofrontal cortex and piriform cortex accompany an associative learning deficit. *PLoS One* 8, e56191.
- Gillette, R., Miller-Crews, I., Skinner, M.K., Crews, D., 2015. Distinct actions of ancestral vinclozolin and juvenile stress on neural gene expression in the male rat. *Front. Genet.* 6, 56.
- Goble, K.H., Bain, Z.A., Padow, V.A., Lui, P., Klein, Z.A., Romeo, R.D., 2011. Pubertal-related changes in hypothalamic-pituitary-adrenal axis reactivity and cytokine secretion in response to an immunological stressor. *J. Neuroendocrinol.* 23, 129–135.
- Goel, N., Workman, J.L., Lee, T.T., Innala, L., Viau, V., 2014. Sex differences in the HPA axis. *Comp. Physiol.* 4, 1121–1155.
- Goldman, L., Winget, C., Hollingshead, G.W., Levine, S., 1973. Postweaning development of negative feedback in the pituitary-adrenal system of the rat. *Neuroendocrinology* 12, 199–211.
- Granholm, L., Roman, E., Nylander, I., 2015. Single housing during early adolescence causes time-, area- and peptide-specific alterations in endogenous opioids of rat brain. *Br. J. Pharmacol.* 172, 606–614.
- Grant, K.E., Compas, B.E., Stuhlmacher, A.F., Thurm, A.E., McMahon, S.D., Halpert, J.A., 2003. Stressors and child and adolescent psychopathology: moving from markers to mechanisms of risk. *Psychol. Bull.* 129, 447–466.
- Green, M., McCormick, C., 2016. Sex and stress steroids in adolescence: gonadal regulation of the hypothalamic-pituitary-adrenal axis in the rat. *Gen. Comp. Endocrinol.* 234, 110–116.
- Green, M.R., Barnes, B., McCormick, C.M., 2013. Social instability stress in adolescence increases anxiety and reduces social interactions in adulthood in male Long-Evans rats. *Dev. Psychobiol.* 55, 849–859.
- Green, M.R., McCormick, C.M., 2013a. Effects of social instability stress in adolescence on long-term, not short-term, spatial memory performance. *Behav. Brain Res.* 256, 165–171.
- Green, M.R., McCormick, C.M., 2013b. Effects of stressors in adolescence on learning and memory in rodent models. *Horm. Behav.* 64, 364–379.
- Green, M.R., McLaren, M., Hodges, T.E., McCormick, C.M., 2015. Effects of Testosterone on Hypothalamic-pituitary-adrenal Function before, during, and after Puberty in Male Rats. Abstract. Society for Neuroscience, Chicago.
- Green, M.R., Notterdorff, R., Simone, J.J., McCormick, C.M., 2016. Glucocorticoid receptor translocation and expression of relevant genes in hippocampus in adolescent and adult male rats. *Psychoneuroendocrinology* 73, 32–41.
- Gerry, J.D., Hastings, P.D., 2011. In search of HPA axis dysregulation in child and adolescent depression. *Clin. Child. Fam. Psychol. Rev.* 14, 135–160.
- Gunnar, M.R., Wewerka, S., Frenn, K., Long, J.D., Griggs, C., 2009. Developmental changes in hypothalamus–pituitary–adrenal activity over the transition to adolescence: normative changes and associations with puberty. *Dev. Psychopathol.* 21, 69–85.
- Haller, J., Fuchs, E., Halasz, J., Makara, G.B., 1999. Defeat is a major stressor in males while social instability is stressful mainly in females: towards the development of a social stress model in female rats. *Brain Res. Bull.* 50, 33–39.
- Hammels, C., Pishva, E., De Vry, J., van den Hove, D.L., Prickaerts, J., van Winkel, R., Selten, J.P., Lesch, K.P., Daskalakis, N.P., Steinbusch, H.W., van Os, J., 2015. Defeat stress in rodents: from behavior to molecules. *Neurosci. Biobehav. Rev.* 59, 111–140.
- Hammerslag, L.R., Gulley, J.M., 2014. Age and sex differences in reward behavior in adolescent and adult rats. *Dev. Psychobiol.* 56, 611–621.
- Handa, R.J., Weiser, M.J., 2014. Gonadal steroid hormones and the hypothalamo–pituitary–adrenal axis. *Front. Neuroendocrinol.* 35, 197–220.
- Handy, C., Yanaga, S., Reiss, A., Zona, N., Robinson, E., Saxton, K.B., 2016. Stress during adolescence alters palatable food consumption in a context-dependent manner. *PLoS One* 11, e0148261.
- Hankin, B.L., Badanes, L.S., Smolen, A., Young, J.F., 2015. Cortisol reactivity to stress among youth: stability over time and genetic variants for stress sensitivity. *J. Abnorm. Psychol.* 124, 54–67.
- Harrell, C.S., Burgado, J., Kelly, S.D., Johnson, Z.P., Neigh, G.N., 2015. High-fructose diet during periadolescent development increases depressive-like behavior and remodels the hypothalamic transcriptome in male rats. *Psychoneuroendocrinology* 62, 252–264.
- Harrell, C.S., Hardy, E., Boss-Williams, K., Weiss, J.M., Neigh, G.N., 2013. Sex and lineage interact to predict behavioral effects of chronic adolescent stress in rats. *Behav. Brain Res.* 248, 57–61.
- Hegab, I.M., Wei, W., 2014. Neuroendocrine changes upon exposure to predator odors. *Physiol. Behav.* 131, 149–155.
- Heine, V.M., Maslam, S., Zareno, J., Joels, M., Lucassen, P.J., 2004. Suppressed proliferation and apoptotic changes in the rat dentate gyrus after acute and chronic stress are reversible. *Eur. J. Neurosci.* 19, 131–144.
- Hellemans, K.G.C., Benge, L.C., Olmstead, M.C., 2004. Adolescent enrichment partially reverses the social isolation syndrome. *Dev. Brain Res.* 150, 103–115.
- Hernandez-Gonzalez, M., 2000. Prepubertal genital grooming and penile erections in relation to sexual behavior of rats. *Physiol. Behav.* 71, 51–56.
- Herzog, C.J., Czeh, M., Wuttke, W., Schulz-Herbruggen, O., Hellweg, R., Fkugge, G., Fuchs, E., 2009. Chronic social instability stress in female rats: a potential animal model for female depression. *Neuroscience* 159, 982–992.
- Hetzell, A., Rosenkranz, J.A., 2014. Distinct effects of repeated restraint stress on basolateral amygdala neuronal membrane properties in resilient adolescent and adult rats. *Neuropsychopharmacology* 39, 2114–2130.
- Hill, R.A., Von Soly, S.K., Ratnayake, U., Klug, M., Binder, M.D., Hannan, A.J., van den Buuse, M., 2014. Long-term effects of combined neonatal and adolescent stress on brain-derived neurotrophic factor and dopamine receptor expression in the rat forebrain. *Biochim. Biophys. Acta BBA-Mol. Basis Dis.* 1842, 2126–2135.
- Hodges, T.E., Green, M.R., Simone, J.J., McCormick, C.M., 2014. Effects of social context on endocrine function and Zif268 expression in response to an acute stressor in adolescent and adult rats. *Int. J. Dev. Neurosci.* 35, 25–34.
- Hodges, T.E., McCormick, C.M., 2015. Adolescent and adult rats habituate to repeated isolation, but only adolescents sensitize to partner unfamiliarity. *Horm. Behav.* 69, 16–30.
- Hol, T., Van Den Berg, C.L., van Ree, J.M., Spruijt, B.M., 1999. Isolation during the play period in infancy decreases adult social interactions in rats. *Behav. Brain Res.* 100, 91–97.
- Hong, S., Flashner, B., Chiu, M., ver Hoeve, E., Luz, S., Bhatnagar, S., 2012. Social isolation in adolescence alters behaviors in the forced swim and sucrose preference tests in female but not in male rats. *Physiol. Behav.* 105, 269–275.
- Hori, M., Yamada, K., Ohnishi, J., Sakamoto, S., Furue, H., Murakami, K., Ichitani, Y., 2014. Tickling during adolescence alters fear-related and cognitive behaviors in rats after prolonged isolation. *Physiol. Behav.* 131, 62–67.
- Horton, J.A., Barateau, J.T., Loomis, R.M., Strauss, J.A., Damron, T.A., 2008. Ontogeny of skeletal maturation in the juvenile rat. *Anat. Rec.* 291, 283–292.
- Juraska, J.M., Willing, J., 2016. Pubertal onset as a critical transition for neural development and cognition. *Brain Res.* <http://dx.doi.org/10.1016/j.brainres.2016.04.012i> (ePub ahead of print).
- Kabbaj, M., Isgor, C., Watson, S.J., Akil, H., 2002. Stress during adolescence alters behavioral sensitivity to amphetamine. *Neuroscience* 113, 395–400.
- Kajantie, E., Phillips, D.J., 2006. The effects of sex and hormonal status on the physiological response to acute psychosocial stress. *Psychoneuroendocrinology* 31, 151–178.
- Kaplowitz, E.T., Savenkova, M., Karatsoreos, I.N., Romeo, R.D., 2016. Somatic and neuroendocrine changes in response to chronic corticosterone exposure during adolescence in male and female rats. *J. Neuroendocrinol.* 28, 10.111/jne.12336.
- Karanges, E.A., Ramos, L., Damphney, B., Suraev, A.S., Ki, K.M., McGregor, I.S., Hunt, G.E., 2016. Contrasting regional Fos expression in adolescent and young adult rats following acute administration of the antidepressant paroxetine. *Brain Res. Bull.* 121, 246–254.
- Karkhanis, A.N., Alexander, N.J., McCool, B.A., Weiner, J.L., Jones, S.R., 2015. Chronic social isolation during adolescence augments catecholamine response to acute ethanol in the basolateral amygdala. *Synapse* 69, 385–395.
- Keeley, R.J., Trow, J., McDonald, R.J., 2015. Strain and sex differences in puberty onset and the effects of THC administration on weight gain and brain volumes. *Neuroscience* 305, 328–342.
- Kelly, S.D., Harrell, C.S., Neigh, G.N., 2014. Chronic stress modulates regional cerebral glucose transporter expression in an age-specific and sexually-dimorphic manner. *Physiol. Behav.* 126, 39–49.
- Kendig, M.D., Bowen, M.T., Kemp, A.H., McGregor, I.S., 2011. Predatory threat

- induces huddling in adolescent rats and residual changes in early adulthood suggestive of increased resilience. *Behav. Brain Res.* 225, 405–414.
- Korenbrot, C.C., Huhtaniemi, I.T., Weiner, R.I., 1977. Preputial separation as an external sign of pubertal development in the male rat. *Biol. Reprod.* 17, 298–303.
- Kusek, M., Tokarski, K., Hess, G., 2013. Repeated restraint stress enhances glutamatergic transmission in the paraventricular nucleus of the rat hypothalamus. *J. Physiol. Pharmacol.* 64, 546–570.
- Lee, H., Noh, J., 2015. Social exclusion intensifies anxiety-like behavior in adolescent rats. *Behav. Brain Res.* 284, 112–117.
- Lee, J.M., Kaciroti, N., Appugliese, D., Corwyn, R.F., Bradley, R.H., Lumeng, J.C., 2010. Body mass index and timing of pubertal initiation in boys. *Arch. Pediatr. Adolesc. Med.* 164, 139–144.
- Lee, P.R., Brady, D., Koenig, J.I., 2003. Corticosterone alters N-methyl-D-aspartate receptor subunit mRNA expression before puberty. *Mol. Brain Res.* 115, 55–62.
- Lee, T.T.Y., Hill, M.N., 2013. Age of stress exposure modulates the immediate and sustained effects of repeated stress on corticolimbic cannabinoid CB 1 receptor binding in male rats. *Neuroscience* 249, 106–114.
- Li-Tempel, T., Larra, M.F., Sandt, E., Mériaux, S.B., Schote, A.B., Schächinger, H., Müller, C.P., Turner, J.D., 2016. The cardiovascular and hypothalamus-pituitary-adrenal axis response to stress is controlled by glucocorticoid receptor sequence variants and promoter methylation. *Clin. Epigenetics* 8, 12.
- Lohmiller, J.J., Sonya, P., 2006. Reproduction and breeding. In: Suckow, M.A., Weisbroth, S.H., Franklin, C.L. (Eds.), *The Laboratory Rat*, second ed. Elsevier, San Diego, pp. 146–164.
- Lopez-Duran, N.L., McGinnis, E., Kuhlman, K., Geiss, E., Vargas, I., Mayer, S., 2015. HPA-axis stress reactivity in youth depression: evidence of impaired regulatory processes in depressed boys. *Stress* 18, 545–553.
- Lui, P., Padov, V.A., Franco, D., Hall, B.S., Park, B., Klein, Z.A., Romeo, R.D., 2012. Divergent stress-induced neuroendocrine and behavioral responses prior to puberty. *Physiol. Behav.* 107, 104–111.
- Lukkes, J.L., Burke, A.R., Zelin, N.S., Hale, M.W., Lowry, C.A., 2012. Post-weaning social isolation attenuates c-Fos expression in GABAergic interneurons in the basolateral amygdala of adult female rats. *Physiol. Behav.* 107, 719–725.
- Lukkes, J.L., Mokin, M.V., Scholl, J.L., Forster, G.L., 2009a. Adult rats exposed to early-life social isolation exhibit increased anxiety and conditioned fear behavior, and altered hormonal stress responses. *Horm. Behav.* 55, 248–256.
- Lukkes, J.L., Norman, K.J., Meda, S., Andersen, S.L., 2016. Sex differences in the ontogeny of CRF receptors during adolescent development in the dorsal raphe nucleus and ventral tegmental area. *Synapse* 70, 126–133.
- Lukkes, J.L., Summers, C.H., Scholl, J.L., Renner, K.J., Forster, G.L., 2009b. Early life social isolation alters corticotropin-releasing factor responses in adult rats. *Neuroscience* 158, 845–855.
- Lukkes, J.L., Vuong, S., Scholl, J.L., Ikiver, H., Forster, G.L., 2009c. Corticotropin-releasing factor receptor antagonism within the dorsal raphe nucleus reduces social anxietylike behavior after early-life social isolation. *J. Neurosci.* 29, 9955–9960.
- Luo, X.M., Yuan, S.N., Guan, X.T., Xie, X., Shao, F., Wang, W.W., 2014. Juvenile stress affects anxiety-like behavior and limbic monoamines in adult rats. *Physiol. Behav.* 135, 7–16.
- Lussier, A.L., Caruncho, H.J., Kalynchuk, L.E., 2009. Repeated exposure to corticosterone, but not restraint, decreases the number of reelin-positive cells in the adult rat hippocampus. *Neurol. Lett.* 460, 170–174.
- MacKay, J.C., 2016. Long-term Behavioral and Neuroendocrine Consequences of Early Adversity (Juvenile Stressor Exposure), and the Buffering Effects of 'comfort' food. Doctoral dissertation. Université d'Ottawa/University of Ottawa, Psychology, University of Ottawa, Ottawa, p. 289.
- Marshall, A.D., 2016. Developmental timing of trauma exposure relative to puberty and the nature of psychopathology among adolescent girls. *J. Am. Acad. Child. Adolsc. Psychiatr.* 55, 25–32.
- Mathews, I.Z., Brudzynski, S.M., McCormick, C.M., 2011. Heightened locomotor-activating effects of amphetamine administered into the nucleus accumbens in adolescent rats. *I. J. Dev. Neurosci.* 29, 501–507.
- Mathews, I.Z., Mills, R.G., McCormick, C.M., 2008a. Chronic social stress in adolescence influenced both amphetamine conditioned place preference and locomotor sensitization. *Dev. Psychobiol.* 50, 451–459.
- Mathews, I.Z., Waters, P., McCormick, C.M., 2009. Changes in hyporesponsiveness to acute amphetamine and age differences in tyrosine hydroxylase immunoreactivity in the brain over adolescence in male and female rats. *Dev. Psychobiol.* 51, 417–428.
- Mathews, I.Z., Wilton, A., Styles, A., McCormick, C.M., 2008b. Heightened neuroendocrine function in males to a heterotypic stressor and increased depressive behaviour in females after adolescent social stress in rats. *Behav. Brain Res.* 190, 33–40.
- McCormick, C.M., 2010. An animal model of social instability stress in adolescence and risk for drugs of abuse. *Physiol. Behav.* 99, 194–203.
- McCormick, C.M., Green, M.R., 2013. From the stressed adolescent to the anxious and depressed adult: investigations in rodent models. *Neuroscience* 249, 242–257.
- McCormick, C.M., Green, M.R., Cameron, N.M., Nixon, F., Levy, M.J., Clark, R.A., 2013a. Deficits in male sexual behavior in adulthood after social instability stress in adolescence in rats. *Horm. Behav.* 63, 5–12.
- McCormick, C.M., Hodges, T.E., Simone, J.J., 2015. Peer pressures: social instability stress in adolescence and social deficits in adulthood in an animal model. *Dev. Cognit. Neurosci.* 11, 2–11.
- McCormick, C.M., Mathews, I.Z., 2010. Adolescent development, hypothalamic-pituitary-adrenal function, and programming of adult learning and memory. *Prog. Neuro-Psychopharmacol. Biol. Psychiatr.* 34, 756–765.
- McCormick, C.M., Merrick, A., Sezen, J., Helmreich, D.L., 2007. Social instability in adolescence alters the central and peripheral hypothalamic-pituitary-adrenal responses to a repeated homotypic stressor in male and female rats. *J. Neuroendocrinol.* 19, 116–126.
- McCormick, C.M., Mongillo, D.L., Simone, J.J., 2013b. Age and adolescent social stress effects on fear extinction in female rats. *Stress* 16, 678–688.
- McCormick, C.M., Nixon, F., Thomas, C., Lowie, B., Dyck, J., 2010. Hippocampal cell proliferation and spatial memory performance after social instability stress in adolescence in female rats. *Behav. Brain Res.* 208, 23–29.
- McCormick, C.M., Robarts, D., Gleason, E., Kelsey, J.E., 2004. Stress during adolescence enhances locomotor sensitization to nicotine in adulthood in female, but not male, rats. *Horm. Behav.* 46, 458–466.
- McCormick, C.M., Robarts, D., Kopeckina, K., Kelsey, J.E., 2005. Long-lasting, sex-and age-specific effects of social stress on corticosterone responses to restraint and locomotor responses to psychostimulants in rats. *Horm. Behav.* 48, 64–74.
- McCormick, C.M., Smith, C., Mathews, I.Z., 2008. Effects of chronic social stress in adolescence on anxiety and neuroendocrine response to mild stress in male and female rats. *Behav. Brain Res.* 187, 228–238.
- McCormick, C.M., Thomas, C.M., Sheridan, C.S., Nixon, F., Flynn, J.A., Mathews, I.Z., 2012. Social instability stress in adolescent male rats alters hippocampal neurogenesis and produces a deficit in spatial location memory in adulthood. *Hippocampus* 22, 1300–1312.
- Meaney, M.J., Szyf, M., Seckl, J.R., 2007. Epigenetic mechanisms of perinatal programming of hypothalamic-pituitary-adrenal function and health. *Trends Mol. Med.* 13, 269–277.
- Montagu, A., 1977. Sociogenic brain damage. In: Arieti, S., Chrzanowski, G. (Eds.), *New Dimensions in Psychiatry*: a World View, vol. 2. Wiley, New York, pp. 4–25.
- Morrissey, M.D., Mathews, I.Z., McCormick, C.M., 2011. Enduring deficits in contextual and auditory fear conditioning after adolescent, not adult, social instability stress in male rats. *Neurobiol. Learn. Mem.* 95, 46–56.
- Nacher, J., Pham, K., Gil-Fernandez, V., McEwen, B.S., 2004. Chronic restraint stress and chronic corticosterone treatment modulate differentially the expression of molecules related to structural plasticity in the adult rat piriform cortex. *Neuroscience* 126, 503–509.
- Negrón-Oyarzo, I., Dagnino-Subiabre, A., Carvajal, P.M., 2015. Synaptic impairment in layer 1 of the prefrontal cortex induced by repeated stress during adolescence is reversed in adulthood. *Front. Cell. Neurosci.* 9, 442.
- Negrón-Oyarzo, I., Pérez, M.A., Terreros, G., Muñoz, P., Dagnino-Subiabre, A., 2014. Effects of chronic stress in adolescence on learned fear, anxiety, and synaptic transmission in the rat prelimbic cortex. *Behav. Brain Res.* 259, 342–353.
- Novick, A.M., Mears, M., Forster, G.L., Lei, Y., Tejani-Butt, S.M., Watt, M.J., 2016. Adolescent social defeat alters N-methyl-D-aspartic acid receptor expression and impairs fear learning in adulthood. *Behav. Brain Res.* 304, 51–59.
- O'Shea, M., Singh, M.E., McGregor, I.S., Mallet, P.E., 2004. Chronic cannabinoid exposure produces lasting memory impairment and increased anxiety in adolescent but not adult rats. *J. Psychopharmacol.* 18, 502–508.
- Okuda, S.R., Urbanski, H.F., 1994. Puberty in the rat. In: Knobil, E., Neill, J.D. (Eds.), *The Physiology of Reproduction*. Raven Press, New York, pp. 363–410.
- Oldehinkel, A.J., Bouma, E.M., 2011. Sensitivity to the depressogenic effect of stress and HPA-axis reactivity in adolescence: a review of gender differences. *Neurosci. Biobehav. Rev.* 35, 1757–1770.
- Padival, M.A., Blume, S.R., Vantreas, J.E., Rosenkranz, J.A., 2015. Qualitatively different effect of repeated stress during adolescence on principal neuron morphology across lateral and basal nuclei of the rat amygdala. *Neuroscience* 291, 128–145.
- Panagiotakopoulos, L., Kelly, S., Neigh, G.N., 2015. HIV-1 proteins accelerate HPA axis habituation in female rats. *Physiol. Behav.* 180, 8–15.
- Panksepp, J., Beatty, W.W., 1980. Social deprivation and play in rats. *Behav. Neural Biol.* 30, 197–206.
- Parent, A.S., Franssen, D., Fudvoye, J., Gerard, A., Bourguignon, J.P., 2015. Developmental variations in environmental influences including endocrine disruptors on pubertal timing and neuroendocrine control: revision of human observations and mechanistic insight from rodents. *Front. Neuroendocrinol.* 38, 12–36.
- Parent, A.S., Teilmann, G., Juul, A., Skakkebaek, N.E., Toppari, J., Bourguignon, J.P., 2003. The timing of normal puberty and the age limits of sexual precocity: variations around the world, secular trends, and changes after migration. *Endocr. Rev.* 24, 688–693.
- Pignatelli, D., Xiao, F., Gouveia, A.M., Ferreira, J.G., Vinson, G.P., 2006. Adrenarche in the rat. *J. Endocrinol.* 191, 301–308.
- Plant, T.M., 2015. Neuroendocrine control of the onset of puberty. *Front. Neuroendocrinol.* 38, 73–88.
- Post, G.B., Kemper, H.C., 1993. Nutrient intake and biological maturation during adolescence. The Amsterdam growth and health longitudinal study. *Eur. J. Clin. Nutr.* 47, 400–408.
- Pryce, C.R., 2008. Postnatal ontogeny of expression of the corticosteroid receptor genes in mammalian brains: inter-species and intra-species differences. *Brain Res. Rev.* 57, 596–605.
- Pyter, L.M., Kelly, S.D., Harrell, C.S., Neigh, G.N., 2013. Sex differences in the effects of adolescent stress on adult brain inflammatory markers in rats. *Brain Behav. Immun.* 30, 88–94.
- Qiao, H., Li, M.X., Xu, C., Chen, H.B., An, S.C., Ma, X.M., 2016. Dendritic spines in depression: what we learned from animal models. *Neurol. Plast.* <http://>

- dx.doi.org/10.1155/2016/8056370.
- Raab, D.K., Morin, B., Medici, C.N., Smith, R.F., 2000. Adolescent cocaine and injection stress effects on the estrous cycle. *Physiol. Behav.* 70, 417–424.
- Radley, J.J., Rocher, A.B., Janssen, W.G., Hof, P.R., McEwen, B.S., Morrison, J.H., 2005. Reversibility of apical dendritic retraction in the rat medial prefrontal cortex following repeated stress. *Exp. Neurol.* 196, 199–203.
- Radtke, K.M., Schauer, M., Gunter, H.M., Ruf-Leuschner, M., Sill, J., Meyer, A., Elbert, T., 2015. Epigenetic modifications of the glucocorticoid receptor gene are associated with the vulnerability to psychopathology in childhood maltreatment. *Transl. Psychiatr.* 5, e571.
- Rafatgianni, A., Diamantopoulou, A., Alikaridis, F., Stamatakis, A., Stylianopoulou, F., 2012. Effects of interaction of an early experience of reward through maternal contact or its denial with social stress during adolescence on the serotonergic system and the stress responsiveness of adult female rats. *Neuroscience* 209, 84–96.
- Rana, S., Nam, H., Glover, M.E., Akil, H., Watson, S.J., Clinton, S.M., Kerman, I.A., 2016. Protective effects of chronic mild stress during adolescence in the low-novelty responder rat. *Stress* 19, 133–138.
- Rau, A.R., Chappell, A.M., Butler, T.R., Ariwodola, O.J., Weiner, J.L., 2015. Increased basolateral amygdala pyramidal cell excitability may contribute to the anxiogenic phenotype induced by chronic early-life stress. *J. Neurosci.* 35, 9730–9740.
- Reich, C.G., Iskander, A.N., Weiss, M.S., 2013. Cannabinoid modulation of chronic mild stress-induced selective enhancement of trace fear conditioning in adolescent rats. *J. Psychopharmacol.* 27, 947–955.
- Romeo, R.D., Lee, S.J., Chhua, N., McPherson, C.R., McEwen, B.S., 2004a. Testosterone cannot activate an adult-like stress response in prepubertal male rats. *Neuroendocrinology* 79, 125–132.
- Romeo, R.D., Lee, S.J., McEwen, B.S., 2004b. Differential stress reactivity in intact and ovariectomized prepubertal and adult female rats. *Neuroendocrinology* 80, 387–393.
- Ruploch, T., Bischof, H.-J., von Engelhardt, N., 2014. Social experience during adolescence influences how male zebra finches (*Taeniopygia guttata*) group with conspecifics. *Behav. Ecol. Sociobiol.* 68, 537–549.
- Saalfeld, J., Spear, L., 2016. The ontogeny of ethanol aversion. *Physiol. Behav.* 156, 164–176.
- Sachser, N., 1992. The ability to arrange with conspecifics depends on social experiences around puberty. *Physiol. Behav.* 53, 539–544.
- Saul, M.L., Tylee, D., Becoats, K.T., Guerrero, B.G., Sweeney, P., Helmreich, D.L., Fudge, J.L., 2012. Long-term behavioral consequences of stress exposure in adolescent versus young adult rats. *Behav. Brain Res.* 229, 226–234.
- Schramm-Saptya, N.L., Cha, Y.M., Chaudhry, S., Wilson, W.A., Swartzwelder, H.S., Kuhn, C.M., 2007. Differential anxiogenic, aversive, and locomotor effects of THC in adolescent and adult rats. *Psychopharmacology* 191, 867–877.
- Shulman, E.P., Smith, A.R., Silva, K., Icenogle, G., Duell, N., Chein, J., Steinberg, L., 2016. The dual systems model: review, reappraisal, and reaffirmation. *Dev. Cogn. Neurosci.* 17, 103–117.
- Sisk, C.L., Foster, D.L., 2004. The neural basis of puberty and adolescence. *Nat. Neurosci.* 7, 1040–1047.
- Skelly, M.J., Chappell, A.E., Carter, E., Weiner, J.L., 2015. Adolescent social isolation increases anxiety-like behavior and ethanol intake and impairs fear extinction in adulthood: possible role of disrupted noradrenergic signaling. *Neuropharmacology* 97, 149–159.
- Snyder, K., Barry, M., Plona, Z., Ho, A., Zhang, X.Y., Valentino, R.J., 2015a. The impact of social stress during adolescence or adulthood and coping strategy on cognitive function of female rats. *Behav. Brain Res.* 286, 175–183.
- Snyder, K.P., Barry, M., Valentino, R.J., 2015b. Cognitive impact of social stress and coping strategy throughout development. *Psychopharmacology* 232, 185–195.
- Sonei, N., Amiri, S., Jafarian, I., Anoush, M., Rahimi-Baleai, M., Bergen, H., Haj-Mirzaian, A., Hosseini, M.J., 2016. Mitochondrial dysfunction bridges negative affective disorders and cardiomyopathy in socially isolated rats: pros and cons of fluoxetine. *World J. Biol. Psychiatr.* 1–15.
- Spear, B.A., 2002. Adolescent growth and development. *J. Acad. Nutr. Diet.* March S23–S29.
- Spear, L.P., 2000. Neurobehavioral changes in adolescence. *Curr. Dir. Psychol. Sci.* 9, 111–114.
- Spear, L.P., 2015. Adolescent alcohol exposure: are there separable vulnerable periods within adolescence? *Physiol. Behav.* 148, 122–130.
- Staples, L.G., 2010. Predator odor avoidance as a rodent model of anxiety: learning-mediated consequences beyond the initial exposure. *Neurobiol. Learn. Mem.* 94, 435–445.
- Stephens, M.A.C., Mahon, P.B., McCaul, M.E., Wand, G.S., 2016. Hypothalamic–pituitary–adrenal axis response to acute psychosocial stress: effects of biological sex and circulating sex hormones. *Psychoneuroendocrinology* 66, 47–55.
- Sternier, E.Y., Kalyanchuk, L.E., 2010. Behavioral and neurobiological consequences of prolonged glucocorticoid exposure in rats: relevance to depression. *Prog. Neuro-Psychopharm. Biol. Psychiatr.* 34, 777–790.
- Stevens, H.E., Vaccarino, F.M., 2015. How animal models inform child and adolescent psychiatry. *J. Am. Acad. Child. Adolesc. Psychiatr.* 54, 352–359.
- Suo, L., Zhao, L., Si, J., Liu, J., Zhu, W., Chai, B., Zhang, Y., Feng, J., Ding, Z., Luo, Y., Shi, H., 2013. Predictable chronic mild stress in adolescence increases resilience in adulthood. *Neuropsychopharmacology* 38, 1387–1400.
- Takahashi, L.K., 2015. Olfactory systems and neural circuits that modulate predator odor fear. *Front. Behav. Neurosci.* 8, 72.
- Taylor, S.B., Taylor, A.R., Koenig, J.I., 2013. The interaction of disrupted Type II Neuregulin 1 and chronic adolescent stress on adult anxiety- and fear-related behaviors. *Neuroscience* 249, 31–42.
- Tirelli, E., Laviola, G., Adriani, W., 2003. Ontogenesis of behavioral sensitization and conditioned place preference induced by psychostimulants in laboratory rodents. *Neurosci. Biobehav. Rev.* 27, 163–178.
- Toledo-Rodriguez, M., Sandi, C., 2007. Stress before puberty exerts a sex- and age-related impact on auditory and contextual fear conditioning in the rat. *Neuroplast.* 2007, 71203.
- Toledo-Rodriguez, M., Sandi, C., 2011. Stress during adolescence increases novelty seeking and risk-taking behavior in male and female rats. *Front. Behav. Neurosci.* 5, 17.
- Torregrossa, M.M., Xie, M., Taylor, J.R., 2012. Chronic corticosterone exposure during adolescence reduces impulsive action but increases impulsive choice and sensitivity to yohimbine in male Sprague-Dawley rats. *Neuropharmacology* 57, 1656–1670.
- Torres, O.V., Tejeda, H.A., Natividad, L.A., O'Dell, L.E., 2008. Enhanced vulnerability to the rewarding effects of nicotine during the adolescent period of development. *Pharmacol. Biochem. Behav.* 90, 658–663.
- Toth, I., Neumann, I.D., 2013. Animal models of social avoidance and social fear. *Cell Tissue Res.* 354, 107–118.
- Traslañá, G.A.A., de Oliveira, F.L., Franci, C.R., 2014. Early adolescent stress alters behavior and the HPA axis response in male and female adult rats: the relevance of the nature and duration of the stressor. *Physiol. Behav.* 133, 178–189.
- Tsai, S.F., Huang, T.Y., Chang, C.Y., Hsu, Y.C., Chen, S.J., Yu, L., Kuo, Y.M., Jen, C.J., 2014. Social instability stress differentially affects amygdalar neuron adaptations and memory performance in adolescent and adult rats. *Front. Behav. Neurosci.* 8, 27.
- Tsoory, M., Richter-Levin, G., 2006. Learning under stress in the adult rat is differentially affected by 'juvenile' or 'adolescent' stress. *Int. J. Neuropharmacol.* 8, 1–16.
- van den Berg, C.L., Hol, T., Van Ree, J.M., Spruijt, B.M., Everts, H., Koolhaas, J.M., 1999. Play is indispensable for an adequate development of coping with social challenges in the rat. *Dev. Psychobiol.* 34, 129–138.
- van Haaster, L.H., de Rooij, D.G., 1993. Spermatogenesis is accelerated in the immature Djungarian and Chinese hamster and rat. *Biol. Reprod.* 49, 1229–1235.
- Vandenbergh, J.G., 1976. Acceleration of sexual maturation in female rats by male stimulation. *J. Reprod. Fertil.* 46, 451–453.
- Varlinskaya, E.I., Truxell, E.M., Spear, L.P., 2013. Repeated restraint stress alters sensitivity to the social consequences of ethanol differentially in early and late adolescent rats. *Pharmacol. Biochem. Behav.* 113, 38–45.
- Vazquez, D.M., Akil, H., 1993. Pituitary-adrenal response to ether vapor in the weanling animal: characterization of the inhibitory effect of glucocorticoids on adrenocorticotropin secretion. *Pediatr. Res.* 34, 646–653.
- Veenit, V., Cordero, M.I., Tzanoulinou, S., Sandi, C., 2013. Increased corticosterone in peripubertal rats leads to long-lasting alterations in social exploration and aggression. *Front. Behav. Neurosci.* 7.
- Veenit, V., Riccio, O., Sandi, C., 2014. CRHR1 links peripuberty stress with deficits in social and stress-coping behaviors. *J. Psychiatr. Res.* 53, 1–7.
- Ver Hoeve, E.S., Kelly, C., Luz, S., Ghanshani, S., Bhatnagar, S., 2013. Short-term and long-term effects of repeated social defeat during adolescence or adulthood in female rats. *Neuroscience* 249, 63–73.
- Vetter-O'Hagen, C.S., Spear, L.P., 2012. Hormonal and physical markers of puberty and their relationship to adolescent-typical novelty-directed behavior. *Dev. Psychobiol.* 54, 523–535.
- Viau, V., Bingham, B., Davis, J., Lee, P., Wong, M., 2005. Gender and puberty interact on the stress-induced activation of parvocellular neurosecretory neurons and corticotropin-releasing hormone messenger ribonucleic acid expression in the rat. *Endocrinology* 146, 137–146.
- Vidal, J., Buwalda, B., Koolhaas, J.M., 2011a. Differential long-term effects of social stress during adolescence on anxiety in Wistar and wild-type rats. *Behav. Process* 87, 176–182.
- Vidal, J., Buwalda, B., Koolhaas, J.M., 2011b. Male Wistar rats are more susceptible to lasting social anxiety than Wild-type Groningen rats following social defeat stress during adolescence. *Behav. Process* 88, 76–80.
- Vidal, J., de Bie, J., Granneman, R.A., Wallinga, A.E., Koolhaas, J.M., Buwalda, B., 2007. Social stress during adolescence in Wistar rats induces social anxiety in adulthood without affecting brain monoaminergic content and activity. *Physiol. Behav.* 92, 824–830.
- Waters, P., McCormick, C.M., 2011. Caveats of chronic exogenous corticosterone treatments in adolescent rats and effects on anxiety-like and depressive behaviour and HPA function. *Biol. Mood Anxiety Disord.* 1, 1–13.
- Watt, M.J., Burke, A.R., Renner, K.J., Forster, G.L., 2009. Adolescent male rats exposed to social defeat exhibit altered anxiety behavioral and limbic monoamines as adults. *Behav. Neurosci.* 123, 564–576.
- Watt, M.J., Roberts, C.L., Scholl, J.L., Meyer, D.L., Müller, L.C., Barr, J.L., Novick, A.M., Forster, G.L., 2014. Decreased prefrontal cortex dopamine activity following adolescent social defeat in male rats: role of dopamine D2 receptors. *Psychopharmacology* 231, 1627–1636.
- Weintraub, A., Singaravelu, J., Bhatnagar, S., 2010. Enduring and sex-specific effects of adolescent social isolation in rats on adult stress reactivity. *Brain Res.* 1343, 83–92.
- Wilkin, M.M., Waters, P., McCormick, C.M., Menard, J.L., 2012. Intermittent physical stress during early- and mid-adolescence differentially alters rats' anxiety- and depression-like behaviours in adulthood. *Behav. Neurosci.* 126, 344–360.

- Willey, A.R., Anderson, R.I., Morales, M., Ramirez, R.L., Spear, L.P., 2012. Effects of ethanol administration on corticosterone levels in adolescent and adult rats. *Alcohol* 46, 29–36.
- Wright, K.M., DiLeo, A., McDannald, M.A., 2015. Early adversity disrupts the adult use of aversive prediction errors to reduce fear in uncertainty. *Front. Behav. Neurosci.* 9, 227.
- Wright, L.D., Hebert, K.E., Perrot-Sinal, T.S., 2008. Periadolescent stress exposure exerts long-term effects on adult stress responding and expression of prefrontal dopamine receptors in male and female rats. *Psychoneuroendocrinology* 33, 130–142.
- Wright, L.D., Muir, K.E., Perrot, T.S., 2012. Enhanced stress responses in adolescent versus adult rats exposed to cues of predation threat, and peer interaction as a predictor of adult defensiveness. *Dev. Psychobiol.* 54, 47–69.
- Wright, L.D., Muir, K.E., Perrot, T.S., 2013. Stress responses of adolescent male and female rats exposed repeatedly to cat odor stimuli, and long-term enhancement of adult defensive behaviors. *Dev. Psychobiol.* 55, 551–557.
- Wulsin, A.C., Wick-Carlson, D., Packard, B.A., Morano, R., Herman, J.P., 2016. Adolescent chronic stress causes hypothalamo–pituitary–adrenocortical hypo-responsiveness and depression-like behavior in adult female rats. *Psychoneuroendocrinology* 65, 109–117.
- Xu, L.Z., Liu, L.J., Yuan, M., Li, S.X., Yue, X.D., Lai, J.L., Lu, L., 2016. Short photoperiod condition increases susceptibility to stress in adolescent male rats. *Behav. Brain Res.* 300, 38–44.
- Zaidan, H., Gaisler-Salomon, I., 2015. Prereproductive stress in adolescent female rats affects behavior and corticosterone levels in second-generation offspring. *Psychoneuroendocrinology* 58, 120–129.
- Zapatero-Caballero, H., Sanchez-Franco, F., Fernandez-Mendez, C., Frutos, M.G.S., Botella-Cubells, L.M., Fernandez-Vazquez, G., 2004. Gonadotropin-releasing hormone receptor gene expression during pubertal development of female rats. *Biol. Reprod.* 70, 348–355.
- Zapatero-Caballero, H., Sanchez-Franco, F., Guerra-Perez, N., Fernandez-Mendez, C., Fernandez-Vazquez, G., 2003. Gonadotropin-releasing hormone receptor gene expression during pubertal development of male rats. *Biol. Reprod.* 68, 1764–1770.
- Zhang, J., Lam, S.P., Kong, A.P., Ma, R.C., Li, S.X., Chan, J.W., Yu, M.W., Zhou, J., Chan, M.H., Ho, C., Li, A.M., Tang, X.Y.W., 2016. Family conflict and lower morning cortisol in adolescents and adults: modulation of puberty. *Sci. Rep.* 6, 22531.
- Zhang, T.Y., Labonté, B., Wen, X.L., Turecki, G., Meaney, M.J., 2013. Epigenetic mechanisms for the early environmental regulation of hippocampal glucocorticoid receptor gene expression in rodents and humans. *Neuropharmacology* 38, 111–123.
- Zhang, W., Rosenkranz, J.A., 2016. Effects of repeated stress on age-dependent GABAergic regulation of the lateral nucleus of the amygdala. *Neuropharmacology* 41, 2309–2323.
- Zitnik, G.A., Curtis, A.L., Wood, S.K., Arner, J., Valentino, R.J., 2015. Adolescent social stress produces an enduring activation of the rat locus coeruleus and alters its coherence with the prefrontal cortex. *Neuropharmacology* 41, 1376–1385.