

Inducing cellular senescence *in vitro* by using genetically encoded photosensitizers

Nadezhda V. Petrova¹, Artem V. Luzhin¹, Ekaterina O. Serebrovskaya², Alina P. Ryumina², Artem K. Velichko¹, Sergey V. Razin^{1,3,4}, Omar L. Kantidze^{1,4}

¹Institute of Gene Biology, Russian Academy of Sciences, 119334 Moscow, Russia

²Shemyakin-Ovchinnikov Institute of Bioorganic Chemistry, Russian Academy of Sciences, 117997 Moscow, Russia

³Department of Molecular Biology, Lomonosov Moscow State University, 119991 Moscow, Russia

⁴LIA 1066 French-Russian Joint Cancer Research Laboratory, 94805 Villejuif, France

Correspondence to: Omar Kantidze; Sergey Razin **email:** kantidze@gmail.com; sergey.v.razin@usa.net

Keywords: KillerRed, miniSOG, DNA damage, optogenetics, cellular senescence

Received: May 18, 2016 **Accepted:** September 29, 2016 **Published:** October 14, 2016

ABSTRACT

Cellular senescence, a form of cell cycle arrest, is one of the cellular responses to different types of exogenous and endogenous damage. The senescence phenotype can be induced *in vitro* by oncogene overexpression and/or DNA damage. Recently, we have reported a novel mechanism of cellular senescence induction by mild genotoxic stress. Specifically, we have shown that the formation of a small number of DNA lesions in normal and cancer cells during S phase leads to cellular senescence-like arrest within the same cell cycle. Here, based on this mechanism, we suggest an approach to remotely induce premature senescence in human cell cultures using short-term light irradiation. We used the genetically encoded photosensitizers, tandem KillerRed and miniSOG, targeted to chromatin by fusion to core histone H2B to induce moderate levels of DNA damage by light in S phase cells. We showed that the cells that express the H2B-fused photosensitizers acquire a senescence phenotype upon illumination with the appropriate light source. Furthermore, we demonstrated that both chromatin-targeted tandem KillerRed (produces O₂⁻) and miniSOG (produces ¹O₂) induce single-stranded DNA breaks upon light illumination. Interestingly, miniSOG was also able to induce double-stranded DNA breaks.

INTRODUCTION

Cellular senescence is growth arrest characterized by complex phenotypic changes and loss of re-proliferative potential [1, 2]. To study cellular senescence, one can transfect cells with a plasmid that encodes a constitutively active oncogene, thus inducing the so-called oncogene-induced senescence (OIS) [3, 4]. It is thought that in course of OIS cells, which are cell cycle-arrested due to oncogene-induced DNA damage, undergo geroconversion stimulated by activation of mTOR pathway [1, 5]. Mechanistically, oncogene-induced DNA damage may result from DNA hyper-replication [6], replication fork reversal [7], the depletion of nucleotide pools [8] and/or increased levels of reactive oxygen species (ROS) [9]. OIS can be stimulated in a DNA damage response (DDR)-independent manner as well, specifically through the

activation of CDKN2A genomic loci that code for p16^{INK4a} and ARF [10, 11]. Obviously, the expression of activated oncogenes induces an excessively complex composition of senescence-inducing stimuli, making OIS a difficult-to-interpret model [4, 12]. Cellular senescence can be induced by sublethal concentrations of DNA-damaging agents as well; however, in this case, an extremely long incubation time (from hours to days) is usually needed [13]. It is also worth noting that low molecular weight compounds may act in many (sometimes unexpected) ways. For instance, a topoisomerase II inhibitor, doxorubicin, which is known to induce premature senescence, stimulates ROS overproduction and histone eviction, as well as topoisomerase-dependent DNA damage [14-16]. Thus, it is reasonable to develop a simple and versatile method to induce cellular senescence in normal and cancer cell lines.

Here, we propose a novel approach to induce cellular senescence *in vitro* using the genetically encoded photosensitizers tandem KillerRed (tKR, a modified version of KillerRed [17, 18]) and miniSOG (mini singlet oxygen generator) [19, 20]). It is generally thought that upon illumination, photosensitizers produce ROS by a Type I (superoxide anion radicals, hydrogen peroxides and hydroxyl radicals) or Type II (singlet oxygen) photosensitization reaction [21]. Irradiation of the photosensitizers can induce different types of cell death (apoptosis, necrosis, or autophagy), depending on the cellular compartment to which they are targeted [21]. There are only a few genetically encoded photosensitizers [18, 19, 22]; among them, KR is the first developed and the best-studied one, and miniSOG is the most promising, due to its enhanced efficiency of ROS production and relatively low molecular weight [23]. It was proven that KR activation results in superoxide anion radical production (O_2^-), while miniSOG activation results in singlet oxygen production (1O_2) [23-26]. The cell killing induced by photosensitizers usually depends on their ROS-mediated proteotoxic effects [21]. However, it has recently been shown that KR activation in the cell nucleus can lead to DNA damage and may be used for light-induced inhibition of cell division [17] and for localized induction of oxidative DNA lesions [27]. Nevertheless, it is still debatable whether O_2^- and 1O_2 can induce single- and/or double-stranded DNA breaks (SSBs and DSBs), as well as the oxidation of DNA bases [28, 29]. In the

present study, we analyzed the DNA-damaging effects of tKR and miniSOG, which were targeted to the cell nucleus by fusing them to the core histone H2B. We found that although both photosensitizers effectively stimulate SSB formation upon light irradiation, miniSOG can additionally induce a small number of DSBs. Based on our recently reported mechanism of mild genotoxic stress-dependent cellular senescence [30], we employed these genetically encoded photosensitizers to induce a cellular senescence-like state.

RESULTS AND DISCUSSION

Overview of the approach

Recently, we reported a novel mechanism of cellular senescence induction by mild genotoxic stress [30, 31]. Specifically, we showed that formation of a small number of DNA lesions in normal and cancer cells during S phase leads to cellular senescence-like arrest within the same cell cycle. The mechanism of this arrest includes DNA strand breaking in S-phase cells, the collision of replication forks with the breaks, and the formation of difficult-to-repair DSBs [30]. Subsequently, persistent DDR results in proliferation arrest with a cellular senescence phenotype (Figure 1A). This mechanism is noteworthy because it utilizes extremely low concentrations of DNA-damaging agents (e.g., nanomolar concentrations of camptothecin was applied to the cells for 30-60 minutes) to induce cellular




Figure 1. Overview of the method used to optogenetically induce cellular senescence in vitro. (A) Model illustrating how mild genotoxic stress can induce cellular senescence-like proliferation arrest (according to [30]). (B) Overview of the method for inducing cellular senescence using the genetically encoded photosensitizers tandem KillerRed (tKR) and miniSOG that were targeted to chromatin.

senescence [30]. Based on this mechanism, we developed an approach to remotely induce premature senescence in human cell cultures using short-term light irradiation. We employed the genetically encoded photosensitizers tKR and miniSOG and targeted them to chromatin to induce DNA lesions, which, in turn, induced difficult-to-repair DSBs, persistent DDR, and, subsequently, the development of the cellular senescence phenotype (Figure 1B). Briefly, the procedure used to induce cellular senescence includes the following steps: i) establishment of the cell line that transiently or stably expresses either tKR or miniSOG fused to core histone H2B to direct them to chromatin, ii) synchronization of the cells in S phase, and iii) light illumination of the cells (Figure 1B). In the current study, we generally used human HeLa Kyoto cell lines that stably express H2B-tKR or H2B-miniSOG, which were previously established by lentiviral transduction with corresponding constructs [17, 32]. We should mention that the expression levels of H2B-tKR (in contrast to H2B-miniSOG) fusion protein vary significantly across the population of stably transfected cells. This follows from a high cellular heterogeneity of H2B-tKR fusion protein fluorescence (data not shown). This observation is in agreement with the results of quantitative RT-PCR showing that the expression of H2B-miniSOG in the HeLa Kyoto cell line, which stably expresses this fusion protein, is approximately sevenfold higher than the expression of H2B-tKR in the corresponding cell line (data not shown). It was shown earlier that the fusion of tKR or miniSOG to core histone H2B effectively targets them to chromatin but does not induce cell killing or cell cycle alterations until the cells are illuminated with a specific wavelength of light [17, 19]. To synchronize the cells in S phase, we performed a double thymidine block, although one can use any other synchronization technique. Moreover, an asynchronous cell population can also be used; in this case, only the S-phase cells will senesce. To activate H2B-miniSOG or H2B-tKR, we illuminated the corresponding cell lines with blue light (465/95 nm, 65 mW/cm²) for 5 minutes or with green light (540/80 nm, 200 mW/cm²) for 15 minutes, respectively. As it will be shown below such illumination conditions were sufficient to induce cellular senescence but did not lead to cell killing.

DNA damage induced by miniSOG and tKR

First, we analyzed whether genetically encoded photosensitizers targeted to chromatin could induce DNA strand breaks. Photosensitizers produce different ROS upon light-induced activation: O₂⁻, hydrogen peroxides and hydroxyl radicals by the Type I photosensitization reaction and ¹O₂ by the Type II reaction. It is generally thought that these ROS can induce DNA damage. Indeed, it is quite well defined

that hydrogen peroxide (H₂O₂) and hydroxyl (⁻OH) stimulate DNA strand breaks, along with base and sugar oxidation [33]. It is much more complicated to determine the DNA-damaging effects of O₂⁻ and ¹O₂. Although O₂⁻ does not interact with undamaged DNA, it was shown that it could react with oxidatively generated DNA base radicals [34, 35]. Furthermore, O₂⁻ can give rise to H₂O₂ and ⁻OH radicals via a two-stage reaction [28]. ¹O₂ specifically reacts with guanines, thus producing 8-oxoguanines in DNA; however, the question of whether ¹O₂ can induce DNA strand breaks is still open [36, 37]. It should also be mentioned that the DNA-damaging effects of O₂⁻ and ¹O₂ have been studied *in vitro* using free DNA in aqueous solutions. Therefore, it is questionable whether these ROS react with chromatin in living cells in a similar fashion. It is unclear what type of DNA damage can be induced by the genetically encoded photosensitizers, such as miniSOG and tKR. It had been only reported that the chromatin-targeted tKR could oxidize DNA bases [17, 27]. It is established that light-illumination of miniSOG leads to ¹O₂ production [19, 26], but the activation of tKR predominantly results in the formation of O₂⁻; however, the possibility that tKR also produces ¹O₂ was not fully excluded [24, 25, 38]. Here, we investigated whether the chromatin-targeted miniSOG and tKR could induce DNA strand breaks upon activation by light illumination. For this purpose, we used the single-cell gel electrophoresis (SCGE) technique, also known as the “comet assay” [39]. The tail moment, the most meaningful parameter of the comet, which represents the tail length multiplied by the fraction of DNA in the tail, was chosen as a criterion for the degree of DNA breakage.

First, we analyzed SSB generation in asynchronous HeLa cells expressing either H2B-miniSOG or H2B-tKR that were illuminated with the corresponding light source. We used an alkaline modification of the comet assay to perform this analysis. As a positive control for the presence of SSB, we used H₂O₂-treated cells. Blue- or green-light irradiation by itself did not induce any detectable DNA damage in control cells (Figure 2A). However, both chromatin-targeted miniSOG and tKR induced a significant number of SSBs (Figure 2A). As expected, miniSOG known to produce ¹O₂ [19, 26], had a much more pronounced DNA-damaging effect than tKR; it was comparable to the effects of high concentrations of H₂O₂ (200 μM, 1 h). However, this may also be due to several-fold higher level of expression of H2B-miniSOG fusion relative to H2B-tKR fusion (data not shown). It is interesting that although a significant portion of the SSBs induced by miniSOG were repaired within 30 minutes after illumination (Figure 2A, “BL+rec”), the tKR-induced lesions remained unresolved at this time point (Figure

2A, “GL+rec”). The mechanism of tKR-dependent SSB formation is elusive; the only possible (known) way for O_2^- to induce DNA strand breaks is to be converted into H_2O_2 and $^{\cdot}OH$ in a two-step reaction utilizing superoxide dismutase (SOD) and active metal ions [28]. However, the presence of SOD in the nuclei of the untreated cells is controversial – it was recently shown that Sod1 was only translocated from the cytoplasm to the nucleus only upon oxidative stress (treatment of cells with H_2O_2) [40]. It may be that tKR still generates a number of 1O_2 that is responsible for SSB generation in this case.

We next analyzed DSB generation in asynchronous HeLa

cells expressing either H2B-miniSOG or H2B-tKR. For this purpose, we used the neutral comet assay; as a positive control for the presence of DSBs, we used cells that had been treated with the topoisomerase II poison etoposide (VP16; 10 μ g/ml, 1 hr). Similar to SSBs, DSBs were not induced in response to illumination in HeLa cells that did not express the photosensitizers (Figure 2B). DSBs were only generated in HeLa cells expressing H2B-miniSOG that were illuminated with blue light (Figure 2B). Collectively, these results suggest that although both chromatin-targeted photosensitizers effectively stimulate SSB formation upon light irradiation, miniSOG can also produce DSBs.




Figure 2. DNA damage induced by the activation of miniSOG and tKR targeted to chromatin. (A-B) Asynchronous H2B-miniSOG expressing HeLa cells, along with their non-expressing counterparts, were either blue-light irradiated (“BL”; 465–495 nm, 65 mW/cm², 5 min) or light irradiated and recovered for 30 min (“BL+rec”). Asynchronous H2B-tKR expressing HeLa cells, along with their non-expressing counterparts, were either green-light irradiated (“GL”; 540–580 nm, 200 mW/cm², 15 min) or light irradiated and recovered for 30 min (“GL+rec”). Alkaline (A) and neutral (B) comet assays were performed. Non-illuminated cells were used (“control”) as a negative control and cells treated with H_2O_2 (“H2O2”; 200 μ M, 1 hr) were used as a positive control in the alkaline comet assay (A), and cells treated with the topoisomerase II poison etoposide (“VP16”; 10 μ g/ml, 1 hr) were used as a positive control in the neutral comet assay (B). Box plots show the tail moments. The boxed region represents the middle 50% of the tail moments, the horizontal lines represent the medians, and the black crosses indicate the means. * $P < 0.0001$ (two-tailed t -test, $n > 70$), # $P < 0.0001$ (two-tailed t -test, $n > 150$), n.s. – not significant. The results of one of four experiments are shown.




Figure 3. Activated genetically encoded photosensitizers can induce cellular senescence. (A-B) The HeLa Kyoto cell line and its derivatives expressing either H2B-miniSOG or H2B-tKR were synchronized in S phase, illuminated with blue (465-495 nm, 65 mW/cm², 5 min) or green (540-580 nm, 200 mW/cm², 15 min) light, allowed to recover for 48 hr, and stained for γ H2AX (A) or SA- β -gal (B). Control represents the cells that were synchronized and released for 48 hr (non-illuminated). The DNA was stained with DAPI in (A). Scale bar: 50 μ m. (C) Senescent HeLa cells stained for SA- β -gal. Cellular senescence was induced by treatment of S-phase HeLa cells with a DNA topoisomerase I inhibitor camptothecin (1 μ M, 1 h). (D) The HeLa Kyoto cell line and its derivatives expressing either H2B-miniSOG or H2B-tKR were synchronized in S phase, illuminated with corresponding light, allowed to recover for 48 hr, and stained with DAPI. Segmentation of cell nuclei was performed using CellProfiler. Boxplots show nuclear area in each case (*P=0.0001, two-tailed *t*-test).

Light-induced cellular senescence

Cellular senescence may be recognized by the manifestation of several typical signs, including cell and nuclear enlargement, senescence-associated β -galactosidase activity (SA- β -gal), the formation of senescence-associated heterochromatin foci (SAHF) and persistent DDR foci, senescence-associated secretory phenotype (SASP), increased expression of cyclin-dependent kinase inhibitors, etc. [41, 42]. The composition of these markers in a particular cellular senescence state greatly depends on the cell type and on the senescence-inducing stimulus. To test whether light irradiation of the chromatin-targeted photosensitizers induces cellular senescence, we first investigated DDR focus formation. For this purpose, the “parent” HeLa Kyoto cell line and its derivatives that stably express H2B-miniSOG or H2B-tKR were synchronized in S phase, light irradiated (465/95 and 540/80 nm), incubated for 48 hours, and immunostained with an antibody against γ H2AX (phosphorylated at serine-139 variant histone H2AX), which is an ubiquitous DDR marker [43, 44]. It is evident that extensive DDR foci formation was only observed in the photosensitizer-expressing HeLa cells that had been illuminated with the relevant light (blue or green) (Figure 3A and 4A). It should be highlighted that γ H2AX foci were not formed in the non-irradiated cells or in cells that were illumina-

ted with an inappropriate light (blue light for tKR, and green light for miniSOG). This apparently means that the light-irradiation conditions (wavelength, power, and time) are not toxic to the cells by themselves. It is also worth noting that the cells exhibiting extensive γ H2AX staining possessed enlarged nuclei, which is another senescence biomarker (Figure 3D).

To ascertain that the light irradiation-induced state does represent cellular senescence, we assayed the cells that were treated as described above for SA- β -gal activity, the most universal feature of cellular senescence [45]. As expected, only HeLa cells that expressed the photosensitizers and were irradiated with the appropriate light exhibited increased SA- β -gal activity (Figure 3B-C and 4B).

To investigate the temporal kinetics of cellular senescence we assessed the presence of γ H2AX foci in HeLa cells expressing either H2B-miniSOG or H2B-tKR that were light irradiated and recovered for different time periods (0, 3, 6 and 24 hours) (Figure 5A-B). We found that γ H2AX foci were formed during the first hours, but not immediately after light-illumination, and did not disappear, thus, forming persistent DDR foci (Figure 5A-B). Western blot analysis of γ H2AX fully confirmed the results obtained using indirect immunofluorescence (Figure 5A-B). Together, these




Figure 4. Locally activated H2B-miniSOG can induce cellular senescence. (A-B) HeLa cells expressing H2B-miniSOG were synchronized in S phase, illuminated with blue (465-495 nm, 65 mW/cm², 5 min) light, allowed to recover for 24 hr, and stained for γ H2AX (A) or SA- β -gal (B). Only part of each specimen was illuminated. Dashed line shows the boundary between illuminated and non-illuminated parts of the specimens. Scale bars: 100 μ m (A) and 80 μ m (B).

observations support our model of delayed replication-dependent induction of difficult-to-repair DSBs [30].

It is well known that the senescence state is maintained by either p16^{INK4A}- or p21^{CIP1}-dependent signaling pathways [41]. In DNA damage-induced cellular senescence, the expression of p21^{CIP1} is usually increased. Using quantitative reverse transcription-PCR (qRT-PCR) and western blotting, we found that p21^{CIP1} but not p16^{INK4A} was upregulated in response to light irradiation of HeLa cells that express either H2B-miniSOG or H2B-tKR (Figure 6A-B). Interestingly, p21^{CIP1} expression was upregulated only after a protracted recovery period (24 h) and not immediately after illumination (Figure 6A-B).

Finally, we decided to test whether the approach proposed is effectively compatible with a transient expression of these genetically encoded photosensitizers. For this purpose, we analyzed the induction of cellular senescence in HeLa cells that were transiently transfected with either H2B-miniSOG or

H2B-tKR constructs, and then illuminated with an appropriate light. The results obtained clearly show that one can use transient transfection with genetically encoded photosensitizers to induce cellular senescence (Supplementary Figure 1).

In summary, we conclude that the HeLa cells that express H2B-fused photosensitizers acquire a senescence phenotype upon illumination with the appropriate light source in early S phase. This phenotype is characterized by nuclear enlargement, the appearance of DDR foci, increased SA- β -gal activity, and the expression of the CDK inhibitor p21^{CIP1} (Figures 3 and 4). Interestingly, premature senescence was induced much more effectively in HeLa cells that expressed H2B-miniSOG. This is apparent from the fact that, in contrast to the H2B-miniSOG-expressing HeLa cells, not all of the H2B-tKR-expressing cells senesce in response to light illumination (Figure 2A and B). This difference may reflect the higher level of H2B-miniSOG expression and increased efficiency of miniSOG in inducing ROS production.




Figure 5. Temporal kinetics of the formation of the persistent DNA damage response foci induced by activation of miniSOG or tKR. (A-B) HeLa cells that express H2B-miniSOG (A) or H2B-tKR (B) were synchronized in S phase, illuminated with blue (465-495 nm, 65 mW/cm², 5 min) or green (540-580 nm, 200 mW/cm², 15 min) light, allowed to recover for the indicated time intervals (0, 3, 6 and 24 hr). Histone γ H2AX was analyzed by indirect immunofluorescence or WB. Negative control represents the cells that were synchronized but not light illuminated; positive control represents the cells treated with DNA topoisomerase II inhibitor etoposide (VP16; 10 μ g/ml, 1 hr). The DNA was stained with DAPI. Scale bar: 20 μ m.




Figure 6. Analysis of the expression of p21 and p16 CDK inhibitors in HeLa cells expressing genetically encoded photosensitizers. HeLa cells that express H2B-miniSOG or H2B-tKR were synchronized in S phase, illuminated with blue (465-495 nm, 65 mW/cm², 5 min) or green (540-580 nm, 200 mW/cm², 15 min) light, allowed to recover for the indicated time intervals (0, 6, 24 and 48 hr), and subjected to gene expression analysis using qRT-PCR and WB. Control (“C”) represents the non-illuminated cells. The expression of p21^{CIP1} and p16^{INK4a} was analyzed using EvaGreen-based qRT-PCR. The amplification levels of the cDNA were normalized to the level of the GAPDH cDNA. The results of one representative experiment are shown. WB was performed with an antibody against p21; GAPDH was used as the loading control.

CONCLUSIONS

Cellular senescence is one of the stress response pathways, in addition to apoptosis, autophagy, necrosis, etc. Although the contribution of cellular senescence to organismal ageing is still debatable [46], it is clear that cellular senescence has well-defined physiological functions [47, 48]. One of the most studied, physiologically relevant roles of senescence is cancer prevention [48]. The senescence phenotype can be induced *in vitro* by oncogene overexpression and/or DNA damage induction. Here, we suggest an approach to remotely activate the cellular senescence program. We used the genetically encoded photosensitizers tKR and miniSOG and targeted them to chromatin by fusing them to the core histone H2B to induce moderate levels of DNA damage using light (Figure 1). To induce senescence, cells expressing these modified photosensitizers should be synchronized in S phase and illuminated with an appropriate light source for a short time. The advantages of our approach are the following: i) the DNA damage is induced remotely without interfering with the cell culture; ii) the resulting DNA damage is adjustable by varying the light irradiation, time, and/or power; and iii) all types of fluorescent microscopes are appropriate for this method. The most useful feature of the approach is the possibility to

remotely induce cellular senescence in a cell population or even in individual cells using light-illumination. This is applicable in studies of cellular senescence using fast-evolving lab-on-a-chip technologies [49], specifically, to investigate cell-to-cell communication or the cellular secretome.

Here, we also shed light on the question of whether genetically encoded photosensitizers can induce DNA damage, particularly DNA strand breaking. We showed that both chromatin-targeted tKR and miniSOG induced SSBs upon light activation; however, only miniSOG was able to induce DSBs (Figure 4). This difference may be related to the fact that miniSOG produces highly active ¹O₂, whereas tKR produces O₂⁻ [19, 25]. Nevertheless, this is the first direct demonstration of DNA strand breaks produced by genetically encoded photosensitizers localized in the cell nucleus, which allows them to be used for DNA damage studies. These findings can stimulate the development of new strategies for using photosensitizers in translational applications. It is also worth noting that we could investigate the effects of different ROS - ¹O₂ and O₂⁻ - on DNA using tKR and miniSOG targeted to chromatin. Nevertheless, the question of whether and how different ROS induce DNA damage is far from being fully understood.

METHODS

Cell culture, synchronization and transfection

HeLa cell line and HeLa Kyoto cell lines that stably express either H2B-tKR or H2B-miniSOG and their non-expressing counterparts were used. The cells were cultured in DMEM (PanEco) supplemented with 10% fetal bovine serum (FBS; HyClone/GE Healthcare), 10 U/ml penicillin (PanEco) and 10 µg/ml streptomycin (PanEco). The cells were grown at 37°C and 5% CO₂ in a conventional humidified CO₂ incubator. For synchronization by double-thymidine block, 30% confluent cells were incubated with 2 mM thymidine for 16 h, released from the block for 9 h, and then incubated with thymidine for an additional 16 h. To release the cells from thymidine, they were washed twice with phosphate-buffered saline (PBS) and incubated in fresh culture medium.

HeLa cells were transfected with 2 µg of either H2B-miniSOG or H2B-tKR construct using Xfect transfection reagent (Clontech) following the manufacturer's instructions. After 24 h, cells were illuminated to induce cellular senescence. The efficiency of transfection was estimated by fluorescence of genetically encoded photosensitizers transfected; usually more than 80% of cells were transfected.

Cell illumination

An inverted Nikon Eclipse Ti-E fluorescence microscope equipped with a Nikon Intensilight C-HGFI light source and a Nikon Plan Fluor 4x/0.13 objective was used to illuminate the cells. A standard filter set was used to acquire the relevant fluorescence signals and to illuminate the cells. The FITC filter set was used for blue-light illumination (excitation band pass (BP) 465/95 nm, emission BP 515/55 nm), and the Tx Red filter set was used for green-light illumination (excitation BP 540/80 nm, emission BP 600/60 nm). A Laser Power Meter LP1 (Sanwa) was used to measure the total power of the excitation light. The light power density (W/cm²) was estimated by dividing the total power by the area of the illuminated region. The cells stably expressing H2B-miniSOG were illuminated for 5 min with blue light (65 mW/cm² light power density), and the H2B-tKR expressing cells were illuminated for 15 min with green light (200 mW/cm² light power density).

Indirect immunofluorescence

Cells were grown on microscope slides, illuminated, fixed, and permeabilized in CSK buffer (10 mM PIPES, pH 7.0, 100 mM NaCl, 1.5 mM MgCl₂, and 300 mM

sucrose) supplemented with 1% PFA and 2.5% Triton X-100 for 15 min at room temperature. The fixed cells were washed three times for 5 min in PBS. After washing, the cells were preincubated with 1% BSA in PBS for 30 min and then incubated with a primary antibody against γH2AX (Upstate/Millipore, #05-636) in PBS supplemented with 1% BSA for 1 h at room temperature. The cells were washed three times for 5 min with PBS. To visualize the primary antibodies, the samples were incubated with Alexa Fluor 488- or Alexa Fluor 594-conjugated secondary antibodies (Molecular Probes/Life Technologies). The DNA was counterstained with the fluorescent dye 4,6-diamino-2-phenylindole (DAPI) in PBS for 5 min at room temperature. After washing in PBS and distilled water, the samples were mounted using Dako fluorescent mounting medium (Life Technologies). The immunostained samples were analyzed using a Zeiss AxioScope A.1 fluorescence microscope (objective: Zeiss N-Achroplan 40x/0.65; camera: Zeiss AxioCam MRm; acquisition software: Zeiss AxioVision Rel. 4.8.2; Jena, Germany). The images were processed using ImageJ software (version 1.44) and Adobe Photoshop CS6.

SA-β-galactosidase assay

The cells were fixed in 2% formaldehyde and 0.2% glutaraldehyde in PBS for 5 min at room temperature. Then, the cells were washed three times in PBS for 5 min and incubated in staining solution (1 mg/ml 5-bromo-4-chloro-3-indolyl β-D-galactopyranoside (X-gal; Sigma-Aldrich), 40 mM citric acid/sodium phosphate buffer, pH 6.0, 150 mM NaCl, 2 mM MgCl₂, 5 mM K₃Fe(CN)₆, and 5 mM K₄Fe(CN)₆) for 18 h at 37°C. Then, the samples were washed in PBS for 5 min and fixed with methanol for 5 min at room temperature.

Neutral comet assay

After illumination, the cells were immersed in Versen solution and incubated at 37°C for 5 min. The cell suspension was then mixed in a 1:1 ratio with a Trevigen LMAgarose (#4250-050-02) at 37°C. The mixture was pipetted onto comet slides (Trevigen, #3950-300-02) that had been precoated with a 1% normal melting point agarose base layer. The drop containing the cells was covered with a glass cover slip and incubated at 4°C for 5 min. The cover slips were removed, and the slides were then immersed in lysis solution (30 mM EDTA, 0.5% SDS, and 10 mM Tris-HCl, pH 8.0, supplemented with 200 mg/ml proteinase K (Sigma-Aldrich)) at 37°C for 1 h. After lysis, the slides were washed three times for 5 min in PBS and incubated in 1x TBE for 20 min at 4°C. Electrophoresis was performed in a Trevigen electrophoresis system (#4250-050-ES) for 20 min at 4°C and 1 V/cm in 1x

TBE. After washing with PBS, the slides were stained with a 1:3000 dilution of SYBR Green (Thermo Scientific, #S7563). The comets were visualized at 4x magnification using an inverted Nikon Eclipse Ti-E fluorescence microscope equipped with a Nikon Intensilight C-HGFI light source (objective: Nikon Plan Fluor 4x/0.13; camera: DS-Qi2). The images of the comets were analyzed with the CometScore software. Statistical analysis was performed using IBM SPSS Statistics 20.

Alkaline comet assay

The preparation of the samples and slides was similar to the neutral comet assay. The slides were immersed in lysis solution (10 mM Tris-base, 2.5 M NaCl, 100 mM EDTA, 1% Triton X-100, and 10% DMSO, pH 10.0) for 1 h at 4°C in the dark. Then, the slides were immersed in unwinding buffer (300 mM NaOH and 1 mM EDTA, pH 13.0) for 20 min at 4°C. The slides were subjected to electrophoresis in a Trevigen electrophoresis system (#4250-050-ES) for 20 min at 4°C and 0.7 V/cm in unwinding buffer. After electrophoresis, the slides were neutralized with 500 mM Tris-HCl, pH 8.0, and stained with a 1:3000 dilution of SYBR Green (Thermo Scientific, #S7563). The comets were visualized at 4x magnification using an inverted Nikon Eclipse Ti-E fluorescence microscope equipped with a Nikon Intensilight C-HGFI light source (objective: Nikon Plan Fluor 4x/0.13; camera: DS-Qi2). The images of the comets were analyzed with the CometScore software. Statistical analysis was performed using IBM SPSS Statistics 20.

RNA isolation and reverse transcription quantitative PCR

The total RNA was extracted from the cells using TRIzol reagent (Life Technologies), and cDNA synthesis was performed at 42°C for 1 h using 0.5 µg of the total RNA as a template, 0.4 µg of random hexamer primers and 200 U of reverse transcriptase (Fermentas) in the presence of 20 U of ribonuclease inhibitor (Fermentas). The obtained cDNAs were analyzed by quantitative polymerase chain reaction (PCR) using the CFX96 real-time PCR detection system (Bio-Rad Laboratories). Each reaction contained 50 mM Tris-HCl, pH 8.6, 50 mM KCl, 1.5 mM MgCl₂, 0.1% Tween-20, 0.5 µM each primer, 0.2 mM dNTPs, 0.6 µM EvaGreen (Biotium), 0.75 U of Hot Start Taq Polymerase (Sibenzyme) and 50 ng of cDNAs. The PCR cycling conditions were as follows: initial denaturation for 5 min at 94°C, 40 cycles of 15 s at 94°C, 30 s at 65°C and 15 s at 72°C. The gene-specific primer pairs were: GAPDH - aaactgtggcgtgatggc and cagtggggacacggaagg; p16 - acaactgccccgccacaac and

acagtgaaggcagaagcggtg; p21 -
aaggcagggggaaggtgggg and gggggaggacagcagcaga;
H2B-tKR - cagccaccacacctacgag and
tgaagccgatgaaggccag; H2B-miniSOG -
gctttgtgattaccgatccgc and atttctgcacggtgcttg.

Whole-cell extracts preparation and immunoblotting

HeLa cells were lysed by incubation in RIPA buffer (150 mM NaCl, 1% Triton X-100, 0.5% sodium deoxycholate, 0.1% SDS, 50 mM Tris-HCl (pH 8.0), 1 mM dithiothreitol, and 1 mM PMSF) supplemented with Protease Inhibitor Cocktail (Sigma-Aldrich) and Phosphatase Inhibitor Cocktail 2 (Sigma-Aldrich) for 30 min on ice. Next, the cell extracts were sonicated with a VirSonic 100 ultrasonic cell disrupter and stored at -70°C. The protein concentration was measured by the Bradford assay. Aliquots of each sample were separated by SDS-PAGE and blotted onto PVDF membranes (Amersham/GE Healthcare). The membranes were blocked for 1 hr in 2% ECL Advance blocking reagent (GE Healthcare) in PBS containing 0.1% Tween 20 (PBS-T) followed by overnight incubation with a primary antibody diluted in PBS-T containing 2% blocking reagent. After three washes with PBS-T, the membranes were incubated for 1 hr with the secondary antibodies (horseradish peroxidase-conjugated anti-rabbit or anti-mouse IgG) in PBS-T containing 2% blocking agent. The immunoblots were visualised using a Pierce ECL plus western blotting substrate.

FUNDING

This work was supported by a Russian Science Foundation (grant number 14-24-00022). This study was performed using the equipment of the Shared Resource Facility «Live cell biology and biomedical nanotransporters for drug delivery» of the Institute of Gene Biology, Russian Academy of Sciences.

CONFLICTS OF INTEREST

The authors have no conflict of interests to declare.

REFERENCE

1. Blagosklonny MV. Cell cycle arrest is not yet senescence, which is not just cell cycle arrest: terminology for TOR-driven aging. *Aging (Albany NY)*. 2012; 4:159–65. doi: 10.18632/aging.100443
2. Blagosklonny MV. Cell cycle arrest is not senescence. *Aging (Albany NY)*. 2011; 3:94–101. doi: 10.18632/aging.100281
3. Serrano M, Lin AW, McCurrach ME, Beach D, Lowe SW. Oncogenic ras provokes premature cell

- senescence associated with accumulation of p53 and p16INK4a. *Cell*. 1997; 88:593–602. doi: 10.1016/S0092-8674(00)81902-9
4. Bianchi-Smiraglia A, Nikiforov MA. Controversial aspects of oncogene-induced senescence. *Cell Cycle*. 2012; 11:4147–51. doi: 10.4161/cc.22589
 5. Leontieva OV, Blagosklonny MV. Tumor promoter-induced cellular senescence: cell cycle arrest followed by geroconversion. *Oncotarget*. 2014; 5:12715–27. doi: 10.18632/oncotarget.3011
 6. Di Micco R, Fumagalli M, Cicalese A, Piccinin S, Gasparini P, Luise C, Schurra C, Garre' M, Nuciforo PG, Bensimon A, Maestro R, Pelicci PG, d'Adda di Fagagna F. Oncogene-induced senescence is a DNA damage response triggered by DNA hyper-replication. *Nature*. 2006; 444:638–42. doi: 10.1038/nature05327
 7. Neelsen KJ, Zanini IM, Herrador R, Lopes M. Oncogenes induce genotoxic stress by mitotic processing of unusual replication intermediates. *J Cell Biol*. 2013; 200:699–708. doi: 10.1083/jcb.201212058
 8. Mannava S, Moparthy KC, Wheeler LJ, Natarajan V, Zucker SN, Fink EE, Im M, Flanagan S, Burhans WC, Zeitouni NC, Shewach DS, Mathews CK, Nikiforov MA. Depletion of deoxyribonucleotide pools is an endogenous source of DNA damage in cells undergoing oncogene-induced senescence. *Am J Pathol*. 2013; 182:142–51. doi: 10.1016/j.ajpath.2012.09.011
 9. Lee AC, Fenster BE, Ito H, Takeda K, Bae NS, Hirai T, Yu ZX, Ferrans VJ, Howard BH, Finkel T. Ras proteins induce senescence by altering the intracellular levels of reactive oxygen species. *J Biol Chem*. 1999; 274:7936–40. doi: 10.1074/jbc.274.12.7936
 10. Ribeiro JD, Morey L, Mas A, Gutierrez A, Luis NM, Mejetta S, Richly H, Benitah SA, Keyes WM, Di Croce L. ZRF1 controls oncogene-induced senescence through the INK4-ARF locus. *Oncogene*. 2013; 32:2161–68. doi: 10.1038/onc.2012.241
 11. Sreeramaneni R, Chaudhry A, McMahon M, Sherr CJ, Inoue K. Ras-Raf-Arf signaling critically depends on the Dmp1 transcription factor. *Mol Cell Biol*. 2005; 25:220–32. doi: 10.1128/MCB.25.1.220-232.2005
 12. Courtois-Cox S, Jones SL, Cichowski K. Many roads lead to oncogene-induced senescence. *Oncogene*. 2008; 27:2801–09. doi: 10.1038/sj.onc.1210950
 13. Petrova NV, Velichko AK, Razin SV, Kantidze OL. Small molecule compounds that induce cellular senescence. *Aging Cell*. 2016. doi: 10.1111/acel.12518. Epub ahead of print
 14. Kim SY, Kim SJ, Kim BJ, Rah SY, Chung SM, Im MJ, Kim UH. Doxorubicin-induced reactive oxygen species generation and intracellular Ca²⁺ increase are reciprocally modulated in rat cardiomyocytes. *Exp Mol Med*. 2006; 38:535–45. doi: 10.1038/emmm.2006.63
 15. Chang BD, Xuan Y, Broude EV, Zhu H, Schott B, Fang J, Roninson IB. Role of p53 and p21waf1/cip1 in senescence-like terminal proliferation arrest induced in human tumor cells by chemotherapeutic drugs. *Oncogene*. 1999; 18:4808–18. doi: 10.1038/sj.onc.1203078
 16. Pang B, Qiao X, Janssen L, Velds A, Groothuis T, Kerkhoven R, Nieuwland M, Ovaa H, Rottenberg S, van Tellingen O, Janssen J, Huijgens P, Zwart W, Neefjes J. Drug-induced histone eviction from open chromatin contributes to the chemotherapeutic effects of doxorubicin. *Nat Commun*. 2013; 4:1908. doi: 10.1038/ncomms2921
 17. Serebrovskaya EO, Gorodnicheva TV, Ermakova GV, Solovieva EA, Sharonov GV, Zagaynova EV, Chudakov DM, Lukyanov S, Zaraisky AG, Lukyanov KA. Light-induced blockage of cell division with a chromatin-targeted phototoxic fluorescent protein. *Biochem J*. 2011; 435:65–71. doi: 10.1042/BJ20101217
 18. Bulina ME, Chudakov DM, Britanova OV, Yanushevich YG, Staroverov DB, Chepurnykh TV, Merzlyak EM, Shkrob MA, Lukyanov S, Lukyanov KA. A genetically encoded photosensitizer. *Nat Biotechnol*. 2006; 24:95–99. doi: 10.1038/nbt1175
 19. Shu X, Lev-Ram V, Deerinck TJ, Qi Y, Ramko EB, Davidson MW, Jin Y, Ellisman MH, Tsien RY. A genetically encoded tag for correlated light and electron microscopy of intact cells, tissues, and organisms. *PLoS Biol*. 2011; 9:e1001041. doi: 10.1371/journal.pbio.1001041
 20. Souslova EA, Mironova KE, Deyev SM. Applications of genetically encoded photosensitizer miniSOG: from correlative light electron microscopy to immunophotosensitizing. *J Biophotonics*. 2016. doi: 10.1002/jbio.201600120. Epub ahead of print
 21. Buytaert E, Dewaele M, Agostinis P. Molecular effectors of multiple cell death pathways initiated by photodynamic therapy. *Biochim Biophys Acta*. 2007; 1776:86–107. doi: 10.1016/j.bbcan.2007.07.001
 22. Sarkisyan KS, Zlobovskaya OA, Gorbachev DA, Bozhanova NG, Sharonov GV, Staroverov DB, Egorov ES, Ryabova AV, Solntsev KM, Mishin AS, Lukyanov KA. KillerOrange, a Genetically Encoded Photosensitizer Activated by Blue and Green Light. *PLoS One*. 2015; 10:e0145287. doi: 10.1371/journal.pone.0145287
 23. Wojtovich AP, Foster TH. Optogenetic control of ROS production. *Redox Biol*. 2014; 2:368–76.

- doi: 10.1016/j.redox.2014.01.019
24. Vegh RB, Solntsev KM, Kuimova MK, Cho S, Liang Y, Loo BL, Tolbert LM, Bommarius AS. Reactive oxygen species in photochemistry of the red fluorescent protein "Killer Red". *Chem Commun (Camb)*. 2011; 47:4887–89. doi: 10.1039/c0cc05713d
 25. Pletnev S, Gurskaya NG, Pletneva NV, Lukyanov KA, Chudakov DM, Martynov VI, Popov VO, Kovalchuk MV, Wlodawer A, Dauter Z, Pletnev V. Structural basis for phototoxicity of the genetically encoded photosensitizer KillerRed. *J Biol Chem*. 2009; 284:32028–39. doi: 10.1074/jbc.M109.054973
 26. Ruiz-González R, Cortajarena AL, Mejias SH, Agut M, Nonell S, Flors C. Singlet oxygen generation by the genetically encoded tag miniSOG. *J Am Chem Soc*. 2013; 135:9564–67. doi: 10.1021/ja4020524
 27. Lan L, Nakajima S, Wei L, Sun L, Hsieh CL, Sobol RW, Bruchez M, Van Houten B, Yasui A, Levine AS. Novel method for site-specific induction of oxidative DNA damage reveals differences in recruitment of repair proteins to heterochromatin and euchromatin. *Nucleic Acids Res*. 2014; 42:2330–45. doi: 10.1093/nar/gkt1233
 28. Cadet J, Loft S, Olinski R, Evans MD, Bialkowski K, Richard Wagner J, Dedon PC, Møller P, Greenberg MM, Cooke MS. Biologically relevant oxidants and terminology, classification and nomenclature of oxidatively generated damage to nucleobases and 2-deoxyribose in nucleic acids. *Free Radic Res*. 2012; 46:367–81. doi: 10.3109/10715762.2012.659248
 29. Cadet J, Wagner JR. DNA base damage by reactive oxygen species, oxidizing agents, and UV radiation. *Cold Spring Harb Perspect Biol*. 2013; 5:a012559. doi: 10.1101/cshperspect.a012559
 30. Velichko AK, Petrova NV, Razin SV, Kantidze OL. Mechanism of heat stress-induced cellular senescence elucidates the exclusive vulnerability of early S-phase cells to mild genotoxic stress. *Nucleic Acids Res*. 2015; 43:6309–20. doi: 10.1093/nar/gkv573
 31. Petrova NV, Velichko AK, Razin SV, Kantidze OL. Early S-phase cell hypersensitivity to heat stress. *Cell Cycle*. 2016; 15:337–44. doi: 10.1080/15384101.2015.1127477
 32. Ryumina AP, Serebrovskaya EO, Shirmanova MV, Snopova LB, Kuznetsova MM, Turchin IV, Ignatova NI, Klementieva NV, Fradkov AF, Shakhov BE, Zagaynova EV, Lukyanov KA, Lukyanov SA. Flavoprotein miniSOG as a genetically encoded photosensitizer for cancer cells. *Biochim Biophys Acta*. 2013; 1830:5059–67. doi: 10.1016/j.bbagen.2013.07.015
 33. Balasubramanian B, Pogozelski WK, Tullius TD. DNA strand breaking by the hydroxyl radical is governed by the accessible surface areas of the hydrogen atoms of the DNA backbone. *Proc Natl Acad Sci USA*. 1998; 95:9738–43. doi: 10.1073/pnas.95.17.9738
 34. Wagner JR, van Lier JE, Johnston LJ. Quinone sensitized electron transfer photooxidation of nucleic acids: chemistry of thymine and thymidine radical cations in aqueous solution. *Photochem Photobiol*. 1990; 52:333–43. doi: 10.1111/j.1751-1097.1990.tb04189.x
 35. Miaszsek R, Crean C, Joffe A, Geacintov NE, Shafirovich V. Oxidative DNA damage associated with combination of guanine and superoxide radicals and repair mechanisms via radical trapping. *J Biol Chem*. 2004; 279:32106–15. doi: 10.1074/jbc.M313904200
 36. Ravanat JL, Sauvaigo S, Caillat S, Martinez GR, Medeiros MH, Di Mascio P, Favier A, Cadet J. Singlet oxygen-mediated damage to cellular DNA determined by the comet assay associated with DNA repair enzymes. *Biol Chem*. 2004; 385:17–20. doi: 10.1515/BC.2004.003
 37. Cadet J, Ravanat JL, Martinez GR, Medeiros MH, Di Mascio P. Singlet oxygen oxidation of isolated and cellular DNA: product formation and mechanistic insights. *Photochem Photobiol*. 2006; 82:1219–25. doi: 10.1562/2006-06-09-IR-914
 38. Serebrovskaya EO, Edelweiss EF, Stremovskiy OA, Lukyanov KA, Chudakov DM, Deyev SM. Targeting cancer cells by using an antireceptor antibody-photosensitizer fusion protein. *Proc Natl Acad Sci USA*. 2009; 106:9221–25. doi: 10.1073/pnas.0904140106
 39. Olive PL, Banáth JP. The comet assay: a method to measure DNA damage in individual cells. *Nat Protoc*. 2006; 1:23–29. doi: 10.1038/nprot.2006.5
 40. Tsang CK, Liu Y, Thomas J, Zhang Y, Zheng XF. Superoxide dismutase 1 acts as a nuclear transcription factor to regulate oxidative stress resistance. *Nat Commun*. 2014; 5:3446. doi: 10.1038/ncomms4446
 41. Campisi J. Aging, cellular senescence, and cancer. *Annu Rev Physiol*. 2013; 75:685–705. doi: 10.1146/annurev-physiol-030212-183653
 42. Salama R, Sadaie M, Hoare M, Narita M. Cellular senescence and its effector programs. *Genes Dev*. 2014; 28:99–114. doi: 10.1101/gad.235184.113
 43. Paull TT, Rogakou EP, Yamazaki V, Kirchgessner CU, Gellert M, Bonner WM. A critical role for histone H2AX in recruitment of repair factors to nuclear foci after DNA damage. *Curr Biol*. 2000; 10:886–95. doi: 10.1016/S0960-9822(00)00610-2

44. Rogakou EP, Pilch DR, Orr AH, Ivanova VS, Bonner WM. DNA double-stranded breaks induce histone H2AX phosphorylation on serine 139. *J Biol Chem.* 1998; 273:5858–68. doi: 10.1074/jbc.273.10.5858
45. Dimri GP, Lee X, Basile G, Acosta M, Scott G, Roskelley C, Medrano EE, Linskens M, Rubelj I, Pereira-Smith O. A biomarker that identifies senescent human cells in culture and in aging skin in vivo. *Proc Natl Acad Sci USA.* 1995; 92:9363–67. doi: 10.1073/pnas.92.20.9363
46. van Deursen JM. The role of senescent cells in ageing. *Nature.* 2014; 509:439–46. doi: 10.1038/nature13193
47. Muñoz-Espín D, Serrano M. Cellular senescence: from physiology to pathology. *Nat Rev Mol Cell Biol.* 2014; 15:482–96. doi: 10.1038/nrm3823
48. Burton DG, Krizhanovsky V. Physiological and pathological consequences of cellular senescence. *Cell Mol Life Sci.* 2014; 71:4373–86. doi: 10.1007/s00018-014-1691-3
49. Ertl P, Sticker D, Charwat V, Kasper C, Lepperdinger G. Lab-on-a-chip technologies for stem cell analysis. *Trends Biotechnol.* 2014; 32:245–53. doi: 10.1016/j.tibtech.2014.03.004