

Letter to the editor:

CHLOROGENIC ACID AND ITS ROLE IN BIOLOGICAL FUNCTIONS: AN UP TO DATE

Jae Kwang Kim¹, Sang Un Park^{2*}

¹ Division of Life Sciences and Convergence Research Center for Insect Vectors, Incheon National University, Incheon 22012, Korea

² Department of Crop Science, Chungnam National University, 99 Daehak-ro, Yuseong-gu, Daejeon, 34134, Korea

* Corresponding author: E-mail: supark@cnu.ac.kr, Tel.: +82-42-821-5730, Fax: +82-42-822-2631

<http://dx.doi.org/10.17179/excli2019-1404>

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>).

Dear Editor,

Chlorogenic acid (CGA; (1S,3R,4R,5R)-3-[[[(2Z)-3-(3,4-dihydroxyphenyl)prop-2-enyl]oxy]-1,4,5-trihydroxycyclohexanecarboxylic acid) is an ester formed from caffeic acid and quinic acid and works as an intermediate in lignin biosynthesis (Abrankó and Clifford, 2017). CGA, one of the most abundant polyphenol compounds in the human diet, consists of a group of phenolic secondary metabolites isolated from the leaves and fruits of dicotyledonous plants, which is an important component of coffee. CGA has the capacity to manipulate the taste of coffee by modifying astringent, sweet, and sour tastes, which change with the concentration (Tajik et al., 2017).

CGAs are biosynthetically derived from phenylalanine following the phenylpropanoid reaction pathway, which is responsible for the synthesis of several important compounds, like flavonoids, isoflavonoid phytoalexins, coumarins, and lignin (Clifford et al., 2017). There are three possible pathways projected from the p-coumaroyl CoA. Each pathway engages with the same types of enzymatic reactions, such as esterification and hydroxylation (Zhao et al., 2018). These compounds protect plant tissues from damage by oxidative stress, pathogen infection, and wounds. They are also used to mediate animal health (Telles et al., 2017). CGA has a broader range of potential biological properties for health benefits, which might provide non-pharmacological and non-invasive hepatoprotective, antioxidant, anti-diabetic, antimicrobial, anticarcinogenic, anti-inflammatory, and anti-obesity strategies (Maalik et al., 2016; Santana-Gálvez et al., 2017; Tošović et al., 2017; Naveed et al., 2018). Here, we summarize recent literature on the effects of CGAs on different features of health (Table 1).

Table 1: Recent studies of the biological and pharmacological activities of chlorogenic acid

Key findings	Reference
A recent study has shown that CGA can effectively act against Al (III)-stimulated cytotoxicity in RAW264.7 cells through the mitogen-activated protein kinase (MAPK)/Akt-mediated caspase pathways.	Cheng et al., 2019
By suppressing the activation of pro-inflammatory and apoptotic signaling pathways, CGA may be very effective against inflammation and attenuating colitis severity.	Vukelić et al., 2018
CGA, which might be used as a potential therapeutic compound for bovine mastitis, had anti-inflammatory activity by inhibiting nuclear factor kappa beta (NF-κB) activation.	Gao et al., 2018
After treatment with CGA, liver injury and insulin resistance could be relieved by suppressing autophagy through the inactivation of the C-Jun N-terminal kinase pathway in a non-alcoholic fatty liver disease (NAFLD) rat model. Based on this finding, CGA could potentially be used for the treatment of NAFLD.	Yan et al., 2018
Crude extract of <i>Aster koraiensis</i> acted as a potent antiangiogenic agent. The antiangiogenic activity might be attributable to the bioactive component, CGA.	Kim et al., 2018
CGA, when treated with Regorafenib, improved the anti-cytotoxicity action of Regorafenib in hepatocarcinoma (HCC) cells. This combined treatment as a drug might be a safe and more effective option in HCC therapy.	Refolo et al., 2018
Continuously taking CGA for a period of 6-months improved attention, executive, and memory functions in the elderly who complained of subjective memory loss.	Kato et al., 2018
CGAs improve quality of coffee beverages by decreasing the concentration of 5-O-CGA, while maintain the aroma and taste profile. This might be useful to people with sensitive stomachs.	Siebert et al., 2018
CGA has a significant antidepressant response in sleep-deprived rats. The mechanism of action of CGA may involve the regulation of abnormal pathways, including metabolism of nicotinate and nicotinamide; metabolism of glyoxylate and dicarboxylate; metabolism of glycine, serine, and threonine; and metabolism of arginine and proline.	Ma et al., 2018
CGA stimulates bacterial apoptosis in <i>E. coli</i> and the depletion of intracellular ROS (reactive oxygen species), which might regulate the progression of bacterial apoptosis-like death.	Lee and Lee, 2018
Dietary supplementation with CGA enhances the growth performance and reduces the diarrhea incidences of the weaned pigs. It is suggested that CGA improved the antioxidant capacity and enhanced the intestinal digestion and absorption function.	Chen et al., 2018
Semen was effectively stored in 3.2 mg/mL of CGA at 15 °C for up to 24 hr. From this finding, it can be assumed that semen could be stored for longer durations if the CGA concentration increased (6.0 mg/ml or more).	Pereira et al., 2018
CGA, combined with phospholipid complex (PC), improved the post-myocardial infarction inflammatory response in aging hearts, making PC a promising candidate for the clinical development of CGA.	Li et al., 2018
CGA assuaged hepatotoxicity and cholestasis by reducing the uptake and synthesis of bilirubin and bile acids, and finally enhancing the metabolism and efflux of bilirubin and bile acids.	Zhu et al., 2018
By increasing cyclic adenosine monophosphate (cAMP) levels via adenylyl cyclase activation, CGA significantly reduced the carbachol-induced contractions of the mouse urinary bladder.	Kaneda et al., 2018
CGA reduces human lens epithelial cells (hLECs) apoptosis and protects lenses from opacity induced by H ₂ O ₂ via the BCL2 (B-cell lymphoma 2) associated X apoptosis regulator (Bax)/ BCL2 apoptosis regulator (Bcl-2) signaling pathway. CGA shows effective defenses against oxidative stress, which could treat a variety of diseases in clinical practice.	Song et al., 2018

Key findings	Reference
CGA treatment could inhibit the retinal neoangiogenesis during diabetic retinopathy by nullifying high glucose (HG)-induced hypoxia-inducible factor 1-alpha (HIF-1 α)-mediated paracrine vascular endothelial growth factor (VEGF) expression in microglia cells and reducing VEGF-induced angiogenesis in retinal endothelial cells.	Mei et al., 2018
CGA is considered as a therapeutic agent for the treatment of diabetic nephropathy, showing antioxidant and anti-inflammatory effects that are activated via the modulation of the nuclear factor erythroid-derived 2-related factor 2 (Nrf2)/heme oxygenase-1 (HO-1) and NF- κ B pathways.	Bao et al., 2018
CGA prevents acetaminophen (APAP)-induced hepatotoxicity through the Nrf2 antioxidative signaling pathway by blocking the binding of Nrf2 to its deterrence protein, Keap1. Extracellular regulated protein kinase (ERK) 1/2 play a vital role in regulating CGA-induced Nrf2 transcriptional activation. CGA could be utilized as a promising therapeutic agent for the detoxification of APAP-induced hepatotoxicity.	Wei et al., 2018
CGA influences the expression of apoptosis genes related to the oxidative stress and p38 MAP-dependent pathways and the expression of genes encoding stem cell markers. CGA may show polyphenolic anti-cancer effects through the consumption of vegetables and fruits.	Yamagata et al., 2018
By reducing the apoptosis of neuronal cells, CGA mitigates the neuronal damage induced in alcohol exposed neonatal rat.	Guo and Li, 2017
CGA has the potential to relieve liver fibrosis through the miR-21-regulated TGF- β 1/Smad7 signaling pathway, which suggests that CGA could act as a new anti-fibrosis agent against liver fibrosis.	Yang et al., 2017
By administering appropriate amounts of myricetin-3-O- β -rhamnoside and CGA synergistically, the fibroblastic and remodeling phases of wound repair are accelerated.	Moghadam et al., 2017
CGA controls several mechanisms of action for the development of highly valuable therapeutics for metabolic diseases, such as type 2 diabetes.	Sanchez et al., 2017
CGA shows antifungal activity against phytopathogenic fungi relevant to horticulture and agriculture, suggesting potential biofungicides activity in CGA-enriched wastes and by-products.	Martínez et al., 2017
CGA can modify the gut microbial community structure in the intestines, exhibiting lower intestinal and systemic inflammation, and also enhancing the course of the dextran sulfate sodium-induced colitis, which is linked to a proportional increase in <i>Akkermansia</i> .	Zhang et al., 2017
CGA has similar responses to caffeic acid. <i>In vitro</i> maturation (IVM) with 50 μ M of CGA is particularly useful to the <i>in vitro</i> production (IVP) of porcine embryos; 50 μ M of CGA prevents oocytes from DNA damage induced by oxidative stress. The addition of CGA to the maturation medium has the potential to enhance the porcine IVP system.	Nguyen et al., 2017
CGA prevents the interleukin (IL)-1 β -induced inflammatory response, through the NF- κ B signaling pathway. CGA could be a suitable candidate for the treatment of osteoarthritis.	Liu et al., 2017
CGA prevents focal cerebral ischemia reperfusion injury in a rat model by maintaining the inflammatory factor, hypoxia factor, and nerve growth factor.	Miao et al., 2017
CGA prevents the progression of hepatocellular carcinoma via multiple pathways. CGA could be an effective chemopreventive agent for hepatocellular carcinoma.	Yan et al., 2017
CGA restrained benign prostatic hyperplasia in model animals, and this mechanism could protect type II 5-alpha reductase activity.	Huang et al., 2017
Consumption of CGA for a period of 5 days significantly enhances fat oxidation during sleep, indicating that beverages containing CGA may be useful to lessen body fat and protect from obesity. Taking CGA also minimizes sleep latency without any adverse effects on sleep quality.	Park et al., 2017

Key findings	Reference
CGA can be used as a neuraminidase blocker to prevent influenza A virus both in cellular and animal models, revealing a potential utility for the treatment of the influenza virus infection.	Ding et al., 2017
A self-microemulsifying drug delivery system (SMEDDS) promotes the oral bio-availability of CGA by enhancing its absorption and slowing the metabolism of absorbed CGA by changing its distribution from the liver to the kidney. SMEDDS is an increasingly utilized carrier for the oral delivery of CGA.	Chen et al., 2017
CGA significantly protects mice from Con A-induced hepatitis by reducing the activation of Toll-like receptor (TLR) 4 signaling. This response lowers the expression of adhesion molecules and enhances the infiltration and activation of hepatic leukocytes and the production of pro-inflammatory cytokines.	Yuan et al., 2017
CGA reduces glioma growth by enhancing M1-polarized macrophages and protecting M2 phenotypic macrophages as a potential therapeutic approach.	Xue et al., 2017
CGA promotes potassium efflux, resulting in an apoptotic volume decrease (AVD) and G2/M cell cycle arrest in <i>Candida albicans</i> . Thus, potassium efflux via potassium channels improves CGA-induced apoptosis, stimulating several apoptotic processes.	Yun and Lee, 2017
CGA prevents Iso-induced vascular smooth muscle cells (VSMC) damage through the suppression of ROS generation. This finding suggests that CGA could be a novel treatment for vascular diseases.	Wang et al., 2016
CGA prevents both α -naphthylisothiocyanate-induced intrahepatic cholestasis and liver injury. This protective mechanism is modulated through the down-regulation of signal transducer and activator of transcription 3 (STAT3) and NF- κ B signaling.	Tan et al., 2016
CGA mitigates the oxidative stress in streptozocin-induced diabetic nephropathy (DN) rats. Its molecular mechanism could prevent the endoplasmic reticulum-stress response in DN.	Ye et al., 2016
CGA is a novel therapeutic agent for preventing inflammatory hyperplasia of the synovium by enhancing synoviocyte apoptosis in patients with rheumatoid arthritis.	Lou et al., 2016
CGA significantly increases hepatic lipid dysregulation in rats by activating fatty acid metabolism enzymes, enhancing AMP-activated protein kinase activation, and modulating levels of hepatic fatty acids.	Zhou et al., 2016

Acknowledgements

This work was supported by a grant from the Next-Generation BioGreen 21 Program (SSAC, Project #. PJ013328)" Rural Development Administration, Republic of Korea.

Conflict of interest

The authors declare no conflict of interest.

REFERENCES

Abrankó L, Clifford MN. An unambiguous nomenclature for the acyl-quinic acids commonly known as chlorogenic acids. *J Agric Food Chem*. 2017;65:3602-8.

Bao L, Li J, Zha D, Zhang L, Gao P, Yao T, et al. Chlorogenic acid prevents diabetic nephropathy by inhibiting oxidative stress and inflammation through modulation of the Nrf2/HO-1 and NF- κ B pathways. *Int Immunopharmacol*. 2018;54:245-53.

Chen J, Li Y, Yu B, Chen D, Mao X, Zheng P, et al. Dietary chlorogenic acid improves growth performance of weaned pigs through maintaining antioxidant capacity and intestinal digestion and absorption function. *J Anim Sci*. 2018;96:1108-18.

Chen L, Liu CS, Chen QZ, Wang S, Xiong YA, Jing J, et al. Characterization, pharmacokinetics and tissue distribution of chlorogenic acid-loaded self-microemulsifying drug delivery system. *Eur J Pharm Sci*. 2017;100:102-8.

- Cheng D, Zhang X, Tang J, Kong Y, Wang X, Wang S. Chlorogenic acid protects against aluminum toxicity via MAPK/Akt signaling pathway in murine RAW264.7 macrophages. *J Inorg Biochem.* 2019;190:113-20.
- Clifford MN, Jaganath IB, Ludwig IA, Crozier A. Chlorogenic acids and the acyl-quinic acids: discovery, biosynthesis, bioavailability and bioactivity. *Nat Prod Rep.* 2017;34:1391-421.
- Ding Y, Cao Z, Cao L, Ding G, Wang Z, Xiao W. Antiviral activity of chlorogenic acid against influenza A (H1N1/H3N2) virus and its inhibition of neuraminidase. *Sci Rep.* 2017;7:45723.
- Gao R, Yang H, Jing S, Liu B, Wei M, He P, et al. Protective effect of chlorogenic acid on lipopolysaccharide-induced inflammatory response in dairy mammary epithelial cells. *Microb Pathog.* 2018;124:178-82.
- Guo Z, Li J. Chlorogenic acid prevents alcohol-induced brain damage in neonatal rat. *Transl Neurosci.* 2017;8:176-81.
- Huang Y, Chen H, Zhou X, Wu X, Hu E, Jiang Z. Inhibition effects of chlorogenic acid on benign prostatic hyperplasia in mice. *Eur J Pharmacol.* 2017;809:191-5.
- Kaneda T, Sasaki N, Urakawa N, Shimizu K. Effects of chlorogenic acid on carbachol-induced contraction of mouse urinary bladder. *J Pharmacol Sci.* 2018;136:26-30.
- Kato M, Ochiai R, Kozuma K, Sato H, Katsuragi Y. Effect of chlorogenic acid intake on cognitive function in the elderly: a pilot study. *Evid Based Complement Alternat Med.* 2018;2018:8608497.
- Kim J, Lee YM, Jung W, Park SB, Kim CS, Kim JS. *Aster koraiensis* extract and chlorogenic acid inhibit retinal angiogenesis in a mouse model of oxygen-induced retinopathy. *Evid Based Complement Alternat Med.* 2018;2018:6402650.
- Lee B, Lee DG. Depletion of reactive oxygen species induced by chlorogenic acid triggers apoptosis-like death in *Escherichia coli*. *Free Radic Res.* 2018;52:605-15.
- Li Y, Ren X, Lio C, Sun W, Lai K, Liu Y, et al. A chlorogenic acid-phospholipid complex ameliorates post-myocardial infarction inflammatory response mediated by mitochondrial reactive oxygen species in SAMP8 mice. *Pharmacol Res.* 2018;130:110-22.
- Liu CC, Zhang Y, Dai BL, Ma YJ, Zhang Q, Wang Y, et al. Chlorogenic acid prevents inflammatory responses in IL-1 β -stimulated human SW-1353 chondrocytes, a model for osteoarthritis. *Mol Med Rep.* 2017;16:1369-75.
- Lou L, Zhou J, Liu Y, Wei YI, Zhao J, Deng J, et al. Chlorogenic acid induces apoptosis to inhibit inflammatory proliferation of IL-6-induced fibroblast-like synoviocytes through modulating the activation of JAK/STAT and NF- κ B signaling pathways. *Exp Ther Med.* 2016;11:2054-60.
- Ma WN, Zhou MM, Gou XJ, Zhao L, Cen F, Xu Y, et al. Urinary metabolomic study of chlorogenic acid in a rat model of chronic sleep deprivation using gas chromatography-mass spectrometry. *Int J Genomics.* 2018;2018:1361402.
- Maalik A, Bukhari SM, Zaidi A, Shah KH, Khan FA. Chlorogenic acid: A pharmacologically potent molecule. *Acta Pol Pharm.* 2016;73:851-4.
- Martínez G, Regente M, Jacobi S, Del Rio M, Pinedo M, de la Canal L. Chlorogenic acid is a fungicide active against phytopathogenic fungi. *Pestic Biochem Physiol.* 2017;140:30-5.
- Mei X, Zhou L, Zhang T, Lu B, Sheng Y, Ji L. Chlorogenic acid attenuates diabetic retinopathy by reducing VEGF expression and inhibiting VEGF-mediated retinal neoangiogenesis. *Vascul Pharmacol.* 2018;101:29-37.
- Miao M, Cao L, Li R, Fang X, Miao Y. Protective effect of chlorogenic acid on the focal cerebral ischemia reperfusion rat models. *Saudi Pharm J.* 2017;25:556-63.
- Moghadam SE, Ebrahimi SN, Salehi P, Moridi Farihani M, Hamburger M, Jabbarzadeh E. Wound healing potential of chlorogenic acid and myricetin-3-O- β -rhamnoside isolated from *Parrotia persica*. *Molecules.* 2017;22(9):1501.
- Naveed M, Hejazi V, Abbas M, Kamboh AA, Khan GJ, Shumzaid M, et al. Chlorogenic acid (CGA): A pharmacological review and call for further research. *Biomed Pharmacother.* 2018;97:67-74.
- Nguyen TV, Tanihara F, Do L, Sato Y, Taniguchi M, Takagi M, et al. Chlorogenic acid supplementation during in vitro maturation improves maturation, fertilization and developmental competence of porcine oocytes. *Reprod Domest Anim.* 2017;52:969-75.

- Park I, Ochiai R, Ogata H, Kayaba M, Hari S, Hibi M, et al. Effects of subacute ingestion of chlorogenic acids on sleep architecture and energy metabolism through activity of the autonomic nervous system: a randomised, placebo-controlled, double-blinded cross-over trial. *Br J Nutr.* 2017;117:979-84.
- Pereira BA, Chaves BR, Teles MC, Pontelo TP, Oliveira CR, de Souza RV, et al. Chlorogenic acid improves the quality of boar semen subjected to cooled storage at 15°C. *Andrologia.* 2018;50(5):e12978.
- Refolo MG, Lippolis C, Carella N, Cavallini A, Messa C, D'Alessandro R. Chlorogenic acid improves the regorafenib effects in human hepatocellular carcinoma cells. *Int J Mol Sci.* 2018;19(5):1518.
- Sanchez MB, Miranda-Perez E, Verjan JCG, de Los Angeles Fortis Barrera M, Perez-Ramos J, Alarcon-Aguilar FJ. Potential of the chlorogenic acid as multi-target agent: Insulin-secretagogue and PPAR α/γ dual agonist. *Biomed Pharmacother.* 2017;94:169-75.
- Santana-Gálvez J, Cisneros-Zevallos L, Jacobo-Velázquez DA. Chlorogenic acid: Recent advances on its dual role as a food additive and a nutraceutical against metabolic syndrome. *Molecules.* 2017;22(3):358.
- Siebert M, Berger RG, Nieter A. Enzymatic mitigation of 5-O-chlorogenic acid for an improved digestibility of coffee. *Food Chem.* 2018;258:124-8.
- Song J, Guo D, Bi H. Chlorogenic acid attenuates hydrogen peroxide-induced oxidative stress in lens epithelial cells. *Int J Mol Med.* 2018;41:765-72.
- Tajik N, Tajik M, Mack I, Enck P. The potential effects of chlorogenic acid, the main phenolic components in coffee, on health: a comprehensive review of the literature. *Eur J Nutr.* 2017;56:2215-44.
- Tan Z, Luo M, Yang J, Cheng Y, Huang J, Lu C, et al. Chlorogenic acid inhibits cholestatic liver injury induced by α -naphthylisothiocyanate: involvement of STAT3 and NF κ B signalling regulation. *J Pharm Pharmacol.* 2016;68:1203-13.
- Telles AC, Kupski L, Furlong EB. Phenolic compound in beans as protection against mycotoxins. *Food Chem.* 2017;214:293-9.
- Tošović J, Marković S, Dimitrić Marković JM, Mojović M, Milenković D. Antioxidative mechanisms in chlorogenic acid. *Food Chem.* 2017;237:390-8.
- Vukelić I, Detel D, Pučar LB, Potočnjak I, Buljević S, Domitrović R. Chlorogenic acid ameliorates experimental colitis in mice by suppressing signaling pathways involved in inflammatory response and apoptosis. *Food Chem Toxicol.* 2018;121:140-50.
- Wang J, Li J, Liu J, Xu M, Tong X, Wang J. Chlorogenic acid prevents isoproterenol-induced DNA damage in vascular smooth muscle cells. *Mol Med Rep.* 2016;14:4063-8.
- Wei M, Zheng Z, Shi L, Jin Y, Ji L. Natural polyphenol chlorogenic acid protects against acetaminophen-induced hepatotoxicity by activating ERK/Nrf2 antioxidative pathway. *Toxicol Sci.* 2018;162:99-112.
- Xue N, Zhou Q, Ji M, Jin J, Lai F, Chen J, et al. Chlorogenic acid inhibits glioblastoma growth through repolarizing macrophage from M2 to M1 phenotype. *Sci Rep.* 2017;7:39011.
- Yamagata K, Izawa Y, Onodera D, Tagami M. Chlorogenic acid regulates apoptosis and stem cell marker-related gene expression in A549 human lung cancer cells. *Mol Cell Biochem.* 2018;441:9-19.
- Yan H, Gao YQ, Zhang Y, Wang H, Liu GS, Lei JY. Chlorogenic acid alleviates autophagy and insulin resistance by suppressing JNK pathway in a rat model of nonalcoholic fatty liver disease. *J Biosci.* 2018;43:287-94.
- Yan Y, Liu N, Hou N, Dong L, Li J. Chlorogenic acid inhibits hepatocellular carcinoma in vitro and in vivo. *J Nutr Biochem.* 2017;46:68-73.
- Yang F, Luo L, Zhu ZD, Zhou X, Wang Y, Xue J, et al. Chlorogenic acid inhibits liver fibrosis by blocking the miR-21-regulated TGF- β 1/Smad7 signaling pathway in vitro and in vivo. *Front Pharmacol.* 2017;8:929.
- Ye HY, Li ZY, Zheng Y, Chen Y, Zhou ZH, Jin J. The attenuation of chlorogenic acid on oxidative stress for renal injury in streptozotocin-induced diabetic nephropathy rats. *Arch Pharm Res.* 2016;39:989-97.
- Yuan Y, Gong X, Zhang L, Jiang R, Yang J, Wang B, et al. Chlorogenic acid ameliorated concanavalin A-induced hepatitis by suppression of Toll-like receptor 4 signaling in mice. *Int Immunopharmacol.* 2017;44:97-104.
- Yun J, Lee DG. Role of potassium channels in chlorogenic acid-induced apoptotic volume decrease and cell cycle arrest in *Candida albicans*. *Biochim Biophys Acta.* 2017;1861:585-92.
- Zhang Z, Wu X, Cao S, Cromie M, Shen Y, Feng Y, et al. Chlorogenic acid ameliorates experimental colitis by promoting growth of *Akkermansia* in mice. *Nutrients.* 2017;9(7):677.
- Zhao L, Wang D, Liu J, Yu X, Wang R, Wei Y, et al. Transcriptomic analysis of key genes involved in chlorogenic acid biosynthetic pathway and characterization of MaHCT from *Morus alba* L. *Protein Expr Purif.* 2018;156:25-35.

Zhou Y, Ruan Z, Wen Y, Yang Y, Mi S, Zhou L, et al. Chlorogenic acid from honeysuckle improves hepatic lipid dysregulation and modulates hepatic fatty acid composition in rats with chronic endotoxin infusion. *J Clin Biochem Nutr.* 2016;58:146-55.

Zhu L, Wang L, Cao F, Liu P, Bao H, Yan Y, et al. Modulation of transport and metabolism of bile acids and bilirubin by chlorogenic acid against hepatotoxicity and cholestasis in bile duct ligation rats: involvement of SIRT1-mediated deacetylation of FXR and PGC-1 α . *J Hepatobiliary Pancreat Sci.* 2018;25:195-205.