

ORAL PRESENTATION

Open Access

Local control of non-local information flow in oscillatory neuronal networks

Christoph Kirst^{1,2*}, Marc Timme^{1,2}, Demian Battaglia^{1,2}

From Twentieth Annual Computational Neuroscience Meeting: CNS*2011
Stockholm, Sweden. 23-28 July 2011

Control of information flow between neurons or groups of neurons is essential in a functional brain, e.g. for context and brain state dependent processing. In line with recent experimental and theoretical studies [1-5] we show that phase relations between synchronized oscillatory local circuits or brain areas may dynamically create information channels and induce changes in the effective connectivity.

Reducing neuronal oscillatory dynamics to a phase - amplitude description [6,7], we show how alternative phase shifts between different neurons or groups of neurons result in different effective connectivities. In particular, to quantify the information flow, we analytically calculate the time delayed mutual information and transfer entropy between oscillators in a phase locked state. We further present a theoretical framework to predict phase lag patterns within and between groups of oscillators in hierarchical networks. Combining both results we derive the information flow between the oscillators as a function of structural and dynamical network parameter.

We use our results to reveal how effective connectivity is controlled by the underlying physical connectivity and the intrinsic single oscillation frequencies. Interestingly, we find that local changes in the strength of a single link can remotely control the effective connectivity between two different physically unchanged oscillators. Similarly, local inputs modulating the intrinsic frequencies can dynamically and remotely change the information flow between distal nodes.

We link our results to biophysically more realistic networks of spiking neurons. In a clustered network of groups of type I neurons exhibiting gamma oscillations emanating from a PING mechanism [8], we numerically show that local changes of the connectivity or the inputs strengths

within a cluster can non-locally control the phase relations and the information flow between distant clusters.

Conclusion

Our findings reveal that local changes, e.g. in the physical strength of a *local link* or in the *local frequency* due to variation in the *local inputs*, can *remotely and dynamically control* the direction of *non-local global information flow* between distal nodes/clusters in a network. This might provide an efficient local mechanism to control global information processing in neuronal systems and to account for contextual and attentional modulation.

Author details

¹Max Planck Institute for Dynamics and Self-Organization, Göttingen, 37073, Germany. ²Bernstein Center for Computational Neuroscience, Göttingen, 37073, Germany.

Published: 18 July 2011

References

1. Womelsdorf T, Schöffelen JM, Oostenveld R, Singer W, Desimone R, Engel AK, Andreas K, Fries P: Modulation of Neuronal Interactions Through Neuronal Synchronization. *Science* 2007, 316:1609-1612.
2. Buehmann A, Deco G: Optimal Information Transfer in the Cortex through Synchronization. *PLoS Comput Biol* 2010, 6:e1000934.
3. Besserve M, Murayama Y, Schölkopf B, Logothetis NK, Panyeri S: High frequency phase-spike synchronization of extracellular signals modulates causal interactions in monkey primary visual cortex. *Neuroscience Meeting Planner* Society for Neuroscience; 2010, Online.
4. Fries P: Neuronal Gamma-Band Synchronization as a Fundamental Process in Cortical Computation. *Ann Rev Neurosci* 2009, 32:209-224.
5. Battaglia D, Witt A, Geisel T, Wolf F: Dynamic transitions in the effective connectivity of interacting cortical areas. *FENS 2010 abstract* [http://fens2010.neurosciences.asso.fr/abstracts/f5/a130_4.html].
6. Kuramoto Y: *Chemical Oscillations, Waves and Turbulence*. Springer; 1994.
7. Kawamura Y, Nakao H, Arai K, Kori H, Kuramoto Y: Collective Phase Sensitivity. *Phys Rev Lett* 2008, 101:024101.
8. Börgers C, Kopell N: Effects of Noisy Drive on Rhythms in Networks of Excitatory and Inhibitory Neurons. *Neural Comp* 2005, 3:557-608.

doi:10.1186/1471-2202-12-S1-O15

Cite this article as: Kirst et al.: Local control of non-local information flow in oscillatory neuronal networks. *BMC Neuroscience* 2011 12(Suppl 1): O15.

* Correspondence: ckirst@nld.ds.mpg.de

¹Max Planck Institute for Dynamics and Self-Organization, Göttingen, 37073, Germany

Full list of author information is available at the end of the article