

# *N*-Ethylmaleimide Uncouples Muscarinic Receptors from Acetylcholine-sensitive Potassium Channels in Bullfrog Atrium

T. NAKAJIMA, H. IRISAWA, and W. GILES

From the Departments of Medical Physiology and Medicine, University of Calgary School of Medicine, Calgary, Alberta, Canada T2N 4N1

**ABSTRACT** The effect of *N*-ethylmaleimide (NEM), a sulphhydryl alkylating agent, on the acetylcholine-activated K<sup>+</sup> current,  $I_{K(ACh)}$ , has been studied in single cells from bullfrog atrium using a tight-seal, whole-cell voltage clamp technique. Addition of NEM ( $5 \times 10^{-5}$  M) produced a time-dependent complete block of  $I_{K(ACh)}$ . Dialysis of guanosine-5'-*O*-( $\beta$ -thiotriphosphate) (GTP $\gamma$ S,  $5-10 \times 10^{-4}$  M), a nonhydrolyzable GTP analogue, into the myoplasm from the recording pipette gradually activated  $I_{K(ACh)}$  even in the absence of acetylcholine. This effect is thought to be due to a GTP $\gamma$ S-induced dissociation of GTP-binding proteins ( $G_i$  and/or  $G_o$ ) into subunits that can directly activate these K<sup>+</sup> channels. When NEM ( $5 \times 10^{-5}$  M) was applied after the GTP $\gamma$ S effect had fully developed, it failed to inhibit the GTP $\gamma$ S-induced K<sup>+</sup> current, indicating that the NEM effect is unlikely to be on the dissociated subunits of the GTP-binding protein(s) or on the K<sup>+</sup> channels. In contrast, pretreatment with NEM before GTP $\gamma$ S application markedly reduced the muscarinic K<sup>+</sup> current, suggesting that NEM can block this K<sup>+</sup> current by inhibition of the dissociation of the GTP-binding proteins into functional subunits. In NEM-treated cells the stimulatory effect of isoproterenol on  $I_{Ca}$  was present, but the inhibitory action of ACh on  $I_{Ca}$  was completely abolished. These results demonstrated that NEM can preferentially inhibit muscarinic receptor-effector interactions, probably by alkylating the GTP-binding proteins that are essential for these responses.

## INTRODUCTION

The transmembrane channels through which ionic currents flow are integral membrane proteins that can be modified by direct actions of certain proteolytic and chemical agents (Armstrong et al., 1973; Stimers et al., 1985; Gonoi and Hille, 1987; Duncan et al., 1988; Hescheler and Trautwein, 1988). Some of these channels can also be regulated by various second messengers or intramembrane processes, which in turn can be indirectly modified by chemical and proteolytic agents.

In atrial cells from both amphibian and mammalian hearts, muscarinic acetylcholine (ACh) receptors are linked to a specific population of K<sup>+</sup> channels ( $I_{K(ACh)}$ ) via

Address reprint requests to Dr. W. Giles, Departments of Medical Physiology and Medicine, University of Calgary School of Medicine, Calgary, Alberta, Canada T2N 4N1.

GTP-binding proteins (Breitwieser and Szabo, 1985; Pfaffinger et al., 1985; Kurachi et al., 1986*a, b, c*; Nakajima et al., 1990). Some of the strongest evidence that G proteins are involved comes from experiments using the islet-activating protein, pertussis toxin (IAP), which ADP-ribosylates  $\alpha$ -subunits of GTP-binding proteins ( $G_i$  and/or  $G_o$ ) and selectively blocks this muscarinic response. In cardiac muscle activation of the GTP-binding proteins by agonists (e.g., ACh) or nonhormonal agents (e.g., nonhydrolyzable GTP analogues) results in dissociation of G proteins into subunits ( $\alpha$  and  $\beta\gamma$ ), which then can directly activate, for example,  $K^+$  channels (for review see Brown and Birnbaumer, 1988; Neer and Clapham, 1988). However, IAP also blocks muscarinic receptor-mediated inhibition of adenylate cyclase, thereby reversing the anti- $\beta$ -adrenergic (inhibitory) effect of ACh on the calcium current,  $I_{Ca}$  (Hescheler et al., 1986; Nakajima et al., 1990).

*N*-Ethylmaleimide (NEM), a sulphydryl alkylating agent, can abolish hormone-induced inhibition of adenylate cyclase by alkylating the  $\alpha$ -subunits of GTP-binding proteins, e.g.,  $G_i$  and/or  $G_o$  (Aktories et al., 1982; Jakobs et al., 1982; Ukena et al., 1984; Wong et al., 1985; Asano and Ogasawara, 1986; Winslow et al., 1987). Although it is known that NEM can inhibit the electrophysiological effects of ACh in intact atrial preparations (Roberts and Konjovic, 1969; Doods et al., 1986; Braun and Sperelakis, 1988), the mechanisms of this effect remain unclear.

Our main goal was to learn more about the molecular pharmacology of NEM-induced changes in both the direct and indirect effects of ACh in bullfrog atrium. The direct effects were studied by recording the  $K^+$  current activated by muscarinic receptors,  $I_{K(ACh)}$ . The indirect effects were investigated by measuring the effects of ACh on the calcium current,  $I_{Ca}$ , after it had been enhanced by the  $\beta$ -adrenergic agonist, isoproterenol (ISO). The results show that NEM can preferentially block certain muscarinic receptor-effector interactions, probably by alkylating the G proteins ( $G_i$  and/or  $G_o$ ) that are involved in these direct responses. However, NEM has no effect on the G protein ( $G_s$ ) involved in the indirect muscarinic response, which consists of an inhibition of  $I_{Ca}$ .

#### METHODS

Single cells were isolated from the atria of adult bullfrog hearts by enzymatic dissociation as described previously (Hume and Giles, 1981; Giles and Shibata, 1985; Nakajima et al., 1990*a, b*). The transmembrane ionic currents were recorded using a whole-cell voltage clamp technique (Hamill et al., 1981; Hume and Giles, 1983). Details of microelectrode fabrication and electronic circuits for suction microelectrode voltage clamp have been described (Giles and Shibata, 1985; Clark and Giles, 1987). Membrane currents were monitored on a storage oscilloscope and stored in a microcomputer (IBM-AT) using an analogue-to-digital conversion board controlled by our acquisition and display software (Robinson and Giles, 1986). The solution used to fill the micropipette contained (in mM): K-aspartate 90, KCl 20, EGTA-KOH 0.77,  $MgCl_2$  1.0,  $CaCl_2$  0.045,  $Na^{2+}$ -ATP 3, HEPES 5, buffered to pH 7.2–7.3 with KOH.  $P_{Ca}$  of the pipette solution was calculated to be 8.0 (Fabiato and Fabiato, 1979). The cells were superfused with a HEPES-buffered Ringer's solution having the following composition (in mM): NaCl 110, KCl 2.5,  $MgCl_2$  5.0,  $CaCl_2$  2.5, glucose 5, HEPES 5, buffered to pH 7.4 with NaOH. In some experiments GTP $\gamma$ S (guanosine 5'-*O*-(3-thiotriphosphate) tetralithium salt (Boehringer Mannheim Biochemicals, Indianapolis, IN), or Gpp(NH)p (Sigma Chemical Co., St. Louis, MO) were added to the pipette-filling solution. Acetylcholine chloride, atropine

sulfate, and NEM (all from Sigma Chemical Co.) were added to the superfusing Ringer's solution from freshly prepared stock solutions. All experiments were performed at room temperature (22–24°C).

## RESULTS

### *Effect of NEM on the Acetylcholine-activated $K^+$ Current*

Fig. 1 shows the effects of NEM on the  $K^+$  current ( $I_{K(ACh)}$ ) activated by muscarinic receptors in single atrial cells. In control conditions (panel A, part a), a TTX-resistant inward current,  $I_{Ca}$ , is recorded positive to  $-30$  mV.  $I_{Ca}$  is followed by slower activation of  $I_K$ , the delayed rectifier  $K^+$  current. Upon repolarization to the holding potential,  $I_K$  deactivates and corresponding tail currents are recorded. An inwardly rectifying background  $K^+$  current,  $I_{K1}$ , can be elicited by hyperpolarizing pulses. After the application of ACh ( $10^{-6}$  M; panel A, part b), both the holding current at  $-40$  mV and the currents during the depolarizing pulses increase in the outward direction, and concomitantly the inward current during the hyperpolarizing pulses negative to  $-90$  mV also increases. Fig. 1 B shows the corresponding current–voltage relationships, obtained by measuring the currents at the end of the voltage clamp pulses. The ACh-induced currents, defined as the difference between the currents recorded before and after the application of ACh, exhibited inward rectification, reversed near  $-85$  mV, and showed time-dependent relaxations in response to the hyperpolarizing pulses. These ACh effects were blocked completely by atropine ( $10^{-6}$  M), demonstrating that they are mediated by muscarinic receptors (results not shown). These results confirm that these ACh-induced current changes are due to the activation of  $I_{K(ACh)}$  (Momose et al., 1984; Simmons and Hartzell, 1987; Breitwieser and Szabo, 1988; Clark et al., 1990; Nakajima et al., 1990).

After the application of NEM ( $5 \times 10^{-5}$  M) for  $\sim 3$  min, the muscarinic effects of ACh were almost completely abolished (Fig. 1 A, part c). In contrast, neither  $I_{K1}$  nor the  $I_K$  tails were changed significantly, indicating that these ACh-independent  $K^+$  currents were not affected by NEM. These results demonstrate that NEM selectively inhibits  $I_{K(ACh)}$ .

To obtain additional information concerning the time course of development of the inhibitory actions of NEM on  $I_{K(ACh)}$ , NEM ( $5 \times 10^{-5}$  M) and ACh ( $10^{-5}$  M) were applied simultaneously (Fig. 2). At first, the holding current at  $-40$  mV increased in the outward direction. At the same time, the inward current activated by the command pulse to  $-110$  mV also increased. Later  $I_{K(ACh)}$  gradually decreased, in spite of the continuous application of ACh, and 5 min after the recording began  $I_{K(ACh)}$  had declined to near zero. Complete inhibition of  $I_{K(ACh)}$  was obtained 3–6 min after the application of  $5 \times 10^{-5}$  M NEM ( $n = 4$ ). Note that this time-dependent inhibition of  $I_{K(ACh)}$  by NEM is much different than the time course of the inhibition of  $I_{K(ACh)}$  by atropine. When atropine ( $10^{-6}$  M) and ACh ( $10^{-5}$  M) are applied simultaneously,  $I_{K(ACh)}$  failed to develop. The marked differences in these time courses of the inhibition of  $I_{K(ACh)}$  suggest that NEM does not block  $I_{K(ACh)}$  by a simple ligand–receptor interaction, as atropine does.




FIGURE 1. NEM-induced inhibition of the effects of ACh on muscarinic receptor-activated  $K^+$  current,  $I_{K(ACh)}$ . The membrane potential was held at  $-40$  mV and 200-ms command pulses to various membrane potentials were applied at 0.2 Hz. The current traces shown in *A* were recorded in control (*a*), in  $10^{-6}$  M ACh (*b*), and in  $10^{-6}$  M ACh plus  $5 \times 10^{-5}$  M NEM (*c*). The zero current level is indicated by the solid triangle. The current traces during the depolarizing ( $-30$  to  $+30$  mV) and hyperpolarizing pulses ( $-50$  to  $-120$  mV) are shown in the upper and lower parts, respectively. *B* shows current-voltage relationships measured at the end of the pulses (control [○],  $10^{-6}$  M ACh [●], and  $10^{-6}$  M ACh plus  $5 \times 10^{-5}$  M NEM [■]).

#### Effect of NEM on the GTP $\gamma$ S-enhanced $K^+$ Current

Intracellular dialysis from the recording pipette with a nonhydrolyzable GTP analogue, GTP $\gamma$ S ( $5 \times 10^{-5}$  M), gradually increased  $I_{K(ACh)}$  even in the absence of ACh (Fig. 3). Time-dependent relaxations of  $I_{K(ACh)}$  during applied hyperpolarizations and after return to  $-40$  mV were also observed (Fig. 3 *A*, part *b*) after GTP $\gamma$ S was dialyzed into the cell. These current changes could not be blocked by atropine

( $10^{-6}$  M) (Fig. 3 A, part c) and were probably due to direct activation of GTP-binding proteins (Breitwieser and Szabo, 1985; Kurachi et al., 1986a, b, c; Nakajima et al., 1990). Fig. 3, A and B shows that this GTP $\gamma$ S-enhanced  $K^+$  current could not be inhibited by NEM when NEM was applied after GTP $\gamma$ S. Furthermore, as illustrated in Fig. 3 C, the activation of  $I_{K(ACh)}$  by GTP $\gamma$ S was enhanced and accelerated by




FIGURE 2. Time-dependent inhibition by NEM of the ACh-activated  $K^+$  current,  $I_{K(ACh)}$ . 400-ms hyperpolarizing pulses to  $-110$  mV from a holding potential of  $-40$  mV were applied at 0.2 Hz. ACh ( $10^{-5}$  M) and NEM ( $5 \times 10^{-5}$  M) were both applied simultaneously. The current traces shown in A were recorded in control (a), or in ACh ( $10^{-5}$  M) plus NEM ( $5 \times 10^{-5}$  M) (b, c) at the times indicated by the corresponding letters in B. B shows the time course of the changes in the holding current (+, top trace) and the inward current measured at the end of the command pulse of  $-110$  mV ( $\square$ , bottom trace). The current changes were measured from the zero current level, indicated by the solid triangles.

simultaneous superfusion with ACh ( $10^{-6}$  M). After the increase in  $I_{K(ACh)}$  had reached a steady state, atropine ( $10^{-6}$  M) failed to inhibit it, indicating that the  $K^+$  channels were uncoupled from the muscarinic receptors and had been activated irreversibly by GTP $\gamma$ S. Fig. 3 C also shows that NEM ( $5 \times 10^{-5}$  M) failed to inhibit the GTP $\gamma$ S-enhanced  $K^+$  current. These results suggest that NEM does not affect




FIGURE 3. Lack of effect of NEM on the GTP $\gamma$ S-induced increase in K<sup>+</sup> current. GTP $\gamma$ S ( $5 \times 10^{-4}$  M) was included in the recording pipette. The membrane potential was held at  $-40$  mV and hyperpolarizing pulses to  $-110$  mV were applied at 0.2 Hz. The time course of the changes in the holding current (+) and the current measured at the end of the clamp pulses ( $\square$ ) are plotted in B, beginning immediately after the rupture of the membrane. The current traces shown in A were recorded (a) in control (i.e., immediately after the rupture of the membrane), (b) after the application of NEM ( $5 \times 10^{-5}$  M), and (c) after NEM ( $5 \times 10^{-5}$  M) plus atropine ( $10^{-6}$  M) treatment. Zero current levels are shown by solid triangles. The holding current change was measured from zero current level. C shows data from a different cell. ACh ( $10^{-6}$  M) was added after the rupture of the membrane, the time course of the changes in the holding current ( $\square$ ) was plotted, and GTP $\gamma$ S was applied intracellularly from the recording pipette. Finally, ACh ( $10^{-6}$  M) and atropine ( $10^{-6}$  M) plus NEM ( $5 \times 10^{-5}$  M) were added to the bath at the times indicated in the top part of C. The holding current change was measured from zero current level.

either the  $K^+$  channels or the dissociated subunits of the GTP-binding proteins which are essential for activation of  $I_{K(ACh)}$  (Brown and Birnbaumer, 1988; Neer and Clapham, 1988). In addition, NEM has no direct effects on the other types of  $K^+$  channels ( $I_{K1}$  and  $I_K$ ) present in bullfrog atrium (shown in Fig. 1). Thus, NEM appears to act on ACh-sensitive  $K^+$  channels at sites located either on the muscarinic receptors or on the GTP-binding proteins ( $G_i$  and/or  $G_o$ ) that are coupled to these  $K^+$  channels.

#### *Effect of GTP $\gamma$ S on Cells Pretreated with NEM*

To obtain additional evidence concerning whether the site of NEM action involves GTP-binding protein(s), atrial cells were treated with NEM before the application of GTP $\gamma$ S. To ensure that the effects of NEM on  $I_{K(ACh)}$  had developed fully, NEM ( $5 \times 10^{-5}$  M) was added to the superfusate  $\sim 10$  min before the dialysis of GTP $\gamma$ S was started (Fig. 4), i.e., before the rupture of the membrane. After this pretreatment with NEM, intracellular application of GTP $\gamma$ S ( $10^{-3}$  M) by dialysis from the recording pipette failed to activate  $I_{K(ACh)}$ , in contrast to the results shown in Fig. 3. This result was also obtained in cells treated with NEM only; thus, it was not related to the presence or absence of ACh, to the concentration of GTP $\gamma$ S ( $5$ – $10 \times 10^{-4}$  M), or to the choice of the nonhydrolyzable GTP analogue, since experiments in which Gpp(NH)p (2 mM) was used also consistently yielded negative results (not shown). In combination, these findings show that in cells pretreated with NEM ( $5 \times 10^{-5}$  M) the activation of  $I_{K(ACh)}$  by GTP $\gamma$ S is reduced significantly, and they suggest that NEM can inhibit the function of the GTP-binding proteins, e.g., their dissociation, without affecting the properties of the active dissociated subunits themselves.

#### *Effect of NEM on the Inhibitory Action of ACh on $I_{Ca}$*

To test whether the actions of NEM on the GTP-binding proteins ( $G_i/G_o$ ) were selective, the effects of NEM on  $I_{Ca}$  were also examined. Fig. 5 shows  $I_{Ca}$  traces recorded before and after the application of NEM ( $5 \times 10^{-5}$  M). The corresponding peak current–voltage relationships are shown below. The membrane potential was held at  $-40$  mV and command pulses were applied at 0.2 Hz. NEM ( $5 \times 10^{-5}$  M) increased  $I_{Ca}$  markedly with no significant change in its voltage dependence. Once again,  $I_{K1}$  was not changed by NEM. Fig. 5 B shows that NEM ( $5 \times 10^{-5}$  M) very gradually increased  $I_{Ca}$ ; the steady-state effect was recorded  $\sim 3$  min after its application. Subsequent addition of ISO ( $10^{-6}$  M) further increased  $I_{Ca}$ , indicating that the stimulatory GTP-binding protein(s) ( $G_s$ ) remained functional after NEM treatment.

The histograms in Fig. 6 summarize the observed effects of NEM on the indirect inhibitory actions of ACh on  $I_{Ca}$ . As shown in the previous paper (Nakajima et al., 1990), ACh ( $10^{-5}$  M) can completely inhibit the effects of ISO ( $10^{-6}$  M) on  $I_{Ca}$  in bullfrog atrium (Fig. 6, *left*). However, this inhibitory effect of ACh ( $10^{-5}$  M) on  $I_{Ca}$  was completely abolished (Fig. 6, *right*) by NEM, even though NEM by itself increased  $I_{Ca}$ , and application of ISO ( $10^{-6}$  M) after NEM further increased  $I_{Ca}$ .

In bullfrog atrial cells, the effects of ACh on both  $I_{Ca}$  and  $I_{K(ACh)}$  are mediated by a

biochemical pathway involving both muscarinic receptors and GTP-binding proteins ( $G_i$  and/or  $G_o$ ) (Nakajima et al., 1990). Our results suggest that NEM can block this pathway. To determine whether NEM selectivity affects  $G_i$  or  $G_o$  rather than  $G_s$ ,  $GTP\gamma S$  ( $5 \times 10^{-4}$  M) was applied to a cell that had been pretreated with NEM




FIGURE 4. Block of the effect of  $GTP\gamma S$  on  $I_{K(ACh)}$  after pretreatment with NEM.  $GTP\gamma S$  ( $10^{-3}$  M) was added intracellularly after making an impalement with a patch pipette. 400-ms command pulses to  $-110$  mV from a holding potential of  $-40$  mV were applied at 0.2 Hz beginning immediately after the rupture of the membrane. The current traces (*a*, *b*) shown in *A* were recorded at times indicated by the corresponding letters in *B*. *B* shows the time courses of the changes in the holding current (+, top trace) and the inward current measured at the end of the command pulse of  $-110$  mV ( $\square$ , bottom trace). Zero current levels are shown by solid triangles.

( $5 \times 10^{-5}$  M) for  $\sim 10$  min (as shown in Fig. 7). In response to  $GTP\gamma S$  dialysis into the cell,  $I_{Ca}$  gradually increased, probably due to the direct activation of the GTP-binding proteins ( $G_s$ ). ISO ( $10^{-6}$  M) further increased  $I_{Ca}$ , suggesting that the activation of  $G_s$  by  $GTP\gamma S$  is enhanced by ISO. Thus, after NEM treatment the






FIGURE 5. Effects of NEM on the  $Ca^{2+}$  current in single bullfrog atrial cells. The membrane potential was held at  $-40$  mV and command pulses were applied at  $0.2$  Hz. The original current traces obtained by depolarizing pulses between  $-30$  mV and  $+20$  mV are illustrated (a) in control and (b) after NEM ( $5 \times 10^{-5}$  M) treatment. Zero current levels are shown by solid triangles. The initial (peak) current-voltage relationships before (○) and after (●) the application of NEM ( $5 \times 10^{-5}$  M) are shown in c. B illustrates the time course of the changes in the  $Ca^{2+}$  current. NEM ( $5 \times 10^{-5}$  M) and ISO ( $10^{-6}$  M) were added at the times indicated. These data were obtained from a different cell than those in A.

functions of  $G_s$  remain intact. The sites of NEM action therefore appear to be located on the GTP-binding proteins ( $G_i$  and/or  $G_o$ ).

#### *Effect of NEM on $Ca^{2+}$ Current Enhanced by ISO and Cyclic AMP*

NEM by itself can increase  $I_{Ca}$  substantially, from approximately  $-100$  pA to  $-750$  pA (mean value,  $n = 4$ ; Figs. 1, 5, and 6). It is likely that one of the underlying mechanisms of NEM actions on  $I_{Ca}$  involves inactivation of  $G_i$ , which could increase basal levels of cyclic AMP (cAMP). We therefore examined the effects of NEM on  $I_{Ca}$

after increasing  $I_{Ca}$  with the  $\beta$ -agonist, ISO, or with intracellular cAMP. As shown in Fig. 8 A, NEM enhanced  $I_{Ca}$  even after maximal effects of ISO ( $10^{-6}$  M) had developed.  $LaCl_3$  ( $5 \times 10^{-5}$  M), a potent blocker of  $I_{Ca}$ , completely blocked the effects of both ISO and NEM on  $I_{Ca}$ . In addition, in experiments in which intracellular application of cAMP ( $2 \times 10^{-4}$  M) had increased  $I_{Ca}$  maximally, and




FIGURE 6. Summary of the NEM-induced inhibition of ACh effects on the calcium current,  $I_{Ca}$ . 200-ms depolarizing pulses to 0 mV from a holding potential of  $-40$  mV were applied at 0.2 Hz. Peak  $I_{Ca}$  was measured from zero current level.  $n$  = number of cells tested. Mean values and standard deviations are illustrated by the open columns and bars, respectively.

subsequent application of ISO ( $10^{-6}$  M) failed to increase  $I_{Ca}$  any further, NEM ( $5 \times 10^{-5}$  M) consistently increased  $I_{Ca}$ . These results suggest that NEM may affect the  $Ca^{2+}$  channel directly, in addition to inactivating GTP-binding proteins.

#### DISCUSSION

Our results show that NEM can inhibit the muscarinic receptor-activated  $K^+$  current,  $I_{K(ACh)}$ , as well as block the inhibitory effect of ACh on the calcium current,  $I_{Ca}$ , in single myocytes from bullfrog atrium. These inhibitory actions of this sulphhydryl alkylating agent are consistent with previous findings showing that NEM

attenuates the negative chronotropic effects of muscarinic agonists in rat and guinea pig hearts (Robert and Konjovic, 1969; Doods et al., 1986; Braun and Sperelakis, 1988).

To determine whether GTP-binding proteins are involved in these actions of




FIGURE 7. Inhibition of the effects of GTP $\gamma$ S on  $I_{K(ACh)}$  and  $I_{Ca}$  after pretreatment with NEM ( $5 \times 10^{-5}$  M) for  $\sim 10$  min. GTP $\gamma$ S ( $5 \times 10^{-4}$  M) was applied intracellularly from the patch pipette. 200-ms command pulses to 0 mV from a holding potential of  $-40$  mV were applied at 0.2 Hz, beginning immediately after the rupture of the membrane. The current traces shown in *A* were recorded at the times indicated by the corresponding letters in *B*. *B* shows the time courses of the changes in the holding current ( $\square$ , top trace) and the  $Ca^{2+}$  current ( $+$ , bottom trace). The zero current levels are illustrated by solid triangles.

NEM, GTP $\gamma$ S was used to activate  $I_{K(ACh)}$ . In cells pretreated with NEM, GTP $\gamma$ S effects were blocked (Figs. 4 and 9), indicating that NEM eliminated the function of GTP-binding proteins, perhaps by inhibiting the binding of GTP $\gamma$ S to GTP-binding proteins or preventing their dissociation into active subunits. However, NEM was




FIGURE 8. Effect of NEM on  $I_{Ca}$  after ISO or cAMP had been applied. Each cell was held at  $-40$  mV and the depolarizing pulses to 0 mV were applied at 0.2 Hz. The changes in the amplitude of  $I_{Ca}$  measured from the zero current level are plotted on the right side of A and B. A, Effect of NEM on ISO-enhanced  $I_{Ca}$ . The current traces shown in A were recorded in (a) control, (b) ISO ( $10^{-6}$  M), (c) ISO ( $10^{-6}$  M) plus NEM ( $5 \times 10^{-5}$  M), and (d) after the addition of  $LaCl_3$  ( $5 \times 10^{-5}$  M) to c. B shows the effects of NEM on cAMP-enhanced  $I_{Ca}$ . cAMP ( $2 \times 10^{-4}$  M) was applied intracellularly from the patch pipette. The current traces shown in B were recorded (a) immediately after the rupture of the membrane, (b) before, and (c) after the application of ISO ( $10^{-6}$  M), and (d) NEM ( $5 \times 10^{-5}$  M) was added. The diagrams at the right sides of both panels show peak  $I_{Ca}$  measured at the times indicated by the corresponding letters on the raw data. Zero current levels are shown by solid triangles.

not effective when applied after GTP $\gamma$ S and activated  $I_{K(ACh)}$  (Figs. 3 and 9). These results indicate that the site of action of NEM is not on the  $K^+$  channel itself or on the dissociated subunits of the GTP-binding proteins.

Previous work has shown that NEM can selectively abolish hormone- or transmitter-induced inhibition of adenylate cyclase in various preparations, including heart cells (Aktories et al., 1982; Jakobs et al., 1982, 1983; Ukena et al., 1984).




FIGURE 9. Effect of NEM on muscarinic receptor-regulated  $K^+$  current. *a*, The current changes at a holding potential of  $-40$  mV after the application of ACh ( $10^{-5}$  M) (*left*) and 3–5 min after the additional application of NEM ( $5 \times 10^{-5}$  M) (*right*) were measured. *n* = number of cells tested. *b*, The effect of NEM on GTP $\gamma$ S-activated  $K^+$  current. GTP $\gamma$ S ( $5 \times 10^{-4}$  M) was applied intracellularly from the patch pipette. In these experiments, after the rupture of the membrane ACh ( $10^{-6}$  M) was added to ensure that GTP $\gamma$ S activated the GTP-binding proteins maximally. The left bar shows the current changes measured when the ACh-induced current reached a steady state, compared with the control value immediately after the rupture of the membrane. The right bar corresponds to the current changes measured at 5–7 min after the addition of atropine ( $10^{-6}$  M) and NEM ( $5 \times 10^{-5}$  M). *c*, The effect of GTP $\gamma$ S in cells pretreated with NEM; i.e., after 7–10 min of NEM treatment, GTP $\gamma$ S ( $5$ – $10 \times 10^{-4}$  M) was applied intracellularly from the recording pipettes. The current changes were measured 5–7 min after the rupture of the membrane, and were compared with the holding current immediately after the rupture of the membrane. In all panels the mean value and the standard deviation are shown in open columns and bars, respectively.

Our electrophysiological data indicate that the observed effects of NEM on  $I_{K(ACh)}$  are quite specific. Thus, two other  $K^+$  currents that are present in these myocytes,  $I_{K1}$  and  $I_K$ , are not significantly changed by NEM ( $5 \times 10^{-5}$  M) treatment. The specificity of NEM effects may arise from selective actions on different GTP-binding proteins.

G proteins are heterotrimers composed of  $\alpha$ -,  $\beta$ -, and  $\gamma$ -subunits (for review see Gilman, 1987). Recently, Winslow et al. (1987) demonstrated that NEM alkylates only specific sites on  $\alpha$ -subunits of  $G_i$  and/or  $G_o$ ; the  $\beta\gamma$ -subunits of these G proteins are not affected by NEM (Asano and Ogasawara, 1986). It is possible, therefore, that the NEM effects we have observed are due to alkylation of sulphhydryl groups only on the  $\alpha$ -subunits of  $G_i$  and/or  $G_o$ .

Previous biochemical studies have shown that NEM can affect several enzymatic responses, even in a particular cell type. For example, Aktories et al. (1982) observed that treatment of intact adipocytes from hamsters with NEM abolished both the receptor-mediated inhibition of adenylate cyclase and the inhibition of adenylate cyclase activity by GTP or nonhydrolyzable GTP analogues (GTP $\gamma$ S and GMP-P(NH)P). In human platelets NEM not only abolished the  $\alpha_2$ -induced inhibition of the adenylate cyclase, but also caused an increase in the basal, prostaglandin  $E_1$ -stimulated activity of this enzyme (Jakobs et al., 1982). Our results are consistent with these biochemical data, since in myocytes from bullfrog atrium NEM removed the inhibitory effect of ACh on  $I_{Ca}$  (Fig. 7), as well as inhibiting  $I_{K(ACh)}$ .

IAP uncouples muscarinic receptors from the GTP-binding proteins ( $G_i$  and/or  $G_o$ ) (Katada and Ui, 1982; Katada et al., 1986; for review, see Gilman, 1987). We observed that receptor-independent, GTP $\gamma$ S-induced activation of these G proteins and their dissociation into subunits remained unchanged after IAP treatment (Cote et al., 1984; Katada et al., 1986). Previous electrophysiological studies (Logothetis et al., 1987; Nakajima et al., 1990) have also shown that GTP $\gamma$ S can activate  $I_{K(ACh)}$  in cells that have been pretreated with IAP. These results suggest that the ability of the GTP-binding proteins to couple to receptors is selectively blocked by IAP. In contrast with this action of IAP, our data suggest that NEM prevents the GTP $\gamma$ S-induced dissociation of the GTP-binding proteins. Thus, it is likely that different sites on the GTP-binding proteins are modified by NEM and IAP (cf. Neer and Clapham, 1988).

Winslow et al. (1987) have reported that NEM alkylates three cysteine residues on the GTP-binding proteins ( $G_o$  and  $G_i$ ). One of these sites is the one that ADP-ribosylated by IAP and is involved in the coupling of the receptors to G proteins. This effect of NEM can explain previous findings showing that this agent reduces the affinity of the receptor for its agonist, an effect that mimics one of the actions of GTP analogues (Wei and Sulakhe, 1980; Harden et al., 1982; Korn et al., 1983; Flynn and Potter, 1985). The other sites are involved in GTP binding, but at these sites the action of NEM is different from that of IAP. After  $I_{K(ACh)}$  was activated by GTP $\gamma$ S, NEM failed to inhibit it, which may suggest that the binding of GTP $\gamma$ S can prevent the effect of NEM. Thus, NEM seems to be able to act on several sites on GTP-binding proteins.

One implication of the effects of NEM that we have observed is that sulphhydryl groups on the sarcolemma of atrial cells can modulate certain ionic channels. One of the mechanisms underlying the effect of NEM on  $I_{Ca}$  may involve the inactivation of  $G_i$ , and a resultant increase in basal levels of cAMP (Jakobs et al., 1982). However, the effect of NEM on  $I_{Ca}$  cannot be explained only by the inactivation of  $G_i$ .

In summary, NEM inhibits the effect of ACh on  $I_{K(ACh)}$  and inhibitory action of ACh on  $I_{Ca}$ , probably by alkylating the GTP-binding proteins ( $G_i$  and/or  $G_o$ ) that are

essential for these responses. Our results provide the first direct evidence that sulphhydryl groups on GTP-binding proteins are essential for these responses, and suggest that NEM could be a very useful tool for studying the molecular pharmacology of certain G protein-linked responses.

The authors wish to thank Mr. Keith Robinson for assistance with data analysis and Drs. R. B. Clark, M. Kaibara, and A. Braun for valuable discussion. We are grateful to Ms. Lenore Doell for skilled secretarial assistance.

*Original version received 1 March 1989 and accepted version received 9 March 1990.*

#### REFERENCES

- Aktories, K., G. Schultz, and K. H. Jakobs. 1982. Inactivation of a guanine nucleotide regulatory site mediating inhibition of adenylate cyclase in hamster adipocytes. *Naunyn-Schmiedeberg's Archives of Pharmacology*. 321:247–252.
- Armstrong, C. M., F. Bezanilla, and E. Rojas. 1973. Destruction of sodium conductance inactivation in squid axons perfused with pronase. *Journal of General Physiology*. 62:375–391.
- Asano, T., and N. Ogasawara. 1986. Uncoupling of  $\gamma$ -aminobutyric acid receptors from GTP-binding proteins by N-ethylmaleimide: effect of N-ethylmaleimide on purified GTP-binding proteins. *Molecular Pharmacology*. 29:244–249.
- Braun, A. P., and N. C. Sperelakis. 1988. Attenuation by N-ethylmaleimide treatment of the cholinergically induced shortening of action potential duration in guinea pig right atrium. *Biochemical Pharmacology*. 37:4577–4581.
- Breitwieser, G. E., and G. Szabo. 1985. Uncoupling of cardiac muscarinic and  $\beta$ -adrenergic receptors from ion channels by a guanine nucleotide analogue. *Nature*. 317:538–540.
- Breitwieser, G. E., and G. Szabo. 1988. Mechanism of muscarinic receptor-induced K channel activation as revealed by hydrolysis-resistant GTP analogues. *Journal of General Physiology*. 91:469–493.
- Brown, A. M., and L. Birnbaumer. 1988. Direct G protein gating of ion channels. *American Journal of Physiology*. 254:H401–410.
- Clark, R. B., and W. R. Giles. 1987.  $Na^+$  current in single cells from bullfrog atrium: voltage dependence and ion transfer properties. *Journal of Physiology*. 391:235–265.
- Clark, R. B., T. Nakajima, W. Giles, K. Kanai, Y. Momose, and G. Szabo. 1990. Two distinct types of inwardly rectifying  $K^+$  channels in bull-frog atrial myocytes. *Journal of Physiology*. 424:229–251.
- Cote, T. E., E. A. Frey, and R. D. Sekura. 1984. Altered activity of the inhibitory guanyl nucleotide-binding component ( $N_i$ ) induced by pertussis toxin: uncoupling of  $N_i$  from receptor with continued coupling of  $N_i$  to the catalytic unit. *Journal of Biological Chemistry*. 259:8693–8698.
- Doods, H. N., D. Davidesko, M. J. Mathy, H. D. Batink, A. de Jonge, and P. A. van Zwieten. 1986. Discrimination by N-ethylmaleimide between the chronotropic and inotropic response to muscarinic receptor stimulation in rat atrium. *Naunyn-Schmiedeberg's Archives of Pharmacology*. 333:182–185.
- Duncan, G., N. J. Emptage, and K. R. Hightower. 1988. P-Chloro-mercuriphenyl sulphonate activates a guanine-sensitive potassium conductance in frog lens. *Journal of Physiology*. 404:637–648.
- Fabiato, A., and F. Fabiato. 1979. Calculator programs for computing the composition of solutions containing multiple metals and ligands used for experiments in skinned muscle cells. *Journal de Physiologie*. 75:463–503.
- Flynn, D. D., and L. T. Potter. 1985. Different effects of N-ethylmaleimide on  $M_1$  and  $M_2$  muscarinic receptors in rat brain. *Proceedings of the National Academy of Sciences USA*. 82:580–583.

- Giles, W. R., and E. F. Shibata. 1985. Voltage clamp of bullfrog cardiac pacemaker cells: a quantitative analysis of potassium currents. *Journal of Physiology*. 368:265–292.
- Gilman, A. G. 1987. G proteins: transducers of receptor-generated signals. *Annual Review of Biochemistry*. 56:615–649.
- Gonoi, T., and B. Hille. 1987. Gating of Na channels. Inactivation modifiers discriminate among models. *Journal of General Physiology*. 89:253–274.
- Hamill, O. P., A. Marty, E. Neher, B. Sakmann, and F. J. Sigworth. 1981. Improved patch clamp technique for high-resolution current recording from cells and cell-free membrane patches. *Pflügers Archiv*. 391:85–100.
- Harden, T. K., A. G. Scheer, and M. M. Smith. 1982. Differential modification of the interaction of cardiac muscarinic cholinergic and beta-adrenergic receptors with a guanine nucleotide binding component(s). *Molecular Pharmacology*. 21:570–580.
- Hartzell, H. C. 1988. Regulation of cardiac ion channels by catecholamines, acetylcholine and second messenger systems. *Progress in Biophysics and Molecular Biology*. 52:165–247.
- Hescheler, J., M. Kameyama, and W. Trautwein. 1986. On the mechanism of muscarinic inhibition of the cardiac Ca current. *Pflügers Archiv*. 407:182–189.
- Hescheler, J., and W. Trautwein. 1988. Modification of L-type calcium current by intracellularly applied trypsin in guinea-pig ventricular myocytes. *Journal of Physiology*. 404:259–274.
- Hume, J. R., and W. Giles. 1981. Active and passive electrical properties of single bullfrog atrial cells. *Journal of General Physiology*. 78:18–43.
- Hume, J. R., and W. Giles. 1983. Ionic currents in single isolated bullfrog atrial cells. *Journal of General Physiology*. 81:153–194.
- Jakobs, K. H., K. Aktories, and G. Schultz. 1983. Inhibitory coupling of hormone and neurotransmitter receptors to adenylate cyclase. *Journal of Receptor Research*. 3:137–149.
- Jakobs, K. H., P. Lasch, M. Minuth, K. Aktories, and G. Schultz. 1982. Uncoupling of  $\alpha$ -adrenoceptor-mediated inhibition of human platelet adenylate cyclase by N-ethylmaleimide. *Journal of Biological Chemistry*. 257:2829–2833.
- Katada, T., M. Oinuma, and M. Ui. 1986. Mechanisms for inhibition of the catalytic activity of adenylate cyclase by guanine nucleotide-binding proteins serving as the substrate of the islet-activating protein, pertussis toxin. *Journal of Biological Chemistry*. 261:5215–5221.
- Katada, T., and M. Ui. 1982. Direct modification of the membrane adenylate cyclase system by islet-activating protein due to ADP-ribosylation of a membrane protein. *Proceedings of the National Academy of Sciences USA*. 79:3129–3133.
- Korn, S. J., M. W. Martin, and T. K. Harden. 1983. N-Ethylmaleimide-induced alteration in the interaction of agonists with muscarinic cholinergic receptors of rat brain. *Journal of Pharmacology and Experimental Therapeutics*. 224:118–126.
- Kurachi, Y., T. Nakajima, and T. Sugimoto. 1986a. Acetylcholine activation of K channels in cell-free membrane of atrial cells. *American Journal of Physiology*. 251:H681–H684.
- Kurachi, Y., T. Nakajima, and T. Sugimoto. 1986b. On the mechanism of activation of muscarinic K channels by adenosine in isolated atrial cells: involvement of GTP-binding proteins. *Pflügers Archiv*. 407:264–274.
- Kurachi, Y., T. Nakajima, and T. Sugimoto. 1986c. Role of intracellular  $Mg^{2+}$  in the activation of muscarinic  $K^+$  channel in cardiac atrial membrane. *Pflügers Archiv*. 407:572–574.
- Logothetis, D. E., Y. Kurachi, J. Galper, E. J. Neer, and D. E. Clapham. 1987. The  $\beta\gamma$  subunits of GTP-binding protein activate the muscarinic K channel in heart. *Nature*. 325:321–326.
- Momose, Y., W. Giles, and G. Szabo. 1984. Acetylcholine-induced K current in amphibian atrial cells. *Biophysical Journal*. 45:20–22.



- Nakajima, T., S. Wu, H. Irisawa, and W. Giles. 1990. Mechanism of acetylcholine-induced inhibition of Ca current in bullfrog atrial myocytes. *Journal of General Physiology*. 96:865–885.
- Neer, E. J., and D. E. Clapham. 1988. Roles of G protein subunits in transmembrane signalling. *Nature*. 333:129–134.
- Pfaffinger, P. J., J. M. Martin, D. D. Hunter, N. M. Nathanson, and B. Hille. 1985. GTP-binding proteins couple cardiac muscarinic receptors to a K channel. *Nature*. 317:536–538.
- Roberts, C. H., and J. Konjovic. 1969. Differences in the chronotropic and inotropic response of the rat atrium to choline esters, cholinesterase inhibitors and certain blocking agents. *Journal of Pharmacology and Experimental Therapeutics*. 169:109–119.
- Robinson, K., and W. Giles. 1986. Menu-driven A/D conversion, display and plotting programs for the IBM PC in electrophysiological studies. *Computer Methods and Programs in Biomedicine*. 23:319–329.
- Simmons, M. A., and H. C. Hartzell. 1987. A quantitative analysis of the acetylcholine-activated potassium current in single cells from frog atrium. *Pflügers Archiv*. 409:454–461.
- Stimers, J., F. Bezanilla, and R. Taylor. 1985. Sodium channel activation in the squid giant axon. *Journal of General Physiology*. 85:65–82.
- Ukena, D., E. Poeschla, E. Huttemann, and U. Schwabe. 1984. Effects of N-ethylmaleimide on adenosine receptors of rat fat cells and human platelets. *Naunyn-Schmiedeberg's Archives of Pharmacology*. 327:247–253.
- Wei, J. W., and P. V. Wulakhe. 1980. Requirement for sulphhydryl groups in the differential effects of magnesium ion and GTP on agonist binding to muscarinic cholinergic receptor sites in rat atrial membrane fraction. *Naunyn-Schmiedeberg's Archives of Pharmacology*. 327:247–253.
- Winslow, J. W., J. D. Bradley, J. A. Smith, and E. J. Neer. 1987. Reactive sulphhydryl groups of  $\alpha_{59}$ , a guanine nucleotide-binding protein from brain. *Journal of Biological Chemistry*. 262:4501–4507.
- Wong, S. R., R. Martin, and A. M. Tolkovsky. 1985. Pertussis-toxin substrate in guanosine 5'-[ $\beta$ -thio]diphosphate-, N-ethylmaleimide-,  $Mg^{2+}$ - and temperature-sensitive GTP-binding protein. *Biochemical Journal*. 232:191–197.