



Current and Possible Future Therapeutic Options for Huntington's Disease

Mackenzie W. Ferguson^{1,†}, Connor J. Kennedy^{1,†},
Thulani H. Palpagama¹, Henry J. Waldvogel¹, Richard L. M. Faull¹
and Andrea Kwakowsky^{1,2} 

Journal of Central Nervous System Disease
Volume 14: 1–27
© The Author(s) 2022
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/11795735221092517


¹Centre for Brain Research, Department of Anatomy and Medical Imaging, Faculty of Medical and Health Sciences, University of Auckland, Auckland, New Zealand. ²Pharmacology and Therapeutics, School of Medicine, Galway Neuroscience Centre, National University of Ireland Galway, Galway, Ireland.

ABSTRACT

Huntington's disease (HD) is an autosomal neurodegenerative disease that is characterized by an excessive number of CAG trinucleotide repeats within the huntingtin gene (*HTT*). HD patients can present with a variety of symptoms including chorea, behavioural and psychiatric abnormalities and cognitive decline. Each patient has a unique combination of symptoms, and although these can be managed using a range of medications and non-drug treatments there is currently no cure for the disease. Current therapies prescribed for HD can be categorized by the symptom they treat. These categories include chorea medication, antipsychotic medication, antidepressants, mood stabilizing medication as well as non-drug therapies. Fortunately, there are also many new HD therapeutics currently undergoing clinical trials that target the disease at its origin; lowering the levels of mutant huntingtin protein (mHTT). Currently, much attention is being directed to antisense oligonucleotide (ASO) therapies, which bind to pre-RNA or mRNA and can alter protein expression via RNA degradation, blocking translation or splice modulation. Other potential therapies in clinical development include RNA interference (RNAi) therapies, RNA targeting small molecule therapies, stem cell therapies, antibody therapies, non-RNA targeting small molecule therapies and neuroinflammation targeted therapies. Potential therapies in pre-clinical development include Zinc-Finger Protein (ZFP) therapies, transcription activator-like effector nuclease (TALEN) therapies and clustered regularly interspaced short palindromic repeats (CRISPR)/CRISPR-associated system (Cas) therapies. This comprehensive review aims to discuss the efficacy of current HD treatments and explore the clinical trial progress of emerging potential HD therapeutics.

KEYWORDS: chorea, antipsychotic medication, antidepressants, mood stabilizers, antisense oligonucleotides, RNAi therapies, Huntington's disease

RECEIVED: July 6, 2021. ACCEPTED: March 21, 2022.

TYPE: Review

DECLARATION OF CONFLICTING INTERESTS: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

FUNDING: The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by Oakley Mental Health Research Foundation (AK, MF, TP, HJW, RLMF; 3725204),

Alzheimers New Zealand Charitable Trust (AK; 370836), Alzheimers New Zealand (AK; 3718869), Freemasons New Zealand (AK; 3719321), Aotearoa Foundation, Centre for Brain Research and University of Auckland (AK; 3705579), Neurological Foundation of New Zealand (AK, TP; 848010), Brain Research New Zealand (AK, HJW, RLF), Health Research Council of New Zealand (RLF and HJW; 3627373).

CORRESPONDING AUTHOR: Andrea Kwakowsky, Centre for Brain Research, Department of Anatomy and Medical Imaging, Faculty of Medical and Health Sciences, University of Auckland, 85 Park Rd, Grafton, Auckland 1142, New Zealand. Email: a.kwakowsky@auckland.ac.nz

Introduction

HD is an autosomal dominant neurodegenerative disease that currently has no approved cure. HD is characterized by an increased number of CAG trinucleotide repeats in the huntingtin gene (*HTT*) on the short arm of chromosome 4. A higher number of CAG repeats in the huntingtin gene is directly associated with a greater probability of HD onset. Individuals with greater than 36 CAG repeats have an increased risk of HD, while patients with over 40 repeats are predestined to experience disease onset within their life span.^{1–5} The *HTT* gene which possesses the CAG repeats is transcribed into the huntingtin protein (HTT).⁶ A high number of CAG repeats results in a dysfunctional mHTT protein, forming protein aggregates within the neurons and other cell types.⁷ These aggregates, among other mechanisms, are hypothesized to cause

mitochondrial dysfunction, neuronal stress, excitotoxicity and neuroinflammation (Figure 1).^{8–12} The onset of HD typically occurs at middle age, however, HD symptoms can develop at any time during the lifespan.^{13,14} Worldwide, the prevalence of HD is 2.71 per 100,000, however, this number increases to 5.70 per 100,000 in westernised countries such as Europe, the United States and Australia.¹⁵ This has been postulated to be the result of a higher frequency of the mutant HD gene in western haplotypes.¹⁶

A variety of symptoms have been recorded with the majority resulting in the deterioration of cognitive ability, motor function, a change in behaviour or a combination of these symptoms.^{17,18} At present, there are no approved treatments that directly target the mutated HTT gene or corresponding mHTT aggregates. Current approved HD drugs address the range



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/ham/open-access-at-sage>).




Figure 1. Mechanisms of Huntington's disease pathogenesis.

of symptoms that arise as a consequence of the disease, but they do not address the disease at its origin.¹⁹ Each patient's symptoms can be unique, hence there is no standardized treatment with medication prescribed on a case-by-case basis and an extensive range of medications used to treat each of the unique symptoms.^{13,20} Side effects from medications used to treat HD can also affect each patient differently.¹⁸

Most HD symptoms can be classified under motor, cognitive or behavioural categories.^{14,18} Motor symptoms include chorea during the early and mid-stages of disease onset, however, in later stages of disease progression movements become slower (bradykinetic) and less frequent (hypokinetic).²¹ Alternatively, behavioural symptoms include depression, apathy and anxiety.^{22,23} In 1988, a survey of 2835 HD patients discovered that 40% of patients had depression at the time of the study, while 50% had sought clinical help for depression in the past.²⁴ Other less common psychiatric symptoms include obsessive-compulsive behaviour, psychosis and aggression. Memory deficits and retrogression of thought processing and organization can arise as cognitive symptoms.^{21,24} Other symptoms can also present, such as unintentional weight loss and sleep difficulties.¹⁴

The field of potential future HD therapies is advancing rapidly and holds an exciting future. Currently, attention is particularly focussed on lowering mHTT through RNA and

DNA targeting therapies.²⁵ Of these, antisense oligonucleotides are the most rapidly advancing potential therapies.²⁶ Currently, Tominersen has one of the largest phase 3 clinical trials, and results from this trial were expected in 2022.^{27,28} However, the dosing in this trial has been prematurely discontinued and the results so far are being assessed.²⁹ Most RNA and DNA targeting therapies are delivered invasively via intrathecal or intracranial injection.³⁰ Furthermore, DNA based therapies are all in pre-clinical stages and have added ethical complications due to their potential to edit germline DNA.³¹ Other less invasive therapies are therefore being investigated, including stem cell therapies, antibody therapies (both mostly delivered intravenously) and small molecule therapies (delivered orally).^{25,32,33} All of these concepts hold promise and have their own strengths and weaknesses. Many are in clinical trials with manifest HD participants,²⁷ however, there is still significant development of these drugs needed until new therapies are approved and available to HD patients. Accessible new therapies will likely take many years to eventuate.³⁴ This comprehensive review will aim to compare the relative effectiveness of currently approved medications for HD and explore new treatment possibilities, with a special focus on those undergoing clinical trials. We intend to provide a broad overview of the field of current and potential future HD therapies as a forefront resource, rather than focussing on specific therapy types as many

previous reviews have done.³⁵ A particular emphasis is placed on currently researched therapies, but therapies investigated in the past 5 to 10 years were included if they had been comprehensively studied or discussed in their subfield.

Current Therapeutic Options for Huntington's Disease

Chorea Medications

The most commonly prescribed HD medications are targeted to reduce chorea (Table 1). One of the major pathological hallmarks of HD is the degeneration of the basal ganglia and particularly the striatum, which has been linked to the development of chorea. The basal ganglia contributes to the initiation of movement and suppression of unwanted movements. Refinement of movement by basal ganglia structures occur through the direct and indirect pathways.³⁶⁻³⁹ In HD, it is hypothesized that the early loss of the enkephalin containing medium spiny neurons result in reduced inhibition of the globus pallidus externus (GPe) and therefore increased inhibition of the subthalamic nucleus (STN). This in turn leads to decreased activation of the GPe and the substantia nigra pars reticulata (SNr) resulting in decreased inhibition of the thalamus. An increase in the resulting excitation of the cortex through excitatory thalamocortical projections can lead to excess uncontrolled movements causing chorea.³⁷⁻³⁹ Although, chorea is the hallmark symptom of HD, affective and cognitive symptoms can be observed decades before the onset of chorea.⁴⁰

Tetrabenazine. Tetrabenazine (TBZ) was the first drug approved to treat Huntington's disease-associated chorea.⁴¹ Although the direct link between its mechanism of action and overall pharmacodynamic effect of reducing chorea is not fully understood, its pharmacological action is thought to selectively deplete central monoamines from the nerve terminal by

reversibly inhibiting the human vesicular monoamine transporter 2 (VMAT2).⁴² To understand the effectiveness of the drug, a randomized control trial was completed by the Huntington's Study Group where 84 patients received TBZ (n = 54) or placebo (n = 30) over 12 weeks.⁴³ The dose was gradually increased over the first 7 weeks to either a maximum of 100 mg/day, until desired anti-chorea effects were achieved, or until side effects prevented continued participation of the patient. The outcome was measured via a change in chorea score of the Unified Huntington's Disease Rating Scale (UHDRS).⁴⁴ Following the 12 weeks, it was concluded that TBZ significantly reduced chorea severity by 5.0 UHDRS units relative to only 1.5 UHDRS units in the placebo treatment. Four subjects did not complete the treatment due to adverse side effects. Following on from this, another 15-year longitudinal study examined patients taking TBZ and found that 17.2% of HD patients experienced adverse side effects including drowsiness, parkinsonism, depression, insomnia, anxiety and akathisia. However, the majority of these side effects were eliminated with a reduction in dose.⁴⁵

Deutetrabenazine. A similar medication to TBZ used to treat HD-associated chorea is deutetrabenazine (DBZ). DBZ is an isotopic isomer of TBZ, where 6 hydrogen atoms have been replaced by deuterium atoms. Therefore, DBZ has the same chemical function as TBZ, however, the deuterium atoms extend the half-life of the compound, in turn allowing for less frequent administration.^{20,46} A clinical trial, completed by Frank et al. (2016) randomized 90 HD patients between being administered DBZ (n = 45) or a placebo (n = 45) over a 12-week period (NCT01795859). Patient chorea scores were both measured at baseline and following the treatment intervention. HD patients receiving DBZ all experienced significant improvements in their chorea scores at the end of the 12 weeks relative to the control patients.⁴⁷

Table 1. Currently used medications for HD treatment.

CLASS OF MEDICATION	MEDICATION	MECHANISM OF ACTION
Chorea medication	Tetrabenazine	Depletes central monoamines by reversibly inhibiting VMAT2
	Deutetrabenazine	Depletes central monoamines by reversibly inhibiting VMAT2, however, has a longer half-life
Antipsychotic medication	Olanzapine	Inhibits dopamine receptors, serotonin receptors, histamine receptors as well as α_1 -adrenergic and muscarinic receptors
	Risperidone	Selectively inhibiting serotonin and dopamine-D2 receptors
Antidepressants (SSRIs)	Citalopram	Inhibit the reuptake of 5-HT into the presynaptic nerve terminal
	Fluoxetine	Inhibit the reuptake of 5-HT into the presynaptic nerve terminal
	Sertraline	Inhibit the reuptake of 5-HT into the presynaptic nerve terminal
Mood stabilizers	Lamotrigine	Blockade of voltage-gated sodium ion channels, however, also suppresses excitatory neurotransmitters glutamate and aspartate
	Carbamazepine	Blockade of voltage-gated sodium ion channels

Table 2. Overview of potential future HD treatments.

THERAPY	SPONSOR	DEVELOPMENT	ADMINISTRATION	ALLELE SPECIFICITY	DOSING FREQUENCY
RNA targeting therapies					
ASO					
<i>Tominersen</i>	Hoffmann-La Roche	Phase 3	Intrathecal injection	Allele non-specific	Multiple doses
<i>WVE-120102/120101</i>	Wave life Sciences	Phase 1b/2a	Intrathecal injection	Allele-specific WVE-120101: rs362307 (SNP1) WVE-120102: rs362331 (SNP2)	Multiple doses
<i>WVE-003</i>	Wave life Sciences	Pre-clinical	Undisclosed	Allele-specific: SNP3 (undisclosed)	Multiple doses
<i>(CUG)7</i>	BioMarin	Pre-clinical	Intracerebroventricular infusion (mice)	Allele-specific: CAG repeats	Multiple doses
<i>TTX-3360</i>	Triplet Therapeutics	Pre-clinical	Intracerebroventricular infusion (mice)	Not applicable	Unknown
RNAi					
<i>AMT-130</i>	UniQure	Phase 1b/2a	Intrastriatal injection (delivered by AAV5)	Allele non-specific	One dose
<i>AAV.shHD2.1^a</i>	Spark	Pre-clinical	Intracranial injection (delivered by AAV1)	Allele non-specific	One dose
<i>VY-HTT01</i>	Voyager	Pre-clinical	Intracranial injection (delivered by AAV1)	Allele non-specific	One dose
Small molecules					
<i>Branaplam</i>	Novartis Pharmaceutical	Pre-clinical	Oral	Allele non-specific	Multiple doses (weekly)
<i>PTC518</i>	PTC Therapeutics	Phase 1	Oral	Allele non-specific	Multiple doses
<i>Unnamed small molecule</i>	Nuredis	Pre-clinical	Gene deletion in animal model, intracerebroventricular bolus injection	Allele-specific: elongation cofactors required for expanded CAG repeat transcription	Single dose
DNA targeting therapies					
Zinc protein fingers					
<i>TAK-686</i>	Takeda and Sangamo	Pre-clinical	Intrastriatal injection	Allele-specific: expanded CAG repeats	Single dose
<i>ZF-KOX1</i>	European Research Council (undertaken by Imperial College London/ Fingers4Cure)	Pre-clinical	Intraventricular injection	Allele-specific: expanded CAG repeat	Single dose
CRISPR/Cas9					
<i>Unnamed CRISPR/Cas9</i>	Harvard University	Pre-clinical	NA (in cell lines) – theoretically intracranial injection	Allele-specific: SNPs related to mHTT	Single dose

(Continued)

Table 2. Continued.

THERAPY	SPONSOR	DEVELOPMENT	ADMINISTRATION	ALLELE SPECIFICITY	DOSING FREQUENCY
<i>Unnamed CRISPR/Cas9</i>	NIH and National Natural Science Foundation of China (undertaken by Emory University)	Pre-clinical	Intrastriatal injection	Allele non-specific	Single dose
Stem cell therapies					
<i>Cellavita HD</i>	Azidus Brasil	Phase 2/3	Intravenous infusion	Not applicable	Multiple doses
<i>Foetal stem cell transplant</i>	Health and Care Research Wales (undertaken by Cardiff University)	Phase 1	Intrastriatal injection	Not applicable	Single dose
<i>Autologous stem/stromal cells</i>	Regeneris Medical	NA (in clinical trial)	Intravenous injection	Not applicable	Unknown
Antibody therapies					
<i>ANX005</i>	Annexon, Inc	Phase 3	Intravenous infusion	Not applicable	Multiple doses
<i>VX15/2503</i>	Vaccinex, Inc	Phase 2	Intravenous infusion	Not applicable	Multiple doses
<i>W20</i>	National Natural Science Foundation of China, National Science and Technology Major Projects of New Drugs	Pre-clinical	Intracerebroventricular injection (mice)	Allele-specific: does not impact wtHTT	Multiple doses
<i>INT41</i>	Vybion Inc	Pre-clinical	Intrastriatal injection (mice)	Allele-specific: binds to mHTT fragments	Single dose
<i>C6-17</i>	AFFiRiS	Pre-clinical	Unknown	Allele-specific – binds to HTT protein near the aa586 caspase-6 cleavage region	Unknown
Other small molecule therapies					
<i>Pridopidine</i>	Prilenia Therapeutics	Phase 3	Oral	Not applicable	Multiple doses
<i>Laquinimod</i>	Teva Pharmaceutical Industries Ltd	Phase 2 (in HD)	Oral	Not applicable	Multiple doses
<i>Fenofibrate</i>	University of California, Irvine	Phase 2	Oral	Not applicable	Multiple doses
<i>Neflamapimod</i>	EIP Pharma Inc	Phase 2	Oral	Not applicable	Multiple doses
<i>Nilotinib</i>	Georgetown University	Phase 1	Oral	Not applicable	Multiple doses
<i>SRX246</i>	Azevan Pharmaceuticals ^a	Phase 2	Oral	Not applicable	Multiple doses
<i>Varenicline</i>	University of Auckland and University of Otago	Open label study	Oral	Not applicable	Multiple doses
<i>SAGE-718</i>	Sage Therapeutics	Phase 1	Oral	Not applicable	Multiple doses
<i>PBT2</i>	Prana Biotechnology Limited	Phase 2	Oral	Not applicable	Multiple doses

^aAAV.shHD2.1 is also known as AAV.shHD2.

Both TBZ and DBZ prescriptions are labelled with warnings about the side effects of depression and suicidality.⁴⁸ However, individuals with HD taking DBZ had a significantly lower risk of developing neuropsychiatric adverse outcomes (agitation, depression, drowsiness, insomnia and parkinsonism) relative to individuals with HD taking tetrabenazine.⁴⁹ Since the

induction of the drug in a clinical setting various studies have been completed analysing the link between suicidality and TBZ/DBZ. Experiments from both Dorsney et al. (2013), and Shultz et al. (2018), both found that TBZ use was not associated with an increased incidence of depression or suicidality.^{50,51} The contrast in findings could be due to the unaccounted increase in

depression and suicidality from the HD disease alone.²⁴ It is known that depression significantly increases in HD patients with 10% to 25% attempting suicide at least once.^{24,52} These studies highlight the need for further research to be conducted to understand whether the increase in depression and suicidality is a result of TBZ/DBZ or if HD patients already have an underlying susceptibility to these symptoms.

Dopamine antagonists (Antipsychotics or neuroleptics)

Dopamine antagonists are used as second-line agents for chorea relief. These medications are primarily used for the treatment of psychosis, however, they also help to relieve chorea symptoms due to their interaction with the D2 receptor. Specific medications that antagonise dopamine are discussed in further detail below under antipsychotic medication⁵³⁻⁵⁵.

Anti-glutamatergics

There has also been some research into the role that glutamate excitotoxicity plays in HD-associated chorea. Hence, medications such as amantadine and riluzole have been trialed as anti-glutamatergics to reduce chorea symptoms.⁵⁶

Amantadine

Amantadine is a non-competitive N-methyl-D-aspartate (NMDA) receptor antagonist. However, it is hypothesized to affect the dopaminergic system and is considered the most effective agent to treat levodopa-induced dyskinesia in patients with Parkinson's disease.⁵⁷ Verhagen Metman et al. trialed Amantadine to treat HD-associated chorea. To evaluate Amantadine's effects, 24 HD patients entered a double blinded study where the treatment group received 400 mg/day of Amantadine. Over a 2-week treatment period, chorea scores were lower in the treatment group, with the median decrease in extremity chorea at rest being 36% ($P = .04$).⁵⁸

Riluzole

Riluzole is another anti-glutamatergic that has also been trialed to treat Huntington's associated chorea. One study by Rosas et al. treated 8 HD patients with 50 mg of riluzole twice a day over 6 weeks. They found that the patients' chorea rating score improved by 35% ($P = .013$) and regressed back to their original scores once discontinued ($P = .026$).⁵⁹ Alternatively, a similar study treated 9 HD patients with an identical dosing regime, but over 12 months. They discovered that patients had improved chorea scores at 3 months, however, this was sustained at 12 months.⁶⁰ Therefore, the long-term efficacy of riluzole to treat HD-specific chorea is still in question.

Antipsychotic Medication

Olanzapine. Olanzapine is the second most commonly prescribed drug for HD²⁰ (Table 1). Olanzapine is traditionally used as an antipsychotic drug that is primarily used to treat symptoms associated with schizophrenia.⁶¹ Olanzapine is now prescribed to treat many HD patients expressing behavioural symptomology. A study on the short-term effects of olanzapine with HD patients ($n = 11$) found that 6 months following administration their behavioural assessment scores within the UHDRS significantly improved.⁶² Chorea scores also improved in these patients, however, were not significant. This suggests that olanzapine may be a more suitable medication for HD patients with behavioural symptoms, while TBZ/DBZ would better address HD-associated chorea. Patients who present with multiple symptom groups are often prescribed multiple medications, taking into account possible negative drug-drug interactions.²¹ This example again illustrates that there is no universal approach to treating HD and highlights the need for each patient to be assessed individually.

Olanzapine inhibits dopamine receptors (D1, D2 and D4), serotonin receptors (5-HT_{2A}, 5-HT_{2C}), histamine receptors (H₁) as well as α 1-adrenergic and muscarinic receptors.⁵⁴ Unlike the majority of other antipsychotic medications, olanzapine has a higher affinity for 5-HT receptors compared to D2 receptors, therefore, being classed as an atypical antipsychotic.⁵⁴

Side effects of this medication include excess weight gain and dyslipidemia. This has been observed in rat studies as well as clinical trials.^{63,64} To overcome this, it is important to take into account the patient's current metabolic health before prescribing the medication. A study by Ball et al., (2001) found that by placing olanzapine patients on a weight loss program they could mitigate some of the weight gain effects. However, sustained weight management was only observed in males.⁶⁵

Risperidone. Another traditionally used antipsychotic medication that is prescribed to HD patients is risperidone.²⁰ Similar to olanzapine, risperidone has antagonistic properties, selectively inhibiting serotonin-5HT₂ (5-HT) and dopamine-D2 receptors⁵⁵ and is commonly used to treat schizophrenia.⁶⁶ Risperidone is an atypical antipsychotic drug. An atypical antipsychotic is a drug that has a higher binding affinity to 5-HT receptors (e.g. olanzapine), while alternatively, a typical antipsychotic has a higher binding affinity to D2 receptors.⁶⁷ However, it is important to note that while risperidone is classified as an atypical antipsychotic drug it binds to both 5-HT₂ and D2 receptors.⁶⁸

Research completed by Duff et al. (2008) administered risperidone to 17 HD patients over 14 months. They found that patients taking risperidone medication had significantly improved behavioural and psychiatric assessment scores compared to controls ($n = 12$).⁶⁹ Additionally, motor symptoms of the treatment group stabilized while the control group's motor symptoms deteriorated⁶⁹ (NCT04201834). This suggests that risperidone may have dual benefits for both the treatment of

motor and behavioural symptoms associated with HD. However, further evidence supporting this dual benefit of risperidone is minimal, therefore, larger randomized control trials must be completed in the future to understand the drug's full efficacy.

A randomized control trial was completed by Conley & Mahmoud, comparing the relative efficacy of risperidone and olanzapine at reducing adverse cognitive and mood behaviours in Schizophrenic patients ($n = 377$) over 8 weeks.⁷⁰ Both treatments were efficacious and well tolerated by patients. The only difference was that a greater proportion of patients taking olanzapine (27%) experienced a significant increase in body weight percentage relative to those taking risperidone (12%).⁷⁰ This suggests that weight gain is an area of concern when prescribing antipsychotic drugs to HD patients, and highlights risperidone as a potentially more suitable option.

Antidepressants

Approximately 40% of HD patients experience partial or ongoing HD-associated depression and 1 in 10 attempt suicide post-disease diagnosis.^{24,52,71} Antidepressant drugs are often prescribed to HD patients suffering from depression, however, these do not address other psychotic symptoms nor the Huntington's associated chorea. The majority of antidepressants used to treat HD-associated depression are selective serotonin reuptake inhibitors (SSRIs) (Table 1). SSRIs preferentially inhibit the reuptake of 5-HT into the presynaptic nerve terminal. This increases the concentration of 5-HT in the extracellular environment, increasing the amount of 5-HT available to bind to receptors on the post-synaptic cleft,⁷²⁻⁷⁴ which leads to desensitization of the receptors. Desensitization of serotonin receptors may contribute to the therapeutic actions of SSRIs and also account for the development of tolerance to acute side effects of these drugs. SSRIs might result in strong but slow disinhibition of 5-HT neurotransmission. Treatment with antidepressants also causes a downregulation of 5-HT₂ receptors.⁷⁵⁻⁷⁷

SSRIs also have several undesirable side effects such as sexual dysfunction and weight gain.⁷⁸ These symptoms typically arise following a month of treatment and can be prevented if diagnosed early.⁷⁸ Some studies have also found an increased risk of suicide in adolescents taking SSRIs, however, this may be less of an issue in HD patients as HD onset does not usually occur until middle age. This same risk factor has not been observed in adults using SSRI medication.^{79,80} Overall, the literature suggests an inadequate knowledge to properly manage how antidepressants are prescribed to HD patients.⁸¹ Further research is required to understand which SSRIs are most suitable for HD patients and whether these are the best medication to address the high rates of suicidality.

Citalopram. A common SSRI that is prescribed to HD patients is citalopram. A randomized control trial by Beglinger et al., (2014) administered HD patients with citalopram and measured any relative changes in executive function. Thirty-three

adult HD patients were either administered with 20 mg of citalopram or a placebo pill daily over 20 weeks. Although there were no significant changes in executive function, patients showed a significant improvement in depression symptoms on the Hamilton Depression Rating Scale (HAM-D).⁸²

Fluoxetine. A second SSRI that is prescribed to HD patients is fluoxetine. This drug is used primarily to treat HD-associated unipolar depression.²¹ One case study found that 2 non-depressed HD patients who took fluoxetine experienced reduced obsessive compulsive disorder (OCD) symptoms and mood improvements.⁸³ Fluoxetine acts to reduce depression in the short-term, however, there are limited clinical trials examining its effectiveness in HD patients with a need for further studies becoming evident.⁸⁴

Sertraline. Another SSRI that is often prescribed to HD patients is sertraline. This medication not only reduces depression but has also been recorded to cause a complete cessation of aggression and OCD symptoms associated with HD.^{85,86}

Mood Stabilizing Medications

HD patients with behavioural symptoms can also present with symptoms such as OCD, aggression and bipolar disorder. To stabilize these mood disorders, patients can be prescribed a range of anticonvulsants such as sodium valproate, lamotrigine and carbamazepine.²¹ Anticonvulsants have been reported to be effective at minimizing the severity of these symptoms in numerous studies.⁸⁷⁻⁹²

Traditionally, anticonvulsants were commonly used for treating seizures.⁹³ Seizures are caused by synchronized neuron populations firing abnormally and excessively.^{94,95} Anticonvulsants reduce this synchrony via 3 main mechanisms: (1) modulation of voltage-gated ion channels; (2) upregulation of inhibitory neurotransmitters; (3) downregulation of excitatory neurotransmitters.⁹⁶ However, anticonvulsants like sodium valproate and carbamazepine are also involved in longer term mechanisms such as modulating cellular plasticity and resilience. This is achieved through regulation of inositol biosynthesis (MIP synthase), cyclic adenosine monophosphate (c-AMP) response element binding protein, brain-derived neurotrophic factor (BDNF), mitogen-activated protein kinases and arachidonic acid pathway regulation, among others.⁹⁷ Furthermore, sodium valproate inhibits histone deacetylase, creating long-term gene expression changes thought to produce long-term mood stabilizing effects.⁹⁸

The exact pharmacodynamics of anticonvulsants is specific to each drug and can include several of the aforementioned mechanisms.⁹⁶ Nonetheless, all drugs have similar effects on stabilizing mood in HD patients.^{21,99-101}

Lamotrigine. Lamotrigine also blocks voltage-gated sodium ion channels, however, it also suppresses excitatory

neurotransmitters glutamate and aspartate.¹⁰² Although lamotrigine is used as a mood stabilizer in HD, some studies suggest its suppression of excitatory neurotransmitters may reduce neuron excitotoxicity, thus, mitigating one of the pathological symptoms in HD.^{21,103} A double blind, placebo-controlled study was completed on 64 early diagnosed HD patients (<5 years) over 30 months. The effectiveness of treatment was analysed using a quantified neurological examination and a set of cognitive and motor tests. Lamotrigine did not significantly slow the progression of HD over the 30 months. Therefore, although beneficial against 1 symptom, lamotrigine does not possess neuroprotective attributes.^{103,104}

Carbamazepine. Carbamazepine's mechanism of action is solely through the blockade of voltage-gated sodium ion channels. However, the exact pharmacodynamics of the drug has not been fully elucidated and there is still some debate within the literature around this.^{105,106} Although, carbamazepine is prescribed as a mood stabilizing drug for HD, no clinical research was found on its specific benefits to HD patients.²¹

Anticonvulsant drugs exhibit a number of side effects. General side effects can include hypersensitivity reactions, blood dyscrasias, dizziness, gastrointestinal disturbances, depression and hyponatremia.²¹ Aside from these side effects, carbamazepine also has the potential to cause serious skin conditions such as Stevens-Johnson syndrome or toxic epidermal necrolysis.¹⁰⁷ This could suggest why carbamazepine is prescribed less often than other anticonvulsants.²¹

Drug treatment can be beneficial to HD patients, although prescribed drugs should be based on concurrent symptoms as each case is unique. Patients are required to be closely monitored when taking medications and management should be patient specific. Additionally, the side effects of each drug mentioned also need to be carefully recorded and ensured they are balanced with their benefits.

Non-Invasive Strategies and Life-Style Adaptations

The overall treatment of HD requires a holistic approach to health, involving a range of health professionals.¹⁰⁸ Non-invasive strategies and life-style adaptations can also play a vital role in the patient's management of HD. Chorea is the primary symptom that can severely impair balance and gait. Physiotherapists can support patients by extending the proper function of these actions as well as assessing when walking aids or wheelchairs are required.^{109,110} Similarly, occupational therapists can assist HD patients through completing home assessments and installing bathroom adaptations or handrails so patients can prolong living in the comfort of their own home.¹¹¹

Patients who exhibit motor symptoms can also develop speech impairments such as hyperkinetic dysarthria. Hyperkinetic dysarthria involves involuntary muscle movements in the mouth, throat and respiratory system, resulting in voice

hoarseness and abnormal prosody.¹¹² For these cases, patients can receive speech and language therapy to optimize their verbal communication for as long as possible. If a patient presents with an inability to speak, speech and language therapists can provide communication charts or electronic communicating devices, enabling patients to continue conversing with friends and family.^{113,114}

Malnutrition is another major issue for HD patients, with the majority having a lower than average body weight.¹¹⁵ Dieticians can help patients in these circumstances, providing them with diet plans and methods to overcome the chorea-induced eating deficits such as blending foods or providing liquid food substitutes.¹¹⁵ A Percutaneous Endoscopic Gastroscopy (PEG) may be performed for some patients in which swallowing is severely affected.¹¹⁶

To alleviate the psychological and cognitive symptoms associated with HD, sessions with a psychologist can be used alongside the medications provided. This may not only improve the patient's mental health, but also can be used to monitor how the patient is responding to each medication they are receiving.²¹

Surgical Treatments

An alternative surgical treatment is deep brain stimulation.^{117,118} This intervention is primarily used to improve chorea symptoms.^{119,120} A study from Gonzalez et al. (2014), found deep brain stimulation to be effective at reducing chorea in pharmacologically resistant chorea HD patients. It did not, however, reduce dystonia or bradykinesia.¹²¹ As pharmacological resistant HD is rare, the use of this intervention is rare. The infrequent use of this treatment can also be attributed to the additional risks associated with the invasive procedure.¹²²

Overall, treatment of HD should not solely be the clinician's responsibility and the medications they prescribe. For optimal treatment, a range of health care professionals should be involved with an extensive network of communication between them.¹²³

Potential Future Therapeutic Options for Huntington's Disease

Antisense Oligonucleotide (ASO) Therapies

During the past decade, attention has moved from targeting broad spectrum and downstream processes to targeting root causes for potential HD therapies [Table 2](#), such as lowering mHTT ([Figure 1](#)).³⁰ ASO therapies hold significant attention in potential HD therapeutics, with multiple ASOs currently in clinical trials.²⁵ ASOs are single stranded oligonucleotide analogues.³¹ They bind to pre-mRNA or mRNA and can act through multiple mechanisms, including RNA degradation, blocking translation and splice modulation¹²⁴; these ultimately alter protein expression. ASOs can be allele-specific, indicating they specifically target mHTT, or allele non-specific, indicating they target both wild type HTT (wtHTT) and mHTT. There




Figure 2. Tominersen and WVE-120101/WVE-120102 therapy development timeline.

are 3 ASOs in clinical trials currently for HD – Tominersen (an allele non-specific ASO), WVE-120101 and WVE-120102 (allele-specific ASOs)³⁴ (Figure 2).

Tominersen, WVE-120101 and WVE-120102 function by complementary base pairing to target pre-mRNA, inducing RNase H1-mediated degradation. They reduce mHTT mRNA levels, therefore, lowering mHTT protein synthesis. Administration occurs via intrathecal injection into the cerebrospinal fluid (CSF) via lumbar puncture, allowing distribution to the central nervous system.^{125,126}

Tominersen. Hoffman-La Roche's RO7234292 or Tominersen (previously called IONIS-HTTRx/RG6042; IONIS Pharmaceuticals) is an ASO that binds to wild type HTT (wtHTT) and mHTT mRNA, inducing degradation.¹²⁶ It is currently the most developed HD ASO therapeutic.²⁷

Tominersen distributed well in mice and non-human primate brains during pre-clinical experiments. It lowered mHTT mRNA and mHTT protein in a dose-dependent, sustained fashion. Tominersen reversed the HD phenotype, improved survival and reduced brain atrophy in mice.¹²⁷ These experiments informed the following phase 1b/2a clinical trial design.

In the initial randomized, double blinded phase 1b/2a clinical trial (NCT02519036), participants (n = 46) were divided into 5 groups of ascending doses and a placebo group. Tominersen was given every 28 days in 4 doses, and participants were followed up after 4 months.¹²⁸ No serious adverse events occurred, and minor adverse event incidence was similar for

Tominersen and placebo groups. CSF mHTT was reduced by 40% on average at 90 mg and 120 mg doses and Tominersen showed dose dependency.^{129,130}

Multiple trials have commenced since NCT02519036 began. A randomized phase 2 open label extension (OLE) trial (NCT03342053) further assessed adverse events, recruiting people (n = 46) that had participated in the NCT02519036 trial.^{131,132} Participants were given monthly or bimonthly Tominersen 120 mg doses for 15 months.¹²⁹ The trial concluded in 2019, with results yet to be published.

GENERATION HD1 (NCT03761849) is a randomized, double blinded phase 3 trial to measure HD phenotype changes using the Composite Unified Huntington's Disease Rating Scale (cUHDRS) and total functional capacity (UHDRS-TFC), among other measurements. Researchers will divide participants (n = 909) with manifest HD into 3 groups – placebo, and Tominersen doses (120 mg) every 8 or 16 weeks. Dosing was planned to occur for 25 months, and follow up would have been 29 months.^{28,133} GENERATION HD1 was expected to conclude in 2022,²⁸ however, was suspended upon independent data review in early 2021 due to Tominersen's benefit to risk ratio over time. No new safety signals arose during this review.²⁹

A randomized phase 3 OLE clinical trial (NCT03842969/GEN-EXTEND) including previous Tominersen trial participants (N = 1100) will measure long-term tolerability, behaviour changes and adverse event incidence.^{131,134} Participants will receive Tominersen every 8 or 16 weeks, depending on the dose timing of their previous trial. It was expected to conclude in

June 2024,¹³⁴ but has been placed on hold due to another trial's benefit to risk ratio.²⁹

GEN-PEAK (NCT04000594) is a non-randomized, open label phase 1 trial investigating Tominersen's pharmacokinetics. It will conclude in 2021. Participants (N = 20 in total) in 3 groups of ascending doses will receive 2 doses 28 days apart. CSF Tominersen and mHTT concentrations, and plasma Tominersen concentrations will be measured.¹³⁵

WVE-120101 and WVE-120102. Wave Life Sciences' WVE-120101 and WVE-120102 are allele-specific, stereopure ASOs. They target single nucleotide polymorphisms (SNPs) specific to HD genotypes – WVE-120101 targets rs362307 (SNP1), and WVE-120102 targets rs362331 (SNP2).¹³⁶ They are allele-specific, which is thought to potentially avoid long-term negative effects since they do not lower wtHTT.¹³⁷ As they target SNPs, WVE-120101 and WVE-120102 cannot treat all HD patients – but combined use could target 80% of European HD patients.¹²⁵ Unpublished pre-clinical experiments showed they reduced mHTT mRNA without significantly altering wtHTT mRNA and protein in patient derived cell lines.¹³⁸ This led to randomized, double blinded phase 1b/2a clinical trials (PRECISION-HD1 and PRECISION-HD2; NCT03225833, NCT03225846), which recruited participants with early manifest HD and rs362307 (SNP1) or rs362331 (SNP2). Participants (n = 60 per trial) were divided into 6 groups – 2 mg, 4 mg, 8 mg, 16 mg or 32 mg WVE-120101/WVE-120102 doses or placebo (.9% NaCl).^{126,136} The parallel studies intended to measure adverse events, tolerability, pharmacokinetics and pharmacodynamics.¹³⁶

Preliminary PRECISION-HD2 results showed WVE-120102 reduced CSF mHTT protein by 12.4% compared to placebo across all groups (n = 44). This was statistically significant in higher doses, and indicated dose dependency. Total HTT did not change between groups. There were no serious adverse events, and less minor and moderate adverse events for WVE-120102 (72%) than placebo (83%).¹²⁶ However, in early 2021 Wave Life Sciences announced PRECISION-HD2 trial data showed WVE-120102 caused no change in mHTT compared to placebo. PRECISION-HD1 data showed similar results, therefore, both trials were stopped.¹³⁹

OLE trials began for patients who participated in the PRECISION trials (NCT04617847 for WVE-120101, NCT04617860 for WVE-120102). They received multiple 8 mg or 16 mg doses of WVE-120101/WVE-120102.^{126,138} OLE results for PRECISION-HD2 indicated slight but inconsistent mHTT lowering, without the ability to consistently lower mHTT with increased doses. Dosing in OLEs has stopped for WVE-120101 and WVE-120102 given their inability to significantly lower mHTT and therefore be of clinical benefit.¹³⁹

Wave Life Science's WVE-003 is a pre-clinical ASO that targets an unspecified SNP (SNP3) in mHTT mRNA.¹⁴⁰ WVE-003 reduced mHTT mRNA in vitro and achieved

mHTT mRNA knockdown in vivo.¹⁴¹ Application for a clinical trial was submitted in late 2020.¹⁴⁰

Other ASOs. BioMarin's (CUG)7 is a pre-clinical allele-specific ASO which targets expanded CAG repeats in mHTT mRNA.³¹ After promising results in HD patient derived fibroblasts,¹⁴² researchers found CUG (7) reduced mHTT protein concentration in R6/2 and Q175 HD model mice by 15-60%, improved motor symptoms and increased cortical volume.¹⁴³ R6/2 mice are transgenic HD models created by expressing exon one of the human *mHTT* gene, containing 150 CAG repeats.¹⁴⁴ Q175 HD knock-in mice express human mHTT within the mouse HD gene, containing 179 CAG repeats.¹⁴⁵

A number of further pre-clinical HD ASO therapeutics are currently in development. Some groups are investigating ASOs that target SNPs associated with HD alleles, therefore, aiming to selectively suppress mHTT expression.^{146,147} Results in HD mice model neurons,¹⁴⁶ and in vivo using HD model mice¹⁴⁷ were positive. Total HTT reducing and mHTT specific ASOs have improved cognitive and behavioural deficits in HD model mice with safety and tolerability, and reduced HTT in the cortex and limbic structures in non-human primate brains (where cognitive and behavioural symptoms largely originate).¹⁴⁸ Another ASO induces exon 12 skipping in HTT pre-mRNA, preventing cleavage sites being expressed and therefore 586 amino acid N-terminal huntingtin fragments being created. Rodent studies showed successful exon skipping, a shorter HTT protein and reduced 586 amino acid N-terminal huntingtin fragments.¹⁴⁹

Some ASOs are non-HTT-targeting, such as Triplet Therapeutics' TTX-3360. It targets MSH3 within the DNA Damage Response pathway, which is often involved in repeat expansion. TTX-3360 is safe and tolerable in HD model mice, and 50% MSH3 knockdown reduced or inhibited mHTT repeat expansion. An Investigational New Drug/Clinical Trial Application submission is planned for late 2021.¹⁵⁰

RNA Interference (RNAi) Therapies

RNAi therapies use transgenes that express RNA molecules such as micro-RNA (miRNA), short hairpin RNA (shRNA) and short interfering RNA (siRNA).^{26,31} Cellular machinery processes these molecules, and they label their target mRNA for degradation. Degradation occurs via enzymes in the RNA-induced silencing complex (RISC),³¹ which stops mRNA translation and therefore protein synthesis.²⁶ RNAi therapies are delivered by a viral vector via intracranial injection.³¹ There are advantages and disadvantages associated with RNAi therapies. For example, shRNA therapies can result in difficulty controlling shRNA expression, but shRNA therapies also tend to produce less off target effects than other RNAi therapies and last longer.¹⁵¹⁻¹⁵³ These factors, and other factors associated

with miRNA and siRNA, must be considered when developing RNAi therapies.

AMT-130. AMT-130 (UniQure Biopharma B.V., named AAV5-miHTT in pre-clinical studies) is a RNAi based gene therapy. It is the only gene therapy currently in clinical trials for HD.²⁷ AMT-130 contains a gene encoding for a miRNA, which is delivered via adeno-associated virus vector serotype 5 (AAV5). AMT-130 activates RISC, which stops mHTT (and wtHTT) protein synthesis. AMT-130 is administered via intrastriatal injection.²⁶

In an HD rat model, Hu128/21 HD mice and HD patient derived neurons, AAV5-miHTT reduced mHTT mRNA and protein levels.¹⁵⁴⁻¹⁵⁶ Hu128/21 HD mice result from crossing an HD mouse model, YAC128, with a wtHTT model, BAC21, giving them 2 full-length human HTT transcripts that are heterozygous for the HD mutation.¹⁵⁷ Potent mHTT suppression was sustained for 7 months post-injection in Hu128/21 HD mice.¹⁵⁸ In Q175 HD mice, AAV5-miHTT lowered human HTT protein and striatal (39%) and cortical mHTT aggregates 12 months post-injection. R6/2 HD mice showed motor symptom improvement and prolonged survival.¹⁵⁹ mHTT aggregates were suppressed in an HD rat model – reducing neuronal dysfunction.¹⁵⁵ AAV5-miHTT lowered human mHTT mRNA (72.8%) and protein (85.3%) in the minipig brain, and indicated dose dependency.¹⁶⁰ AAV5-miHTT did not create off target effects caused by gene dysregulation or incorrect processing in HD patient derived neurons,¹⁵⁶ but at high doses caused astrogliosis in Hu128/21 mice.¹⁵⁸ AMT-130 showed safety, tolerability and widespread distribution in the brain of non-human primates.¹⁶¹

Phase 1b/2a clinical trials will test AMT-130's safety and proof of concept in 26 manifest HD patients (NCT04120493). Some will be given a low or high dose via MRI guided bilateral intrastriatal injection, and others will be given placebo (imitation surgery).²⁷ Researchers are measuring adverse effects, CSF biomarkers, CSF AMT-130, mHTT concentration and clinical scale changes (like UHDRS) during the core study period and follow up.¹⁶² Data from the initial 6 months of dosing in 2 patients, and 90 day data from the next 2 patients indicated AMT-130 was safe.¹⁶³ This prompted complete enrolment of the first dosing cohort (n = 10). The study is expected to be completed in 2026, with an OLE beginning in late 2021.¹⁶⁴

Other RNAi therapies. Voyager Therapeutics' VY-HTT01 is an allele non-specific RNAi based gene therapy. An miRNA is expressed and delivered via adeno-associated virus vector serotype 1 (AAV1), targeting HTT mRNA degradation. It is administered via intracranial injection.³¹ In YAC128 transgenic HD mice, which express full-length human HTT with 128 CAG repeats, VY-HTT01 reduced striatal HTT protein by

40%, partially corrected motor and behavioural symptoms, reduced HTT aggregates and showed effective striatal distribution.^{165,166} Unpublished results showed VY-HTT01 lowered striatal (67%) and thalamic (73%) HTT mRNA in non-human primates.¹⁶⁷ Clinical trial applications were submitted in late 2020, but are on hold to resolve chemistry, manufacturing and control issues.¹⁶⁸

Spark Therapeutics are also developing an RNAi based drug, using an AAV1 delivered shRNA (AAV.shHD2.1)³¹. Rodent studies showed motor symptom improvement, reduced human mHTT expression¹⁶⁹ and neuroprotection on administration with AAV.shHD2.1.¹⁷⁰ In non-human primates, lowering HTT expression via RNAi did not lead to neuronal degeneration or an immune response for 6 weeks post-injection into the putamen, strengthening RNAi as a potential HD therapy.¹⁷¹ Clinical trials have not yet been initiated.³⁴

RNA Targeting Small Molecule Therapies

Many small molecule therapies alter pre-mRNA to include or exclude target exons, leading to an altered protein (also called splice modulation). They are administered orally, which is advantageous compared to other invasive administration methods. However, this also increases the risk of adverse effects due to their systemic distribution and lack of specificity.²⁵ All HD RNA targeting small molecule therapies are either pre-clinical or in phase 1 clinical trials currently, with no clinical trial data yet published.³⁴

PTC Therapeutics' PTC518 is a RNA splice modulator that induces a pseudoexon inclusion containing a premature stop codon. This enhances mHTT mRNA degradation – resulting in lowered mHTT protein. Unpublished pre-clinical data showed PTC518 is systemically distributed (including widespread central nervous system distribution) and indicated dose dependency. In HD patients' isolated cells PTC518 lowered mHTT mRNA and protein, and in transgenic BACHD HD mice, which express the full-length human mHTT with 97 mixed CAA-CAG repeats within exon 1, PTC518 systemically lowered HTT protein.¹⁷²⁻¹⁷⁴ A phase 1 trial has commenced with healthy participants, using ascending single and multiple doses. It is expected to indicate correct dosing, proof of concept and safety. Proof of concept was shown in healthy volunteers with dose-dependent HTT mRNA splicing. It was also well tolerated and showed predictable pharmacology.^{173,175}

Novartis Pharmaceuticals' Branaplam (also called LMI070) is an oral, weekly administered small molecule RNA splice modulator. It was initially designed to treat spinal muscular atrophy (SMA),¹⁷⁶ but researchers found it lowered mHTT protein in SMA and HD animal models. It also lowered HTT mRNA in HD models. A phase 2b clinical trial with manifest HD participants is planned for in 2021.¹⁷⁷

Nuredis are developing an RNA splice modulator that specifically targets transcription elongation cofactors such as Spt4.^{34,178}

These are required for expanded CAG repeat transcription. Therefore, Spt4 inhibition would result in mHTT protein reduction. Pre-clinical data showed Spt4 inhibition using small molecules reduced expanded polyQ proteins in neurons, while restoring neuronal function and regional enzymatic activity. SUPT4H (a mammalian Spt4 ortholog) also reduced mHTT protein, toxicity and mHTT aggregates in neurons.¹⁷⁸ In R6/2 HD mice, SUPT4H reduction prolonged survival, reduced motor symptoms and reduced mHTT aggregation. Further studies conducted in zQ175 HD mice showed SUPT4H reduced mHTT mRNA and protein.¹⁷⁹

There are other groups that are searching for and developing RNA targeting small molecules to treat HD – including molecules modulating protein disulfide isomerase in the endoplasmic reticulum to reduce mHTT toxicity,¹⁸⁰ molecules altering degradation or autophagy mechanisms¹⁸¹ and molecules targeting expanded CAG repeat RNA to reduce mHTT protein synthesis.¹⁸²⁻¹⁸⁴ CAG lengths can be somatically unstable in different tissue types. The mechanism underpinning this is unknown. Understanding and targeting this somatic instability may be a pathway to future therapies.¹⁸⁵

Zinc-Finger Protein (ZFP) Therapies

DNA targeting therapies affect the mHTT gene itself by introducing protein coding sequences to the brain parenchyma that bind to DNA regions. They require a viral vector for delivery and are administered intracranially.³¹ ZFPs are synthetically made sequences that mimic naturally occurring motifs. They contain DNA binding components that often include multiple zinc finger peptides, which can bind to 3 to 5 target DNA nucleotides. The major ZFPs currently being developed for HD bind to expanded CAG repeats, preventing the mHTT gene's transcription. This lowers mHTT protein levels without altering the gene itself.²⁵ However, zinc finger nucleases (ZFNs) can alter genetic material, allowing for the mHTT gene to be corrected or disrupted.³⁴ ZFPs have successfully reduced mHTT protein by 60% in the R6/2 HD mice model brain, reduced aggregation and improved motor symptoms.¹⁸⁶

There are 2 major ZFPs in development for HD – TAK-686 and ZF-KOX1.¹⁸⁶⁻¹⁸⁸ These therapies are both in pre-clinical stages currently.²⁵ TAK-686, which targets expanded CAG repeats, has shown success in reducing mHTT in knock-in HdhQ50 mice containing an HD allele with 50 CAG repeats. It also reduced mHTT by 62% and improved behavioural symptoms in R6/2 HD mice.^{188,189} A modified version of ZF-KOX1 (for mouse compatibility) delayed HD symptoms and inhibited mHTT expression by 77% in HD model mice.¹⁸⁷

Transcription Activator-like Effector Nuclease (TALEN) Therapies

TALENs have DNA binding domains that contain repeating peptides which bind to DNA nucleotides.³⁴ They cause double

strand breaks via artificial nucleases, allowing for segment deletion or correction.¹⁹⁰ All TALENs in development for HD therapy are in pre-clinical or drug discovery stages. Their impacts on phenotype, toxicity and an effective delivery system have not been established yet. TALENs also have not been investigated in HD animal models.¹⁹¹ However, TALENs have successfully removed expanded CAG repeats in yeast cells by inducing double strand breaks, resulting in gene conversion. They did not increase mutation rate or cause genomic rearrangement.¹⁹²

Lowered mHTT gene expression and protein aggregation was also achieved in HD patient derived fibroblasts using a TALEN and SNP specific transcription activator-like effector (TALE-SNP). The TALE-SNP prevented transcription of the mHTT gene by targeting SNPs associated with it, leaving wtHTT gene expression unaltered. TALENs were used for gene correction by creating double strand breaks which were repaired – deleting the expanded CAG repeats. Results showed that some of the TALE-SNPs reduced mHTT expression by up to 20% without altering normal HTT expression, indicating proof of concept.^{25,191}

Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR)/CRISPR-Associated System (Cas) Therapies

Cas9 (CRISPR associated protein 9) is a naturally occurring immune enzyme found in some bacteria, which has been utilized as a genome editing tool for the past decade. It functions via a single-guide RNA (sgRNA) binding to its target DNA, and Cas9 nucleases inducing double strand breaks.³⁴ Double strand breaks are repaired by error-prone non-homologous end joining (NHEJ), causing frameshifts which can impair gene expression.^{190,193} All CRISPR approaches are delivered with viral vectors and intracranial injection.³⁴ In HD therapeutics, CRISPR/Cas9 therapies may remove expanded CAG repeats to correct HD alleles, inactivate HD-associated alleles or target the HTT gene itself, non-specifically reducing HTT. Multiple groups are investigating CRISPR pre-clinically for HD.¹⁹³⁻¹⁹⁸

CRISPR/Cas9's efficiency has been assessed in cell lines. A paired Cas9 Nickase strategy successfully excised expanded CAG repeats in exon 1 of the HTT gene in 3 distinct HD patient derived fibroblast lines, inactivating *HTT*. Nickases cause single strand breaks, reducing off target effects due to increased specificity. Each cell line treated with Cas9 Nickases had different CAG repeat lengths and showed approximately 70% reduction in HTT protein.¹⁹³ A strategy using CRISPR-Cas9 and *piggyBac* transposon was also successful in HD human induced pluripotent stem cells, correcting the mHTT allele and amending HD mediated phenotypic defects.¹⁹⁴

A PAM-altering, mHTT associated SNP based CRISPR/Cas9 approach is also in development. This therapy induces a deletion mutation that inactivates the mHTT allele, leaving an intact wtHTT allele. In HD patient derived cell lines, excision of CAG repeats associated with mHTT alleles was successful.¹⁹⁵

Some groups have investigated CRISPR in HD animal models. Deletion of the mHTT polyQ domain using an allele non-specific CRISPR/Cas9 based strategy successfully repressed human mHTT expression in HD140Q-knock-in mice's striatum, which express full-length human mHTT with exon 1 containing 140 CAG repeats. CRISPR was administered via striatal injection. This approach reduced mHTT expression, mHTT aggregates, astrocyte reactivity and improved motor symptoms.¹⁹⁶ Another group injected SpCas9 or sgRNAs delivered by recombinant AAVs (abbreviated to rAAV.SpCas9 and rAAV.sgHD1/i3, respectively) into the right hemisphere of BACHD mice. Human HTT exon 1 was deleted, reducing human mHTT mRNA by 40%. The left hemisphere was used as a control, and did not demonstrate lowered mHTT expression.¹⁹⁸ Another group investigated in vivo striatal injection of CRISPR/Cas9 therapies (AAV1-SaCas9-HTT to target human HTT, or AAV1-SaCas9-mRosa26 as a control) in R6/2 HD mice. They found AAV1-SaCas9-HTT had reduced mHTT protein inclusions by 40%, reduced total mHTT protein by 50%, and it improved lifespan and motor symptoms.¹⁹⁷

Stem Cell Therapies

Stem cell therapies are another candidate for future HD therapeutics (Figure 3). Their main advantages are their

potential ability to replace neurons lost due to HD pathology, improve regeneration and provide pro-survival factors. Furthermore, theoretically some long-term benefits would remain due to stem cell incorporation into the patient's brain micro-environment. However, there are also concerns these therapies may induce immune reactions or rejection.¹⁹⁹

Cellavita HD. Cellavita HD is the most developed stem cell therapy for HD currently²⁷ (Figure 3). It is a mesenchymal stem cell therapy, and is administered via intravenous injection.²⁰⁰

Azidus Brasil began a non-randomized, open label phase 1 clinical trial in 2017 to test safety and effectiveness of their stem cell therapy for HD, with completion expected in 2023 (NCT02728115/SAVE-DH).^{27,200} Participants (n = 6 total) with manifest HD will be given 3 low or high monthly Cellavita HD doses. They will be followed up for 5 years to track adverse events and efficacy through the UHDRS scale and inflammatory markers, among other measurements.²⁰⁰

A randomized, triple blind phase 2 clinical trial began in 2018 to test Cellavita HD's dose-response relationship (NCT03252535/ADORE-DH).^{27,201} Thirty-five participants with manifest HD will be given 9 total low or high doses of Cellavita HD or placebo. Researchers are primarily measuring




Figure 3. Cellavita HD, foetal striatal stem CRT and autologous stromal/adipose-derived stem cell therapy development timeline.

UHDRS to assess effective dosing. This trial will conclude in 2022.²⁰¹

An OLE phase 2/3 trial has also been registered, but is not yet recruiting (NCT04219241/ADORE-EXT). It will assess longer term safety and efficacy in 35 manifest HD participants who took part in the ADORE-DH trial.^{27,202} They will receive multiple doses over 24 months. UHDRS, TFC, neurological image improvement and clinical worsening will be measured, among other factors. It will conclude in 2022.²⁰²

Foetal striatal stem cell replacement therapy. Foetal striatal stem cell replacement therapy was first investigated in HD patients over 20 years ago. Multiple early studies showed foetal striatal cell replacement therapy (CRT) was safe but relatively ineffective at altering patient symptoms. A phase 1, randomized controlled trial using unilateral intrastriatal grafts in 4 HD patients indicated no adverse events but no HD symptom improvement (ISRCTN36485475).²⁰³ Similarly, a later trial using bilateral foetal striatal CRT grafts showed the UHDRS score was similar across those with (n = 5) and without (n = 12) striatal grafts. The grafts did not cause significant adverse events.²⁰⁴

Some trials have shown HD symptom improvements. In 5 HD patients, asynchronous bilateral intrastriatal transplants of foetal striatal neuroblasts created increased striatal metabolic activity, cognitive and motor improvements in 3 of the 5 patients (compared to 22 control HD patients). However, the other 2 patients' condition declined similarly to the control group.²⁰⁵ Another group also found mild short-term symptom improvement in 6 patients, but this only persisted in 3 and did not alter overall HD progression.²⁰⁶

A randomized, open label, phase 1 clinical trial using a relatively higher CRT dose to previous studies was initiated in 2017. CRT will be administered to 5 HD patients via intrastriatal injection and patients will receive immunosuppressant medication (ISRCTN52651778/TRIDENT).^{27,207} TRIDENT aims to assess safety at 4 weeks, and the effectiveness of an increased CRT dose. The trial will conclude in 2023.²⁰⁷

However, foetal striatal CRT also raises concerns. Immunosuppressants used to reduce graft rejection have been problematic due to side effects and lowered compliance.²⁰⁸ Immune responses can occur due to foetal striatal stem cell grafts – over a decade after the ISRCTN36485475 trial, 1 of the 4 patients showed strong microglial activation, immune cell infiltration and mHTT protein aggregates at the graft site.²⁰⁹ However, while some groups believe immunosuppressants are required,²¹⁰ others have shown patients' immune responses after CRT grafts are unpredictable – therefore excessive immunosuppressant therapy is undesirable.²¹¹ Some groups have found that striatal foetal neural grafts can only provide clinical improvement for a few years, after which patients decline. Therefore, CRT may not be viable as a permanent cure.²¹² Ethical concerns also persist due to the use of foetal cells,

and graft survival is uncertain due to findings showing reduced vascularization, and lowered astrocytes and pericyte numbers.²¹³

Autologous stem cell therapies. Autologous stromal cells are also being investigated by Regeneris Medical to treat a range of neurological disorders in a current clinical trial (n = 300), including HD (NCT03297177).²⁷ Autologous stem cell transplantation requires reprogramming mutated cells or isolating cells with the wild type allele.²¹⁴ Pre-clinical data for autologous adipose derived stem cells transplanted bilaterally into the striatum were positive in R6/2 HD model mice. This approach reduced striatal apoptosis, prolonged lifespan, reduced mHTT aggregates and modulated striatal neuronal loss.²¹⁵

A non-randomized, placebo-controlled study will assess the safety and change in neurological function of those given autologous stem/stromal cells compared to normal saline for up to 5 years.²⁷ Autologous cells will be collected from patients' subdermal fat, isolated, concentrated and neutralized before being given intravenously. The study is expected to be completed in 2023.²¹⁶

Mesenchymal stem cell therapies. Mesenchymal stem cells (MSCs) have also been trialled in HD animal models. MSCs originate in the umbilical cord (UC), amniotic fluid, bone marrow and adipose tissue. They are self-renewing, can release neuroprotective factors like BDNF and can differentiate into multiple cell types, potentially including neural cells.^{217,218} No teratomas or overgrowths have been found in MSC animal model studies.²¹⁷ Striatal injection of mice UC-derived MSCs in R6/2 HD mice improved spatial memory and reduced neuropathological deficits, but did not change motor impairments.²¹⁹ Striatal transplantation of UC-derived rat MSCs in 3-nitropropionic acid (3-NP) lesioned HD rats improved motor coordination, reduced striatal atrophy, prevented oxidative stress in PC12 cells and enhanced cell viability.²²⁰ 3-NP HD rats were systemically, subcutaneously administered with 3-NP, a mitochondrial toxin, to produce striatal lesions.²²¹ Another 3-NP HD rat study found injection of bone marrow derived rat MSCs resulted in no significant striatal inflammation, no lateral ventricle enlargement, reduced neurological and behavioural deficits and the presence of striatal neurotrophic factors.²²²

Induced pluripotent stem cell-derived neural stem cells. Induced pluripotent stem cell-derived neural stem cells (iPS-NSCs) are self-renewing, reprogrammed somatic adult cells that can differentiate into many cell types.²²³ Striatal injection of mice iPS-NSCs in an HD rat model was trialled and monitored using serial ¹⁸F-FDG PET/CT. Researchers found that the rats' striatal volume was partially recovered, memory, learning and glucose metabolism improved and iPS-NSCs differentiated into striatal neurons and glia.²²⁴ Quinolinic acid HD rats undergo striatal injections of quinolinic acid to induce an HD phenotype.²²⁵ Another study in a quinolinic acid-induced HD rat model found striatal transplantation with human iPS-NSCs




Figure 4. ANX005 and VX15/2503 antibody therapy development timeline.

reduced behavioural deficits, reduced inflammation, promoted neurogenesis and replaced some neural cells.²²⁶ Striatal injection with control mice iPS-NSCs into YAC128 HD mice resulted in motor improvement, increased striatal BDNF and iPS-NSC differentiation into neural cells.²²⁷

Antibody Therapies

Antibody-based therapies are not only being evaluated for a range of tauopathies and synucleinopathies, but are also emerging as a potential therapy for monogenic disorders of the central nervous system, including HD (Figure 4).

ANX005. Annexon, Inc’s ANX005 is a monoclonal antibody that targets C1q, preventing classical complement cascade initiation.²²⁸ When neuronal stress is already implicated during HD pathogenesis, C1q’s role in synaptic pruning can be disrupted – affecting synapse loss, neurodegeneration and neuroinflammation. Through C1q inhibition, ANX005 is thought to reduce synapse loss and neurodegeneration.^{229,230} ANX005 was initially intended to treat Guillain-Barré Syndrome (GBS) and Alzheimer’s Disease (AD).²²⁸

Pre-clinical studies showed high affinity binding of ANX005 to C1q in multiple species, and successful CSF and serum C1q reduction in monkeys and rats. Complement cascade inhibition in vivo and in vitro in GBS and AD mice models was successful. Safe dosing frequency, dose dependency and tolerability was also established in monkeys and rats.²²⁸ A murine version of ANX005 reduced neurofilament light chain and synapse loss in HD animal models.²²⁹

A randomized, placebo-controlled, double blinded phase 1 clinical trial began in late 2016 in 27 healthy volunteers via an ascending dose schedule (NCT03010046). ANX005 was given intravenously, either on its own or with intravenous immunoglobulin (IVIg). The study aimed to assess safety, pharmacokinetics and pharmacodynamics. It was terminated in 2018 since there were no adverse events at the maximum dose.²³¹ Subsequently, investigators began testing ANX005 for HD, GBS, Amyotrophic Lateral Sclerosis (ALS) and Warm Autoimmune Haemolytic Anaemia (wAIHA) patients.²³²⁻²³⁵

A phase 1b clinical trial in 31 GBS patients showed promising top-line results using an ascending dose schedule. Its aims were to test safety, tolerability and pharmacokinetics. The ANX005 dose was increased after safety was established in up to 100 mg/kg in 19 participants. No significant adverse events occurred, and ANX005 successfully inhibited C1q. ANX005

was associated with early muscle strength improvement, reduced neurofilament light chain and small GBS disability scale improvements.²³⁶ ANX005 is also in a phase 1b, open label clinical trial to assess safety in 12 GBS patients (NCT04035135),²³² and a randomized, double blinded phase 2/3 trial in 180 GBS patients to assess efficacy and safety (NCT04701164).²³⁷

There are also 2 phase 2, open label studies in progress for ALS and wAIHA (NCT04569435, NCT04691570, respectively). ANX005 will be given to 24 participants with ALS for 12 weeks (NCT04569435), and given to 12 patients with wAIHA for 2 doses (NCT04691570). Both studies will test ANX005's safety, tolerability, pharmacodynamics and pharmacokinetics. The ALS trial will end in late 2021, and the wAIHA trial in early 2022.^{234,235}

An open label phase 2a clinical trial will evaluate ANX005's safety, tolerability, pharmacokinetics and pharmacodynamics in HD patients or patients at risk for HD (n = 24) (NCT04514367). Seven ANX005 infusions will be given for up to 10 weeks, then patients will be followed for 3 months. The study will be completed in late 2021.²³³

VX15/2503. Vaccinex's VX15/2503 (also called Pepinemab) is an IgG4 monoclonal antibody that inhibits protein semaphorin 4D (SEMA4D). SEMA4D is a protein associated with neuroinflammation by promoting glial cell activation and activating signalling cascades causing oligodendrocyte and neural precursor cell apoptosis, among other processes.²³⁸

Pre-clinical studies showed that VX15/2503 did not induce toxicity or adverse events in rats or non-human primates. It lowered T-cell SEMA4D and induced long-term saturation with 5 weekly doses. Dose dependency was established.²³⁸ Anti-SEMA4D is not immunosuppressive and improves blood brain barrier integrity.²³⁹ Another study found that anti-SEMA4D improved cortical and striatal atrophy, and cognitive symptoms in YAC128 HD model mice. It did not influence motor symptoms.²⁴⁰

Vaccinex, Inc and the Huntington Study Group conducted a double blinded, phase 2 clinical trial testing VX15/2503's safety, tolerability and pharmacokinetics in late prodromal and early manifest HD patients (NCT02481674/SIGNAL). Patients were divided into 2 cohorts, A (12 months treatment) and B (36 months treatment), and given intravenous monthly VX15/2503 doses or a placebo. Investigators assessed MRI brain volume, motor and cognitive function and CSF biomarkers, among other measures. The study ended in mid-2020.^{241,242}

Preliminary SIGNAL results from Cohort A (n = 36) showed VX15/2503 was tolerable and safe. Reduced metabolic activity and brain volume reduction appeared to be prevented. Cohort A were given VX15/2503 for another 5 months (open label), with a 3 month follow up.²⁴³ However, results from cohort B1 with early manifest HD (n = 179) found that VX15/2503 did not prevent brain volume and metabolic activity reduction, or significantly alter motor, cognitive or behavioural function.²⁴⁴ Trends towards cognitive benefits

were found, and post-hoc analysis using Clinical Global Impression of Change in an advanced HD subpopulation confirmed this. However, these findings require further investigation.²⁴⁵

VX15/2503 is also in clinical trials for AD, head and neck cancer and colorectal cancer,²⁴⁶⁻²⁴⁹ and has been tested in Advanced Solid Tumours, Multiple Sclerosis (MS), melanoma, non-small cell lung cancer.²⁵⁰⁻²⁵² Advanced Solid Tumour (NCT01313065) (N = 42) and MS (NCT01764737) (N = 10) trials found VX15/2503 was safe and tolerable.^{250,251}

Anti-mHTT antibody therapies. Anti-mHTT antibodies have been tested pre-clinically for over 20 years. Their intracellular targets, such as the PolyQ, PolyP and N-Terminal exon 1 domains, are affected using intrabodies.³² Intrabodies are simpler forms of antibodies delivered via viral vectors that are able to block and inactivate proteins, enhance degradation and stop protein misfolding intracellularly.²⁵³

PolyQ targeting intrabodies have had negative results, but PolyP/proline-rich region and N-Terminal exon 1 domain targeting intrabodies have had mixed results. rAAV6-INT41 (developed by Vybrion Inc) is a PolyP/proline-rich region targeting intrabody that can lower gene dysregulation caused by mHTT.^{32,254} It was delivered in R6/2 HD mice by a viral vector (rAAV6) via bilateral intra-striatal injection. INT41 can reduce small mHTT aggregates by 31% and larger aggregates by 16%, while improving cognitive deficits in R6/2 HD mice.²⁵⁴

Intrabodies targeting mHTT aggregates have had positive results, but raise concerns since conformational antibodies can also increase aggregation.³² W20 is an oligomer specific intrabody that targets mHTT aggregates, among other aggregates, such as α -synuclein and amyloid beta.^{32,255} BACHD mice given W20 via intracerebroventricular injection showed mHTT aggregate reduction in the striatum (by 49.5%) and cortex (by 26%), and lowered apoptotic cells in the striatum (by 73.3%) and cortex (by 69.6%). W20 lowered neuroinflammation, proinflammatory cytokine levels and reactive oxygen species, while improving motor and memory deficits. W20 does not affect WT HTT or surveilling microglia or astrocytes.²⁵⁵

Other intrabodies have also been tested pre-clinically, such as HAPP1 and VL12.3. These target the P-rich epitope of the mHTT protein, and the N-terminal 17 amino acids, respectively. They were found to lower HTT exon 1 toxicity and aggregation.²⁵⁶ C4 also targets the N-Terminal of HTT fragments.²⁵⁷ When fused with mouse orthenine decarboxylase including a deleted C-terminal PEST motif (mODC-PEST), HD exon 1 fragments are reduced, along with lowered HD exon 1 fragment toxicity, aggregation and intrabody degradation.²⁵⁸

Arguably antibody therapies' main advantage is their ability to target extracellular mHTT, since most other therapies cannot do so. This is achieved via active (self-produced antibodies) or passive (pre-made antibodies) immunization.²⁵⁹ Active




Figure 5. Non-RNA targeting small molecule (Pridopidine, Laquinimod, VX-745 and Nilotinib) therapy development timeline.

immunization’s functional benefits are not yet known, and raise safety concerns due to increased immune gene dysregulation. Passive immunization has decreased mHTT in YAC128 mice.³² Furthermore, AFFiRiS/ mAB C6-17 effectively bound and demonstrated specificity to human mHTT in cell-based assays and immunohistochemistry. C6-17 targets a site within the HTT protein near the aa586 caspase-6 cleavage region, and has blocked extracellular mHTT uptake. Distribution of C6-17 was quick and widespread in HD model mice, showed reduced mHTT in the central nervous system and improved motor symptoms.^{260,261}

Antibody therapies may be most beneficial in combination with RNA based therapies to target both intracellular and extracellular mHTT.²⁵⁹

Other Small Molecule Therapies

Pridopidine. Pridopidine (also named TV-7820; formerly named ACR16 or Huntexil) is an oral small molecule developed by Prilenia

Therapeutics (previously developed by Teva Pharmaceutical Industries Ltd) for HD motor symptoms²⁶² (Figure 5). It is a dopamine stabilizer that binds to dopamine type 2 receptors, therefore, affecting striatal dopamine pathways involved in motor HD pathophysiology.^{262,263} It also activates the sigma-1 receptor, increasing BDNF production and supporting neuroprotection.^{262,264} Pre-clinical studies in R6/2 HD mice showed pridopidine prevented apoptosis, increased pro-survival molecules and reduced mHTT aggregate size – indicating neuroprotection. It also improved motor function.²⁶⁵ Pridopidine prevented medium spiny neuron loss in YAC128 HD mice.²⁶⁶

Pridopidine was first clinically investigated as an HD therapy in 58 HD patients in a randomized, double blinded trial (ACR16C007). It showed tolerability and safety. While cognitive scores did not differ between placebo and pridopidine, voluntary motor performance improved using pridopidine.²⁶⁷

Further trials assessed efficacy and safety of pridopidine. In a 26-week, double blinded phase 3 clinical trial (NCT00665223/MermaiHD), 437 HD patients were randomized to 45 mg or 90 mg pridopidine, or placebo. Modest motor improvements were found using UHDRS-total motor score (TMS), but this was a tertiary endpoint and other improvements (like the mMS) were not statistically significant.²⁶⁸

A 12-week double blinded phase 2/3 clinical trial (NCT00724048/HART) randomized 227 HD patients to 3 dose groups or placebo. No statistically significant modified motor score (mMS) improvements were found, but trends pointed towards voluntary motor function improvement at 90 mg.²⁶⁹ Both studies found acceptable tolerability and safety. Similarly, a 26-week, double blinded, phase 2 clinical trial found tolerability (NCT02006472/PRIDE-HD). 408 HD patients were randomized to 4 differing pridopidine dose groups or placebo. Investigators found no UHDRS change between placebo and pridopidine groups. A few serious adverse events occurred that likely related to HD symptoms, and 1 death during the trial was possibly affected by pridopidine administration.^{263,270}

The PRIDE-HD trial was extended from 26 to 52 weeks in 262 patients. At 52 weeks, those taking 45 mg of Pridopidine had a lowered TFC compared to placebo, indicating slowed HD progression. Patients administered 45 mg did not experience TFC decline for the entire year they were taking pridopidine. They also found Pridopidine also has a dose-specific effect.²⁷¹

These findings led to an OLE for the MermaiHD trial and for the HART trial (NCT01306929/OPEN-HART). Both studies found long-term tolerability and safety.^{272,273} In OPEN-HART, 225 HD patients were administered 45 mg Pridopidine twice daily over 36 months. The trial found HD progressed similarly between placebo and Pridopidine groups for the first 36 months (as measured by TFC and TMS).²⁷³ However, additional data showed that while TFC declined over the full 60 months, the rate of decline slowed for those taking pridopidine compared to placebo. TFC and TMS also remained stable between 48 and 60 months. However, the sample size had dropped to 33 patients by 60 months which could create confounding due to those remaining in the trial potentially having a less aggressive disease form.²⁷⁴ An OLE study was also initiated for PRIDE-HD participants (NCT02494778/OPEN PRIDE-HD)²⁷⁵

A double blinded, phase 3 clinical trial (NCT04556656/PROOF-HD) will investigate efficacy (via UHDRS-TFC score) and safety in 480 HD patients. Participants will be randomized to 45 mg Pridopidine twice daily or a placebo. The study will end in 2023 and an OLE is planned for PROOF-HD participants.²⁷⁶

Laquinimod. Laquinimod, developed by Teva Pharmaceutical Industries, is an orally administered immunomodulator therapy²⁷⁷ (Figure 5). It likely reduces central nervous system leukocyte infiltration and is neuroprotective by stimulating BDNF production, but its precise mechanism of action is unclear.²⁷⁸ Pre-clinical

studies showed laquinimod decreases microglial activation, reactive astrogliosis, astrocytic NF- κ B activation, proinflammatory cytokine production, proinflammatory changes and oligodendroglial apoptosis, among other neuroinflammatory processes.²⁷⁹ Laquinimod has been tested as an MS therapeutic for over 15 years,²⁷⁸ with phase 3 clinical trials showing positive efficacy results and confirming safety and tolerability (NCT00509145/ALLEGRO).²⁸⁰ YAC128 mice treated with laquinimod displayed improvements in motor function and motor learning and further, displayed a reduction in depressive-like behaviours suggesting it a possible therapy for HD patients.²⁸¹

Laquinimod was tested in a randomized, double blinded, 52-week phase 2 clinical trial assessing efficacy (via UHDRS-TMS and other clinical measurements), safety and caudate volume atrophy (NCT02215616/LEGATO-HD). Participants (N = 352 total) with HD received 0.5 mg, 1 mg or 1.5 mg laquinimod or placebo.^{277,282} The 1.5 mg dose was terminated in 2016 due to safety concerns prompted by a study on laquinimod for MS.²⁷⁷ LEGATO-HD did not show a statistically significant improvement in UHDRS-TMS, but caudate volume significantly improved. Laquinimod was safe and well tolerated.²⁸³ However, further analysis showed that astrogliosis and gliosis was reduced in laquinimod groups.²⁸⁴ Some Q-motor measurements were also statistically significantly improved in 0.5 mg laquinimod, however, multiplicity corrections were not performed.²⁸⁵

VX-745. EIP Pharma's VX-745 (also called Neflamapimod) modulates neuroinflammation and microglial dysfunction, and decreases microglial proinflammatory cytokine upregulation via p38 α MAPK inhibition.^{286,287} VX-745 has been investigated for over 20 years,²⁸⁸ but has only recently been trialed for HD²⁷ (Figure 5). In AD model mice, VX-745 reduced hippocampal interleukin-1 beta (IL-1 β) and improved cognitive performance – but these benefits' peaks occurred at different doses, indicating independent processes were occurring.²⁸⁷

EIP Pharma Inc and Voisin Consulting, Inc will fund a randomized, placebo-controlled, double blinded, 10-week phase 2 clinical trial in patients with early HD using orally administered VX-745 (n = 16) (NCT03980938).²⁷ It will primarily assess hippocampal dysfunction changes after 20 weeks. The study began in 2019 and is currently recruiting.²⁸⁹

VX-745 treatment has also been assessed in AD and Lewy Body Dementia (LBD). A trial in LBD showed cognition and mobility benefits with VX-745 administration,²⁹⁰ but a study on mild AD did not indicate statistically significant clinical benefit.²⁹¹ All studies to date have shown VX-745 is safe and tolerable.^{290,291}

Nilotinib

Nilotinib (also called Tasigna[®] and AMN107) is being investigated as an oral HD therapeutic^{27,292} (Figure 5). It is a

BCR-ABL tyrosine kinase inhibitor that traditionally treats chronic myeloid leukaemia (CML).²⁹² Nilotinib reduced oxidative stress in Parkinson's Disease (PD) model mice via ABL inhibition and improved motor performance.²⁹³ Tyrosine kinase inhibitors also modulated misfolded protein pathophysiology by autophagic clearance in PD model mice, positively affecting neuroinflammation.²⁹⁴ Clinical trials with PD and LBD participants (n = 12) used lower Nilotinib doses than those that treat CML. Investigators found Nilotinib was safe and tolerable, and appeared to inhibit central nervous system ABL. However, the study was small and not placebo-controlled, necessitating further trials.²⁹² Georgetown University are recruiting for an open label phase 1 clinical trial to assess safety, tolerability and CSF biomarkers associated with motor and behavioural HD symptoms (n = 20) (NCT03764215/Tasigna-HD).^{27,295} Ten participants with HD will be given 150 mg for 3 months, and if safe and tolerable another ten participants will be given 300 mg for a further 3 months. There is no control group.²⁹⁵

PF-02545920. Multiple therapies have been discontinued despite promising pre-clinical results. Pfizer's PF-02545920 is an oral phosphodiesterase 10a (PDE10a) inhibitor.^{277,296} It aims to treat HD motor symptoms. PDE10a inhibition improves basal ganglia pathology associated with HD, such as improving spiny projection neuron pathology and responsiveness in HD mice.²⁹⁷ A randomized, double blinded, 26-week phase 2 trial assessing safety and efficacy (via UHDRS-TMS) of PF-02545920 compared 5 mg and 20 mg PF-02545920 with placebo in 272 HD patients with chorea (NCT02197130/Amaryllis).^{277,296} Results showed no change in UHDRS-TMS or secondary endpoints between placebo and PF-02545920, despite PF-02545920 being safe and tolerable. An OLE study for Amaryllis participants was terminated after these results,²⁷⁷ and PF-02545920 was discontinued from Pfizer's drug development pipeline.²⁹⁸

Other small molecule therapies are also being investigated. Fenofibrate is a peroxisome proliferator-activated receptor (PPAR) agonist,²⁷ capable of modifying γ coactivator-1 α (PGC-1 α).²⁹⁹ PGC-1 α is a transcriptional co-regulator involved in mitochondrial malfunction in HD, leading to neuronal degeneration.³⁰⁰ Fenofibrate is in a 6 month, randomized, placebo-controlled, double blinded, phase 2 clinical trial assessing safety and efficacy (NCT03515213).²⁷ Azevan Pharmaceuticals' SRX246 is a vasopressin 1a receptor antagonist that has completed Phase 2 clinical trials (NCT02507284), aimed at treating HD patients with irritability and aggression. It showed safety, tolerability and validity as a treatment candidate in the near future.³⁰¹ Varenicline is a nicotinic acetylcholine receptor agonist targeting cognitive improvements in early HD patients who smoke. HD patients treated with varenicline showed significant executive function and emotional recognition improvements, without clinically significant adverse events.³⁰² Sage Therapeutics' SAGE-718 is an NMDA receptor

positive allosteric modulator (PAM) that targets HD cognitive symptoms. Phase 1 results showed tolerability, no clinically significant adverse events and improved cognitive performance from baseline, leading to planned phase 2 trial initiation for late 2021.^{303,304} PBT2 likely reduces metal induced mHTT aggregation through metal protein attenuation. It has completed phase 2 trials (NCT01590888), showing tolerability and safety but no clear benefit for cognitive scores – warranting further studies.³⁰⁵

Further Neuroinflammation Targeted Therapies for HD

Minocycline. The presence of chronic inflammation in HD has presented a potential therapeutic target with a number of treatments targeting inflammation being proposed. The second-generation tetracycline drug, minocycline, was 1 such proposed therapy that showed anti-inflammatory and anti-apoptotic properties. Mice treated with a low dose of minocycline brought about better behavioural performance than HD mice without treatment (5 mg/kg/day).³⁰⁶ Stabilization of general neuropsychological and motor symptoms along with significant improvements in psychiatric symptoms were seen in patients administered with 100 mg/day of minocycline in a 2 day clinical trial.³⁰⁷ However, a low sample size of 11 patients suggests that these results must be corroborated with similar studies on a larger number of patients. A further clinical trial of minocycline treatment on 30 patients over a 6 month period reported a trend toward symptom improvement on the URHDS.³⁰⁸ The lack of statistical significance in these results were attributed to the short overall treatment period with the study suggesting the need for a period longer than 6 months for effective minocycline treatment. A larger study of 87 HD patients over 18 months assessed the potential benefits of this drug treatment in a phase 3 clinical trial (clinicaltrials.gov ID: NCT00277355) and found that patient improvement did not reach a futility threshold.³⁰⁹ Based on these findings, further trials of minocycline administration at 200 mg/day were not warranted.³⁰⁹

Cannabidiol. Cannabinoids have also been postulated to be potential therapeutic targets due to their anti-inflammatory properties.³¹⁰⁻³¹² A clinical trial of cannabidiol treatment on 15 HD patients showed no improvement in motor function over a 6-week period.³¹³ The low sample size of this study again poses an issue when concluding the effectiveness of cannabidiol as a therapeutic agent. However, a double-blind, randomized, placebo-controlled, cross-over pilot clinical trial with Sativex (®), a botanical extract with an equimolecular combination of delta-9-tetrahydrocannabinol and cannabidiol, conducted over 12 weeks on 24 HD patients, also reported no significant symptomatic effects (clinicaltrials.gov ID: NCT01502046).³¹⁴

The failure of these clinical trials may suggest that global immunosuppression (for example through corticosteroids) or the effects of anti-inflammatory treatments may be too broad

and may also highlight a need to consider the heterogeneity of microglial and astrocytic phenotypes.³¹⁵⁻³¹⁷ A coherent understanding of the diverse characteristics of these cells and their functions in a healthy state, different disease states, infection, trauma, development and ageing is critical for better therapeutic outcomes.

Conclusions

Positive steps are being made in the development of current HD therapeutics. Guidance for clinicians on HD treatment is improving – guidelines were released relatively recently to elevate international standards of HD care.¹¹⁶ Current therapies focus on symptom management via multidisciplinary teams, using pharmacological and non-pharmacological treatment.³¹⁸ However, HD management has not significantly advanced for the past 20 years.⁵² Furthermore, diverse and changeable symptom presentations creates complexity when treating HD.¹⁸ There is no cure for HD, or therapeutic to alter disease onset or progression.¹¹⁶ HD management requires a tailored approach not only due to symptom diversity, but also adverse effects, common contraindications, drug-drug interactions and certain HD medications worsening other symptoms.^{18,52,319} Clinicians must navigate these issues in addition to effectively treating HD symptoms.

Perhaps the most prominent shortcoming of today's HD therapeutics and previous experimental therapeutics is their inability to impact specific HD targets, instead influencing non-specific and downstream processes.³⁰ This limits their efficiency as disease mechanisms will still propagate and cause harm where they are not inhibited. Pursuing HD-specific targets at the root cause of disease is therefore the new and likely future research direction.³⁰ The recent surge in interest to develop therapies that target mHTT RNA and DNA shows this trend.³¹ Additional therapeutic targets are being investigated, including dopaminergic and glutamatergic neurotransmission, synaptic activity, BDNF signalling, mitochondrial function and biogenics, and aggregate inhibition.³²⁰⁻³³⁰ However, several of these targets were primarily researched over 10 years ago, and do not appear to be areas of significant interest in the field currently. Peripheral targets are also being investigated due to peripheral HD symptoms and HTT expression.³³¹⁻³³³ For example, the gut microbiome is a key target due to its bidirectional communication with the brain and alteration in HD.³³⁴⁻³³⁸

RNA therapies, particularly antisense oligonucleotides, have favourable results and are currently of the most promising future HD therapeutics.^{26,339} However, RNA targeting therapies come with their own challenges. It is unknown whether non-selective HTT lowering is safe, if off target effects will occur in larger populations, whether mHTT is the singular cause of HD pathology and if permanent HTT suppression is required. Furthermore, if allele-specific treatments are required, 1 SNP will not treat all HD patients.^{137,340} Many RNA based

therapies also require invasive administration.³¹ Whether this is tolerable and effective in the long term for larger populations will likely be determined by future research. Initial clinical findings from RNAi therapies and RNA targeting small molecules, and further clinical results from antisense oligonucleotides will clarify RNA based therapies' efficacy for HD patients. Particularly, results of larger trials such as PRECISION-HD1 and PRECISION-HD2, and further information on GENERATION-HD1 will likely inform whether antisense oligonucleotides remain the front-runners of future HD therapeutics.

DNA therapies may also enter clinical trials within the next decade, but much work still needs to be done to achieve this goal. Ensuring DNA therapies, particularly CRISPR therapies, are safe for human use will be of primary importance.¹³⁷ Combination therapies may also be explored in future research. Current research suggests combining multiple RNA targeting therapies,³⁴¹ or RNA targeting therapies with antibodies or stem cells^{32,214} may be beneficial.

Work on future HD therapeutics will continue for numerous years to come and resulting treatments for HD patients are likely some time away.³⁴ However, many of the latest developments are encouraging and a new era of HD therapeutics may be approaching.

ORCID iD

Andrea Kwakowsky  <https://orcid.org/0000-0002-3801-4956>

REFERENCES

1. Reiner A, Dragatsis I, Dietrich P. Genetics and neuropathology of Huntington's disease. *Int Rev Neurobiol.* 2011;98:325-372.
2. Gusella JF, Wexler NS, Conneally PM, Naylor SL, Anderson MA, Tanzi RE, et al. A polymorphic DNA marker genetically linked to Huntington's disease. *Nature.* 1983;306(5940):234-238.
3. Nance MA. Genetics of Huntington disease. *Handb Clin Neurol.* 2017;144:3-14.
4. Duyao M, Ambrose C, Myers R, Novelletto A, Persichetti F, Frontali M, et al. Trinucleotide repeat length instability and age of onset in Huntington's disease. *Nat Genet.* 1993;4(4):387-392.
5. The Huntington's Disease Collaborative Research Group. A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's disease chromosomes. The Huntington's disease collaborative research group. *Cell.* 1993; 72(6):971-983.
6. Hoffner G, Djian P. Protein aggregation in Huntington's disease. *Biochimie.* 2002; 84(4):273-278.
7. Scherzinger E, Sittler A, Schweiger K, Heiser V, Lurz R, Hasenbank R, et al. Self-assembly of polyglutamine-containing huntingtin fragments into amyloid-like fibrils: Implications for Huntington's disease pathology. *Proc Natl Acad Sci Unit States Am.* 1999;96(8):4604-4609.
8. Kumar A, Ratan RR. Oxidative stress and Huntington's disease: The good, the bad, and the ugly. *J Huntingt Dis.* 2016;5(3):217-237.
9. Quintanilla RA, Johnson GV. Role of mitochondrial dysfunction in the pathogenesis of Huntington's disease. *Brain Research Bulletin.* 2009;80(4-5):242-247.
10. Tabrizi S, Cleeter M, Xuereb J, Taanman JW, Cooper J, Schapira A. Biochemical abnormalities and excitotoxicity in Huntington's disease brain. *Ann Neurol.* 1999; 45(1):25-32.
11. Palpagama TH, Waldvogel HJ, Faull RLM, Kwakowsky A. The role of microglia and astrocytes in Huntington's disease. *Front Mol Neurosci.* 2019;12:258.
12. Gil JM, Rego AC. Mechanisms of neurodegeneration in Huntington's disease. *Eur J Neurosci.* 2008;27(11):2803-2820.
13. Walker FO. Huntington's disease. *Lancet.* 2007;369(9557):218-228.
14. Roos RAC. Huntington's disease: A clinical review. *Orphanet Journal of Rare Diseases.* 2010;5:40.

15. Pringsheim T, Wiltshire K, Day L, Dykeman J, Steeves T, Jette N. The incidence and prevalence of Huntington's disease: A systematic review and meta-analysis. *Mov Disord*. 2012;27(9):1083-1091.
16. Warby SC, Montpetit A, Hayden AR, Carroll JB, Butland SL, Visscher H, et al. CAG expansion in the Huntington disease gene is associated with a specific and targetable predisposing haplogroup. *Am J Hum Genet*. 2009;84(3):351-366.
17. Mestre T, Ferreira J, Coelho MM, Rosa M, Sampaio C. Therapeutic interventions for symptomatic treatment in Huntington's disease. *Cochrane Database Syst Rev*. 2009(3).
18. Videnovic A. Treatment of Huntington disease. *Curr Treat Options Neurol*. 2013; 15(4):424-438.
19. Frank S. Treatment of Huntington's disease. *Neurotherapeutics*. 2014;11(1): 153-160.
20. Coppin EM, Roos RA. Current pharmacological approaches to reduce chorea in Huntington's disease. *Drugs*. 2017;77(1):29-46.
21. Ghosh R, Tabrizi SJ. Clinical aspects of Huntington's disease. In: *Behavioral Neurobiology of Huntington's Disease and Parkinson's Disease*. Springer; 2013: 3-31.
22. Craufurd D, Snowden J. Neuropsychological and neuropsychiatric aspects of Huntington's disease. *Oxf Monogr Med Genet*. 2002;45:62-94.
23. van Duijn E, Craufurd D, Hubers AA, Giltay EJ, Bonelli R, Rickards H, et al. Neuropsychiatric symptoms in a European Huntington's disease cohort (REGISTRY). *J Neurol Neurosurg Psychiatry*. 2014;85(12):1411-1418.
24. Paulsen JS, Nehl C, Hoth KF, Kanz JE, Benjamin M, Conybeare R, et al. Depression and stages of Huntington's disease. *J Neuropsychiatry Clin Neurosci*. 2005;17(4):496-502.
25. Tabrizi SJ, Ghosh R, Leavitt BR. Huntingtin lowering strategies for disease modification in Huntington's disease. *Neuron*. 2019;101(5):801-819.
26. Evers MM, Konstantinova P. AAV5-miHTT gene therapy for Huntington disease: Lowering both huntingtins. *Expert Opin Biol Ther*. 2020;20(10): 1121-1124.
27. Rodrigues FB, Wild EJ. Huntington's disease clinical trials corner: April 2020. *J Huntingt Dis*. 2020;9:185-197.
28. Hoffmann-La Roche. *A Study to Evaluate the Efficacy and Safety of Intrathecally Administered RO7234292*. (RG6042) in Patients With Manifest Huntington's Disease 2018 [cited 2021 08/01/2021]. Available from: <https://clinicaltrials.gov/ct2/show/NCT03761849>.
29. Roche provides update on tominersen programme in manifest Huntington's disease [press release]. Basel, Switzerland, 22 March 2021 2021.
30. Mestre TA, Sampaio C. Huntington disease: Linking pathogenesis to the development of experimental therapeutics. *Curr Neurol Neurosci Rep*. 2017;17(2):18.
31. Wild EJ, Tabrizi SJ. Therapies targeting DNA and RNA in Huntington's disease. *Lancet Neurol*. 2017;16(10):837-847.
32. Denis HL, David LS, Cicchetti F. Antibody-based therapies for Huntington's disease: Current status and future directions. *Neurobiol Dis*. 2019; 132:104569.
33. Maucksch C, Vazey EM, Gordon RJ, Connor B. Stem cell-based therapy for Huntington's disease. *J Cell Biochem*. 2013;114(4):754-763.
34. Tabrizi SJ, Flower MD, Ross CA, Wild EJ. Huntington disease: New insights into molecular pathogenesis and therapeutic opportunities. *Nat Rev Neurol*. 2020; 16(10):529-546.
35. Kim A, Lalonde K, Truesdell A, Gomes Welter P, Brocardo PS, Rosenstock TR, et al. New avenues for the treatment of Huntington's disease. *Int J Mol Sci*. 2021; 22(16).
36. Smith Y, Bevan MD, Shink E, Bolam JP. Microcircuitry of the direct and indirect pathways of the basal ganglia. *Neuroscience*. 1998;86(2):353-387.
37. Albin RL, Reiner A, Anderson KD, Dure LS, Handelin B, Balfour R, et al. Preferential loss of striato-external pallidal projection neurons in presymptomatic Huntington's disease. *Ann Neurol*. 1992;31(4):425-430.
38. Penney JB Jr., Young AB. Striatal inhomogeneities and basal ganglia function. *Mov Disord*. 1986;1(1):3-15.
39. Reiner A, Albin RL, Anderson KD, D'Amato CJ, Penney JB, Young AB. Differential loss of striatal projection neurons in Huntington disease. *Proc Natl Acad Sci U S A*. 1988;85(15):5733-5737.
40. Kirkwood SC, Su JL, Conneally PM, Foroud T. Progression of symptoms in the early and middle stages of Huntington disease. *Arch Neurol*. 2001;58(2):273-278.
41. Hayden MR, Leavitt BR, Yasothan U, Kirkpatrick P. *Tetrabenazine*. Nature Publishing Group; 2009.
42. Zheng G, Dwoskin LP, Crooks PA. Vesicular monoamine transporter 2: Role as a novel target for drug development. *AAPS J*. 2006;8(4):E682-E92.
43. Group HS. Tetrabenazine as antichorea therapy in Huntington disease: A randomized controlled trial. *Neurology*. 2006;66(3):366-372.
44. Kieburz K, Penney JB, Como P, Ranen N, Shoulson I, Feigin A, et al. Unified Huntington's disease rating scale: Reliability and consistency. *Neurology*. 2001; 11(2):136-142.
45. Jankovic J, Beach J. Long-term effects of tetrabenazine in hyperkinetic movement disorders. *Neurology*. 1997;48(2):358-362.
46. Claassen DO, Carroll B, De Boer LM, Wu E, Ayyagari R, Gandhi S, et al. Indirect tolerability comparison of Deutetrabenazine and Tetrabenazine for Huntington disease. *J Clin Movement Disorder*. 2017;4(1):3.
47. Frank S, Testa CM, Stamler D, Kayson E, Davis C, Edmondson MC, et al. Effect of deutetrabenazine on chorea among patients with Huntington disease: A randomized clinical trial. *JAMA*. 2016;316(1):40-50.
48. Yero T, Rey JA. Tetrabenazine (Xenazine), an FDA-approved treatment option for Huntington's disease-related chorea. *Pharma Therapeutic*. 2008;33(12):690.
49. Claassen DO, Carroll B, De Boer LM, Wu E, Ayyagari R, Gandhi S, et al. Indirect tolerability comparison of Deutetrabenazine and Tetrabenazine for Huntington disease. *J Clin Mov Disord*. 2017;4:3.
50. Schultz JL, Killoran A, Nopoulos PC, Chabal CC, Moser DJ, Kamholz JA. Evaluating depression and suicidality in tetrabenazine users with Huntington disease. *Neurology*. 2018;91(3):e202-e7.
51. Dorsey E, Brocht AF, Nichols PE, Darwin KC, Anderson KE, Beck CA, et al. Depressed mood and suicidality in individuals exposed to tetrabenazine in a large Huntington disease observational study. *J Huntingt Dis*. 2013;2(4): 509-515.
52. Walker FO. Huntington's disease. *Lancet*. 2007;369(9557):218-228.
53. Bashir H, Jankovic J. Treatment options for chorea. *Expert Review of Neurotherapeutics*. 2018;18(1):51-63.
54. Fulton B, Goa KL. Olanzapine. *Drugs*. 1997;53(2):281-298.
55. Janssen P, Niemegeers C, Awouters F, Schellekens K, Megens A, Meert T. Pharmacology of risperidone (R 64 766), a new antipsychotic with serotonin-5₂ and dopamine-D₂ antagonistic properties. *J Pharmacol Exp Therapeut*. 1988; 244(2):685-693.
56. Armstrong MJ, Miyasaki JM. Evidence-based guideline: Pharmacologic treatment of chorea in Huntington disease: Report of the guideline development subcommittee of the American Academy of Neurology. *Neurology*. 2012;79(6):597-603.
57. Vijayakumar D, Jankovic J. Drug-induced dyskinesia, part 1: Treatment of levodopa-induced dyskinesia. *Drugs*. 2016;76(7):759-777.
58. Metman LV, Morris M, Farmer C, Gillespie M, Mosby K, Wu J, et al. Huntington's disease: A randomized, controlled trial using the NMDA-antagonist amantadine. *Neurology*. 2002;59(5):694-699.
59. Rosas HD, Koroshetz WJ, Jenkins BG, Chen YI, Hayden DL, Beal MF, et al. Riluzole therapy in Huntington's disease (HD): Movement disorders. *Official Journal of the Movement Disorder Society*. 1999;14(2):326-330.
60. Seppi K, Mueller J, Bodner T, Brandner E, Benke T, Weirich-Schwaiger H, et al. Riluzole in Huntington's disease (HD): An open label study with one year follow up. *J Neurol*. 2001;248(10):866-869.
61. Duggan L, Fenton M, Rathbone J, Dardennes R, El-Dosoky A, Indran S. Olanzapine for schizophrenia. *Cochrane Database Syst Rev*. 2005(2).
62. Squitieri F, Cannella M, Porcellini A, Brusa L, Simonelli M, Ruggieri S. Short-term effects of olanzapine in Huntington disease. *Cognit Behav Neurol*. 2001;14(1):69-72.
63. Eder U, Mangweth B, Ebenbichler C, Weiss E, Hofer A, Hummer M, et al. Association of olanzapine-induced weight gain with an increase in body fat. *Am J Psychiatr*. 2001;158(10):1719-1722.
64. Goudie AJ, Smith JA, Halford JC. Characterization of olanzapine-induced weight gain in rats. *J Psychopharmacol*. 2002;16(4):291-296.
65. Ball MP, Coons VB, Buchanan RW. A program for treating olanzapine-related weight gain. *Psychiatr Serv*. 2001;52(7):967-969.
66. Marder SR, Meibach RC. Risperidone in the treatment of schizophrenia. *Am J Psychiatr*. 1994.
67. Dazzan P, Morgan KD, Orr K, Hutchinson G, Chitnis X, Suckling J, et al. Different effects of typical and atypical antipsychotics on grey matter in first episode psychosis: The AESOP study. *Neuropsychopharmacology*. 2005;30(4):765-774.
68. Komossa K, Rummel-Kluge C, Schwarz S, Schmid F, Hunger H, Kissling W, et al. Risperidone versus other atypical antipsychotics for schizophrenia. *Cochrane Database Syst Rev*. 2011(1).
69. Duff K, Beglinger LJ, O'Rourke ME, Nopoulos P, Paulson HL, Paulsen JS. Risperidone and the treatment of psychiatric, motor, and cognitive symptoms in Huntington's disease. *Ann Clin Psychiatr*. 2008;20(1):1-3.
70. Conley RR, Mahmoud R. A randomized double-blind study of risperidone and olanzapine in the treatment of schizophrenia or schizoaffective disorder. *Am J Psychiatr*. 2001;158(5):765-774.
71. Naarding P, Kremer H, Zitman F. Huntington's disease: A review of the literature on prevalence and treatment of neuropsychiatric phenomena. *Eur Psychiatr*. 2001; 16(8):439-445.
72. Hyttel J. *Pharmacological Characterization of Selective Serotonin Reuptake Inhibitors (SSRIs)*. International clinical psychopharmacology; 1994.
73. Preskorn SH. Clinically relevant pharmacology of selective serotonin reuptake inhibitors. *Clin Pharmacokinet*. 1997;32(1):1-21.
74. Goodnick PJ, Goldstein BJ. Selective serotonin reuptake inhibitors in affective disorders—I. Basic pharmacology. *J Psychopharmacol*. 1998;12(4_suppl 1):S-S20.
75. Taylor C, Fricker AD, Devi LA, Gomes I. Mechanisms of action of antidepressants: From neurotransmitter systems to signaling pathways. *Cell Signal*. 2005; 17(5):549-557.

76. Chu A, Wadhwa R. Selective Serotonin Reuptake Inhibitors. In: StatPearls. Treasure Island (FL): StatPearls Publishing. Copyright © 2021, StatPearls Publishing LLC.; 2021.
77. Khushboo SB. Antidepressants: Mechanism of action, toxicity and possible amelioration. *J Applied Biotechnol Bioengineer.* 2017;3(5).
78. Hirschfeld R. Long-term side effects of SSRIs: Sexual dysfunction and weight gain. *J Clin Psychiatr.* 2003;64:20-24.
79. Gibbons RD, Brown CH, Hur K, Marcus SM, Bhaumik DK, Erkens JA, et al. Early evidence on the effects of regulators' suicidality warnings on SSRI prescriptions and suicide in children and adolescents. *Am J Psychiatr.* 2007;164(9):1356-1363.
80. Garland EJ, Kutcher S, Virani A, Elbe D. Update on the use of SSRIs and SNRIs with children and adolescents in clinical practice. *J Canad Acad Child Adol Psychiatr.* 2016;25(1):4.
81. Moulton CD, Hopkins C, Bevan-Jones WR. Systematic review of pharmacological treatments for depressive symptoms in Huntington's disease. *Mov Disord.* 2014;29(12):1556-1561.
82. Beglinger LJ, Adams WH, Langbehn D, Fiedorowicz JG, Jorge R, Biglan K, et al. Results of the citalopram to enhance cognition in Huntington disease trial. *Mov Disord.* 2014;29(3):401-405.
83. De Marchi N, Daniele F, Ragone MA. Fluoxetine in the treatment of Huntington's disease. *Psychopharmacology.* 2001;153(2):264-266.
84. Benfield P, Heel RC, Lewis SP. Fluoxetine. *Drugs.* 1986;32(6):481-508.
85. Ranen NG, Lipsey JR, Treisman G, Ross CA. Sertraline in the treatment of severe aggressiveness in Huntington's disease. *J Neuropsychiatry Clin Neurosci.* 1996;8(3):338-340.
86. Patzold T, Brüne M. Obsessive compulsive disorder in Huntington disease: A case of isolated obsessions successfully treated with sertraline. *Cognit Behav Neurol.* 2002;15(3):216-219.
87. Monroe RR. Anticonvulsants in the treatment of aggression. *J Nerv Ment Dis.* 1975;160:119-126.
88. Stanford MS, Helfritz LE, Conklin SM, Villemarette-Pittman NR, Greve KW, Adams D, et al. A comparison of anticonvulsants in the treatment of impulsive aggression. *Exp Clin Psychopharmacol.* 2005;13(1):72.
89. Grunze HC. Anticonvulsants in bipolar disorder. *J Ment Health.* 2010;19(2):127-141.
90. Yatham LN. Newer anticonvulsants in the treatment of bipolar disorder. *J Clin Psychiatr.* 2004;65:28-35.
91. Bruno A, Micò U, Pandolfo G, Mallamace D, Abenavoli E, Di Nardo F, et al. Lamotrigine augmentation of serotonin reuptake inhibitors in treatment-resistant obsessive-compulsive disorder: A double-blind, placebo-controlled study. *J Psychopharmacol.* 2012;26(11):1456-1462.
92. Wang HR, Woo YS, Bahk WM. Potential role of anticonvulsants in the treatment of obsessive-compulsive and related disorders. *Psychiatr Clin Neurosci.* 2014;68(10):723-732.
93. Macdonald R, McLean M. Anticonvulsant drugs: Mechanisms of action. *Adv Neurol.* 1986;44:713.
94. Appleton RE, Freeman A, Cross JH. Diagnosis and management of the epilepsies in children: A summary of the partial update of the 2012 NICE epilepsy guideline. *Arch Dis Child.* 2012;97(12):1073-1076.
95. Hammer GD, McPhee SJ. *Pathophysiology of Disease: An Introduction to Clinical Medicine.* Univerza v Ljubljani, Medicinska fakulteta; 2014.
96. Stefan H, Feuerstein T. Novel anticonvulsant drugs. *Pharmacol Ther.* 2007;113(1):165-183.
97. Grunze HCR. The effectiveness of anticonvulsants in psychiatric disorders. *Dialogues Clin Neurosci.* 2008;10(1):77-89.
98. Gould TD, Chen G, Manji HK. Mood stabilizer psychopharmacology. *Clin Neurosci Res.* 2002;2(3-4):193-212.
99. Ross CA, Tabrizi SJ. Huntington's disease: From molecular pathogenesis to clinical treatment. *Lancet Neurol.* 2011;10(1):83-98.
100. Chiu C-T, Liu G, Leeds P, Chuang D-M. Combined treatment with the mood stabilizers lithium and valproate produces multiple beneficial effects in transgenic mouse models of Huntington's disease. *Neuropsychopharmacology.* 2011;36(12):2406-2421.
101. Scheuing L, Chiu C-T, Liao H-M, Linares GR, Chuang D-M. Preclinical and clinical investigations of mood stabilizers for Huntington's disease: What have we learned? *Int J Biol Sci.* 2014;10(9):1024.
102. Lees G, Leach MJ. Studies on the mechanism of action of the novel anticonvulsant lamotrigine (Lamictal) using primary neuroglial cultures from rat cortex. *Brain Res.* 1993;612(1-2):190-199.
103. Wu J, Tang T, Bezprozvany I. Evaluation of clinically relevant glutamate pathway inhibitors in in vitro model of Huntington's disease. *Neurosci Lett.* 2006;407(3):219-223.
104. Kremer B, Clark C, Almqvist E, Raymond L, Graf P, Jacova C, et al. Influence of lamotrigine on progression of early Huntington disease: A randomized clinical trial. *Neurology.* 1999;53(5):1000.
105. Ambrósio AF, Soares-da-Silva P, Carvalho CM, Carvalho AP. Mechanisms of action of carbamazepine and its derivatives, oxcarbazepine, BIA 2-093, and BIA 2-024. *Neurochem Res.* 2002;27(1-2):121-130.
106. Lang DG, Wang CM, Cooper B. Lamotrigine, phenytoin and carbamazepine interactions on the sodium current present in N4TG1 mouse neuroblastoma cells. *J Pharmacol Exp Therapeut.* 1993;266(2):829-835.
107. Devi K, George S, Criton S, Suja V, Sridevi P. Carbamazepine-The commonest cause of toxic epidermal necrolysis and Stevens-Johnson syndrome: A study of 7 years. *Indian J Dermatol Venereol Leprol.* 2005;71(5):325.
108. Bilney B, Morris ME, Perry A. Effectiveness of physiotherapy, occupational therapy, and speech pathology for people with Huntington's disease: A systematic review. *Neurorehabilitation Neural Repair.* 2003;17(1):12-24.
109. Mirek E, Filip M, Banaszkiwicz K, Rudzińska M, Szymura J, Pasiut S, et al. The effects of physiotherapy with PNF concept on gait and balance of patients with Huntington's disease—pilot study. *Neurol Neurochir Pol.* 2015;49(6):354-357.
110. Quinn L, Busse M. Physiotherapy clinical guidelines for Huntington's disease. *Neurodegener Dis Manag.* 2012;2(1):21-31.
111. Busse ME, Khalil H, Quinn L, Rosser AE. Physical therapy intervention for people with Huntington disease. *Phys Ther.* 2008;88(7):820-831.
112. Barkmeier-Kraemer JM, Clark HM. Speech-language pathology evaluation and management of hyperkinetic disorders affecting speech and swallowing function. *Tremor Hyperkinetic Movement.* 2017;7:489.
113. Butnaru H, Krueger WO. *Communication Device and Method for Deaf and Mute Persons.* Google Patents; 2001.
114. Pennington L, Goldbart J, Marshall J. Speech and language therapy to improve the communication skills of children with cerebral palsy. *Cochrane Database Syst Rev.* 2004(2).
115. Żukiewicz-Sobczak W, Król R, Wróblewska P, Piątek J, Gibas-Dorna M. Huntington disease—principles and practice of nutritional management. *Neurol Neurochir Pol.* 2014;48(6):442-448.
116. Bachoud-Lévi A-C, Ferreira J, Massart R, Yousov K, Rosser A, Busse M, et al. International Guidelines for the Treatment of Huntington's Disease. *Front Neurol.* 2019;10(710).
117. Nagel SJ, Machado AG, Gale JT, Lobel DA, Pandya M. Preserving cortico-striatal function: Deep brain stimulation in Huntington's disease. *Front Syst Neurosci.* 2015;9(32).
118. Sharma M, Deogaonkar M. Deep brain stimulation in Huntington's disease: Assessment of potential targets. *J Clin Neurosci.* 2015;22(5):812-817.
119. Temel Y, Cao C, Vlamings R, Blokland A, Ozen H, Steinbusch HW, et al. Motor and cognitive improvement by deep brain stimulation in a transgenic rat model of Huntington's disease. *Neurosci Lett.* 2006;406(1-2):138-141.
120. Wojtecki L, Groiss SJ, Hartmann CJ, Elben S, Omlor S, Schnitzler A, et al. Deep brain stimulation in Huntington's disease—preliminary evidence on pathophysiology, efficacy and safety. *Brain Sciences.* 2016;6(3):38.
121. Gonzalez V, Cif L, Biolsi B, Garcia-Ptacek S, Seychelles A, Sanrey E, et al. Deep brain stimulation for Huntington's disease: Long-term results of a prospective open-label study. *J Neurosurg.* 2014;121(1):114-122.
122. Fenoy AJ, Simpson RK. Risks of common complications in deep brain stimulation surgery: Management and avoidance. *J Neurosurg.* 2014;120(1):132-139.
123. Ghosh R, Tabrizi SJ. Clinical aspects of Huntington's disease. *Curr Top Behav Neurosci.* 2015;22:3-31.
124. Scoles DR, Pulst SM. Oligonucleotide therapeutics in neurodegenerative diseases. *RNA Biol.* 2018;15(6):707-714.
125. Rodrigues FB, Wild EJ. Huntington's disease clinical trials corner: February 2018. *J Huntingt Dis.* 2018;7(1):89-98.
126. Marxreiter F, Stemick J, Kohl Z. Huntingtin lowering strategies. *Int J Mol Sci.* 2020;21(6):2146.
127. Kordasiewicz HB, Stanek LM, Wancewicz EV, Mazur C, McAlonis MM, Pytel KA, et al. Sustained therapeutic reversal of Huntington's disease by transient repression of huntingtin synthesis. *Neuron.* 2012;74(6):1031-1044.
128. Ionis Pharmaceuticals I. *Safety, Tolerability, Pharmacokinetics, and Pharmacodynamics of ISIS 443139 in Participants with Early Manifest Huntington's Disease.* National Institutes of Health; 2019. [updated May 31 2019; cited 2021 08/01/2021]. Available from: <https://clinicaltrials.gov/ct2/show/NCT02519036>.
129. Tabrizi SJ, Leavitt BR, Landwehrmeyer GB, Wild EJ, Saft C, Barker RA, et al. Targeting Huntingtin expression in patients with Huntington's disease. *N Engl J Med.* 2019;380(24):2307-2316.
130. Dhuri K, Bechtold C, Quijano E, Pham H, Gupta A, Vikram A, et al. Antisense oligonucleotides: An emerging area in drug discovery and development. *J Clin Med.* 2020;9(6):2004.
131. Rodrigues FB, Ferreira JJ, Wild EJ. Huntington's disease clinical trials corner: June 2019. *J Huntingt Dis.* 2019;8(3):363-371.
132. Hoffmann-La Roche. A Study to Evaluate the Safety, Tolerability, Pharmacokinetics, and Pharmacodynamics of RO7234292 (ISIS 443139) in Huntington's Disease Patients Who Participated in Prior Investigational Studies of RO7234292

- (ISIS 443139) 2019 [updated June 9 2020; cited 2021 08/01/2021]. Available from: <https://clinicaltrials.gov/ct2/show/NCT03342053>.
133. Rodrigues FB, Quinn L, Wild EJ. Huntington's disease clinical trials corner: January 2019. *J Huntingt Dis*. 2019;8:115-125.
 134. Hoffmann-La Roche. An Open-Label Extension Study to Evaluate Long-Term Safety and Tolerability of RO7234292 (RG6042) in Huntington's Disease Patients Who Participated in Prior Roche and Genentech Sponsored Studies 2019 [updated December 10 2020; cited 2021 08/01/2021]. Available from: <https://clinicaltrials.gov/ct2/show/NCT03842969>.
 135. Hoffmann-La Roche. A Study to Investigate the Pharmacokinetics and Pharmacodynamics of RO7234292 (RG6042) in CSF and Plasma, and Safety and Tolerability Following Intrathecal Administration in Patients With Huntington's Disease 2019 [cited 2021 08/01/2021]. Available from: <https://clinicaltrials.gov/ct2/show/NCT04000594>.
 136. Hersch S, Claassen D, Edmondson M, Wild E, Guercioli R, Panzara M. Multicenter, randomized, double-blind, placebo-controlled phase 1b/2a studies of WVE-120101 and WVE-120102 in patients with Huntington's disease (P2.006). *Neurology*. 2017;88(16 suppl):P2.006.
 137. Kaemmerer WF, Grondin RC. The effects of huntingtin-lowering: What do we know so far? *Degener Neurol Neuromuscul Dis*. 2019;9:3-17.
 138. Wave Life Sciences Announces Topline Data and Addition of Higher Dose Cohort in Ongoing Phase 1b/2a PRECISION-HD2 Trial in Huntington's Disease [press release]. Cambridge, Massachusetts: Wave Life Sciences, December 30 2019 2019.
 139. Wave Life Sciences Provides Update on Phase 1b/2a PRECISION-HD Trials. [press release]. March 29 2021.
 140. Wave Life Sciences Reports Third Quarter 2020 Financial Results and Provides Business Update [press release]. Cambridge, Massachusetts: Wave Life Sciences, November 9 2020 2020.
 141. Dale E, Frank-Kamenetsky M, Taborn K, Aklilu K, Liu Y, Berkovitch S, et al. Stereopure oligonucleotides for the selective silencing of mutant Huntingtin (4703). *Neurology*. 2020;94(15 suppl):4703.
 142. Evers MM, Pepers BA, van Deutekom JCT, Mulders SAM, den Dunnen JT, Aartsma-Rus A, et al. Targeting several CAG expansion diseases by a single antisense oligonucleotide. *PLoS One*. 2011;6(9):e24308.
 143. Datson NA, González-Barriga A, Kourkouta E, Weij R, van de Giessen J, Mulders S, et al. The expanded CAG repeat in the huntingtin gene as target for therapeutic RNA modulation throughout the HD mouse brain. *PLoS One*. 2017;12(2):e0171127.
 144. Li JY, Popovic N, Brundin P. The use of the R6 transgenic mouse models of Huntington's disease in attempts to develop novel therapeutic strategies. *NeuroRx*. 2005;2(3):447-464.
 145. Menalled LB, Kudwa AE, Miller S, Fitzpatrick J, Watson-Johnson J, Keating N, et al. comprehensive behavioral and molecular characterization of a new knock-in mouse model of Huntington's disease: zQ175. *PLoS One*. 2012;7(12):e49838.
 146. Skotte NH, Southwell AL, Østergaard ME, Carroll JB, Warby SC, Doty CN, et al. Allele-specific suppression of mutant huntingtin using antisense oligonucleotides: Providing a therapeutic option for all Huntington disease patients. *PLoS One*. 2014;9(9):e107434.
 147. Southwell AL, Skotte NH, Kordasiewicz HB, Østergaard ME, Watt AT, Carroll JB, et al. In vivo evaluation of candidate allele-specific mutant huntingtin gene silencing antisense oligonucleotides. *Mol Ther*. 2014;22(12):2093-2106.
 148. Southwell AL, Kordasiewicz HB, Langbehn D, Skotte NH, Parsons MP, Villanueva EB, et al. Huntingtin suppression restores cognitive function in a mouse model of Huntington's disease. *Sci Transl Med*. 2018;10(461):eaar3959.
 149. Evers MM, Tran H-D, Zalachoras I, Meijer OC, den Dunnen JT, van Ommen G-JB, et al. Preventing formation of toxic N-terminal Huntingtin fragments through antisense oligonucleotide-mediated protein modification. *Nucleic Acid Therapeut*. 2014;24(1):4-12.
 150. Triplet Therapeutics Presents Preclinical Data for TTX-3360, Announces Equity Pledge at the CHDI Foundation's 16th Annual HD Therapeutics Conference [press release]. April 27 2021.
 151. McBride JL, Boudreau RL, Harper SQ, Staber PD, Monteys AM, Martins I, et al. Artificial miRNAs mitigate shRNA-mediated toxicity in the brain: Implications for the therapeutic development of RNAi. *Proc Natl Acad Sci USA*. 2008;105(15):5868-5873.
 152. Grimm D, Streetz KL, Jopling CL, Storm TA, Pandey K, Davis CR, et al. Fatality in mice due to oversaturation of cellular microRNA/short hairpin RNA pathways. *Nature*. 2006;441(7092):537-541.
 153. Aguiar S, van der Gaag B, Cortese FAB. RNAi mechanisms in Huntington's disease therapy: siRNA versus shRNA. *Transl Neurodegener*. 2017;6:30.
 154. Miniarikova J, Zanella I, Huseinovic A, van der Zon T, Hanemaaijer E, Martier R, et al. Design, characterization, and lead selection of therapeutic miRNAs targeting Huntingtin for development of gene therapy for Huntington's disease. *Mol Ther Nucleic Acids*. 2016;5(3):e297-e.
 155. Miniarikova J, Zimmer V, Martier R, Brouwers CC, Pythoud C, Richetin K, et al. AAV5-miHTT gene therapy demonstrates suppression of mutant Huntingtin aggregation and neuronal dysfunction in a rat model of Huntington's disease. *Gene Therapy*. 2017;24(10):630-639.
 156. Keskin S, Brouwers CC, Sogorb-Gonzalez M, Martier R, Depla JA, Vallès A, et al. AAV5-miHTT lowers Huntingtin mRNA and protein without Off-target effects in patient-derived neuronal cultures and astrocytes. *Mol Ther Method Clin Dev*. 2019;15:275-284.
 157. Southwell AL, Skotte NH, Villanueva EB, Østergaard ME, Gu X, Kordasiewicz HB, et al. A novel humanized mouse model of Huntington disease for preclinical development of therapeutics targeting mutant huntingtin alleles. *Hum Mol Genet*. 2017;26(6):1115-1132.
 158. Caron NS, Southwell AL, Brouwers CC, Cengio LD, Xie Y, Black HF, et al. Potent and sustained huntingtin lowering via AAV5 encoding miRNA preserves striatal volume and cognitive function in a humanized mouse model of Huntington disease. *Nucleic Acids Res*. 2020;48(1):36-54.
 159. Spronck EA, Brouwers CC, Vallès A, de Haan M, Petry H, van Deventer SJ, et al. AAV5-miHTT gene therapy demonstrates sustained Huntingtin lowering and functional improvement in Huntington disease mouse models. *Mol Therapy Method Clin Dev*. 2019;13:334-343.
 160. Evers MM, Miniarikova J, Juhas S, Vallès A, Bohuslavova B, Juhasova J, et al. AAV5-miHTT gene therapy demonstrates broad distribution and strong human mutant Huntingtin lowering in a Huntington's disease minipig model. *Mol Ther*. 2018;26(9):2163-2177.
 161. Spronck EA, Vallès A, Lampen MH, Montenegro-Miranda PS, Keskin S, Heijink L, et al. Intrastriatal administration of AAV5-miHTT in non-human primates and rats is well tolerated and results in miHTT transgene expression in key areas of Huntington disease pathology. *Brain Sci*. 2021;11(2):129.
 162. UniQure. Safety and Proof-of-Concept (POC) Study With AMT-130 in Adults With Early Manifest Huntington Disease 2019 [updated October 28 2020; cited 2021 09/01/2021]. Available from: <https://clinicaltrials.gov/ct2/show/NCT04120493>.
 163. uniQure Announces Completion of Enrollment in First Cohort of Phase I/II Clinical Trial of AMT-130 for the Treatment of Huntington's Disease [press release]. Amsterdam, The Netherlands, April 5, 2021 2021.
 164. UniQure Announces Positive Recommendation from Data Safety Monitoring Board of Phase I/II Clinical Trial of AMT-130 for the Treatment of Huntington's Disease [press release]. Lexington, Massachusetts; Amsterdam, The Netherlands; UniQure, February 8 2021 2021.
 165. Stanek LM, Sardi SP, Mastis B, Richards AR, Treleven CM, Taksir T, et al. Silencing mutant huntingtin by adeno-associated virus-mediated RNA interference ameliorates disease manifestations in the YAC128 mouse model of Huntington's disease. *Hum Gene Ther*. 2014;25(5):461-474.
 166. Ghilan M, Bostrom CA, Hryciw BN, Simpson JM, Christie BR, Gil-Mohapel J. YAC128 Huntington's disease transgenic mice show enhanced short-term hippocampal synaptic plasticity early in the course of the disease. *Brain Research*. 2014;1581:117-128.
 167. Voyager Therapeutics Announces Preclinical Data For Huntington's Disease And Amyotrophic Lateral Sclerosis Programs At The Congress Of The European Society Of Gene And Cell Therapy [press release]. Cambridge, Massachusetts, October 16 2018.
 168. Voyager Therapeutics Provides Regulatory Update On VY-HTT01 Program [press release]. Cambridge, Massachusetts: Voyager Therapeutics, October 13 2020.
 169. Harper SQ, Staber PD, He X, Eliason SL, Martins IH, Mao Q, et al. RNA interference improves motor and neuropathological abnormalities in a Huntington's disease mouse model. *Proc Natl Acad Sci USA*. 2005;102(16):5820-5825.
 170. Franich NR, Fitzsimons HL, Fong DM, Klugmann M, Daring MJ, Young D. AAV vector-mediated RNAi of mutant huntingtin expression is neuroprotective in a novel genetic rat model of Huntington's disease. *Mol Ther*. 2008;16(5):947-956.
 171. McBride JL, Pitzer MR, Boudreau RL, Dufour B, Hobbs T, Ojeda SR, et al. Preclinical safety of RNAi-mediated HTT suppression in the rhesus macaque as a potential therapy for Huntington's disease. *Mol Ther*. 2011;19(12):2152-2162.
 172. Bhattacharyya A, ed. Identification and development of orally administered, CNS-penetrant small molecules that lower huntingtin protein levels by inducing a novel splicing event that alters the stability of huntingtin mRNA. CHDI HD Therapeutics Conference; 2019; Palm Springs, California.
 173. PTC Therapeutics Announces that PTC518 Has Entered into a Phase 1 Clinical Trial for the Huntington's Disease Program [press release]. South Plainfield, New Jersey 2020.
 174. Gray M, Shirasaki DI, Cepeda C, André VM, Wilburn B, Lu X-H, et al. Full-length human mutant huntingtin with a stable polyglutamine repeat can elicit progressive and selective neuropathogenesis in BACHD mice. *J Neurosci*. 2008;28(24):6182-6195.
 175. Therapeutics PTC. PTC518 Phase 1 Huntington's Disease Program Update 2021 [cited 2022 January]. Available from: <https://ir.ptcbio.com/events/event-details/ptc518-phase-1-huntingtons-disease-program-update>.
 176. Shorrock HK, Gillingwater TH, Groen EJN. Overview of current drugs and molecules in development for spinal muscular atrophy therapy. *Drugs*. 2018;78(3):293-305.

177. Novartis receives US Food and Drug Administration (FDA) Orphan Drug Designation for branaplam (LMI070) in Huntington's disease (HD) [press release]. Basel, Switzerland, October 21 2020.
178. Liu CR, Chang CR, Chern Y, Wang TH, Hsieh WC, Shen WC, et al. Spt4 is selectively required for transcription of extended trinucleotide repeats. *Cell*. 2012;148(4):690-701.
179. Cheng H-M, Chern Y, Chen IH, Liu C-R, Li S-H, Chun SJ, et al. Effects on murine behavior and lifespan of selectively decreasing expression of mutant huntingtin allele by supt4h knockdown. *PLoS Genetics*. 2015;11(3):e1005043.
180. Zhou X, Li G, Kaplan A, Gaschler MM, Zhang X, Hou Z, et al. Small molecule modulator of protein disulfide isomerase attenuates mutant huntingtin toxicity and inhibits endoplasmic reticulum stress in a mouse model of Huntington's disease. *Hum Mol Genet*. 2018;27(9):1545-1555.
181. Bard J, Wall MD, Lazari O, Arjomand J, Munoz-Sanjuan I. Advances in huntingtin disease drug discovery: Novel approaches to model disease phenotypes. *J Biomol Screen*. 2014;19(2):191-204.
182. Khan E, Mishra SK, Mishra R, Mishra A, Kumar A. Discovery of a potent small molecule inhibiting Huntington's disease (HD) pathogenesis via targeting CAG repeats RNA and Poly Q-protein. *Sci Rep*. 2019;9(1):16872.
183. Khan E, Tawani A, Mishra SK, Verma AK, Upadhyay A, Kumar M, et al. Myricetin reduces toxic level of CAG repeats RNA in Huntington's disease (HD) and spino cerebellar ataxia (SCAs). *ACS Chem Biol*. 2018;13(1):180-188.
184. Khan E, Biswas S, Mishra SK, Mishra R, Samanta S, Mishra A, et al. Rationally designed small molecules targeting toxic CAG repeat RNA that causes Huntington's disease (HD) and spinocerebellar ataxia (SCAs). *Biochimie*. 2019;163:21-32.
185. Lee J-M, Zhang J, Su AI, Walker JR, Wiltshire T, Kang K, et al. A novel approach to investigate tissue-specific trinucleotide repeat instability. *BMC Syst Biol*. 2010;4:29.
186. Garriga-Canut M, Agustín-Pavón C, Herrmann F, Sánchez A, Dierssen M, Fillat C, et al. Synthetic zinc finger repressors reduce mutant huntingtin expression in the brain of R6/2 mice. *Proc Natl Acad Sci USA*. 2012;109(45):E3136-E3145.
187. Agustín-Pavón C, Mielcarek M, Garriga-Canut M, Isalan M. Deimmunization for gene therapy: Host matching of synthetic zinc finger constructs enables long-term mutant huntingtin repression in mice. *Mol Neurodegener*. 2016;11(1):64.
188. Zeitler B, Froelich S, Marlen K, Shivak DA, Yu Q, Li D, et al. Allele-selective transcriptional repression of mutant HTT for the treatment of Huntington's disease. *Nat Med*. 2019;25(7):1131-1142.
189. Menalled LB. Knock-in mouse models of Huntington's disease. *NeuroRx*. 2005;2(3):465-470.
190. Nemudnyi AA, Valetdinova KR, Medvedev SP, Zakian SM. TALEN and CRISPR/Cas genome editing systems: Tools of discovery. *Acta naturae*. 2014;6(3):19-40.
191. Fink KD, Deng P, Gutierrez J, Anderson JS, Torrest A, Komarla A, et al. Allele-specific reduction of the mutant huntingtin allele using transcription activator-like effectors in human Huntington's disease fibroblasts. *Cell Transplantation*. 2016;25(4):677-686.
192. Richard G-F, Viterbo D, Khanna V, Mosbach V, Castelain L, Dujon B. Highly specific contractions of a single CAG/CTG trinucleotide repeat by TALEN in yeast. *PLoS One*. 2014;9(4):e95611.
193. Dabrowska M, Juzwa W, Krzyzosiak WJ, Olejniczak M. Precise excision of the CAG tract from the Huntingtin gene by Cas9 nickases. *Front Neurosci*. 2018;12:75.
194. Xu X, Tay Y, Sim B, Yoon S-I, Huang Y, Ooi J, et al. Reversal of phenotypic abnormalities by CRISPR/Cas9-mediated gene correction in Huntington disease patient-derived induced pluripotent stem cells. *Stem Cell Reports*. 2017;8(3):619-633.
195. Shin JW, Kim K-H, Chao MJ, Atwal RS, Gillis T, MacDonald ME, et al. Permanent inactivation of Huntington's disease mutation by personalized allele-specific CRISPR/Cas9. *Hum Mol Genet*. 2016;25(20):4566-4576.
196. Yang S, Chang R, Yang H, Zhao T, Hong Y, Kong HE, et al. CRISPR/Cas9-mediated gene editing ameliorates neurotoxicity in mouse model of Huntington's disease. *J Clin Invest*. 2017;127(7):2719-2724.
197. Ekman F, Ojala D, Adil M, Lopez P, Schaffer D, Gaj T. CRISPR-Cas9-mediated genome editing increases lifespan and improves motor deficits in a Huntington's disease mouse model. *Mol Ther Nucleic Acids*. 2019;17:829-839.
198. Monteyes AM, Ebanks SA, Keiser MS, Davidson BL. CRISPR/Cas9 editing of the mutant Huntingtin Allele In vitro and in vivo. *Mol Ther*. 2017;25(1):12-23.
199. Salado-Manzano C, Perpiña U, Straccia M, Molina-Ruiz FJ, Cozzi E, Rosser AE, et al. Is the immunological response a bottleneck for cell therapy in neurodegenerative diseases? *Front Cell Neurosci*. 2020;14:250.
200. Azidus Brasil. Safety Evaluation of Cellavita HD Administered Intravenously in Participants With Huntington's Disease (SAVE-DH) 2016 [cited 2021 13 January]. Available from: <https://clinicaltrials.gov/ct2/show/NCT02728115>.
201. Azidus Brasil. Dose-response Evaluation of the Cellavita HD Product in Patients With Huntington's Disease (ADORE-DH) 2017. Available from: <https://clinicaltrials.gov/ct2/show/NCT03252535>.
202. Azidus Brasil. Clinical Extension Study for Safety and Efficacy Evaluation of Cellavita-HD Administration in Huntington's Patients. (ADORE-EXT) 2020 [cited 2021 13 January]. Available from: <https://clinicaltrials.gov/ct2/show/NCT04219241>.
203. Rosser AE, Barker RA, Harrower T, Watts C, Farrington M, Ho AK, et al. Unilateral transplantation of human primary fetal tissue in four patients with Huntington's disease: NEST-UK safety report ISRCTN no 36485475. *J Neurol Neurosurg Psychiatr*. 2002;73(6):678-685.
204. Barker RA, Mason SL, Harrower TP, Swain RA, Ho AK, Sahakian BJ, et al. The long-term safety and efficacy of bilateral transplantation of human fetal striatal tissue in patients with mild to moderate Huntington's disease. *J Neurol Neurosurg Psychiatr*. 2013;84(6):657-665.
205. Bachoud-Lévi A-C, Remy P, Nguyen J-P, Brugieres P, Lefaucheur J-P, Bourdet C, et al. Motor and cognitive improvements in patients with Huntington's disease after neural transplantation. *Lancet*. 2000;356(9246):1975.
206. Paganini M, Biggeri A, Romoli AM, Mechi C, Ghelli E, Berti V, et al. Fetal striatal grafting slows motor and cognitive decline of Huntington's disease. *J Neurol Neurosurg Psychiatr*. 2014;85(9):974-981.
207. Drew C, Rosser A, Gray W. A single site, open label, phase I study to assess the safety and feasibility of foetal cell transplants in the striatum of people with Huntington's disease 2017. Available from: <http://www.isrctn.com/ISRCTN52651778>.
208. Bachoud-Lévi AC, Bourdet C, Brugieres P, Nguyen JP, Grandmougin T, Haddad B, et al. Safety and tolerability assessment of intrastriatal neural allografts in five patients with Huntington's disease. *Exp Neurol*. 2000;161(1):194-202.
209. Maxan A, Mason S, Saint-Pierre M, Smith E, Ho A, Harrower T, et al. Outcome of cell suspension allografts in a patient with Huntington's disease. *Ann Neurol*. 2018;84(6):950-956.
210. Krystkowiak P, Gaura V, Labalette M, Rialland A, Remy P, Peschanski M, et al. Alloimmunisation to donor antigens and immune rejection following foetal neural grafts to the brain in patients with Huntington's disease. *PLoS One*. 2007;2(1):e166.
211. Krebs SS, Trippel M, Prokop T, Omer TN, Landwehrmeyer B, Weber WA, et al. Immune response after striatal engraftment of fetal neuronal cells in patients with Huntington's disease: Consequences for cerebral transplantation programs. *Clinic Exp Neuroimmunol*. 2011;2(2):25-32.
212. Bachoud-Lévi AC, Gaura V, Brugieres P, Lefaucheur JP, Boissé MF, Maison P, et al. Effect of fetal neural transplants in patients with Huntington's disease 6 years after surgery: A long-term follow-up study. *Lancet Neurol*. 2006;5(4):303-309.
213. Cisbani G, Freeman TB, Soulet D, Saint-Pierre M, Gagnon D, Parent M, et al. Striatal allografts in patients with Huntington's disease: Impact of diminished astrocytes and vascularization on graft viability. *Brain*. 2013;136(Pt 2):433-443.
214. Cho IK, Hunter CE, Ye S, Pongos AL, Chan AWS. Combination of stem cell and gene therapy ameliorates symptoms in Huntington's disease mice. *Npj Regenerative Medicine*. 2019;4(1):7.
215. Lee ST, Chu K, Jung KH, Im WS, Park JE, Lim HC, et al. Slowed progression in models of Huntington disease by adipose stem cell transplantation. *Ann Neurol*. 2009;66(5):671-681.
216. Welter R. Autologous Stem/Stromal Cells in Neurological Disorders and Disease (NDD) 2017 [cited 2021 20 February]. Available from: <https://clinicaltrials.gov/ct2/show/NCT03297177>.
217. Haddad MS, Wenceslau CV, Pompeia C, Kerkis I. Cell-based technologies for Huntington's disease. *Dem Neuropsychol*. 2016;10(4):287-295.
218. Fink KD, Deng P, Torrest A, Stewart H, Pollock K, Gruenloh W, et al. Developing stem cell therapies for juvenile and adult-onset Huntington's disease. *Regen Med*. 2015;10(5):623-646.
219. Fink KD, Rossignol J, Crane AT, Davis KK, Bombard MC, Bavar AM, et al. Transplantation of umbilical cord-derived mesenchymal stem cells into the striata of R6/2 mice: Behavioral and neuropathological analysis. *Stem Cell Research & Therapy*. 2013;4(5):130.
220. Ebrahimi MJ, Aliaghaei A, Boroujeni ME, Khodaghali F, Meftahi G, Abdollahifar MA, et al. human umbilical cord matrix stem cells reverse oxidative stress-induced cell death and ameliorate motor function and striatal atrophy in rat model of Huntington Disease. *Neurotox Res*. 2018;34(2):273-284.
221. Vis JC, Verbeek MM, De Waal RM, Ten Donkelaar HJ, Kremer HP. 3-Nitropropionic acid induces a spectrum of Huntington's disease-like neuropathology in rat striatum. *Neuropathol Appl Neurobiol*. 1999;25(6):513-521.
222. Rossignol J, Boyer C, Lévêque X, Fink KD, Thinard R, Blanchard F, et al. Mesenchymal stem cell transplantation and DMEM administration in a 3NP rat model of Huntington's disease: Morphological and behavioral outcomes. *Behav Brain Res*. 2011;217(2):369-378.
223. Liu J. Induced pluripotent stem cell-derived neural stem cells: New hope for stroke? *Stem Cell Res Ther*. 2013;4(5):115.
224. Mu S, Wang J, Zhou G, Peng W, He Z, Zhao Z, et al. Transplantation of induced pluripotent stem cells improves functional recovery in Huntington's disease rat model. *PLoS One*. 2014;9(7):e101185.
225. Sanberg PR, Calderon SF, Giordano M, Tew JM, Norman AB. The quinolinic acid model of Huntington's disease: Locomotor abnormalities. *Exp Neurol*. 1989;105(1):45-53.
226. Yoon Y, Kim HS, Hong CP, Li E, Jeon I, Park HJ, et al. Neural transplants from human induced pluripotent stem cells rescue the pathology and behavioral defects in a rodent model of Huntington's disease. *Front Neurosci*. 2020;14:558204.

227. Al-Gharaibeh A, Culver R, Stewart AN, Srinageshwar B, Spelde K, Frollo L, et al. Induced pluripotent stem cell-derived neural stem cell transplantations reduced behavioral deficits and ameliorated neuropathological changes in YAC128 mouse model of Huntington's disease. *Front Neurosci.* 2017;11:628.
228. Lansita JA, Mease KM, Qiu H, Yednock T, Sankaranarayanan S, Kramer S. Nonclinical development of ANX005: A humanized anti-C1q antibody for treatment of autoimmune and neurodegenerative diseases. *Int J Toxicol.* 2017; 36(6):449-462.
229. Keswani SC. Phase 2a Study of ANX005, a Humanized anti-C1q mAb, in Patients with Huntington disease. Huntington Study Group Annual Conference; October 30; Online2020.
230. Cho K. Emerging roles of complement protein C1q in neurodegeneration. *Aging Dis.* 2019;10(3):652-663.
231. Annexon I. Single Dose Study of ANX005 in Healthy Volunteers 2017. Available from: <https://clinicaltrials.gov/ct2/show/NCT03010046>.
232. Annexon I. A Clinical Study of ANX005 and IVIG in Subjects With Guillain Barré Syndrome (GBS) 2019. Available from: <https://clinicaltrials.gov/ct2/show/NCT04035135> <https://clinicaltrials.gov/ct2/show/NCT04035135>.
233. Annexon I. An Open Label Study of ANX005 in Subjects With, or at Risk for, Manifest Huntington's Disease 2020 [cited 2021 26 January]. Available from: <https://clinicaltrials.gov/ct2/show/NCT04514367>.
234. Annexon I. Study of ANX005 in Adults With Amyotrophic Lateral Sclerosis (ALS) 2020. Available from: <https://clinicaltrials.gov/ct2/show/NCT04569435>.
235. Annexon I. Safety, Tolerability, Pharmacokinetics and Pharmacodynamics of ANX005 in Subjects With Warm Autoimmune Hemolytic Anemia (wAIHA) 2020 [cited 2021 28 January]. Available from: <https://clinicaltrials.gov/ct2/show/NCT04691570>.
236. Islam Z, Papri N, Jahan I, Azad KAK, Kroon H, Humphris E, et al. Inhibition of C1q, initiator of the classical complement cascade, by ANX005 for the treatment of Guillain-Barré Syndrome. Results from a Phase 1b study. 2020 American Academy of Neurology (AAN) Annual Meeting; 28 April; Toronto, Canada2020.
237. Annexon I. Efficacy and Safety of ANX005 in Subjects With Guillain-Barré Syndrome 2021. Available from: <https://clinicaltrials.gov/ct2/show/NCT04701164>.
238. Leonard JE, Fisher TL, Winter LA, Cornelius CA, Reilly C, Smith ES, et al. Nonclinical safety evaluation of VX15/2503, a humanized IgG4 Anti-SEMA4D antibody. *Mol Cancer Therapeut.* 2015;14(4):964.
239. Smith ES, Jonason A, Reilly C, Veeraraghavan J, Fisher T, Doherty M, et al. SEMA4D compromises blood-brain barrier, activates microglia, and inhibits remyelination in neurodegenerative disease. *Neurobiol Dis.* 2015;73:254-268.
240. Southwell AL, Franciosi S, Villanueva EB, Xie Y, Winter LA, Veeraraghavan J, et al. Anti-semaphorin 4D immunotherapy ameliorates neuropathology and some cognitive impairment in the YAC128 mouse model of Huntington disease. *Neurobiol Dis.* 2015;76:46-56.
241. Vaccinex I. A Study in Subjects With Late Prodromal and Early Manifest Huntington's Disease (HD) to Assess the Safety, Tolerability, Pharmacokinetics, and Efficacy of Pepinemab (VX15/2503) (SIGNAL) 2015. Available from: <https://clinicaltrials.gov/ct2/show/NCT02481674>.
242. Rodrigues FB, Wild EJ. Huntington's disease clinical trials corner: August 2018. *J Huntingt Dis.* 2018;7(3):279-286.
243. Vaccinex, Inc. Announces Preliminary Data from the SIGNAL Clinical Trial (Investigational Drug VX15/2503 as a Potential Treatment for Huntington's Disease) [press release]. Rochester, New York 2017.
244. Top-Line Results of Phase 2 SIGNAL Study in Huntington's Disease Support Potential for Cognitive Benefit of Pepinemab [press release]. September 22 2020 2020.
245. Vaccinex I. Learnings from the SIGNAL Phase 2 Study of Treatment with Pepinemab Antibody. Huntington Study Group 2020 Medical Conference; 30 October 2020; Virtual2020.
246. University of Utah. Anti-SEMA4D Monoclonal Antibody VX15/2503 With Nivolumab or Ipilimumab in Treating Patients With Stage III or IV Melanoma 2018. Available from: <https://www.clinicaltrials.gov/ct2/show/NCT03425461?term=VX15&draw=5&rank=2>.
247. Vaccinex I. SEMA4D Blockade Safety and Brain Metabolic Activity in Alzheimer's Disease (AD) (SIGNAL-AD) 2020. Available from: <https://www.clinicaltrials.gov/ct2/show/NCT04381468>.
248. Emory University. VX15/2503 and Immunotherapy in Resectable Pancreatic and Colorectal Cancer 2017. Available from: <https://www.clinicaltrials.gov/ct2/show/NCT03373188> <https://www.clinicaltrials.gov/ct2/show/NCT03373188>.
249. Emory University. VX15/2503 in Combination With Ipilimumab or Nivolumab in Patients With Head and Neck Cancer 2018. Available from: <https://www.clinicaltrials.gov/ct2/show/NCT03690986>.
250. LaGanke C, Samkoff L, Edwards K, Jung Henson L, Repovic P, Lynch S, et al. Safety/tolerability of the anti-semaphorin 4D Antibody VX15/2503 in a randomized phase 1 trial. *Neurol Neuroimmunol Neuroinflamm.* 2017;4(4):e367.
251. Patnaik A, Weiss GJ, Leonard JE, Rasco DW, Sachdev JC, Fisher TL, et al. Safety, pharmacokinetics, and pharmacodynamics of a humanized anti-semaphorin 4D antibody, in a first-in-human study of patients with advanced solid tumors. *Clin Cancer Res.* 2016;22(4):827-836.
252. Vaccinex I. VX15/2503 in Combination With Avelumab in Advanced Non-small Cell Lung Cancer (CLASSICAL-Lung) 2017. Available from: <https://www.clinicaltrials.gov/ct2/show/NCT03268057>.
253. Khoshnan A, Southwell AL, Bugg CW, Ko JC, Patterson PH. recombinant intrabodies as molecular tools and potential therapeutics for Huntington's disease. In: Lo DC, Hughes RE, eds *Neurobiology of Huntington's Disease: Applications to Drug Discovery*. Boca Raton (FL: CRC Press/Taylor & Francis; 2011.
254. Amaro IA, Henderson LA. An intrabody drug (rAAV6-INT41) reduces the binding of N-terminal Huntingtin fragment(s) to DNA to basal levels in PC12 cells and delays cognitive loss in the R6/2 animal model. *J Neurodegen Dis.* 2016; 2016:7120753.
255. Zha J, Liu X-M, Zhu J, Liu S-Y, Lu S, Xu P-X, et al. A scFv antibody targeting common oligomeric epitope has potential for treating several amyloidoses. *Sci Rep.* 2016;6:36631.
256. Southwell AL, Khoshnan A, Dunn DE, Bugg CW, Lo DC, Patterson PH. Intrabodies binding the proline-rich domains of mutant huntingtin increase its turnover and reduce neurotoxicity. *J Neurosci.* 2008;28(36):9013-9020.
257. Miller TW, Zhou C, Gines S, MacDonald ME, Mazarakis ND, Bates GP, et al. A human single-chain Fv intrabody preferentially targets amino-terminal huntingtin fragments in striatal models of Huntington's disease. *Neurobiol Dis.* 2005;19(1):47-56.
258. Butler DC, Messer A. Bifunctional anti-huntingtin proteasome-directed intrabodies mediate efficient degradation of mutant huntingtin exon 1 protein fragments. *PLoS One.* 2011;6(12):e29199.
259. Denis HL, Lauriol F, Cicchetti F. Are immunotherapies for Huntington's disease a realistic option?. *Mol Psychiatry.* 2019;24:364-377.
260. Positive preclinical in vivo results with AFFiRS' antibody mAb C6-17 to treat Huntington's disease to be presented at the 16th Annual Huntington's Disease Therapeutics Conference [press release]. April 27 2021.
261. Bartl S, Oueslati A, Southwell AL, Siddu A, Parth M, David LS, et al. Inhibiting cellular uptake of mutant huntingtin using a monoclonal antibody: Implications for the treatment of Huntington's disease. *Neurobiol Dis.* 2020;141:104943.
262. Waters S, Tedroff J, Ponten H, Klammer D, Sonesson C, Waters N. Pridopidine: Overview of pharmacology and rationale for its use in Huntington's disease. *J Huntingt Dis.* 2018;7(1):1-16.
263. Reilmann R, McGarry A, Grachev ID, Savola JM, Borowsky B, Eyal E, et al. Safety and efficacy of pridopidine in patients with Huntington's disease (PRIDE-HD): A phase 2, randomised, placebo-controlled, multicentre, dose-ranging study. *Lancet Neurol.* 2019;18(2):165-176.
264. Geva M, Kusko R, Soares H, Fowler KD, Birnberg T, Barash S, et al. Pridopidine activates neuroprotective pathways impaired in Huntington Disease. *Hum Mol Genet.* 2016;25(18):3975-3987.
265. Squitieri F, Di Pardo A, Favellato M, Amico E, Maglione V, Frati L. Pridopidine, a dopamine stabilizer, improves motor performance and shows neuroprotective effects in Huntington disease R6/2 mouse model. *J Cell Mol Med.* 2015;19(11):2540-2548.
266. Ryskamp D, Wu J, Geva M, Kusko R, Grossman I, Hayden M, et al. The sigma-1 receptor mediates the beneficial effects of pridopidine in a mouse model of Huntington disease. *Neurobiol Dis.* 2017;97(Pt A):46-59.
267. Lundin A, Dietrichs E, Haghighi S, Göller M-L, Heiberger A, Loutfi G, et al. Efficacy and safety of the dopaminergic stabilizer Pridopidine (ACR16) in patients with Huntington's disease. *Clin Neuropharmacol.* 2010;33(5):260-264.
268. de Yebenes JG, Landwehrmeyer B, Squitieri F, Reilmann R, Rosser A, Barker RA, et al. Pridopidine for the treatment of motor function in patients with Huntington's disease (MermaiHD): A phase 3, randomised, double-blind, placebo-controlled trial. *Lancet Neurol.* 2011;10(12):1049-1057.
269. Huntington Study Group HART Investigators. A randomized, double-blind, placebo-controlled trial of pridopidine in Huntington's disease. *Mov Disord.* 2013; 28(10):1407-1415.
270. Prilenia. A Phase 2, to Evaluating the Safety and Efficacy of Pridopidine Versus Placebo for Symptomatic Treatment in Patients With Huntington's Disease 2013. Available from: <https://clinicaltrials.gov/ct2/show/NCT02006472>.
271. McGarry A, Leinonen M, Kiebertz K, Geva M, Olanow CW, Hayden M. Effects of Pridopidine on Functional Capacity in Early-Stage Participants from the PRIDE-HD Study. *J Huntingt Dis.* 2020;9(4):371-380.
272. Squitieri F, Landwehrmeyer B, Reilmann R, Rosser A, de Yebenes JG, Prang A, et al. One-year safety and tolerability profile of pridopidine in patients with Huntington disease. *Neurology.* 2013;80(12):1086-1094.
273. McGarry A, Kiebertz K, Abler V, Grachev ID, Gandhi S, Auinger P, et al. Safety and exploratory efficacy at 36 months in open-HART, an open-label extension study of pridopidine in Huntington's disease. *J Huntingt Dis.* 2017;6:189-199.
274. McGarry A, Auinger P, Kiebertz K, Geva M, Mehra M, Abler V, et al. Additional safety and exploratory efficacy data at 48 and 60 months from open-HART, an open-label extension study of pridopidine in Huntington disease. *J Huntingt Dis.* 2020;9:173-184.
275. Prilenia. A Study Evaluating if Pridopidine is Safe, Efficacious, and Tolerable in Patients With Huntington's Disease (Open PRIDE-HD) 2015. Available from: <https://clinicaltrials.gov/ct2/show/NCT02494778>.
276. Prilenia. PRIDopidine's Outcome On Function in Huntington Disease, PROOF-HD 2020. Available from: <https://clinicaltrials.gov/ct2/show/NCT04556656>.

277. Rodrigues FB, Wild EJ. Clinical trials corner: September 2017. *J Huntingt Dis*. 2017;6(3):255-263.
278. Haggiag S, Ruggieri S, Gasperini C. Efficacy and safety of laquinimod in multiple sclerosis: Current status. *Therap Adv Neurol Dis*. 2013;6(6):343-352.
279. Brück W, Pfortner R, Pham T, Zhang J, Hayardeny L, Piryatinsky V, et al. Reduced astrocytic NF- κ B activation by laquinimod protects from cuprizone-induced demyelination. *Acta Neuropathologica*. 2012;124(3):411-424.
280. Filippi M, Rocca MA, Pagani E, De Stefano N, Jeffery D, Kappos L, et al. Placebo-controlled trial of oral laquinimod in multiple sclerosis: MRI evidence of an effect on brain tissue damage. *J Neurol Neurosurg Psychiatr*. 2014;85(8):851.
281. Garcia-Miralles M, Tan J, Radulescu C, Sidik H, Belinson H, Zach N, et al. Laquinimod treatment improves myelination deficits at the transcriptional and ultrastructural levels in the YAC128 mouse model of Huntington disease. *Mol Neurobiol*. 2019;56(6):4464-4478.
282. Teva Pharmaceutical Industries. A Clinical Study in Participants With Huntington's Disease (HD) to Assess Efficacy and Safety of Three Oral Doses of Laquinimod (LEGATO-HD) 2014 [updated May 4 2020; cited 2021 17 February]. Available from: <https://clinicaltrials.gov/ct2/show/NCT02215616>.
283. Reilmann R, Gordon MF, Anderson KE, Feigin A, Tabrizi SJ, Leavitt BR, et al. The efficacy and safety results of laquinimod as a treatment for Huntington disease (LEGATO-HD) (S16.007). *Neurology*. 2019;92(15 suppl ment):S16.007.
284. Leavitt BR, Reilmann R, Gordon MF, Anderson KE, Feigin A, Tabrizi SJ, et al. eds. Magnetic resonance spectroscopy evaluation of neuronal integrity and astrogliosis in a phase 2 study of laquinimod as a treatment for Huntington disease (LEGATO-HD) *International Congress of Parkinson's Disease and Movement disorders*. Nice, France; 2019.
285. Reilmann R, Gordon MF, Schubert R, Anderson KE, Feigin A, Tabrizi SJ, et al. eds. Quantitative motor (Q-Motor) assessments suggest a beneficial central effect of laquinimod in a phase II study in Huntington disease (LEGATO-HD) *International Congress of Parkinson's Disease and Movement disorders*. NiceFrance2019; 2019.
286. Bachstetter AD, Xing B, de Almeida L, Dimayuga ER, Watterson DM, Van Eldik LJ. Microglial p38 α MAPK is a key regulator of proinflammatory cytokine up-regulation induced by toll-like receptor (TLR) ligands or beta-amyloid (A β). *J Neuroinflammation*. 2011;8:79.
287. Alam JJ. Selective brain-targeted antagonism of p38 MAPK α reduces hippocampal IL-1 β levels and improves Morris Water maze performance in aged rats. *J Alzheim Dis : JAD*. 2015;48(1):219-227.
288. Haddad JJ. VX-745. Vertex pharmaceuticals. *Curr Opin Invest Drugs*. 2001;2(8):1070-1076.
289. EIP Pharma Inc. Within Subject Crossover Study of Cognitive Effects of Neflamapimod in Early-Stage Huntington Disease 2019 [cited 2021 18 February]. Available from: <https://clinicaltrials.gov/ct2/show/NCT03980938>.
290. EIP Pharma Announces Positive Phase 2 Results for Neflamapimod in Mild-to-Moderate Dementia with Lewy bodies (DLB) [press release]. Boston, Massachusetts, 6 October 2020 2020.
291. EIP Pharma Announces Clinical Trial Results of REVERSE-SD, a Phase 2b Study of Neflamapimod in Early-stage Alzheimer's Disease [press release]. Boston, Massachusetts, November 7 2019.
292. Pagan F, Hebron M, Valadez EH, Torres-Yaghi Y, Huang X, Mills RR, et al. Nilotinib effects in Parkinson's disease and dementia with lewy bodies. *J Parkinsons Dis*. 2016;6(3):503-517.
293. Hebron ML, Lonskaya I, Moussa CEH. Nilotinib reverses loss of dopamine neurons and improves motor behavior via autophagic degradation of α -synuclein in Parkinson's disease models. *Hum Mol Genet*. 2013;22(16):3315-3328.
294. Hebron ML, Lonskaya I, Olopade P, Selby ST, Pagan F, Moussa CEH. Tyrosine kinase inhibition regulates early systemic immune changes and modulates the neuroimmune response in α -synucleinopathy. *J Clin Cell Immunol*. 2014;5:259.
295. Anderson KE. Nilotinib in Huntington's Disease (Tasigna HD) 2018 [cited 2021 20 February]. Available from: <https://clinicaltrials.gov/ct2/show/NCT03764215>.
296. Pfizer. Randomized, Placebo Controlled Study Of The Efficacy And Safety Of PF-02545920 In Subjects With Huntington's Disease 2014 [cited 2021 18 February]. Available from: <https://clinicaltrials.gov/ct2/show/NCT02197130>.
297. Beaumont V, Zhong S, Lin H, Xu W, Bradaia A, Steidl E, et al. Phosphodiesterase 10A inhibition improves cortico-basal ganglia function in Huntington's disease models. *Neuron*. 2016;92(6):1220-1237.
298. Pfizer. Product Pipeline for New Pharmaceutical Drugs 2021 [updated February 2 2021; cited 2021 18 February]. Available from: <https://www.pfizer.com/science/drug-product-pipeline>.
299. Liu GZ, Hou TT, Yuan Y, Hang PZ, Zhao JJ, Sun L, et al. Fenofibrate inhibits atrial metabolic remodelling in atrial fibrillation through PPAR- α /sirtuin 1/PGC-1 α pathway. *Br J Pharmacol*. 2016;173(6):1095-1109.
300. Johri A, Chandra A, Flint Beal M. PGC-1 α , mitochondrial dysfunction, and Huntington's disease. *Free Radic Biol Med*. 2013;62:37-46.
301. Brownstein MJ, Simon NG, Long JD, Yankey J, Maibach HT, Cudkowicz M, et al. Safety and tolerability of SRX246, a vasopressin 1a antagonist, in irritable Huntington's disease patients—a randomized phase 2 clinical trial. *J Clin Med*. 2020;9(11):3682.
302. McGregor AL, Dysart J, Tingle MD, Russell BR, Kydd RR, Finucane G. Varenicline improves motor and cognitive symptoms in early Huntington's disease. *Neuropsychiatric Disease and Treatment*. 2016;12:2381-2386.
303. Sage Therapeutics Announces Planned Progression of SAGE-718 to Phase 2 in Huntington's Disease and Presentations at the 2019 Annual Meeting of the American College of Neuropsychopharmacology (ACNP) [press release]. December 10 2019.
304. Sage Therapeutics to Host Sage Science Spotlight: SAGE-718 In Depth [press release]. May 13 2021.
305. Huntington Study Group Reach2HD Investigators. Safety, tolerability, and efficacy of PBT2 in Huntington's disease: A phase 2, randomised, double-blind, placebo-controlled trial. *Lancet Neurol*. 2015;14(1):39-47.
306. Li C, Yuan K, Schluesener H. Impact of minocycline on neurodegenerative diseases in rodents: A meta-analysis. *Rev Neurosci*. 2013;24(5):553.
307. Bonelli RM, Hödl AK, Hofmann P, Kapfhammer H-P. Neuroprotection in Huntington's disease: A 2-year study on minocycline. *Int Clin Psychopharmacol*. 2004;19(6):337.
308. Thomas M, Ashizawa T, Jankovic J. Minocycline in Huntington's disease: A pilot study. *Mov Disord*. 2004;19(6):692-695.
309. Cudkowicz M. A futility study of minocycline in Huntington's disease. *Mov Disord*. 2010;25(13):2219-2224.
310. Möller T. Neuroinflammation in Huntington's disease. *J Neural Transm*. 2010;117(8):1001-1008.
311. Rocha NP, Ribeiro FM, Furr-Stimming E, Teixeira AL. Neuroimmunology of Huntington's Disease: Revisiting Evidence from Human Studies. *Mediat Inflamm*. 2016;2016:8653132.
312. Stella N. Endocannabinoid signaling in microglial cells. *Neuropharmacology*. 2009;56:244-253.
313. Consroe P, Laguna J, Allender J, Snider S, Stern L, Sandyk R, et al. Controlled clinical trial of cannabidiol in Huntington's disease. *Pharmacol Biochem Behav*. 1991;40(3):701-708.
314. López-Sendón Moreno JL, García Caldentey J, Trigo Cubillo P, Ruiz Romero C, García Ribas G, Alonso Arias M, et al. A double-blind, randomized, cross-over, placebo-controlled, pilot trial with Sativex in Huntington's disease. *J Neurol*. 2016;263(7):1390-1400.
315. Masgrau R, Guaza C, Ransohoff RM, Galea E. Should we stop saying 'glia' and 'neuroinflammation'? *Trends Mol Med*. 2017;23(6):486-500.
316. Matias I, Morgado J, Gomes FCA. Astrocyte heterogeneity: Impact to brain aging and disease. *Front Aging Neurosci* 2019.
317. Ransohoff RM. A polarizing question: Do M1 and M2 microglia exist? *Nat Neurosci*. 2016;19(8):987-991.
318. Sturrock A, Leavitt BR. The clinical and genetic features of Huntington disease. *J Geriatr Psychiatr Neurol*. 2010;23(4):243-259.
319. Frank S. Treatment of Huntington's disease. *Neurotherapeutics*. 2014;11(1):153-160.
320. André VM, Cepeda C, Levine MS. Dopamine and glutamate in Huntington's disease: A balancing act. *CNS Neuroscience & Therapeutics*. 2010;16(3):163-178.
321. Chen J, Wang E, Cepeda C, Levine M. Dopamine imbalance in Huntington's disease: A mechanism for the lack of behavioral flexibility. *Front Neurosci*. 2013;7.
322. Miller BR, Bezprozvanny I. Corticostriatal circuit dysfunction in Huntington's disease: Intersection of glutamate, dopamine and calcium. *Future Neurol*. 2010;5(5):735-756.
323. Zuccato C, Cattaneo E. Role of brain-derived neurotrophic factor in Huntington's disease. *Prog Neurobiol*. 2007;81(5):294-330.
324. Xie Y, Hayden MR, Xu B. BDNF overexpression in the forebrain rescues Huntington's disease phenotypes in YAC128 mice. *J Neurosci*. 2010;30(44):14708-14718.
325. Erta M, Quintana A, Hidalgo J, Interleukin-6, a major cytokine in the central nervous system. *Int J Biol Sci*. 2012;8(9):1254-1266.
326. Costa V, Scorrano L. Shaping the role of mitochondria in the pathogenesis of Huntington's disease. *EMBO J*. 2012;31(8):1853-1864.
327. Chaudhary RK, Patel KA, Patel MK, Joshi RH, Roy I. Inhibition of Aggregation of Mutant Huntingtin by Nucleic Acid Aptamers In Vitro and in a Yeast Model of Huntington's Disease. *Mol Ther*. 2015;23(12):1912-1926.
328. Zhang X, Smith DL, Meriin AB, Engemann S, Russel DE, Roark M, et al. A potent small molecule inhibits polyglutamine aggregation in Huntington's disease neurons and suppresses neurodegeneration in vivo. *Proc Natl Acad Sci USA*. 2005;102(3):892.
329. Senior K. Synaptic activity could hold the key to therapy for Huntington disease. *Nat Rev Neurol*. 2010;6(1):1.
330. Sepers MD, Raymond LA. Mechanisms of synaptic dysfunction and excitotoxicity in Huntington's disease. *Drug Discov Today*. 2014;19(7):990-996.
331. Marques Sousa C, Humbert S. Huntingtin: Here, there, everywhere! *J Huntingt Dis*. 2013;2(4):395-403.
332. Carroll JB, Bates GP, Steffan J, Saft C, Tabrizi SJ. Treating the whole body in Huntington's disease. *Lancet Neurol*. 2015;14(11):1135-1142.
333. van der Burg JMM, Björkqvist M, Brundin P. Beyond the brain: Widespread pathology in Huntington's disease. *Lancet Neurol*. 2009;8(8):765-774.
334. Kong G, Ellul S, Narayana VK, Kanojia K, Ha HTT, Li S, et al. An integrated metagenomics and metabolomics approach implicates the microbiota-gut-brain axis in the pathogenesis of Huntington's disease. *Neurobiol Dis*. 2021;148:105199.

335. Du G, Dong W, Yang Q, Yu X, Ma J, Gu W, et al. Altered gut microbiota related to inflammatory responses in patients with Huntington's disease. *Front Immunol.* 2020;11:603594.
336. Stan TL, Soylu-Kucharz R, Burleigh S, Prykhodko O, Cao L, Franke N, et al. Increased intestinal permeability and gut dysbiosis in the R6/2 mouse model of Huntington's disease. *Sci Rep.* 2020;10(1):18270.
337. Wasser CI, Mercieca E-C, Kong G, Hannan AJ, McKeown SJ, Glikmann-Johnston Y, et al. Gut dysbiosis in Huntington's disease: Associations among gut microbiota, cognitive performance and clinical outcomes. *Brain Commun.* 2020;2(2):fcaa110.
338. Kong G, Cao KL, Judd LM, Li S, Renoir T, Hannan AJ. Microbiome profiling reveals gut dysbiosis in a transgenic mouse model of Huntington's disease. *Neurobiol Dis.* 2020;135:104268.
339. McColgan P, Tabrizi SJ. Huntington's disease: A clinical review. *Eur J Neurol.* 2018;25(1):24-34.
340. Harper SQ. Progress and challenges in RNA interference therapy for Huntington disease. *Arch Neurol.* 2009;66(8):933-938.
341. Meghan S, Neil A. Frontiers in neuroscience RNA- and DNA- based therapies for Huntington's disease. In: Lo DC, Hughes RE, eds *Neurobiology of Huntington's Disease: Applications to Drug Discovery*. Boca Raton (FL: CRC Press/Taylor & Francis; 2011:.