

MEETING ABSTRACT

Open Access

Identification and characterization of small molecular NPR-B receptor antagonists

Henriette Andresen^{1,2}, Lise Román Moltzau^{1*}, Steffen Fagerheim^{1,3}, David Gloriam⁴, Trond Bach^{1,3}, Finn Olav Levy^{1,3}

From 7th International Conference on cGMP Generators, Effectors and Therapeutic Implications Trier, Germany. 19-21 June 2015

Background

Natriuretic peptides increase in heart failure. Atrial natriuretic peptide (ANP) and brain natriuretic peptide (BNP) activate the natriuretic peptide receptor (NPR)-A and C-type natriuretic peptide (CNP) the NPR-B, causing production of cyclic 3',5'-guanosine monophosphate (cGMP). Our group has previously shown that CNP potentiates β_1 -adrenoceptor-mediated inotropic response by increasing cAMP levels. This is explained by a cross-talk between cGMP and cAMP, where cGMP produced by NPR-B inhibits phosphodiesterase (PDE) 3 from degrading cAMP. Increased β_1 -adrenoceptor signalling is harmful in heart failure which is the basis for use of β -blockers in heart failure therapy. Further, PDE3 inhibition has been associated with increased mortality in heart failure. Thus, an NPR-B antagonist could eliminate the unwanted effects due to PDE3 inhibition and increased β_1 -adrenoceptor signalling by CNP. There are no selective NPR-B antagonists available, and our aim is to identify and characterize novel non-peptide small molecular selective NPR-B antagonists for potential use in heart failure therapy.

Methods

Small molecular compounds were screened for activity and selectivity towards NPR-A and NPR-B, using a cGMP assay based on AlphaScreen technology. Based on this technology, we can identify compounds that interfere with BNP- and CNP-stimulated cGMP production through the NPR-A and NPR-B receptor, respectively. Antagonism properties can also be characterized.

Results and conclusion

Potential NPR-B antagonists were identified by high throughput screening of about 20,000 compounds. Hits were tested against the NPR-A to determine selectivity, and characterized as non-competitive and reversible inhibitors of NPR-B response in HEK293 cells, displaying properties of negative allosteric modulators [1]. Information on structure and activity relationship (SAR), was collected and over 50 new compounds, clustered in five different groups of chemically related compounds, were designed *in silico* and were tested for activity and selectivity towards NPR-A and NPR-B. From this testing, three out of five groups were found interesting. The ligands with the best antagonistic property from each of the 3 groups were used to select 60 more compounds by *in silico* design for further testing.

Authors' details

¹Department of Pharmacology, University of Oslo and Oslo University Hospital, Oslo, Norway. ²School of Pharmacy, University of Oslo, Oslo, Norway. ³K.G. Jebsen Cardiac Research Centre and Center for Heart Failure Research, University of Oslo, Oslo, Norway. ⁴Department of Drug Design and Pharmacology, Faculty of Health and Medical Sciences, University of Copenhagen, Copenhagen, Denmark.

Published: 2 September 2015

Reference

1. Bach T, Bergholtz S, Riise J, Qvigstad E, Skomedal T, Osnes JB, et al: Identification of small molecule NPR-B antagonists by high throughput screening - potential use in heart failure. *Naunyn-Schmiedeberg's Arch Pharmacol* 2014, **387**(1):5-14.

doi:10.1186/2050-6511-16-S1-A70

Cite this article as: Andresen et al.: Identification and characterization of small molecular NPR-B receptor antagonists. *BMC Pharmacology and Toxicology* 2015 **16**(Suppl 1):A70.

* Correspondence: l.r.moltzau@medisin.uio.no

¹Department of Pharmacology, University of Oslo and Oslo University Hospital, Oslo, Norway

Full list of author information is available at the end of the article