

RESEARCH ARTICLE

Open Access

The role of iron uptake in pathogenicity and symbiosis in *Photorhabdus luminescens* TT01

Robert J Watson¹, Peter Millichap¹, Susan A Joyce^{1,2}, Stuart Reynolds¹ and David J Clarke^{*1,2}

Abstract

Background: *Photorhabdus* are Gram negative bacteria that are pathogenic to insect larvae whilst also having a mutualistic interaction with nematodes from the family *Heterorhabditis*. Iron is an essential nutrient and bacteria have different mechanisms for obtaining both the ferrous (Fe²⁺) and ferric (Fe³⁺) forms of this metal from their environments. In this study we were interested in analyzing the role of Fe³⁺ and Fe²⁺ iron uptake systems in the ability of *Photorhabdus* to interact with its invertebrate hosts.

Results: We constructed targeted deletion mutants of *exbD*, *feoABC* and *yfeABCD* in *P. luminescens* TT01. The *exbD* mutant was predicted to be crippled in its ability to obtain Fe³⁺ and we show that this mutant does not grow well in iron-limited media. We also show that this mutant was avirulent to the insect but was unaffected in its symbiotic interaction with *Heterorhabditis*. Furthermore we show that a mutation in *feoABC* (encoding a predicted Fe²⁺ permease) was unaffected in both virulence and symbiosis whilst the divalent cation transporter encoded by *yfeABCD* is required for virulence in the Tobacco Hornworm, *Manduca sexta* (Lepidoptera) but not in the Greater Wax Moth, *Galleria mellonella* (Lepidoptera). Moreover the Yfe transporter also appears to have a role during colonization of the IJ stage of the nematode.

Conclusion: In this study we show that iron uptake (via the TonB complex and the Yfe transporter) is important for the virulence of *P. luminescens* to insect larvae. Moreover this study also reveals that the Yfe transporter appears to be involved in Mn²⁺-uptake during growth in the gut lumen of the IJ nematode. Therefore, the Yfe transporter in *P. luminescens* TT01 is important during colonization of both the insect and nematode and, moreover, the metal ion transported by this pathway is host-dependent.

Background

Photorhabdus is a genus of Gram negative bioluminescent bacteria that are members of the Enterobacteriaceae and are therefore close relatives of important mammalian pathogens such as *Escherichia coli* and *Salmonella*. *Photorhabdus* have a complex life-style that involves a pathogenic interaction with insect larvae and a mutualistic interaction with nematodes from the family *Heterorhabditis* (for recent reviews see [1,2]). The bacteria can be normally found colonizing the gut of the infective juvenile (IJ) stage of the nematode. The IJ is a free-living, soil-dwelling stage of the nematode whose role is to seek out and infect susceptible insect larvae. Once inside the insect the IJ regurgitate their bacterial symbionts into the insect hemolymph and, here, the bacteria divide expo-

nentially [3,4]. The bacteria produce a range of activities, including hydrolytic enzymes, that contribute to the efficient conversion of the insects internal organs and tissues into bacterial biomass and the insect eventually dies of septicemia 48-72 hours post-infection [5]. At this point the IJ recovers to become an adult hermaphrodite that feeds on the bacterial biomass and lays eggs that develop through juvenile stages (L1-L4) before adulthood. After 2-3 rounds of nematode reproduction uncharacterized environmental signals stimulate the formation of an alternative L3 stage nematode called the IJ. The IJ is initially colonized by 1-2 *Photorhabdus* cells in a complex transmission process that has only recently been phenomenologically described [6]. These founder cells grow and divide resulting in a final population of *Photorhabdus* in the IJ of between 50-100 colony forming units (CFU). The IJs then emerge from the insect cadaver ready to search for more susceptible insect larvae.

* Correspondence: david.clarke@ucc.ie

¹ Department of Biology and Biochemistry, University of Bath, Bath BA2 7AY, UK
Full list of author information is available at the end of the article

The *Heterorhabditis* nematode is bacteriophorous and, during growth and development, the nematode feeds on the bacterial biomass present within the cadaver. Therefore the *Photorhabdus* cells must be able to satisfy the nutritional requirements of the nematode population. The genetic basis of the nutritional interaction between *Photorhabdus* and *Heterorhabditis* is not well understood. There is some evidence that crystalline inclusion proteins (encoded by *cipA* and *cipB*) produced by *Photorhabdus* have a role in nematode nutrition. Mutations in the *cipA* and *cipB* genes of *Photorhabdus luminescens* NC1 resulted in a strain that was unable to support nematode growth and development [7]. However the *cipA* and *cipB* mutations were pleiotropic making it difficult to confidently define a role for these proteins in nematode nutrition. Nonetheless it has been shown that overproduction of CipA and CipB in *E. coli* can improve the growth and development of *Steinernema* nematodes implying some role for these proteins in nematode nutrition [8]. A mutation in another gene, *ngrA*, encoding a phosphopantetheinyl (P'ant) transferase, was also shown to prevent nematode growth and development [9]. The *ngrA* gene was shown to be required for the production of small bioactive molecules such as siderophores and antibiotics [9]. Interestingly the stilbene antibiotic produced by all strains of *Photorhabdus* (3,5-dihydroxy-4-isopropylstilbene (ST)) has been shown to be important as a signal for the nematode and is involved in stimulating the recovery of the IJ to the adult hermaphrodite [10]. Moreover we have also recently shown that a mutation in the *exbD* gene of *Photorhabdus temperata* K122 was unable to support the growth and development of its nematode partner, *H. downesi* [11]. The *exbD* gene encodes a component of the TonB complex which is important in mediating the active uptake of siderophore-iron complexes via their cognate outer membrane receptors [12,13]. The defect in symbiosis of the K122 *exbD* mutant was rescued by the addition of FeCl_3 to the media suggesting that siderophore-mediated iron uptake was important for nematode growth and development [11].

Iron is an essential nutrient that is generally found in the insoluble ferric (Fe^{3+}) form [14]. Many bacteria produce siderophores, molecules with very high affinities for Fe^{3+} , in order to be able to successfully compete for Fe^{3+} in their environments [15,16]. The siderophores bind the Fe^{3+} and then bind to specific receptors on the surface of the bacteria. The siderophore-iron complex is then transported into the cell before the Fe^{3+} is reduced to Fe^{2+} and stored as a complex with iron-binding proteins such as bacterioferritin or used for the assembly of important cofactors such as Fe-S clusters [14,17]. Bacteria also have mechanisms to transport the low levels of ferrous (Fe^{2+}) iron that may be available in their environments. These


transport pathways include the FeoABC permease and the YfeABCD divalent cation transporter [14,18]. In this study we wanted to undertake a comprehensive analysis of the role of iron in the symbiosis between the sequenced strain of *Photorhabdus* (*P. luminescens* TT01) and its invertebrate hosts i.e. the insect and the nematode partner, *H. bacteriophora*. Therefore we constructed targeted mutants in genes predicted to play important roles in the uptake of both Fe^{3+} and Fe^{2+} and we tested these mutants for their ability to interact with the different invertebrate partners of *Photorhabdus*.

Results

Genetic analysis of iron uptake systems in *P. luminescens*

TT01

We have previously shown that the *exbD* gene is important for both virulence and symbiosis in *P. temperata* (Pt) K122 [11]. The *exbD* gene encodes a component of the TonB complex (containing TonB, ExbD and ExbB) that is required for siderophore-mediated ferric (Fe^{3+}) iron uptake in many bacteria [13]. The genome sequence of *P. luminescens* (Pl) TT01 has been available since 2003 at which time it was noted that the genome contained the largest known set of iron, heme, hemin and siderophore receptors [19]. This suggested an important role for iron acquisition in the life cycle of *P. luminescens* and we decided to undertake an analysis of the role of iron uptake in the sequenced strain. *In silico* analysis of the genome sequence of Pl TT01 identified a single *tonB* gene (*plu2485*) and a single genetic locus containing *exbD* (*plu3940*) and *exbB* (*plu3941*) (Figure 1A). To compare the role of the TonB complex in both Pl and Pt we constructed a deletion mutation in the *exbD* gene of Pl TT01 (the same gene that was mutated in Pt K122). It would be expected that the $\Delta exbD$ mutant strain would be crippled for iron uptake via any siderophore-mediated pathway. In Pt K122 the *exbD::Km* mutation resulted in an increase in the size of the halo produced on CAS indicator agar indicating accumulation of a siderophore in the agar ([11] and Figure 1B). We have previously shown that this siderophore is likely to be photobactin, a catechol siderophore that was originally identified in *P. luminescens* NC1 [11,20]. Although the Pl TT01 genome is predicted to encode a variety of siderophores, it is interesting that the *phb* genes, encoding the proteins required for photobactin biosynthesis, are not present [19]. Moreover, the Pl TT01 $\Delta exbD$ mutation was observed to have no effect on siderophore production as observed by no change in halo size on CAS agar (Figure 1B). Therefore, Pl TT01 does not appear to be limited for iron during growth on LB agar. Nonetheless we would expect that the $\Delta exbD$ mutant would be limited in its ability to scavenge for iron under iron-limiting conditions. To test this we cultured Pl



TT01 and the $\Delta exbD$ mutant in LB supplemented with 50 μ M 2'-2'-dipyridyl (DIP), an iron chelator, and measured growth (Figure 1C). In the absence of DIP, the growth curves of both the WT and the $\Delta exbD$ mutant were identical. However, in the presence of DIP, it was clear that the $\Delta exbD$ mutant grew at a slower rate than the WT con-

firming that the $\Delta exbD$ mutant was less efficient at scavenging iron.

Bacteria can also utilize the small amounts of soluble ferrous (Fe^{2+}) iron that are present in their environments, usually in a manner that is independent of the TonB complex. We identified genes encoding two potential TonB-independent Fe^{2+} uptake systems, the FeoABC system and the YfeABCD system in the PI TT01 genome (see Table 1 and Figure 2). The FeoABC system is encoded by the *feoABC* operon in which FeoB is predicted to be a GTPase directly involved in Fe^{2+} transport [21]. On the other hand YfeABCD is an ABC transporter that mediates uptake of divalent cations, including Fe^{2+} [18,22]. To test for the role of these genes in PI TT01 we constructed $\Delta feoABC$ and $\Delta yfeABCD$ mutant strains (Δfeo and Δyfe respectively). We also combined mutations to produce the double mutants $\Delta feo \Delta yfe$, $\Delta exbD \Delta yfe$ and $\Delta exbD \Delta feo$ and an $\Delta exbD \Delta yfe \Delta feo$ triple mutant. These iron transport mutants were then tested for their ability to grow on iron-restricted medium i.e. LB agar supplemented with increasing levels of DIP. All strains could grow equally well in the absence of DIP and, as expected, all strains carrying the $\Delta exbD$ allele showed reduced growth, compared to the WT, on media containing 100 μ M DIP (Figure 3). In addition, the *yfeABCD* locus may also play an important role in iron uptake as the Δyfe mutant did not grow as well as WT in the presence of 150 μ M DIP. Moreover the affects of the Δyfe and $\Delta exbD$ mutations appear to be additive confirming that the Yfe ABC transporter and the TonB complex function independently (Figure 3). On the other hand, the Δfeo mutant was unaffected at all concentrations of DIP suggesting that this system does not play a significant role in iron scavenging under these conditions. Interestingly the $\Delta exbD \Delta yfe \Delta feo$ triple mutant was still able to grow on LB agar plates (even in the presence of 50 μ M DIP) suggesting that PI TT01 has additional mechanisms for scavenging iron.

Role of iron uptake in pathogenicity

To determine the affect of the iron transport mutations on virulence we injected approximately 200 CFU of each strain into 10 *Galleria mellonella* larvae. PI TT01 killed the insects in around 48 h, as did both the Δyfe and Δfeo mutant strains (data not shown). On the other hand no insects injected with the $\Delta exbD$ mutant died over the 168 h period of the experiment (data not shown). The $\Delta exbD$ mutant was also avirulent when injected into larvae of another insect model, the Tobacco Hornworm, *Manduca sexta* (Figure 4). Importantly, in *Manduca*, the virulence of the $\Delta exbD$ mutant could be rescued by the pre-injection of 5 mM $FeCl_3$ into the insect (Figure 4). We have shown that the injection of 5 mM $FeCl_3$ was not toxic to

Table 1: Iron transport genes in *P. luminescens* TT01 analyzed in this study.

gene	PI annotation	score	Best hit
<i>tonB</i>	<i>plu2485</i>	4e-27	PMI1355 <i>tonB</i> <i>P. mirabilis</i> HI4320 TonB protein
<i>exbD</i>	<i>plu3940</i>	5e-68	YpsIP31758_0592 <i>exbD</i> <i>Y. pseudotuberculosis</i> IP 31758
<i>exbB</i>	<i>plu3941</i>	1e-79	ECA0358 <i>exbB</i> <i>E. carotovora</i> SCRI1043 Biopolymer transport
<i>feoA</i>	<i>plu0209</i>	8e-27	b3408 <i>feoA</i> <i>E. coli</i> K12 Ferrous iron transport protein A
<i>feoB</i>	<i>plu0208</i>	0.0	b3409 <i>feoB</i> <i>E. coli</i> K12 Ferrous iron transport protein B
<i>feoC</i>	<i>plu0207</i>	2e-20	ef ZP_04612647.1 <i>Yersinia rohdei</i> ATCC 43380 FeoC
<i>yfeA</i>	<i>plu2672</i>	1e-136	YpsIP31758_1705 <i>yfeA</i> <i>Y. pseudotuberculosis</i> IP 31758
<i>yfeB</i>	<i>plu2673</i>	1e-139	PMI1026 <i>sitB</i> <i>P. mirabilis</i> HI4320 Iron ABC transporter
<i>yfeC</i>	<i>plu2674</i>	1e-124	YpsIP31758_1703 <i>yfeC</i> <i>Y. pseudotuberculosis</i> IP 31758
<i>yfeD</i>	<i>plu2675</i>	1e-125	YpsIP31758_1702 <i>yfeD</i> <i>Y. pseudotuberculosis</i> IP 31758


the insect (data not shown). Remarkably, whilst the Δfeo mutant was equally as virulent as the WT in *Manduca*, the Δyfe mutant was avirulent in this insect host (Figure 4). This suggests that the requirement of the *yfeABCD* operon as a virulence factor is dependent on the insect host. Moreover virulence of the Δyfe mutant could be rescued by the pre-injection of $FeCl_3$ confirming that the ability to scavenge for iron is an important virulence factor in PI TT01 (Figure 4).


Role of iron uptake in symbiosis

In this study we wanted to determine the affect of the iron homeostasis mutations in PI TT01 on nematode growth and development. Therefore lipid agar plates were inoculated with the strains to be tested (PI TT01, $\Delta exbD$, Δyfe , Δfeo , $\Delta exbD \Delta yfe \Delta feo$) and, 4 days later, the

bacterial biomass was seeded with 40 surface-sterilised *H. bacteriophora* IJs. We observed that all of the PI TT01 mutants, even the $\Delta exbD \Delta yfe \Delta feo$ triple mutant, were as competent as, if not better than, the WT in their ability to support the growth and development of their nematode partner as measured by the IJ yield i.e. total number of IJs collected/number of IJs inoculated (see Figure 5). This is in sharp contrast to what we had previously observed with Pt K122 *exbD::Km* where we reported that *H. downesi* nematodes failed to reproduce on the mutant bacteria [11].

Nematode development culminates in the formation of a new generation of IJs that are colonized by the bacteria on which the nematodes have been cultured. Therefore, in order to ensure the symbiotic cycle had been completed, the IJs recovered from these symbiosis assays were





surface sterilised, crushed individually and the lysate was spread onto LB agar. In this way it was determined that there were, on average, 42 CFU of PI TT01 present in the gut of each IJ (Figure 6A). Moreover the *ΔexbD* and *Δfeo* mutant strains were able to colonize the IJ as well as the WT (Figure 6A). However the *Δyfe* and *ΔexbD Δyfe Δfeo* mutants appeared to colonize the nematodes at a level that was significantly lower than WT ($P < 0.0001$) suggesting that the *yfeABCD* locus may be important during colonization of the IJ (Figure 6A).

YfeABCD (also known as SitABCD) is an ABC divalent cation transporter that has been shown to transport both Fe²⁺ and Mn²⁺ [18,23,24]. In addition, both YfeABCD and Mn²⁺ have been implicated in resistance to reactive oxygen species (ROS) [22,25]. *Photorhabdus* have been reported to be very sensitive to the low levels of ROS (particularly H₂O₂) generated in LB agar plates after exposure of the plates to fluorescent light [26]. Therefore the low numbers of CFU obtained with the *Δyfe* mutant could be explained by poor plating efficiencies due to an

increased sensitivity to ROS. To test this we crushed IJs grown on either PI TT01 or *Δyfe* and plated the lysate on LB agar supplemented with 0.1% (w/v) pyruvate (a known scavenger of H₂O₂). There was no difference in the number of WT PI TT01 recovered from IJs when the lysate was plated on either LB agar or LB agar supplemented with pyruvate (Figure 6B). On the other hand, the number of CFU recovered from IJs grown on the *Δyfe* mutant increased to WT levels when the lysate was plated on LB agar supplemented with pyruvate (see Figure 6B). Similar results were obtained when the LB agar plates were supplemented with catalase (28 U ml⁻¹) or if the plates were stored in the dark before use (data not shown). Therefore the *Δyfe* mutant does colonize the IJ to the same level as PI TT01 although the *Δyfe* mutant appears to be more sensitive to ROS than the WT. Interestingly we did not see any difference in the sensitivity of WT or the *Δyfe* mutant to ROS when the strains were grown on LB agar and exposed to 30% (v/v) H₂O₂ (data not shown). Therefore the *Δyfe* mutant is not inherently more sensitive to oxidative stress and the increased sensitivity to ROS appears to be dependent on growth within the IJ, suggesting a role for the YfeABCD transporter in this environment.

Bioassays using *H. downesi* reveals symbiosis defect in PI TT01 *DexbD*

We had previously shown that the *exbD* gene in Pt K122 was required for the growth and development of *H. downesi* [11]. In this study we report that *H. bacteriophora* grows normally on the equivalent mutation in PI TT01 (Figure 5). Therefore is the *H. downesi* nematode more sensitive to the *exbD* mutation or is the Pt K122 *exbD::Km* mutant less capable of supporting nematode growth and development in general? To test this we set up a set of bioassays whereby PI TT01 *ΔexbD* and Pt K122 *exbD::Km* were incubated separately with their cognate nematode partner or the nematode partner of the other bacterium. For 14 days after inoculation we monitored nematode growth and reproduction and observed that *H. downesi* did not grow on the Pt K122 *exbD::Km* mutant, as expected, but did grow normally when cultured with Pt K122, PI TT01 or PI TT01 *ΔexbD* (Table 2). In contrast *H. bacteriophora* grew well on all strains tested suggesting that Pt K122 *exbD::Km* is not generally compromised in its ability to support nematode growth and reproduction. Therefore it does appear that the *H. downesi* nematode has a more stringent requirement for iron compared to *H. bacteriophora*.

Discussion

In this study we have genetically tested the role of iron uptake in the interactions between *Photorhabdus* and its invertebrate hosts. We have constructed targeted dele-




Figure 4 Virulence of the $\Delta exbD$ and $\Delta yfeABCD$ mutants can be rescued by $FeCl_3$. Overnight cultures were prepared and 1000 CFU of WT (diamonds), $\Delta exbD$ (squares) and $\Delta yfeABCD$ (triangles) were injected into 5th instar *M. sexta* larvae (open symbols) or larvae pre-injected with 10 μ l of a solution containing 5 mM $FeCl_3$ (filled symbols). The data shown is the mean of at least 2 independent experiments (with n = 10 insects/experiment). For clarity the standard deviations are not shown but these values were within expected limits (0-35%).

tions of genes on the *P. luminescens* TT01 genome that are predicted to be important in both ferric (Fe^{3+}) and ferrous (Fe^{2+}) iron uptake and we have tested these mutants for phenotypes associated with virulence against insect larvae and symbiosis with *H. bacteriophora* nematodes. Our results confirm that iron uptake is important during virulence of the insect and also reveal some interesting features of the role of divalent cation uptake during the pathogenic and mutualistic interactions of *Photographus*.

In this study we have shown that the TT01 $\Delta exbD$ mutation is avirulent in the two different insect models that were tested. The *exbD* gene encodes for a protein that is part of the TonB complex that is found in many Gram negative bacteria. This inner membrane protein complex (composed of ExbD, ExbB and TonB) effectively transduces energy (using the proton motive force) from the inner membrane, across the periplasm, to the outer membrane [13,27]. The TonB complex interacts with outer membrane proteins (such as siderophore receptors)

and the energy is used to facilitate the uptake of molecules through these outer membrane proteins. Bioinformatics can be used to identify proteins that interact with TonB based on the presence of a specific amino acid sequence called the TonB box. In this way 12 TonB-dependent receptors, the majority of which (75%) are predicted to be involved in iron uptake, have been identified in TT01 [27]. In this study we have shown that the lack of virulence associated with the $\Delta exbD$ mutation was due to the inability of this mutant to scavenge iron within the insect environment as virulence could be rescued by the pre-injection of $FeCl_3$. Circulating iron in the insect is bound to transferrin and it has been shown that the transcription of the transferrin gene is increased in *M. sexta* after a microbial challenge suggesting that reducing the availability of iron is part of the insect innate immune response (P. Millichap, unpublished data). We have previously shown that a mutation in the *exbD* gene of Pt K122 was attenuated in virulence due to a decrease in the growth rate in the insect [11]. Our data suggests the PI




Figure 5 The IJ yield after growth on *P. luminescens*. The different bacterial strains were inoculated onto lipid agar plates, incubated for 3-4 days at 30°C and 40 surface-sterilised *H. bacteriophora* IJs were added to the biomass. The plates were incubated for 21 days at 25°C and the IJ yields were determined (i.e. total number of IJs/50). For each experiment 5 plates were analyzed for each strain and the experiment was repeated 3 times. Therefore the data shown is the mean ± standard deviation of n = 15 plates for each strain. Statistical significance was determined using a T-test and IJ yields significantly different ($P < 0.01$) to those obtained using TT01 are indicated with an asterisk.

TT01 $\Delta exbD$ mutant strain is unable to grow in the insect implying that Pt K122 is better at scavenging iron in the insect. Although we have not investigated the reasons for this difference we have confirmed that, similar to what has been reported in other pathogens, TonB complex-mediated iron-uptake is critical for the virulence of *Photorhabdus*.

Nutritional interactions are one of the major driving forces in symbiotic associations [28-31] and our data suggests that iron is an important nutrient in *Photorhabdus-Heterorhabditis* interactions. During growth and development the nematodes feed on the bacterial biomass implying that this biomass must be able to satisfy all of the nematodes nutritional requirements, including the requirement for iron. We have previously shown that iron uptake in Pt K122 is required for the normal growth and development of Hd nematodes [11]. Therefore the Pt K122 $exbD::Km$ mutant was not able to support Hd growth and development but this defect could be rescued by the addition of Fe^{3+} to the media [11]. However, in contrast to this previous work, we have now shown that the $exbD$ gene in PI TT01 is not required for the normal growth and development of the Hb nematode. Cross-feeding experiments, where the Hb nematode was grown on Pt K122 and the Hd nematode was grown on PI TT01, suggested that the nematode was responsible for this difference in iron dependency as the Hb nematode grew equally well on the Pt K122 $exbD::Km$ mutant and the PI TT01 $exbD$ mutant. In addition, although the Hd nematode was observed to grow and develop on both PI TT01

and the PI TT01 $exbD$ mutant, we did observe that the development of Hd IJ nematodes growing on the PI TT01 $exbD$ mutant was significantly delayed compared to Hb growing on the same bacteria (data not shown). This suggests that the Hd nematode might be more sensitive to the presence of the $exbD$ mutation (and therefore iron levels) in their symbiotic bacteria. Such differences in sensitivity to iron levels may be one of the driving forces in the evolution and diversification of the *Photorhabdus-Heterorhabditis* system.

The FeoB protein is an inner membrane Fe^{2+} permease that requires the FeoA-dependent hydrolysis of GTP [21]. The Feo transporter is present in many bacteria and has been reported to have a role in the anaerobic-microaerophilic environment of the gastrointestinal tract of mammals. In this study we show that the FeoABC transporter has no apparent role in either the pathogenic or mutualistic life-styles of *Photorhabdus*. The $yfeABCD$ operon (also found in *Yersinia* and annotated as $sitABCD$ in *Salmonella*, *Shigella* and avian pathogenic *Escherichia coli* (APEC) and $afeABCD$ in *Actinobacillus*) encodes an ATP-dependent divalent cation transporter with affinity for Fe^{2+} and Mn^{2+} [32-36]. The Yfe/Sit/Afe transporter has been shown to have an important role in the virulence of many pathogens and a recent survey in *E. coli* has revealed a strong correlation between the presence of the $yfeABCD$ operon and virulence [35]. In this study we have shown that the $yfeABCD$ operon is important for the virulence of *P. luminescens* in some insect hosts. Therefore the Δyfe mutant was as virulent as the WT bacteria in one lepidopteran insect host, *G. mellonella*, but was completely avirulent in another lepidopteran host, *M. sexta*. This implicates the $yfeABCD$ operon as a possible host-range determining locus in *P. luminescens*. The defect in virulence observed with the Δyfe mutant was rescued by the pre-loading the insect with Fe^{3+} but not Mn^{2+} suggesting that the role of the Yfe transporter in insect virulence is associated with iron homeostasis (data not shown).

In this study we have also shown that the Yfe transporter may have a role during the symbiotic interaction with the nematode, in particular during the colonization of the IJ. We observed that the Δyfe mutant has a very low plating efficiency, compared to WT, on LB agar when isolated directly from the IJ nematode. This low plating efficiency was rescued by the addition of either pyruvate or catalase, known scavengers of H_2O_2 , to the LB agar plates. Therefore the Δyfe mutant appears more sensitive to H_2O_2 than the WT bacteria. The Yfe transporter can mediate the uptake of Mn^{2+} and it has been shown that Mn^{2+} can protect the cells from ROS [18,22]. Although it was thought that part of this protective effect was due to the ability of Mn^{2+} to act as a chemical scavenger of ROS, recent evidence suggests that the role of Mn^{2+} during oxi-




Figure 6 Colonization of IJ nematodes with TT01 and mutant derivatives. A) Individual IJ nematodes ($n = 10$), grown on the different bacterial strains (as indicated), were crushed and the lysate was plated on LB agar to enumerate the CFU within the nematode. The data is shown as a boxplot where the horizontal line within the box represents the median value. The box extends to the first and third quartiles and the whiskers show the upper and lower limits of the data (as defined by the statistical test). Asterisks represent outliers. The level of colonization of strains carrying the $\Delta yfeABCD$ allele was significantly lower than TT01 ($P < 0.0001$, Mann-Whitney). B) As above except that the lysate from each crushed IJ was plated on LB agar with or without added 0.1% (w/v) pyruvate, as indicated.

Table 2: The growth and development of *Heterorhabditis* nematodes on cognate and non-cognate bacteria.

Bacteria	Nematode growth and reproduction ¹	
	<i>H. downesi</i>	<i>H. bacteriophora</i>
Pt K122	+	+
Pt K122 <i>exbD</i> ::Km	-	+
PI TT01	+	+
PI TT01 Δ <i>exbD</i>	+	+

¹presence (+) or absence (-) of nematode growth and reproduction after 14 days

ductive stress in *E. coli* is as an enzyme co-factor (i.e. replacing the Fe²⁺ in Fe-S clusters that are sensitive to oxidative stress) [25]. Many bacteria contain a dedicated Nramp-like Mn²⁺ transporter called MntH [18,37]. In *E. coli* the expression of *mntH* can be induced by oxidative stress and it has been reported that *mntH yfe* double mutants in *Salmonella*, APEC and *Shigella* are sensitive to H₂O₂ [38-40]. Therefore Mn²⁺ uptake appears to be critical in some cells for their ability to survive exposure to H₂O₂. Interestingly analysis of the PI TT01 genome reveals that there is no *mntH* homologue in PI TT01 and, therefore, the Yfe transporter is the only means by which PI TT01 is predicted to be able to obtain Mn²⁺ from the environment. However we could not detect any inherent increase in the sensitivity of the Δ *yfe* mutant to H₂O₂ during growth on LB agar plates. This suggests that there is something specific about the conditions within the nematode that induces the H₂O₂-sensitive phenotype in PI TT01 Δ *yfe*. Recent studies in the model nematode *Caenorhabditis elegans* (a close relative of *Heterorhabditis*) have shown that this nematode produces 3 intestinally localized Nramp-like proteins that are involved in Mn²⁺ transport from the gut lumen [41,42]. Therefore, the levels of Mn²⁺ available to PI TT01 within the gut of the IJ are likely to be very low. This, combined with the absence of a high-affinity Mn²⁺ transporter, would be expected to result in a significant reduction in the level of intracellular Mn²⁺ in the Δ *yfe* mutant that would presumably also limit the ability of the bacterium to respond to an oxidative stress.

Conclusion

In this study we show that siderophore-mediated iron uptake is important for the virulence of *P. luminescens* to insect larvae. This is similar to what has been reported for other pathogens and further highlights the relevance of *Photorhabdus* as a model for studying bacteria-host

interactions [43]. Moreover, in contrast to what we previously reported in another species of *Photorhabdus* (*P. temperata* K122) [11], we show that siderophore-mediated iron uptake in *P. luminescens* TT01 is not required for the growth and development of the nematode. Therefore it appears that different *Photorhabdus-Heterorhabditis* complexes have specific requirements for iron. In addition we show that the *yfeABCD* operon (encoding the Yfe divalent cation transporter) is required for virulence in some, but not all, insect hosts. Although the Yfe transporter can mediate the uptake of either Fe²⁺ or Mn²⁺ we have shown that this transporter is involved in iron uptake during pathogenicity. On the other hand we present data that suggests that the Yfe transporter may be involved in Mn²⁺-uptake during growth in the gut lumen of the IJ nematode. Therefore, the substrate specificity of the Yfe transporter in *P. luminescens* TT01 appears to be dependent on the invertebrate host colonized by the bacteria.

Methods

Bacterial strains and growth conditions

Strains used in this study are listed in Table 3. *Photorhabdus temperata* K122, *Photorhabdus luminescens* subsp *laumondii* TT01 and *Escherichia coli* strains were routinely cultured in Luria-Bertani (LB) broth or on LB agar and were incubated at 30°C or 37°C respectively. CAS agar, for the detection of siderophores, was prepared by adding CAS solution (1:10 (v:v)) into the LB agar just before pouring. CAS solution was prepared as described previously [11]. When required antibiotics were added at the following final concentrations: kanamycin (Km) 50 µg/ml, ampicillin (Amp) 100 µg/ml, chloramphenicol (Cm) 20 µg/ml and rifampicin (Rif) 100 µg/ml.

Construction of deletions in *exbD*, *feoABC* and *yfeABCD*

Targeted deletion mutants were constructed as previously described [10]. Briefly paired oligonucleotides are designed to amplify approximately 500-600 bp of DNA upstream (oligonucleotide 1 + 2) and downstream (oligonucleotide 3 + 4) of the gene to be deleted. The oligonucleotides used in this study are listed in Table 4. These amplicons, which have homologous terminal regions, are fused in a primerless PCR and amplified using oligonucleotide 1 + 4 and then cloned into the suicide vector pDS132 [44]. After conjugation of the plasmid from *E. coli* S17-1 (λ pir) into *P. luminescens* TT01 exconjugants were selected by growth in the presence of Cm and Rif. Potential mutants were then grown overnight in LB broth and plated on LB agar with 2% sucrose to select for loss of the plasmid via a second recombination event. Sucrose-resistant, chloramphenicol-sensitive colonies were then screened using colony PCR to identify mutants. Normally

Table 3: Bacterial strains used in this study

Strain	Genotype	Reference
<i>Photorhabdus</i>		
<i>P. temperata</i> (Pt) K122	Spontaneous Rif ^R mutant	Joyce and Clarke, 2003
<i>P. luminescens</i> (Pl) TT01	Spontaneous Rif ^R mutant	Bennett and Clarke, 2005
BMM417	K122 <i>exbD</i> ::Km	Watson and Clarke, 2005
BMM430	TT01 Δ <i>exbD</i>	This study
BMM431 (Δ <i>yfe</i>)	TT01 Δ <i>yfe</i> ABCD	This study
BMM432 (Δ <i>feo</i>)	TT01 Δ <i>feo</i> ABC	This study
BMM433	TT01 Δ <i>exbD</i> Δ <i>yfe</i>	This study
BMM434	TT01 Δ <i>exbD</i> Δ <i>feo</i>	This study
BMM435	TT01 Δ <i>feo</i> Δ <i>yfe</i>	This study
BMM436	TT01 Δ <i>exbD</i> Δ <i>feo</i> Δ <i>yfe</i>	This study
E.coli		
S17-1 (λ <i>pir</i>)	lysogenised with λ <i>pir</i> , replication of <i>ori</i> R6K	Laboratory stock

mutants are detected at a frequency of between 10-30% and the amplicons from 2-3 of the colonies are sequenced to confirm the integrity of the deletion.

Virulence assays

The pathogenicity of *P. luminescens* was assessed using final instar *Galleria mellonella* larvae (purchased from Livefood (UK)) and freshly molted 5th instar *Manduca sexta* larvae (cultured at the University of Bath) as the

model insect hosts. Briefly overnight cultures of *P. luminescens* TT01 were washed 3 times in 1 × PBS and the density adjusted appropriately so that 200 CFU or 1000 CFU could be injected into the hemolymph of *G. mellonella* or *M. sexta*, respectively. Insects were incubated at 30°C and monitored for death at regular time intervals. Where appropriate insect were pre-injected with 10 μl of either 5 mM FeCl₃ or 5 mM MnCl₂ at least 30 min before the bacteria were injected.

Table 4: Oligonucleotides used for construction of targeted deletion mutants.

Gene(s)	Sequence 5' to 3' ^{*:}	Name
<i>exbD</i>	1. TTAT <u>G</u> CATGCGGTGATTGCTTCTGTTACTT GG	RJW115
	2. GAATCAGTGACAATTACATAAGTCACCTTGCTCTG	RJW116
	3. CAAGGTGACTTATGTAATTGCACTGATTCTTCC	RJW117
	4. TTAT <u>G</u> AGCTCGCCAACCAATTTGCCTCTGCCCTAC	RJW118
<i>yfe</i> ABCD	1. TTAT <u>G</u> CATGCGGTTATCAATACCTGCCAGATGC	RJW171
	2. CCCTTTTGTACATAAATCAAACC	RJW172
	3. GGTTTGAATTTATGTAACAAAAGGGTTATATCTG	RJW173
	4. TTAT <u>G</u> AGCTCGGTGTTGAAGTTGTTACTTATAGC	RJW174
<i>feo</i> ABC	1. TTAT <u>G</u> CATGCGTAGTAAAAGCGGGTATATCG	RJW167
	2. GCTAATCATTTTCAATTCCTACATATGACCTTCCG	RJW168
	3. CGGAAGGTCATATGTAGGAATTGAAAATGATTAGC	RJW169
	4. TTAT <u>G</u> AGCTCCAAAACGCTTCTCTTAGAAGATGC	RJW170

*: the underlined sequence indicate the restriction endonuclease sites used for cloning the amplicon into pDS132.

Nematode growth and development

To determine the ability of each mutant to support nematode growth and development we carried out in vitro symbiosis assays. Therefore the bacteria were cultured overnight in LB and 50 µl was spread, in a Z pattern, onto the surface of a lipid agar plate (/500 ml: 12.5 g nutrient agar, 5 g corn syrup, 2.5 g yeast extract, 2.5 ml cod liver oil, 1 g MgCl₂·6H₂O) containing Rif and incubated at 30°C for 3-4 days. The bacterial biomass was then seeded with 40 surface-sterilised *H. bacteriophora* IJs and incubated at 25°C for 21 days. The presence of the Rif ensures that any bacteria present in the IJ are not able to compete with the lawn of bacteria present on the lipid agar plate. After 21 days the new generation of IJs had migrated to the lid of the Petri dish and these nematodes were collected in 1 × PBS and enumerated to determine the IJ yield (i.e. total number of IJs collected/number of IJs inoculated).

Colonization assay

To determine colonization levels by each of the mutants IJs collected from the in vitro symbiosis assays were incubated at room temperature for at least 7 days before analysis. This incubation provides the bacteria with the opportunity to reproduce in the IJ gut and form a stable population. The IJs were surface-sterilised by washing in 0.4% (w/v) hyamine and individual IJs were crushed in 100 µl of PBS and the lysate was plated on LB (with or without added pyruvate). The plates were incubated at 30°C and the number of CFU's was determined after 48 h.

Authors' contributions

RJW undertook all of the experiments described in this manuscript with the exception of the virulence assays in *Manduca sexta* (which were carried out by PM). RJW, SAJ and DJC conceived of the study. SAJ, SR and DJC designed the experiments and DJC wrote the manuscript. All authors have read and approved the final manuscript.

Acknowledgements

RJW and PM were supported by studentships from the BBSRC. SAJ was funded through the Exploiting Genomics initiative of the BBSRC (project number: 86/EGA16183 awarded to DJC and SR). Work in the lab of DJC is currently funded by Science Foundation Ireland.

Author Details

¹Department of Biology and Biochemistry, University of Bath, Bath BA2 7AY, UK and ²Department of Microbiology, University College Cork, Cork, Ireland

Received: 30 October 2009 Accepted: 22 June 2010

Published: 22 June 2010

References

1. Waterfield NR, Ciche T, Clarke D: *Photorhabdus* and a host of hosts. *Annu Rev Microbiol* 2009, **63**:557-574.
2. Clarke DJ: *Photorhabdus*: a model for the analysis of pathogenicity and mutualism. *Cell Microbiol* 2008, **10**:2159-2167.
3. Ciche TA, Ensign JC: For the insect pathogen *Photorhabdus luminescens*, which end of a nematode is out? *Appl Environ Microbiol* 2003, **69**:1890-1897.
4. Clarke DJ, Dowds BCA: Virulence mechanisms of *Photorhabdus* sp strain K122 toward wax moth larvae. *J Invert Pathol* 1995, **66**:149-155.
5. Daborn PJ, Waterfield N, Blight MA, ffrench-Constant RH: Measuring virulence factor expression by the pathogenic bacterium *Photorhabdus luminescens* in culture and during insect infection. *J Bacteriol* 2001, **183**:5834-5839.
6. Ciche TA, Kim K, Kaufmann-Daszczuk B, Nguyen KCQ, Hall DH: Cell invasion and matricide during *Photorhabdus luminescens* transmission by *Heterorhabditis bacteriophora* nematodes. *Appl Environ Microbiol* 2008, **74**:2275-2287.
7. Bintrim SB, Ensign JC: Insertional inactivation of genes encoding the crystalline inclusion proteins of *Photorhabdus luminescens* results in mutants with pleiotropic phenotypes. *J Bacteriol* 1998, **180**:1261-1269.
8. You J, Liang S, Cao L, Liu X, Han R: Nutritive significance of crystalline inclusion proteins of *Photorhabdus luminescens* in *Steinernema* nematodes. *FEMS Microbiol Ecol* 2006, **55**:178-185.
9. Ciche TA, Bintrim SB, Horswill AR, Ensign JC: A phosphopantetheinyl transferase homolog is essential for *Photorhabdus luminescens* to support growth and reproduction of the entomopathogenic nematode *Heterorhabditis bacteriophora*. *J Bacteriol* 2001, **183**:3117-3126.
10. Joyce SA, Brachmann AO, Glazer I, Lango L, Schwar G, Clarke DJ, Bode HB: Bacterial biosynthesis of a multipotent stilbene. *Angewandte Chemie* 2008, **47**:1942-1945.
11. Watson RJ, Joyce SA, Spencer GV, Clarke DJ: The *exbD* gene of *Photorhabdus temperata* is required for full virulence in insects and symbiosis with the nematode *Heterorhabditis*. *Mol Microbiol* 2005, **56**:763-773.
12. Braun V: Energy-coupled transport and signal transduction through the gram-negative outer membrane via TonB-ExbB-ExbD-dependent receptor proteins. *FEMS Microbiol Rev* 1995, **16**:295-307.
13. Braun V, Braun M: Active transport of iron and siderophore antibiotics. *Curr Opin Microbiol* 2002, **5**:194-201.
14. Andrews SC, Robinson AK, Rodriguez-Quinones F: Bacterial iron homeostasis. *FEMS Microbiol Rev* 2003, **27**:215-237.
15. Wandersman C, Deleplaire P: Bacterial iron sources: from siderophores to hemophores. *Annu Rev Microbiol* 2004, **58**:611-647.
16. Faraldo-Gomez JD, Sansom MS: Acquisition of siderophores in gram-negative bacteria. *Nat Rev Mol Cell Biol* 2003, **4**:105-116.
17. Andrews SC: Iron storage in bacteria. *Adv Microb Physiol* 1998, **40**:281-351.
18. Papp-Wallace KM, Maguire ME: Manganese transport and the role of manganese in virulence. *Annu Rev Microbiol* 2006, **60**:187-209.
19. Duchaud E, Rusniok C, Frangeul L, Buchrieser C, Givaudan A, Taourit S, Bocs S, Boursaux-Eude C, Chandler M, Charles JF, et al.: The genome sequence of the entomopathogenic bacterium *Photorhabdus luminescens*. *Nat Biotechnol* 2003, **21**:1307-1313.
20. Ciche TA, Blackburn M, Carney JR, Ensign JC: Photobactin: a catechol siderophore produced by *Photorhabdus luminescens*, an entomopathogen mutually associated with *Heterorhabditis bacteriophora* NC1 nematodes. *Appl Environ Microbiol* 2003, **69**:4706-4713.
21. Cartron ML, Maddocks S, Gillingham P, Craven CJ, Andrews SC: Fe²⁺-transport of ferrous iron into bacteria. *Biometals* 2006, **19**:143-157.
22. Kehres DG, Maguire ME: Emerging themes in manganese transport, biochemistry and pathogenesis in bacteria. *FEMS Microbiol Rev* 2003, **27**:263-290.
23. Janakiraman A, Schlauch JM: The putative iron transport system SitABCD encoded on SPI1 is required for full virulence of *Salmonella typhimurium*. *Mol Microbiol* 2000, **35**:1146-1155.
24. Kehres DG, Janakiraman A, Schlauch JM, Maguire ME: SitABCD is the alkaline Mn(2+) transporter of *Salmonella enterica* serovar Typhimurium. *J Bacteriol* 2002, **184**:3159-3166.
25. Anjem A, Varghese S, Imlay JA: Manganese import is a key element of the OxyR response to hydrogen peroxide in *Escherichia coli*. *Mol Microbiol* 2009, **72**:844-858.
26. Xu J, Hurlbert RE: Toxicity of irradiated media for *Xenorhabdus* spp. *Appl Environ Microbiol* 1990, **56**:815-818.
27. Koebnik R: TonB-dependent trans-envelope signalling: the exception or the rule? *Trends Microbiol* 2005, **13**:343-347.
28. Zientz E, Dandekar T, Gross R: Metabolic interdependence of obligate intracellular bacteria and their insect hosts. *Microbiol Mol Biol Rev* 2004, **68**:745-770.

29. Prell J, Poole P: **Metabolic changes of rhizobia in legume nodules.** *Trends Microbiol* 2006, **14**:161-168.
30. Leser TD, Molbak L: **Better living through microbial action: the benefits of the mammalian gastrointestinal microbiota on the host.** *Environ Microbiol* 2009, **11**:2194-2206.
31. Harrison MJ: **Signaling in the arbuscular mycorrhizal symbiosis.** *Annu Rev Microbiol* 2005, **59**:19-42.
32. Rhodes ER, Menke S, Shoemaker C, Tomaras AP, McGillivray G, Actis LA: **Iron acquisition in the dental pathogen *Actinobacillus actinomycetemcomitans*: what does it use as a source and how does it get this essential metal?** *Biometals* 2007, **20**:365-377.
33. Zhou D, Hardt WD, Galan JE: ***Salmonella typhimurium* encodes a putative iron transport system within the centisome 63 pathogenicity island.** *Infect Immun* 1999, **67**:1974-1981.
34. Runyen-Janecky LJ, Reeves SA, Gonzales EG, Payne SM: **Contribution of the *Shigella flexneri* Sit, luc, and Feo iron acquisition systems to iron acquisition in vitro and in cultured cells.** *Infect Immun* 2003, **71**:1919-1928.
35. Sabri M, Leveille S, Dozois CM: **A SitABCD homologue from an avian pathogenic *Escherichia coli* strain mediates transport of iron and manganese and resistance to hydrogen peroxide.** *Microbiology* 2006, **152**:745-758.
36. Perry RD, Mier I Jr, Fetherston JD: **Roles of the Yfe and Feo transporters of *Yersinia pestis* in iron uptake and intracellular growth.** *Biometals* 2007, **20**:699-703.
37. Richer E, Courville P, Bergevin I, Cellier MF: **Horizontal gene transfer of "prototype" Nramp in bacteria.** *J Mol Evol* 2003, **57**:363-376.
38. Kehres DG, Janakiraman A, Slauch JM, Maguire ME: **Regulation of *Salmonella enterica* serovar Typhimurium *mntH* transcription by H₂O₂, Fe(2+), and Mn(2+).** *J Bacteriol* 2002, **184**:3151-3158.
39. Sabri M, Caza M, Proulx J, Lymberopoulos MH, Bree A, Moulin-Schouleur M, Curtiss R, Dozois CM: **Contribution of the SitABCD, MntH, and FeoB metal transporters to the virulence of avian pathogenic *Escherichia coli* O78 strain chi7122.** *Infect Immun* 2008, **76**(2):601-611.
40. Runyen-Janecky L, Dazenski E, Hawkins S, Warner L: **Role and regulation of the *Shigella flexneri* sit and MntH systems.** *Infect Immun* 2006, **74**:4666-4672.
41. Settivari R, Levora J, Nass R: **The divalent metal transporter homologues SMF-1/2 mediates dopamine neuron sensitivity in *Caenorhabditis elegans* models of manganese and Parkinson's disease.** *J Biol Chem* 2009, **284**:35758-68.
42. Bandyopadhyay J, Song HO, Park BJ, Singaravelu G, Sun JL, Ahnn J, Cho JH: **Functional assessment of Nramp-like metal transporters and manganese in *Caenorhabditis elegans*.** *Biochem Biophys Res Commun* 2009, **390**:136-141.
43. Schaible UE, Kaufmann SH: **Iron and microbial infection.** *Nat Rev Microbiol* 2004, **2**:946-953.
44. Philippe N, Alcaraz J-P, Coursange E, Geiselmann J, Schneider D: **Improvement of pCVD442, a suicide plasmid for gene allele exchange in bacteria.** *Plasmid* 2004, **51**:246-255.

doi: 10.1186/1471-2180-10-177

Cite this article as: Watson *et al.*, The role of iron uptake in pathogenicity and symbiosis in *Photobacterium luminescens* TT01 *BMC Microbiology* 2010, **10**:177

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

