

3-Ethylsulfinyl-2-(3-fluorophenyl)-5,6-methylenedioxy-1-benzofuran**Hong Dae Choi,^a Pil Ja Seo^a and Uk Lee^{b*}**

^aDepartment of Chemistry, Dongeui University, San 24 Kaya-dong, Busanjin-gu, Busan 614-714, Republic of Korea, and ^bDepartment of Chemistry, Pukyong National University, 599-1 Daeyeon 3-dong, Nam-gu, Busan 608-737, Republic of Korea

Correspondence e-mail: uklee@pknu.ac.kr


Received 25 October 2012; accepted 30 October 2012

Key indicators: single-crystal X-ray study; $T = 173\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.002\text{ \AA}$; R factor = 0.038; wR factor = 0.101; data-to-parameter ratio = 17.2.

In the title compound, $\text{C}_{17}\text{H}_{13}\text{FO}_4\text{S}$, the 3-fluorophenyl ring makes a dihedral angle of $6.14(5)^\circ$ with the mean plane [r.m.s. deviation = $0.008(1)\text{ \AA}$] of the benzofuran fragment. In the crystal, molecules are linked by weak $\text{C}-\text{H}\cdots\text{F}$, $\text{C}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\pi$ interactions, forming a three-dimensional network. The crystal structure also exhibits slipped $\pi-\pi$ interactions between the 3-fluorophenyl rings of neighbouring molecules [centroid–centroid distance = $3.769(2)\text{ \AA}$ and slippage = $1.684(2)\text{ \AA}$].

Related literature

For background information on and the crystal structures of related benzofuran compounds, see: Choi *et al.* (2010); Seo *et al.* (2011).

**Experimental***Crystal data*

$\text{C}_{17}\text{H}_{13}\text{FO}_4\text{S}$
 $M_r = 332.33$
Monoclinic, $P2_1/c$

$a = 8.8516(2)\text{ \AA}$
 $b = 21.8221(4)\text{ \AA}$
 $c = 7.7228(2)\text{ \AA}$

$\beta = 102.949(1)^\circ$
 $V = 1453.80(6)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 0.25\text{ mm}^{-1}$
 $T = 173\text{ K}$
 $0.39 \times 0.29 \times 0.26\text{ mm}$

Data collection

Bruker SMART APEXII CCD diffractometer
Absorption correction: multi-scan (*SADABS*; Bruker, 2009)
 $T_{\min} = 0.908$, $T_{\max} = 0.937$

14376 measured reflections
3600 independent reflections
3131 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.038$
 $wR(F^2) = 0.101$
 $S = 1.02$
3600 reflections

209 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.45\text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.35\text{ e \AA}^{-3}$

Table 1Hydrogen-bond geometry (\AA , $^\circ$). $Cg1$ is the centroid of the C2–C7 benzene ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
$\text{C9}-\text{H9A}\cdots\text{O4}^i$	0.99	2.45	3.314 (2)	145
$\text{C9}-\text{H9B}\cdots\text{O2}^{ii}$	0.99	2.59	3.519 (3)	156
$\text{Cl6}-\text{H16A}\cdots\text{O4}^{iii}$	0.99	2.29	3.2689 (19)	169
$\text{C6}-\text{H6}\cdots\text{F1}^{iv}$	0.95	2.37	3.2532 (18)	154
$\text{C17}-\text{H17B}\cdots\text{Cg1}^v$	0.98	2.74	3.642 (3)	153

Symmetry codes: (i) $x + 1, -y + \frac{3}{2}, z + \frac{1}{2}$; (ii) $x, -y + \frac{3}{2}, z + \frac{1}{2}$; (iii) $x, -y + \frac{3}{2}, z - \frac{1}{2}$; (iv) $x + 1, y, z + 1$; (v) $x, y, z - 1$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 1998); software used to prepare material for publication: *SHELXL97*.

This work was supported by the Blue-Bio Industry Regional Innovation Center (RIC08-06-07) at Dongeui University as an RIC program under the Ministry of Knowledge Economy and Busan City.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: MW2094).

References

- Brandenburg, K. (1998). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
Bruker (2009). *APEX2*, *SADABS* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
Choi, H. D., Seo, P. J., Son, B. W. & Lee, U. (2010). *Acta Cryst. E66*, o1296.
Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
Seo, P. J., Choi, H. D., Son, B. W. & Lee, U. (2011). *Acta Cryst. E67*, o3504.
Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.

supplementary materials

Acta Cryst. (2012). E68, o3339 [doi:10.1107/S1600536812044832]

3-Ethylsulfinyl-2-(3-fluorophenyl)-5,6-methylenedioxy-1-benzofuran

Hong Dae Choi, Pil Ja Seo and Uk Lee

Comment

As a part of our ongoing study of 5,6-methylenedioxy-1-benzofuran derivatives containing [3-ethylsulfinyl-2-(4-fluorophenyl)] (Choi *et al.*, 2010) and [2-(4-fluorophenyl)-3-isopropylsulfinyl] (Seo *et al.*, 2011) substituents, we report herein the crystal structure of the title compound.

In the title molecule (Fig. 1), the benzofuran unit is essentially planar, with a mean deviation of 0.008 (1) Å from the least-squares plane defined by the nine constituent atoms. The dihedral angle between the 3-fluorophenyl ring and the mean plane of the benzofuran fragment is 6.14 (5)°. In the crystal structure (Figs. 2 & 3), molecules are connected by weak C—H···F, C—H···O and C—H···π interactions (Table 1, Cg1 is the centroid of the C2–C7 benzene ring). The crystal packing (Fig. 3) also exhibits slipped π–π interactions between the 3-fluorophenyl rings of neighbouring molecules, with a Cg2···Cg2^{vi} distance of 3.769 (2) Å and an interplanar distance of 3.372 (2) Å resulting in a slippage of 1.684 (2) Å (Cg2 is the centroid of the C10–C15 3-fluorophenyl ring).

Experimental


3-Chloroperoxybenzoic acid (77%, 224 mg, 1.0 mmol) was added in small portions to a stirred solution of 3-ethylsulfonyl-2-(3-fluorophenyl)-5,6-methylenedioxy-1-benzofuran (284 mg, 0.9 mmol) in dichloromethane (40 mL) at 273 K. After being stirred at room temperature for 4 h, the mixture was washed with saturated sodium bicarbonate solution and the organic layer was separated, dried over magnesium sulfate, filtered and concentrated at reduced pressure. The residue was purified by column chromatography (hexane–ethyl acetate, 1:1 v/v) to afford the title compound as a colorless solid [yield 73%, m.p. 408–409 K; R_f = 0.61 (hexane–ethyl acetate, 1:1 v/v)]. Single crystals suitable for X-ray diffraction were prepared by slow evaporation of a solution of the title compound in acetone at room temperature.

Refinement


All H atoms were geometrically positioned and refined using a riding model, with C—H = 0.95 Å for aryl, 0.99 Å for methylene and 0.95 Å for methyl H atoms, respectively. $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ for aryl and methylene, and $1.5U_{\text{eq}}(\text{C})$ for methyl H atoms. The positions of methyl hydrogens were optimized rotationally.

Computing details


Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT* (Bruker, 2009); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *DIAMOND* (Brandenburg, 1998); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008).

**Figure 1**

The molecular structure of the title compound with the atom numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H atoms are presented as small spheres of arbitrary radius.

**Figure 2**

A view of the C—H···F and C—H···O interactions (dotted lines) in the crystal structure of the title compound. H atoms not participating in hydrogen-bonding were omitted for clarity. [Symmetry codes: (i) $x + 1, -y + 3/2, z + 1/2$; (ii) $x, -y + 3/2, z + 1/2$; (iii) $x, -y + 3/2, z - 1/2$; (iv) $x + 1, y, z + 1$ (vii) $x - 1, -y + 3/2, z - 1/2$; (viii) $x, -y + 3/2, z + 1/2$; (ix) $x, -y + 3/2, z + 1/2$; (x) $x - 1, y, z - 1$.]

**Figure 3**

A view of the C—H \cdots π and π — π interactions (dotted lines) in the crystal structure of the title compound. H atoms not participating in hydrogen-bonding were omitted for clarity. [Symmetry codes: (v) $x, y, z - 1$; (vi) $-x, -y + 1, -z + 1$; (xi) $x, y, z + 1$.]

3-Ethylsulfinyl-2-(3-fluorophenyl)-5,6-methylenedioxy-1-benzofuran

Crystal data

$C_{17}H_{13}FO_4S$
 $M_r = 332.33$
Monoclinic, $P2_1/c$
Hall symbol: -P 2ybc
 $a = 8.8516 (2)$ Å
 $b = 21.8221 (4)$ Å
 $c = 7.7228 (2)$ Å
 $\beta = 102.949 (1)^\circ$
 $V = 1453.80 (6)$ Å³
 $Z = 4$

$F(000) = 688$
 $D_x = 1.518 \text{ Mg m}^{-3}$
Melting point: 408 K
Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å
Cell parameters from 6139 reflections
 $\theta = 2.9\text{--}28.2^\circ$
 $\mu = 0.25 \text{ mm}^{-1}$
 $T = 173$ K
Block, colourless
 $0.39 \times 0.29 \times 0.26$ mm

Data collection

Bruker SMART APEXII CCD
diffractometer
Radiation source: rotating anode
Graphite multilayer monochromator
Detector resolution: 10.0 pixels mm⁻¹
 φ and ω scans

Absorption correction: multi-scan
(SADABS; Bruker, 2009)
 $T_{\min} = 0.908$, $T_{\max} = 0.937$
14376 measured reflections
3600 independent reflections
3131 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.025$

$\theta_{\max} = 28.3^\circ$, $\theta_{\min} = 1.9^\circ$
 $h = -11 \rightarrow 11$

$k = -29 \rightarrow 27$
 $l = -10 \rightarrow 10$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.038$

$wR(F^2) = 0.101$

$S = 1.02$

3600 reflections

209 parameters

0 restraints

Primary atom site location: structure-invariant
direct methods

Secondary atom site location: difference Fourier
map

Hydrogen site location: difference Fourier map

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.0488P)^2 + 0.7371P]$
where $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} = 0.001$

$\Delta\rho_{\max} = 0.45 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.35 \text{ e } \text{\AA}^{-3}$

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R-factor wR and goodness of fit S are based on F^2 , conventional R-factors R are based on F, with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F, and R-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.20748 (4)	0.653717 (15)	0.40724 (5)	0.02313 (11)
F1	-0.09845 (13)	0.46852 (5)	0.11643 (15)	0.0452 (3)
O1	0.44774 (12)	0.52350 (4)	0.70767 (14)	0.0247 (2)
O2	0.76433 (15)	0.73096 (6)	0.93233 (18)	0.0424 (3)
O3	0.86160 (15)	0.64201 (6)	1.07899 (18)	0.0431 (3)
O4	0.18119 (13)	0.70803 (5)	0.51487 (16)	0.0341 (3)
C1	0.33826 (16)	0.60530 (6)	0.55117 (19)	0.0215 (3)
C2	0.47112 (16)	0.62662 (6)	0.68258 (18)	0.0222 (3)
C3	0.54231 (17)	0.68385 (7)	0.7276 (2)	0.0252 (3)
H3	0.5037	0.7207	0.6683	0.030*
C4	0.67160 (18)	0.68235 (7)	0.8635 (2)	0.0271 (3)
C5	0.72968 (17)	0.62887 (7)	0.9530 (2)	0.0278 (3)
C6	0.66316 (17)	0.57266 (7)	0.9140 (2)	0.0271 (3)
H6	0.7020	0.5362	0.9755	0.033*
C7	0.53247 (16)	0.57424 (6)	0.77457 (19)	0.0229 (3)
C8	0.32896 (16)	0.54330 (6)	0.57115 (19)	0.0223 (3)
C9	0.8859 (2)	0.70603 (8)	1.0666 (2)	0.0403 (4)
H9A	0.9872	0.7137	1.0365	0.048*
H9B	0.8866	0.7259	1.1821	0.048*
C10	0.22658 (16)	0.49448 (6)	0.48447 (19)	0.0225 (3)
C11	0.10942 (17)	0.50551 (7)	0.3331 (2)	0.0257 (3)
H11	0.0948	0.5451	0.2812	0.031*
C12	0.01626 (18)	0.45741 (7)	0.2619 (2)	0.0274 (3)
C13	0.03117 (18)	0.39881 (7)	0.3285 (2)	0.0292 (3)

H13	-0.0358	0.3668	0.2743	0.035*
C14	0.14759 (19)	0.38828 (7)	0.4776 (2)	0.0314 (3)
H14	0.1613	0.3483	0.5272	0.038*
C15	0.24452 (18)	0.43523 (7)	0.5557 (2)	0.0280 (3)
H15	0.3238	0.4272	0.6583	0.034*
C16	0.33604 (19)	0.67847 (7)	0.2700 (2)	0.0306 (3)
H16A	0.2947	0.7164	0.2061	0.037*
H16B	0.4390	0.6880	0.3462	0.037*
C17	0.3535 (2)	0.62964 (10)	0.1369 (3)	0.0450 (4)
H17A	0.3961	0.5923	0.2000	0.068*
H17B	0.4237	0.6443	0.0642	0.068*
H17C	0.2519	0.6206	0.0600	0.068*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.02202 (18)	0.01839 (17)	0.02712 (19)	0.00208 (12)	0.00156 (13)	0.00075 (13)
F1	0.0445 (6)	0.0362 (5)	0.0436 (6)	-0.0053 (4)	-0.0138 (5)	-0.0012 (4)
O1	0.0256 (5)	0.0197 (5)	0.0266 (5)	0.0015 (4)	0.0013 (4)	0.0023 (4)
O2	0.0397 (7)	0.0311 (6)	0.0465 (7)	-0.0092 (5)	-0.0113 (6)	-0.0031 (5)
O3	0.0365 (7)	0.0404 (7)	0.0419 (7)	-0.0063 (5)	-0.0133 (5)	0.0026 (5)
O4	0.0370 (6)	0.0226 (5)	0.0417 (7)	0.0079 (4)	0.0069 (5)	-0.0049 (5)
C1	0.0206 (6)	0.0208 (6)	0.0233 (6)	0.0025 (5)	0.0050 (5)	0.0011 (5)
C2	0.0216 (6)	0.0227 (7)	0.0226 (7)	0.0025 (5)	0.0056 (5)	-0.0009 (5)
C3	0.0261 (7)	0.0212 (7)	0.0278 (7)	0.0020 (5)	0.0046 (6)	-0.0011 (5)
C4	0.0264 (7)	0.0257 (7)	0.0288 (7)	-0.0023 (6)	0.0051 (6)	-0.0037 (6)
C5	0.0225 (7)	0.0342 (8)	0.0248 (7)	-0.0003 (6)	0.0011 (6)	-0.0010 (6)
C6	0.0262 (7)	0.0271 (7)	0.0264 (7)	0.0040 (6)	0.0021 (6)	0.0047 (6)
C7	0.0233 (7)	0.0211 (7)	0.0241 (7)	0.0002 (5)	0.0051 (5)	0.0000 (5)
C8	0.0219 (6)	0.0220 (7)	0.0225 (7)	0.0022 (5)	0.0041 (5)	0.0007 (5)
C9	0.0382 (9)	0.0403 (9)	0.0363 (9)	-0.0093 (8)	-0.0048 (7)	-0.0005 (7)
C10	0.0241 (7)	0.0196 (6)	0.0253 (7)	0.0001 (5)	0.0086 (5)	-0.0007 (5)
C11	0.0279 (7)	0.0212 (7)	0.0278 (7)	0.0001 (5)	0.0058 (6)	-0.0005 (5)
C12	0.0271 (7)	0.0274 (7)	0.0270 (7)	0.0003 (6)	0.0045 (6)	-0.0030 (6)
C13	0.0319 (8)	0.0242 (7)	0.0333 (8)	-0.0058 (6)	0.0111 (6)	-0.0059 (6)
C14	0.0391 (9)	0.0202 (7)	0.0360 (8)	-0.0007 (6)	0.0106 (7)	0.0017 (6)
C15	0.0308 (7)	0.0236 (7)	0.0289 (7)	0.0009 (6)	0.0050 (6)	0.0031 (6)
C16	0.0310 (8)	0.0296 (8)	0.0301 (8)	-0.0033 (6)	0.0046 (6)	0.0080 (6)
C17	0.0486 (11)	0.0550 (12)	0.0347 (9)	-0.0016 (9)	0.0162 (8)	-0.0018 (8)

Geometric parameters (\AA , $^\circ$)

S1—O4	1.4954 (11)	C8—C10	1.4602 (19)
S1—C1	1.7651 (14)	C9—H9A	0.9900
S1—C16	1.8029 (16)	C9—H9B	0.9900
F1—C12	1.3568 (18)	C10—C11	1.399 (2)
O1—C7	1.3717 (17)	C10—C15	1.400 (2)
O1—C8	1.3809 (16)	C11—C12	1.372 (2)
O2—C4	1.3733 (18)	C11—H11	0.9500
O2—C9	1.424 (2)	C12—C13	1.373 (2)

O3—C5	1.3724 (18)	C13—C14	1.381 (2)
O3—C9	1.420 (2)	C13—H13	0.9500
C1—C8	1.366 (2)	C14—C15	1.385 (2)
C1—C2	1.4473 (19)	C14—H14	0.9500
C2—C7	1.3905 (19)	C15—H15	0.9500
C2—C3	1.407 (2)	C16—C17	1.513 (3)
C3—C4	1.369 (2)	C16—H16A	0.9900
C3—H3	0.9500	C16—H16B	0.9900
C4—C5	1.395 (2)	C17—H17A	0.9800
C5—C6	1.365 (2)	C17—H17B	0.9800
C6—C7	1.393 (2)	C17—H17C	0.9800
C6—H6	0.9500		
O4—S1—C1	106.63 (7)	O3—C9—H9B	109.9
O4—S1—C16	106.65 (7)	O2—C9—H9B	109.9
C1—S1—C16	97.95 (7)	H9A—C9—H9B	108.3
C7—O1—C8	107.07 (10)	C11—C10—C15	118.99 (13)
C4—O2—C9	105.97 (13)	C11—C10—C8	121.69 (13)
C5—O3—C9	105.99 (13)	C15—C10—C8	119.32 (13)
C8—C1—C2	107.41 (12)	C12—C11—C10	118.03 (14)
C8—C1—S1	128.05 (11)	C12—C11—H11	121.0
C2—C1—S1	124.35 (11)	C10—C11—H11	121.0
C7—C2—C3	120.10 (13)	F1—C12—C11	117.79 (14)
C7—C2—C1	104.88 (12)	F1—C12—C13	117.90 (13)
C3—C2—C1	135.01 (13)	C11—C12—C13	124.31 (14)
C4—C3—C2	114.80 (13)	C12—C13—C14	117.26 (14)
C4—C3—H3	122.6	C12—C13—H13	121.4
C2—C3—H3	122.6	C14—C13—H13	121.4
C3—C4—O2	127.07 (14)	C13—C14—C15	120.90 (14)
C3—C4—C5	123.50 (14)	C13—C14—H14	119.5
O2—C4—C5	109.43 (13)	C15—C14—H14	119.5
C6—C5—O3	126.94 (14)	C14—C15—C10	120.50 (14)
C6—C5—C4	123.35 (14)	C14—C15—H15	119.8
O3—C5—C4	109.70 (14)	C10—C15—H15	119.8
C5—C6—C7	113.02 (13)	C17—C16—S1	111.26 (12)
C5—C6—H6	123.5	C17—C16—H16A	109.4
C7—C6—H6	123.5	S1—C16—H16A	109.4
O1—C7—C2	110.68 (12)	C17—C16—H16B	109.4
O1—C7—C6	124.09 (12)	S1—C16—H16B	109.4
C2—C7—C6	125.22 (13)	H16A—C16—H16B	108.0
C1—C8—O1	109.95 (12)	C16—C17—H17A	109.5
C1—C8—C10	135.85 (13)	C16—C17—H17B	109.5
O1—C8—C10	114.20 (12)	H17A—C17—H17B	109.5
O3—C9—O2	108.90 (13)	C16—C17—H17C	109.5
O3—C9—H9A	109.9	H17A—C17—H17C	109.5
O2—C9—H9A	109.9	H17B—C17—H17C	109.5
O4—S1—C1—C8	134.07 (14)	C1—C2—C7—C6	179.62 (14)
C16—S1—C1—C8	-115.83 (15)	C5—C6—C7—O1	178.29 (14)

O4—S1—C1—C2	−40.34 (14)	C5—C6—C7—C2	−0.7 (2)
C16—S1—C1—C2	69.76 (13)	C2—C1—C8—O1	0.08 (16)
C8—C1—C2—C7	−0.34 (16)	S1—C1—C8—O1	−175.09 (10)
S1—C1—C2—C7	175.05 (11)	C2—C1—C8—C10	−179.64 (16)
C8—C1—C2—C3	178.82 (16)	S1—C1—C8—C10	5.2 (3)
S1—C1—C2—C3	−5.8 (2)	C7—O1—C8—C1	0.22 (16)
C7—C2—C3—C4	0.3 (2)	C7—O1—C8—C10	180.00 (12)
C1—C2—C3—C4	−178.78 (16)	C5—O3—C9—O2	0.8 (2)
C2—C3—C4—O2	178.99 (15)	C4—O2—C9—O3	−1.3 (2)
C2—C3—C4—C5	−0.4 (2)	C1—C8—C10—C11	5.9 (3)
C9—O2—C4—C3	−178.18 (17)	O1—C8—C10—C11	−173.83 (13)
C9—O2—C4—C5	1.29 (19)	C1—C8—C10—C15	−173.27 (17)
C9—O3—C5—C6	178.63 (17)	O1—C8—C10—C15	7.0 (2)
C9—O3—C5—C4	−0.03 (19)	C15—C10—C11—C12	0.5 (2)
C3—C4—C5—C6	0.0 (3)	C8—C10—C11—C12	−178.70 (14)
O2—C4—C5—C6	−179.54 (15)	C10—C11—C12—F1	179.18 (13)
C3—C4—C5—O3	178.68 (15)	C10—C11—C12—C13	−0.6 (2)
O2—C4—C5—O3	−0.81 (19)	F1—C12—C13—C14	−179.45 (14)
O3—C5—C6—C7	−177.91 (15)	C11—C12—C13—C14	0.3 (2)
C4—C5—C6—C7	0.6 (2)	C12—C13—C14—C15	0.1 (2)
C8—O1—C7—C2	−0.45 (16)	C13—C14—C15—C10	−0.2 (2)
C8—O1—C7—C6	−179.59 (14)	C11—C10—C15—C14	−0.1 (2)
C3—C2—C7—O1	−178.82 (13)	C8—C10—C15—C14	179.08 (14)
C1—C2—C7—O1	0.49 (16)	O4—S1—C16—C17	−172.19 (12)
C3—C2—C7—C6	0.3 (2)	C1—S1—C16—C17	77.73 (13)

Hydrogen-bond geometry (Å, °)

Cg1 is the centroid of the C2—C7 benzene ring.

D—H···A	D—H	H···A	D···A	D—H···A
C9—H9A···O4 ⁱ	0.99	2.45	3.314 (2)	145
C9—H9B···O2 ⁱⁱ	0.99	2.59	3.519 (3)	156
C16—H16A···O4 ⁱⁱⁱ	0.99	2.29	3.2689 (19)	169
C6—H6···F1 ^{iv}	0.95	2.37	3.2532 (18)	154
C17—H17B···Cg1 ^v	0.98	2.74	3.642 (3)	153

Symmetry codes: (i) $x+1, -y+3/2, z+1/2$; (ii) $x, -y+3/2, z+1/2$; (iii) $x, -y+3/2, z-1/2$; (iv) $x+1, y, z+1$; (v) $x, y, z-1$.