

Complete Genome Sequences of the Historical *Legionella pneumophila* Strains OLDA and Pontiac

Jeffrey W. Mercante, Shatavia S. Morrison, Brian H. Raphael, Jonas M. Winchell

Division of Bacterial Diseases, National Center for Immunization and Respiratory Diseases, Centers for Disease Control and Prevention, Atlanta, Georgia, USA

Here, we report the complete genome sequences of *Legionella pneumophila* serogroup 1 strains OLDA and Pontiac, which predate the 1976 Philadelphia Legionnaires' disease outbreak. Strain OLDA was isolated in 1947 from an apparent sporadic case, and strain Pontiac caused an explosive outbreak at a Michigan health department in 1968.

Received 24 June 2016 Accepted 29 June 2016 Published 25 August 2016

Citation Mercante JW, Morrison SS, Raphael BH, Winchell JM. 2016. Complete genome sequences of the historical *Legionella pneumophila* strains OLDA and Pontiac. *Genome Announc* 4(4):e00866-16. doi:10.1128/genomeA.00866-16.

Copyright © 2016 Mercante et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Jonas M. Winchell, jwinchell@cdc.gov.

Legionnaires' disease (LD) is a severe and sometimes fatal bacterial pneumonia caused by a growing list of *Legionella* spp., the most clinically relevant is *L. pneumophila*. One of the first high-profile LD outbreaks occurred in 1976 at an American Veterans' convention in Philadelphia (1). Eight years prior to the Philadelphia outbreak, a large epidemic of febrile, nonpneumonic disease, termed "Pontiac Fever," sickened 144 individuals at the Oakland County health department in Pontiac, Michigan (2). The cause of this illness was identified in 1977 when CDC researchers successfully isolated *L. pneumophila* serogroup 1 (sg1) from previously frozen guinea pig tissues experimentally inoculated with the Pontiac agent in 1968 (3). The "rickettsia-like" organism, OLDA, among the earliest cultured legionellae, was originally isolated in 1947 (4) from an apparent sporadic LD case; in 1977, this organism was identified as the same species and serogroup as the Philadelphia outbreak bacterium (5). Both strains were recently propagated from frozen stocks in the CDC archival collection with storage dates of 25 July 1978 for strain OLDA and 10 April 1978 for strain Pontiac.

Pacific Biosciences RSII-compatible (Menlo Park, CA, USA) long-insert DNA libraries were constructed for both strains according to the manufacturer's 10-kb protocol (PN100-286-100-04) and sequenced on single SMRT cells using 240-min movies. Assembly was performed with the Hierarchical Genome Assembly Process 3 (HGAP3) (6) through the PacBio SMRT Analysis System at ~85× coverage using 80,689 and 70,696 reads for the OLDA and Pontiac genomes, respectively. Overlapping contig ends were identified with Gepard version 1.3 (7) and trimmed to produce closed, circular genomes; 2,541,000 (OLDA) and 2,869,376 (Pontiac) paired-end 250-bp Illumina MiSeq (San Diego, CA, USA) reads were mapped to the PacBio assemblies to verify nucleotide accuracy >99.99%. The main OLDA genome is 3,486,082 bp with one circular plasmid, pLP3 (8) (G+C of 37.5%), of 129,883 bp, while strain Pontiac is 3,544,954 bp without extrachromosomal elements; both genomes have a G+C content of 38.4%. The NCBI Prokaryotic Genome Annotation Pipeline (PGAP) (9) identified 3,135 and 3,084 predicted protein coding genes for OLDA and Pontiac, respectively, and nine

rRNAs, 43 tRNAs, and four ncRNAs in both strains; strain OLDA harbored a single CRISPR array. Genome analyses with Genious version 9 (10) revealed that the previously described ~30-kb unstable genetic element (11) was absent from strain OLDA, but a pP36-like element (12) is maintained. A deletion of the *lag-1* O-acetyltransferase also corroborates its monoclonal antibody-2 (mAb2 = mAb3/1) negative phenotype (13, 14). Strain OLDA is sequence type 1 (ST1) (15) and highly similar (≥98.5%) to the ST1 *L. pneumophila* reference strain Paris (NC_006368) (16). Strain Pontiac (ST62) is 93.4% identical to the *L. pneumophila* reference strain L10-023 (NC_002942.5) of the same sequence type, but lacks a 70.5-kb genomic island (position 3,042,379), and exhibits a genomic rearrangement near a *trb/vir* conjugal transfer locus (at position ~2,729,000).

The complete genomes of strains OLDA and Pontiac, among the oldest known clinical and environmental *L. pneumophila* isolates, respectively, will be valuable as reference sequences as we attempt to understand the diversity of this species and its public health importance.

Accession number(s). The whole-genome sequences described here have been deposited at NCBI/GenBank under the accession numbers CP016030 (OLDA chromosome), CP016031 (OLDA plasmid), and CP016029 (Pontiac).

ACKNOWLEDGMENTS

The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

FUNDING INFORMATION

This study was supported, in part, by funds made available through the Office of Advanced Molecular Detection at the Centers for Disease Control and Prevention.

REFERENCES

1. McDade JE, Shepard CC, Fraser DW, Tsai TR, Redus MA, Dowdle WR. 1977. Legionnaires' disease— isolation of a bacterium and demonstration of its role in other respiratory disease. *N Engl J Med* 297:1197–1203. <http://dx.doi.org/10.1056/NEJM197712012972202>.

2. Glick TH, Gregg MB, Berman B, Mallison G, Rhodes WW, Jr, Kasanoff I. 1978. Pontiac fever. An epidemic of unknown etiology in a health department: I. Clinical and epidemiologic aspects. *Am J Epidemiol* 107:149–160.
3. Kaufmann AF, McDade JE, Patton CM, Bennett JV, Skaliy P, Feeley JC, Anderson DC, Potter ME, Newhouse VF, Gregg MB, Brachman PS. 1981. Pontiac fever: isolation of the etiologic agent (*Legionella pneumophila*) and demonstration of its mode of transmission. *Am J Epidemiol* 114:337–347.
4. Jackson EB, Crocker TT, Smadel JE. 1952. Studies on two rickettsia-like agents probably isolated from guinea pigs. *Bacteriol Proc* 52:119.
5. McDade JE, Brenner DJ, Bozeman FM. 1979. Legionnaires' disease bacterium isolated in 1947. *Ann Intern Med* 90:659–661. <http://dx.doi.org/10.7326/0003-4819-90-4-659>.
6. Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <http://dx.doi.org/10.1038/nmeth.2474>.
7. Krumsiek J, Arnold R, Rattei T. 2007. Gepard: a rapid and sensitive tool for creating dotplots on genome scale. *Bioinformatics* 23:1026–1028. <http://dx.doi.org/10.1093/bioinformatics/btm039>.
8. Mikesell P, Ezzell JW, Knudson GB. 1981. Isolation of plasmids in *Legionella pneumophila* and *Legionella*-like organisms. *Infect Immun* 31:1270–1272.
9. Tatusova T, DiCuccio M, Badretdin A. 2013. Prokaryotic genome annotation pipeline. In *The NCBI Handbook*, 2nd ed. NCBI, Bethesda, MD. <http://www.ncbi.nlm.nih.gov/books/NBK174280>.
10. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics* 28:1647–1649. <http://dx.doi.org/10.1093/bioinformatics/bts199>.
11. Lüneberg E, Mayer B, Daryab N, Kooistra O, Zähringer U, Rohde M, Swanson J, Frosch M. 2001. Chromosomal insertion and excision of a 30 kb unstable genetic element is responsible for phase variation of lipopolysaccharide and other virulence determinants in *Legionella pneumophila*. *Mol Microbiol* 39:1259–1271. <http://dx.doi.org/10.1111/j.1365-2958.2001.02314.x>.
12. Doléans-Jordheim A, Akermi M, Ginevra C, Cazalet C, Kay E, Schneider D, Buchrieser C, Atlan D, Vandenesch F, Etienne J, Jarraud S. 2006. Growth-phase-dependent mobility of the *lvh*-encoding region in *Legionella pneumophila* strain Paris. *Microbiology* 152:3561–3568. <http://dx.doi.org/10.1099/mic.0.29227-0>.
13. Petzold M, Thürmer A, Menzel S, Mouton JW, Heuner K, Lück C. 2013. A structural comparison of lipopolysaccharide biosynthesis loci of *Legionella pneumophila* serogroup 1 strains. *BMC Microbiol* 13:198. <http://dx.doi.org/10.1186/1471-2180-13-198>.
14. Joly JR, McKinney RM, Tobin JO, Bibb WF, Watkins ID, Ramsay D. 1986. Development of a standardized subgrouping scheme for *Legionella pneumophila* serogroup 1 using monoclonal antibodies. *J Clin Microbiol* 23:768–771.
15. Gaia V, Fry NK, Afshar B, Lück PC, Meugnier H, Etienne J, Peduzzi R, Harrison TG. 2005. Consensus sequence-based scheme for epidemiological typing of clinical and environmental isolates of *Legionella pneumophila*. *J Clin Microbiol* 43:2047–2052. <http://dx.doi.org/10.1128/JCM.43.5.2047-2052.2005>.
16. Brudno M, Do CB, Cooper GM, Kim MF, Davydov E, Program NCS, Green ED, Sidow A, Batzoglou S. 2003. LAGAN and Multi-LAGAN: efficient tools for large-scale multiple alignment of genomic DNA. *Genome Res* 13:721–731. <http://dx.doi.org/10.1101/gr.926603>.