

trans-Bis(μ -2-hydroxyethanethiolato- κ^2 S:S)bis[dinitrosyliron(II)](Fe—Fe)

Hon Man Lee^a and Show-Jen Chiou^{b*}

^aDepartment of Chemistry, National Changhua University of Education, Changhua, Taiwan 50058, and ^bDepartment of Applied Chemistry, National Chiayi University, 300 Syuefu Road, Chiayi City 60004, Taiwan

Correspondence e-mail: genechiou@mail.ncyu.edu.tw


Received 26 October 2009; accepted 12 November 2009

Key indicators: single-crystal X-ray study; $T = 150$ K; mean $\sigma(C-C) = 0.003$ Å; R factor = 0.028; wR factor = 0.061; data-to-parameter ratio = 19.2.

The title complex, $[Fe_2(C_2H_5OS)_2(NO)_4]$, lies on a crystallographic inversion center. The Fe—Fe distance is characteristic of a metal–metal bond. In the crystal structure, intermolecular O—H···O hydrogen bonds link complex molecules into a two-dimensional network.

Related literature

For iron–nitrosyl complexes, see: Chiang *et al.* (2004); Dillinger *et al.* (2007); Mazany *et al.* (1983).



Experimental

Crystal data

$[Fe_2(C_2H_5OS)_2(NO)_4]$
 $M_r = 385.98$
Monoclinic, $C2/c$
 $a = 16.943$ (3) Å
 $b = 5.0070$ (7) Å
 $c = 14.931$ (2) Å
 $\beta = 94.327$ (3)°

$V = 1263.1$ (3) Å³
 $Z = 4$
Mo $K\alpha$ radiation
 $\mu = 2.65$ mm⁻¹
 $T = 150$ K
 $0.21 \times 0.18 \times 0.02$ mm

Data collection

Bruker SMART APEXII diffractometer
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.606$, $T_{\max} = 0.949$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.028$
 $wR(F^2) = 0.061$
 $S = 0.94$
1597 reflections
83 parameters
H-atom parameters constrained
 $\Delta\rho_{\max} = 0.52$ e Å⁻³
 $\Delta\rho_{\min} = -0.32$ e Å⁻³

Table 1
Selected geometric parameters (Å, °).

Fe1—N1	1.6650 (19)	Fe1—S1	2.2619 (6)
Fe1—S1 ⁱ	2.2555 (7)	Fe1—Fe1 ⁱ	2.7051 (6)
N1—Fe1—N2	117.44 (9)	N2—Fe1—S1	106.00 (7)
N1—Fe1—S1	110.22 (6)		

Symmetry code: (i) $-x + 2, -y, -z$.

Table 2
Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
O3—H3 ⁱⁱ —O3 ⁱⁱ	0.84	1.97	2.7950 (14)	166

Symmetry code: (ii) $-x + \frac{3}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$.

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *DIAMOND* (Brandenburg, 1999).

We are grateful to the National Science Council of Taiwan for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2939).

References

- Brandenburg, K. (1999). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Chiang, C.-Y., Miller, M. L., Reibenspies, J. H. & Daresbourg, M. Y. (2004). *J. Am. Chem. Soc.* **126**, 10867–10867.
- Dillinger, S. A. T., Schmalke, H. W., Berke, T. & Fox, H. (2007). *Dalton Trans.* pp. 3562–3571.
- Mazany, A. M., Fackler, J. P. Jr, Gallagher, M. K. & Seyerth, D. (1983). *Inorg. Chem.* **22**, 2593–2596.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.

supplementary materials

Acta Cryst. (2009). E65, m1600 [doi:10.1107/S1600536809048065]

trans-Bis(μ-2-hydroxyethanethiolato-κ²S:S)bis[dinitrosyliron(II)](Fe—Fe)

H. M. Lee and S.-J. Chiou

Comment

Roussin's red esters (RREs) are photochemically active NO-release compounds and are expected to become potential NO donor drugs (Dillinger *et al.* 2007). The title compound is an air-stable and water soluble RRE compound. The molecular structure of the title complex is shown in Fig. 1. The molecule lies on a crystallographic inversion center. The Fe–Fe distance is characteristic of a M–M bond (Dillinger *et al.* 2007). In the crystal structure, intermolecular O–H···O hydrogen bonds link complex molecules into a two-dimensional network.

Experimental

$\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ (5.0 g, 0.018 mol) was added to H_2SO_4 (4*M*, 50 ml) at 273 K. To the resulting solution, the mixture of NaNO_2 (2.0 g, 0.023 mol) and mercaptoethanol (5.0 ml, 0.071 mol) were added. After stirring for 10 min, the solution was kept in refrigerator (277 K) for one week. Dark red crystals formed and one was used for this structure determination. Yield: 2.1 g (65%). FTIR (THF): 1809 (*w*), 1774 (*vs*), 1748 (*s*) cm^{-1} (*vNO*).

Refinement

All H atoms were positioned geometrically and refined as riding atoms, with $\text{C}_{\text{methylene}}\text{—H}$ = 0.99, $\text{O}\text{—H}$ = 0.84 Å while $U_{\text{iso}}(\text{H})$ = 1.2*U*_{eq}(C) and $U_{\text{iso}}(\text{H})$ = 1.5*U*_{eq}(O) for all the H atoms.

Figures




Fig. 1. The molecular structure of the title complex, showing 50% displacement ellipsoids for non-H atoms. The H atoms are depicted by circles of an arbitrary radius. Unlabeled atoms of the complex are related to labeled atoms by the symmetry operator ($2 - x, -y, -z$).




Fig. 2. Part of the crystal structure showing intermolecular hydrogen bonds as dashed lines.

trans-Bis(μ-2-hydroxyethanethiolato- κ²S:S)bis[dinitrosyliron(II)](Fe—Fe)

Crystal data

$[\text{Fe}_2(\text{C}_2\text{H}_5\text{OS})_2(\text{NO})_4]$

$F_{000} = 776$

supplementary materials

$M_r = 385.98$	$D_x = 2.030 \text{ Mg m}^{-3}$
Monoclinic, $C2/c$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
Hall symbol: -C 2yc	Cell parameters from 1286 reflections
$a = 16.943 (3) \text{ \AA}$	$\theta = 2.4\text{--}26.8^\circ$
$b = 5.0070 (7) \text{ \AA}$	$\mu = 2.65 \text{ mm}^{-1}$
$c = 14.931 (2) \text{ \AA}$	$T = 150 \text{ K}$
$\beta = 94.327 (3)^\circ$	Plate, brown
$V = 1263.1 (3) \text{ \AA}^3$	$0.21 \times 0.18 \times 0.02 \text{ mm}$
$Z = 4$	

Data collection

Bruker SMART APEXII diffractometer	1240 reflections with $I > 2\sigma$
Monochromator: graphite	$R_{\text{int}} = 0.039$
$T = 150 \text{ K}$	$\theta_{\max} = 28.7^\circ$
ω scans	$\theta_{\min} = 2.4^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -22 \rightarrow 22$
$T_{\min} = 0.606, T_{\max} = 0.949$	$k = -6 \rightarrow 6$
5708 measured reflections	$l = -19 \rightarrow 19$
1597 independent reflections	

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.028$	H-atom parameters constrained
$wR(F^2) = 0.061$	$w = 1/[\sigma^2(F_o^2) + (0.0324P)^2]$ where $P = (F_o^2 + 2F_c^2)/3$
$S = 0.94$	$(\Delta/\sigma)_{\max} = 0.003$
1597 reflections	$\Delta\rho_{\max} = 0.52 \text{ e \AA}^{-3}$
83 parameters	$\Delta\rho_{\min} = -0.32 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -

factors(gt) etc. and is not relevant to the choice of reflections for refinement. *R*-factors based on F^2 are statistically about twice as large as those based on F , and *R*-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.89173 (13)	0.1311 (4)	0.14871 (15)	0.0189 (5)
H1A	0.8810	-0.0613	0.1385	0.023*
H1B	0.9190	0.1523	0.2093	0.023*
C2	0.81447 (12)	0.2848 (5)	0.14308 (15)	0.0219 (5)
H2A	0.8253	0.4791	0.1449	0.026*
H2B	0.7836	0.2436	0.0858	0.026*
Fe1	1.062332 (16)	-0.00956 (6)	0.06209 (2)	0.01677 (10)
N1	1.06466 (10)	-0.2257 (4)	0.14697 (12)	0.0200 (4)
N2	1.13967 (11)	0.1950 (4)	0.05687 (13)	0.0216 (4)
O1	1.07764 (10)	-0.3658 (3)	0.20944 (11)	0.0306 (4)
O2	1.19777 (10)	0.3213 (4)	0.06634 (13)	0.0370 (5)
O3	0.77015 (10)	0.2112 (3)	0.21691 (12)	0.0302 (4)
H3	0.7591	0.3487	0.2456	0.045*
S1	0.95465 (3)	0.25677 (10)	0.06417 (4)	0.01729 (13)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0256 (11)	0.0179 (11)	0.0142 (12)	-0.0014 (9)	0.0070 (8)	0.0007 (9)
C2	0.0243 (11)	0.0202 (12)	0.0220 (13)	-0.0032 (9)	0.0071 (9)	0.0006 (10)
Fe1	0.01897 (16)	0.01530 (17)	0.01622 (17)	-0.00062 (13)	0.00241 (11)	-0.00012 (13)
N1	0.0220 (9)	0.0206 (10)	0.0173 (10)	-0.0001 (8)	0.0012 (7)	-0.0025 (8)
N2	0.0225 (9)	0.0206 (10)	0.0224 (11)	-0.0002 (8)	0.0060 (8)	-0.0021 (8)
O1	0.0449 (10)	0.0241 (9)	0.0220 (10)	-0.0008 (8)	-0.0020 (8)	0.0069 (7)
O2	0.0276 (9)	0.0363 (11)	0.0477 (12)	-0.0119 (8)	0.0069 (8)	-0.0058 (9)
O3	0.0378 (9)	0.0235 (9)	0.0322 (11)	-0.0049 (8)	0.0222 (8)	-0.0037 (7)
S1	0.0212 (2)	0.0139 (3)	0.0174 (3)	-0.0006 (2)	0.00546 (19)	-0.0004 (2)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.515 (3)	Fe1—N2	1.6696 (19)
C1—S1	1.824 (2)	Fe1—S1 ⁱ	2.2555 (7)
C1—H1A	0.9900	Fe1—S1	2.2619 (6)
C1—H1B	0.9900	Fe1—Fe1 ⁱ	2.7051 (6)
C2—O3	1.428 (2)	N1—O1	1.174 (2)
C2—H2A	0.9900	N2—O2	1.170 (2)
C2—H2B	0.9900	O3—H3	0.8400
Fe1—N1	1.6650 (19)	S1—Fe1 ⁱ	2.2555 (7)
C2—C1—S1	109.51 (15)	N2—Fe1—S1 ⁱ	110.41 (7)
C2—C1—H1A	109.8	N1—Fe1—S1	110.22 (6)
S1—C1—H1A	109.8	N2—Fe1—S1	106.00 (7)

supplementary materials

C2—C1—H1B	109.8	S1 ⁱ —Fe1—S1	106.43 (2)
S1—C1—H1B	109.8	N1—Fe1—Fe1 ⁱ	121.14 (6)
H1A—C1—H1B	108.2	N2—Fe1—Fe1 ⁱ	121.42 (7)
O3—C2—C1	109.18 (18)	S1 ⁱ —Fe1—Fe1 ⁱ	53.324 (17)
O3—C2—H2A	109.8	S1—Fe1—Fe1 ⁱ	53.106 (18)
C1—C2—H2A	109.8	O1—N1—Fe1	170.09 (16)
O3—C2—H2B	109.8	O2—N2—Fe1	169.24 (18)
C1—C2—H2B	109.8	C2—O3—H3	109.5
H2A—C2—H2B	108.3	C1—S1—Fe1 ⁱ	110.16 (7)
N1—Fe1—N2	117.44 (9)	C1—S1—Fe1	108.68 (7)
N1—Fe1—S1 ⁱ	105.88 (7)	Fe1 ⁱ —S1—Fe1	73.57 (2)


Symmetry codes: (i) $-x+2, -y, -z$.

Hydrogen-bond geometry (\AA , $^\circ$)

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O3—H3 ⁱⁱ —O3 ⁱⁱ	0.84	1.97	2.7950 (14)	166

Symmetry codes: (ii) $-x+3/2, y+1/2, -z+1/2$.

Fig. 1



supplementary materials

Fig. 2

