



# High-Quality Draft Genome Sequence of *Fischerella thermalis* JSC-11, a Siderophilic Cyanobacterium with Bioremediation Potential

Igor Brown,<sup>a\*</sup> Tanja Woyke,<sup>b</sup> Natalia Ivanova,<sup>b</sup> Nicole Shapiro,<sup>b</sup> Jaime Alcorta,<sup>c</sup> Andrei Chistoserdov,<sup>d</sup> Donald Pan,<sup>e</sup> Svetlana Sarkisova,<sup>a</sup> Susannah G. Tringe<sup>b</sup>

<sup>a</sup>Jacobs Engineering/NASA Johnson Space Center, Houston, Texas, USA

<sup>b</sup>Department of Energy Joint Genome Institute, Lawrence Berkeley National Laboratory, Berkeley, California, USA

<sup>c</sup>Department of Molecular Genetics and Microbiology, Biological Sciences Faculty, Pontifical Catholic University of Chile, Santiago, Chile

<sup>d</sup>Department of Biology, University of Louisiana at Lafayette, Lafayette, Louisiana, USA

<sup>e</sup>School of Oceanography, Shanghai Jiao Tong University, Shanghai, China

**ABSTRACT** Here, we report the draft genome sequence of the siderophilic cyanobacterium *Fischerella thermalis* JSC-11, which was isolated from an iron-depositing hot spring. JSC-11 has bioremediation potential because it is capable of both extracellular absorption and intracellular mineralization of colloidal iron. This genomic information will facilitate the exploration of JSC-11 for bioremediation.

We report the high-quality draft genome sequence of the cyanobacterium JSC-11, which was isolated from an iron-depositing mat from Chocolate Pots Hot Springs (Yellowstone National Park, WY, USA) (temperature of 55°C, pH of 5.6, and total soluble iron concentration of ~20 μM) (1). Single-colony isolation was performed as described previously (2, 3). Because optimal growth occurred at 45 to 55°C and 0.4 to 0.6 mM iron, JSC-11 has been characterized as a thermophilic and siderophilic species (4). Its 16S rRNA sequence (GenBank accession number [HM636645.1](#)) shares 100% identity with those of several *Fischerella thermalis* strains, and it shares their characteristic true-branching morphology (5, 6). Therefore, it was classified as *F. thermalis* strain JSC-11 (CCMEE 7001). This strain absorbs iron oxides on its exopolymers and produces intracellular iron deposits (Fig. 1), making it a potential candidate for bioremediation (7).

Genomic DNA of JSC-11 was isolated using the UltraClean microbial DNA isolation kit (MoBio Laboratories, USA), following cultivation in DH medium as described previously (8). The draft genome of JSC-11 was generated at the U.S. Department of Energy (DOE) Joint Genome Institute (JGI) using a combination of four next-generation sequencing (NGS) libraries from the same DNA extract. An Illumina GAIIx (9) library was created using the KAPA-Illumina library creation kit (Kapa Biosystems), yielding 31,933,200 reads of 76-bp average length. One 454 Titanium library was created using the GS FLX Titanium rapid library preparation kit (Roche), generating 267,779 reads of 395-bp average length; two paired-end 454 libraries, with average insert sizes of 4 kb and 8 kb, were created according to the method described by Peng et al. (10), generating 602,963 reads of 160- and 188-bp average length, respectively. Kits were used according to the manufacturers' instructions.

The 454 data were assembled with Newbler v2.3 (9), the Illumina data were assembled with Velvet v1.0.13 (11), and the contigs were computationally shredded into 2-kb and 1.5-kb overlapping fake reads (shreds), which were integrated with the 454 paired-end library reads using parallel Phrap v1.080812 (High Performance Software, LLC). POLISHER (12) was used to correct base errors. Misassemblies were corrected by using Gap Resolution (13), Dupfinisher (14), or sequencing of cloned bridging PCR

**Editor** Julia A. Maresca, University of Delaware

**Copyright** © 2022 Brown et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](#).

Address correspondence to Igor Brown, [igor\\_brown@hotmail.com](mailto:igor_brown@hotmail.com), or Susannah G. Tringe, [sgtringe@lbl.gov](mailto:sgtringe@lbl.gov).


\*Present address: Igor Brown, 3437 Stettler View Rd., Charlotte, North Carolina, USA.

The authors declare no conflict of interest.

**Received** 5 August 2022

**Accepted** 11 October 2022

**Published** 27 October 2022



**FIG 1** Scanning transmission electron microscopy (STEM) views of extracellular and intracellular iron-rich particles found in JSC-11 cells incubated with  $600 \mu\text{M}$   $\text{FeCl}_3 \cdot 6\text{H}_2\text{O}$  and  $0.04 \text{ mM}$   $\text{Na}_2\text{HPO}_4 \cdot 7\text{H}_2\text{O}$ . (Upper, left) Bright-field STEM photograph of a JSC-11 cell encrusted with external Fe-bearing precipitates and containing an electron-dense, internal, Fe-rich particle  $\sim 100 \text{ nm}$  in size. (Upper, right) Dark-field STEM photograph to localize Fe-rich aggregates. (Lower) Quantitative element maps for internal and external particles, showing distribution patterns for phosphorus (P), Fe, and oxygen (O). The electron-dense particles within the JSC-11 cytoplasm contain substantial amounts of P, Fe, and O.

fragments with subcloning. Gaps were closed manually using Consed (15, 16) and 168 PCR and Bubble PCR primer walks.

Annotation was performed with the Oak Ridge National Laboratory annotation pipeline using Prodigal v1.4 (17), followed by manual curation using the JGI GenePRIMP pipeline (18). The JSC-11 genome is 5,380,000 bp long, with a G+C content of 41.0%, and has 99.76% completeness and 0.0% contamination according to CheckM v1.1.3 (19). It contains 4,367 coding genes, 1 rRNA operon, and 43 tRNA genes.

The JSC-11 genome may shed light on specific mechanisms of iron homeostasis in siderophilic cyanobacteria, because it encodes proteins predicted to be involved in iron transport (21 total) or maintaining a balance between intracellular  $\text{Fe}^{2+}$  and  $\text{Fe}^{3+}$  (5 total). Genomic analysis also revealed that JSC-11 may be capable of far-red-light photoacclimation (20).

**Data availability.** The GenBank accession number for the 16S rRNA gene is [HM636645.1](https://www.ncbi.nlm.nih.gov/nuclot/HM636645.1). The GenBank assembly accession number is [GCA\\_000231365.2](https://www.ncbi.nlm.nih.gov/assembly/GCA_000231365.2). The NCBI RefSeq assembly accession number is [GCF\\_000231365.1](https://www.ncbi.nlm.nih.gov/refseq/assembly/GCF_000231365.1). The BioProject accession number is [PRJNA61093](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA61093). The nucleotide GenBank accession number is [AGIZ00000000.1](https://www.ncbi.nlm.nih.gov/nuclot/AGIZ00000000.1). The SRA accession numbers for the raw sequences are [SRR21655585](https://www.ncbi.nlm.nih.gov/sra/SRR21655585), [SRR21655586](https://www.ncbi.nlm.nih.gov/sra/SRR21655586), [SRR21655587](https://www.ncbi.nlm.nih.gov/sra/SRR21655587), and [SRR3927276](https://www.ncbi.nlm.nih.gov/sra/SRR3927276).

#### ACKNOWLEDGMENTS

I.B. and S.S. thank former colleagues at the NASA Johnson Space Center for their help with studies in 2006 to 2010. I.B. extends his personal gratitude to K. Thomas-Keptra for

transmission electron microscopy and energy-dispersive spectroscopy studies. We further thank D. A. Bryant and G. Shen (Pennsylvania State University) for their cooperation in additional purification of isolated DNA and A. Copeland and A. Lapidus for helpful discussions of the manuscript.

The study carried out at Jacobs Engineering/NASA Johnson Space Center was partially supported by the NASA Johnson Space Center Director Discretionary Fund in 2008 and 2009. The work (proposal 10.46936/10.25585/60007313) conducted by the DOE JGI (<https://ror.org/04xm1d337>), a DOE Office of Science User Facility, was supported by the Office of Science of the DOE, operated under contract DE-AC02-05CH11231. J.A. was supported by a Chilean ANID doctoral fellowship (grant 21191763).

## REFERENCES

- Fortney NW, He S, Converse BJ, Boyd ES, Roden EE. 2018. Investigating the composition and metabolic potential of microbial communities in Chocolate Pots hot springs. *Front Microbiol* 9:2075. <https://doi.org/10.3389/fmicb.2018.02075>.
- Brown II, Mummy D, Cooksey KE. 2005. A novel cyanobacterium exhibiting an elevated tolerance for iron. *FEMS Microbiol Ecol* 52:307–314. <https://doi.org/10.1016/j.femsec.2004.11.020>.
- Brown II, Bryant DA, Casamatta D, Thomas-Keprta KL, Sarkisova SA, Shen G, Graham JE, Boyd ES, Peters JW, Garrison DH, McKay DS. 2010. Polyphasic characterization of a thermotolerant siderophilic filamentous cyanobacterium that produces intracellular iron deposits. *Appl Environ Microbiol* 76:6664–6672. <https://doi.org/10.1128/AEM.00662-10>.
- Brown I, Allen C, Mummy DL, Sarkisova S, McKay DS. 2007. Iron-tolerant cyanobacteria, p 425–442. *In* Seckbach J (ed), *Algae and cyanobacteria in extreme environments*. Springer, Dordrecht, Netherlands. [https://doi.org/10.1007/978-1-4020-6112-7\\_23](https://doi.org/10.1007/978-1-4020-6112-7_23).
- Alcorta J, Vergara-Barros P, Antonaru LA, Alcamán-Arias ME, Nürnberg DJ, Diez B. 2019. *Fischerella thermalis*: a model organism to study thermophilic diazotrophy, photosynthesis and multicellularity in cyanobacteria. *Extremophiles* 23:635–647. <https://doi.org/10.1007/s00792-019-01125-4>.
- Rippka R, Deruelles J, Waterbury JB, Herdman M, Stanier RY. 1979. Generic assignments, strain histories and properties of pure cultures of cyanobacteria. *Microbiology* 111:1–61. <https://doi.org/10.1099/00221287-111-1-1>.
- Al-Amin A, Parvin F, Chakraborty J, Kim Y-I. 2021. Cyanobacteria mediated heavy metal removal: a review on mechanism, biosynthesis, and removal capability. *Environ Technol Rev* 10:44–57. <https://doi.org/10.1080/21622515.2020.1869323>.
- Brown I, Tringe SG, Ivanova N, Goodwin L, Shapiro N, Alcorta J, Pan D, Chistoserdov A, Sarkisova S, Woyke T. 2021. High-quality draft genome sequence of the siderophilic and thermophilic *Leptolyngbyaceae* cyanobacterium JSC-12. *Microbiol Resour Announc* 10:e00495-21. <https://doi.org/10.1128/MRA.00495-21>.
- Margulies M, Egholm M, Altman WE, Attiya S, Bader JS, Bemben LA, Berka J, Braverman MS, Chen Y-J, Chen Z, Dewell SB, Du L, Fierro JM, Gomes XV, Godwin BC, He W, Helgesen SHC, Irzyk GP, Jando SC, Alenquer MLI, Jarvie TP, Jirage KB, Kim J-B, Knight JR, Lanza JR, Leamon JH, Lefkowitz SM, Lei M, Li J, Lohman KL, Lu H, Makhijani VB, McDade KE, McKenna MP, Myers EW, Nickerson E, Nobile JR, Plant R, Puc BP, Ronan MT, Roth GT, Sarkis GJ, Simons JF, Simpson JW, Srinivasan M, Tartaro KR, Tomasz A, Vogt KA, Volkmer GA, Wang SH, Wang Y, Weiner MP, Yu P, Begley RF, Rothberg JM. 2005. Genome sequencing in microfabricated high-density picolitre reactors. *Nature* 437:376–380. <https://doi.org/10.1038/nature03959>.
- Peng Z, Zhao Z, Nath N, Froula JL, Clum A, Zhang T, Cheng J-F, Copeland AC, Pennacchio LA, Chen F. 2012. Generation of long insert pairs using a Cre-LoxP inverse PCR approach. *PLoS One* 7:e29437. <https://doi.org/10.1371/journal.pone.0029437>.
- Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res* 18:821–829. <https://doi.org/10.1101/gr.074492.107>.
- LaButti K, Foster B, Lowry S, Trong S, Goltsman E, Lapidus A. 2008. POLISHER: a tool for using ultra short reads in microbial genome finishing. LBNL report LBNL-639E. <https://escholarship.org/uc/item/9wh0d388>.
- Labutti K, Foster B, Lapidus A. 2017. Gap resolution. <https://www.osti.gov/doi/code/biblio/57346>.
- Han C, Chain P. 2006. Finishing repeat regions automatically with Dupfinisher, p 141–146. *In* Arabina HR, Valafar H (ed), *Proceeding of the 2006 International Conference on Bioinformatics & Computational Biology, BIOCOMP 06, Las Vegas, Nevada, USA, June 26–29, 2006*. <https://researchr.org/publication/biocomp%3A2006>.
- Ewing B, Green P. 1998. Base-calling of automated sequencer traces using Phred. II. Error probabilities. *Genome Res* 8:186–194. <https://doi.org/10.1101/gr.8.3.186>.
- Ewing B, Hillier L, Wendl MC, Green P. 1998. Base-calling of automated sequencer traces using Phred. I. Accuracy assessment. *Genome Res* 8:175–185. <https://doi.org/10.1101/gr.8.3.175>.
- Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119. <https://doi.org/10.1186/1471-2105-11-119>.
- Pati A, Ivanova NN, Mikhailova N, Ovchinnikova G, Hooper SD, Lykidis A, Kyrpides NC. 2010. GenePRIMP: a gene prediction improvement pipeline for prokaryotic genomes. *Nat Methods* 7:455–457. <https://doi.org/10.1038/nmeth.1457>.
- Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 25:1043–1055. <https://doi.org/10.1101/gr.186072.114>.
- Gan F, Bryant DA. 2015. Adaptive and acclimative responses of cyanobacteria to far-red light: responses of cyanobacteria to far-red light. *Environ Microbiol* 17:3450–3465. <https://doi.org/10.1111/1462-2920.12992>.