


# Study of the effects of interferon $\beta$ -1a on hospitalized patients with COVID-19: SBMU Taskforce on the COVIFERON study

Mohammad Fallahzadeh<sup>1,2</sup> | Mohamad A. Pourhoseingholi<sup>2,3</sup> |  
 Masoud G. Boroujeni<sup>1,2</sup> | Sajad Besharati<sup>1,2</sup> | Masoud Mardani<sup>1,2,4</sup> |  
 Minoosh Shabani<sup>1,2,4</sup> | Shervin Shokouhi<sup>1,2,4</sup> | Mahdi Amirdosara<sup>2,5</sup> |  
 Mohammadreza Hajjesmaeili<sup>2,5</sup> | Latif Gachkar<sup>1,2,4</sup> | Baran Roshan<sup>2,6</sup> |  
 Masoud Zangi<sup>2,5</sup> | Golshan Mirmomeni<sup>2,7</sup> | Seyed S. N. Irvani<sup>2,8</sup> |  
 Ilad Alavi Darazam<sup>1,2,4</sup> 

<sup>1</sup>Infectious Diseases and Tropical Medicine Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>2</sup>SBMU Taskforce on the COVIFERON Study, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>3</sup>Gastroenterology and Liver Diseases Research Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>4</sup>Department of Infectious Diseases and Tropical Medicine, Loghman Hakim Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>5</sup>Department of Anesthesiology, Loghman Hakim Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran

<sup>6</sup>Department of Clinical Pharmacy, Faculty of Pharmacy, Shahid Beheshti University of Medical Science, Tehran, Iran

<sup>7</sup>Hearing Research Center, Clinical Science Research Institute, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

<sup>8</sup>Independent Senior Researcher, Tabriz, Iran

## Correspondence

Ilad Alavi Darazam, Department of Infectious Diseases, Loghman Hakim Hospital, Shahid Beheshti University of Medical Sciences; Infectious Diseases and Tropical Medicine Research Center, Shahid Beheshti University of Medical Sciences, Velenjak Koodakyar Ave, Tehran, Iran.

Email: [ilad.alavi@sbmu.ac.ir](mailto:ilad.alavi@sbmu.ac.ir) and [ilad13@yahoo.com](mailto:ilad13@yahoo.com)

## Abstract

Interferons are an essential part of the innate immune system and have antiviral and immunomodulatory functions. We studied the effects of interferon  $\beta$ -1a on the outcomes of severe cases of coronavirus disease 2019 (COVID-19). This retrospective study was conducted on hospitalized COVID-19 patients in Loghman-Hakim hospital from February 20, 2020 to April 20, 2020, Tehran, Iran. Patients were selected from two groups, the first group received interferon  $\beta$ -1a in addition to the standard treatment regimen, and the second group received standard care. The clinical progression of two groups during their hospital admission was compared. We studied a total number of 395 hospitalized COVID-19 patients. Out of this number, 111 patients (33.5%) died (31.3% of the interferon  $\beta$ -1a group and 34.1% of the control group). The mortality rate indicated no statistically significant difference between groups ( $p$ -value = 0.348), however for patients who were hospitalized for more than a week, the rate of mortality was lower in the interferon  $\beta$ -1a group ( $p$ -value = 0.014). The median hospital stay was statistically longer for patients treated by interferon  $\beta$ -1a ( $p$ -value < 0.001). The results of this study showed that interferon  $\beta$ -1a can improve the outcomes of hospitalized patients with severe COVID-19, but more adequately-powered randomized controlled trials should be conducted.

## KEYWORDS

COVID-19, efficacy, IFN- $\beta$ 1a, outcome

## 1 | INTRODUCTION

In December 2019, a new virus belonging to the coronaviridae family was reported in Wuhan, China, which was named severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). This virus caused a global outbreak of a respiratory disease known as coronavirus disease 2019 (COVID-19).<sup>1</sup> COVID-19 has a wide range of symptoms including mild self-limited disease to severe progressing pneumonia and multiple organ failure.<sup>2-4</sup> The high rate of transmission and mortality (3.7%) of this disease<sup>5</sup> made it a public health emergency of international concern (PHEIC).<sup>6</sup>

Clinical researchers made efforts to find a specific treatment for COVID-19, and have suggested some kind of medications for the management of this disease.<sup>7,8</sup> The efficacy of different categories of drugs have been evaluated in clinical trials and some medications had relative efficacy.<sup>9</sup> The emergency condition of the COVID-19 pandemic and lack of available drugs of proven efficacy for COVID-19 caused to repurpose drugs for treating COVID-19 which were used for other purposes previously.<sup>10</sup> Various pharmacological interventions were suggested to treat COVID-19, however, their efficiency is questionable. Some antiviral agents, including remdesivir and combination of lopinavir and ritonavir, and immunomodulatory drugs, consisting of corticosteroids, hydroxychloroquine, and interferons, have been used in COVID-19 patents.<sup>11-16</sup>

The innate immune response plays an important role in novel viral infections without previous established adaptive immunity to the pathogen to downturn the severity of the disease.<sup>17</sup> Interferons are an essential part of this immune action and have antiviral and immunomodulatory functions.<sup>17</sup> In a recent clinical trial, interferon beta1-a (IFN- $\beta$ 1a) had significant efficacy in patients with COVID-19.<sup>16</sup> Previous *in vitro* investigations showed the antiviral effect of interferon- $\beta$  against the SARS virus.<sup>18</sup> In this retrospective study, we assessed the clinical results of IFN- $\beta$ 1a in severe cases of COVID-19.

## 2 | MATERIALS AND METHODS

### 2.1 | Study design and setting

This retrospective study was conducted on hospitalized COVID-19 patients in Loghman-Hakim hospital from February 20, 2020 to April 20, 2020, Tehran, Iran. This study started after research ethics committee approval. Patients were selected from two groups with different treatment regimens for COVID-19, which were matched, based on age, sex, and severity of disease regarding arterial oxygen saturation. Patients of the first group received interferon  $\beta$ -1a (IFN- $\beta$ 1a) (Recigen) in addition to standard treatment regimen and patients of the second group received standard care. We obtained demographic characteristics, background health conditions, clinical symptoms, and signs, and laboratory and imaging findings of patients.

### 2.2 | Patients

We studied patients with COVID-19 which was confirmed by positive polymerase chain reactions (PCRs) and positive chest computed tomography (CT) scans. All patients had clinical symptoms accordant with COVID-19. Participants were hospitalized in the ward or intense care unit of Loghman Hospital due to severe COVID-19 pneumonia. Severe COVID-19 was defined as patients with arterial blood oxygen saturation level less than 90 percent in the initial assessment and who suffer from moderate to severe dyspnea, based on patients' assessment. Some of these patients had received IFN- $\beta$ 1a in addition to standard treatment regimen and the rest had received just standard care. Patients of these two groups were matched, based on age, sex, and the severity of the disease. Clinical information of patients has been extracted from Loghman-Hakim hospital central database.

### 2.3 | Treatment

In this study, the patients were selected from two groups. Both groups received medication based on the protocol of the health ministry of Iran for COVID-19 including lopinavir/ritonavir (400 mg/100 mg twice a day for 10 days) (Kaletra) + hydroxychloroquine (400 mg single dose). This treatment regimen was based on Iran's internal guideline for COVID-19 in February 2020, when we initiated this study.<sup>19</sup> Patients of the first group also received IFN- $\beta$ 1a (subcutaneous injections of 44  $\mu$ g (12 000 000 IU) on Days 1, 3, and 5), (Recigen).

### 2.4 | Outcomes

We studied the clinical progression of two groups during their hospital admission. The mortality rate in the early 7 days and in the late phase of admission, duration of hospital admission, arterial blood gas findings, complete blood count findings, C-reactive protein, and erythrocyte sedimentation rate have been compared between two groups.

### 2.5 | Statistical analysis

Frequency rates and percentages were used for categorical variables, and interquartile ranges (IQRs) and median were used for continuous variables. For comparison of the non-normal continuous variables, the Kruskal-Wallis test was used.  $\chi^2$  test was used for comparing the frequency of categorical variables and a logistic regression model was also applied to calculate the odds ratios (ORs) with 95% confidence intervals (CIs). R software version 3.6.1 was used to perform the statistical analyses.

**TABLE 1** Characteristics of the patients at baseline

Parameters	Total (n = 331)	Interferon (n = 64)	Standard Care (n = 267)	p-value
<b>Characteristics</b>				
Age (year)	64.9 (18.2)	62.9 (22.1)	65.4 (17.1)	0.413
Male sex - no. (%)	203 (61.3%)	34 (53.1%)	169 (63.3%)	0.153
<b>Underlying conditions</b>				
Diabetes	107 (32.33%)	17 (26.56%)	90 (33.71%)	0.301
Hypertension	142 (42.90%)	30 (46.88%)	112 (41.95%)	0.485
Cardiovascular disease (CVD)	78 (23.56%)	17 (26.56%)	61 (22.85%)	0.516
Rheumatologic condition	6 (1.81%)	3 (4.69%)	3 (1.12%)	0.089
Asthma	15 (4.53%)	2 (3.13%)	13 (4.87%)	0.744
COPD	23 (6.95%)	5 (7.81%)	18 (6.74%)	0.785
Chronic liver disease	1 (0.3%)	1 (1.56%)	0 (0%)	1.000
Transplant receiver	5 (1.51%)	3 (4.69%)	2 (0.75%)	0.526
Malignancy	7 (2.11%)	1 (1.56%)	6 (2.25%)	0.594
HIV	8 (2.42%)	4 (6.25%)	4 (1.50%)	0.048
Hepatitis B	1 (0.3%)	1 (1.56%)	0 (0%)	1.000
Hypothyroidism	5 (1.51%)	0 (0%)	5 (1.87%)	0.587
<b>Respiratory factors</b>				
Oxygen saturation (SpO <sub>2</sub> ) – median (IQR)	50.1 (35.48–75.2)	57.8 (35.7–83.9)	47.6 (35.3–73.43)	0.157
pH (DISS) - median (IQR)	7.4 (7.37–7.46)	7.4 (7.36–7.47)	7.4 (7.37–7.46)	0.512
PaCO <sub>2</sub> (DISS) - median (IQR)	38.1 (32.08–46.1)	38.2 (30.7–48.13)	39.2 (32.2–46.07)	0.382
PaO <sub>2</sub> (DISS) - median (IQR)	26.9 (21–40.8)	29.5 (20.8–44.2)	26.3 (21–39.23)	0.879
HCO <sub>3</sub> (DISS) - median (IQR)	27.7 (23.5–27.5)	26.3 (24–27.5)	25.5 (23.1–27.5)	0.041
Respiratory rate	19 (17–22)	18 (16–20)	19 (18–22)	<0.001
<b>White blood cell count (×10<sup>9</sup>/L) – median (IQR)</b>				
<4 × 10 <sup>9</sup> /L – no. (%)	23 (7.26%)	4 (6.25%)	19 (7.51%)	0.934
4–10 × 10 <sup>9</sup> /L – no. (%)	200 (63.09%)	42 (65.62%)	158 (62.45%)	0.88
>10 × 10 <sup>9</sup> /L – no. (%)	94 (28.84%)	18 (28.13%)	76 (30.04%)	
<b>Lymphocyte count (×10<sup>9</sup>/L) – median (IQR)</b>				
≥1.0 × 10 <sup>9</sup> /L – no. (%)	139 (44.4%)	24 (38.1%)	115 (46%)	0.653
<1.0 × 10 <sup>9</sup> /L – no. (%)	174 (55.6%)	39 (61.9%)	135 (54%)	0.321
<b>Platelet count (×10<sup>9</sup>/L) – median (IQR)</b>				
≥100 × 10 <sup>9</sup> /L – no. (%)	301 (95.25%)	63 (98.34%)	236 (94.4%)	0.174
<100 × 10 <sup>9</sup> /L – no. (%)	15 (4.75%)	1 (1.56%)	14 (5.6%)	0.180
<b>Neutrophil count (×10<sup>9</sup>/liter) – median (IQR)</b>				
<1.5 × 10 <sup>9</sup> /L – no. (%)	7 (2.36%)	2 (3.23%)	5 (2.13%)	0.654
1.5–8 × 10 <sup>9</sup> /L – no. (%)	193 (64.98%)	41 (66.13%)	152 (64.68%)	0.834
>8 × 10 <sup>9</sup> /L – no. (%)	97 (32.66%)	19 (30.64%)	78 (33.19%)	
<b>Aspartate aminotransferase (AST) (U/L) – median (IQR)</b>				
≤40 U/L – no. (%)	67 (27.34%)	10 (17.24%)	57 (%)	0.915
>40 U/L – no. (%)	178 (72.66%)	48 (82.76%)	130 (%)	0.032

TABLE 1 (Continued)

Parameters	Total (n = 331)	Interferon (n = 64)	Standard Care (n = 267)	p-value
Alanine aminotransferase (ALT) (U/L) - median (IQR)	59 (38–98)	53.4 (37–96)	59.5 (38–99.25)	0.802
≤50 U/L - no. (%)	104 (42.79%)	26 (45.62%)	78 (41.94%)	0.366
>50 U/L - no. (%)	139 (57.21%)	31 (54.38%)	108 (58.06%)	
Lactate Dehydrogenase (LDH) (U/L) - median (IQR)	444.5(301–687.5)	578(383-845)	428(283.5–643)	0.398
≤245 U/L - no. (%)	18 (15.25%)	3 (9.09%)	15 (17.65%)	0.193
>245 U/L - no. (%)	100 (84.75%)	30 (90.91%)	70 (82.35%)	
C-reactive protein (CRP) - median (IQR)	57.05 (29.5–82.9)	48.5 (25.65-69.5)	60 (33.55–83.65)	0.121
CRP < 6 - no. (%)	20 (9.71%)	7 (14.28%)	13 (8.28%)	0.267
CRP > 6 - no. (%)	186 (90.29%)	42 (85.72%)	144 (91.72%)	
Erythrocyte sedimentation rate (ESR) - median (IQR)	48 (25–69)	48 (22–71)	48 (25–68)	0.556
Serum creatinine (μmol/L) - median (IQR)	1.3 (1–1.6)	1.1 (1–1.475)	1.3 (1.1–1.7)	0.412

Abbreviations: COPD, chronic obstructive pulmonary disease; IQR, interquartile range.

TABLE 2 Outcomes

Parameters	Total (n = 331)	Interferon (n = 64)	Standard care (n = 267)	p-value
Mortality	111 (33.5%)	20 (31.3%)	91 (34.1%)	0.768
Early Mortality* - no. (%)	72 (21.75%)	14 (21.87%)	58 (21.72%)	0.348
Late Mortality** - no. (%)	39 (15.06%)	6 (12.00%)	33 (15.78%)	0.014
Hospital stay – median no. of days (IQR)	4 (2–7)	7 (5–9)	4 (2–7)	<0.001

\*Mortality before 7 days of hospitalization.

\*\*Mortality after 7 days of hospitalization.

### 3 | RESULTS

In this retrospective study, which was conducted on hospitalized patients in Loghman-Hakim hospital from February 20, 2020 to April 20, 2020, Tehran Iran with severe COVID-19 pneumonia, was confirmed by positive reverse-transcription polymerase chain reaction and positive chest CT scans. We studied a total number of 395 patients in this study, and from them, 64 patients received IFN-β1a (Recigen) plus the standard care (lopinavir/ritonavir [Kaletra] + hydroxychloroquine) and 331 control patients received standard care.

The mean (±SD) age of total patients was 64.9 (18.2) with the majority being male (61.3%). There were no statistically significant differences in age and sex between the two groups. In Table 1, the demographic and clinical factors across two study groups were presented (Table 1). Although most clinical factors were distributed similarly across the two groups, the rate of HIV infection, HCO<sub>3</sub>, and respiratory rate were statistically different between the two groups (Table 1). Out of the 395 patients under study, 111 patients (33.5%) died (31.3% of the Recigen group and 34.1% of the control group). The mortality rate indicated no difference between the groups, however for patients who were hospitalized for more than a week,

the rate of mortality was lower in the Recigen group, compared to the corresponding group among those who received only the standard care (12.00% mortality in Recigen group vs. 15.78% mortality in the control group, Table 2). Also, the median hospital stay was statistically longer for patients treated by IFN-β1a (median 7 days in the Recigen group vs. 4 days in the control group, p-value < 0.001) (Table 2).

From other clinical factors, the median of white blood cell count, lymphocyte count, and neutrophil count was not significantly different between the Recigen group and the controls. Also, we used a logistic regression model to calculate the odds ratio of mortality for patients who were just treated by the standard care, compare to the Recigen group. According to analysis in a crude model, the odds of death in control groups, compared to the Recigen group was 1.13 (95% CI: 0.63–2.04, p value = 0.67) and the adjusted odds ratio (for respiratory rate, saturation, and HCO<sub>3</sub>) was 1.13 (95% CI: 0.58–2.18, p value = 0.71). As the mortality was lower for the Recigen group in patients with more than one week of hospitalization, we did a subgroup analysis for those patients. The results indicated a substantial significant difference between the two groups; the crude odds ratio was 3.96 (95% CI: 1.37–11.43, p value = 0.011) as similar as the

adjusted odds ratio, which was 4.02 (95% CI: 1.22–13.21,  $p$  value = 0.022), which revealed that patients who were treated with standard care and hospitalized for at least one week were at higher risk of death, compared to same patients who treated with Recigen.

## 4 | DISCUSSION

This study revealed that IFN- $\beta$ 1a decreased the days of hospitalization and the mortality rate in severe COVID-19 patients who had been hospitalized for more than 7 days.

The mortality rate in severe COVID-19 patients was reported between 62% and 81%.<sup>20</sup> A recent study showed COVID-19 patients with more severe disease had significantly decreased interferon activity and SARS-CoV-2 suppressed IFN- $\beta$  release in vitro.<sup>21,22</sup> There is an increased risk of severe lung disease in people with comorbidities, older age, or who receive immunosuppressive medication due to less IFN- $\beta$  production.<sup>23,24</sup>

IFN- $\beta$  is superior to other interferons in inhibition of coronaviruses replication.<sup>25</sup> It should be used in the early phase of viral infection to reach a protective effect and late application of interferon may exacerbate the disease.<sup>26,27</sup> IFN- $\beta$  increases CD73 which plays a role in vascular integrity in hypoxic conditions.<sup>28</sup> A high serum level of interferon is needed to reach an antiviral effect.<sup>27</sup> Long-term safety and tolerability of IFN- $\beta$  are proven due to its application in multiple sclerosis treatment.<sup>29</sup>

In our study, the two groups didn't have a significant difference in mortality rate; however, in patients who were hospitalized for more than a week, the rate of mortality was lower in the Recigen group, compared to those corresponding ones who received standard care. In addition, the median hospital stay was statistically longer for patients treated by IFN- $\beta$ 1a.

In a previous randomized trial, a combination of INF- $\beta$  with lopinavir/ritonavir and ribavirin was safe, alleviated the symptoms, and shortened the duration of hospital stay compared with triple antiviral therapy alone.<sup>30</sup>

In another trial, IFN- $\beta$  significantly improved the discharge rate and lowered the 28-day mortality, especially in patients who received it in the early phase of the disease.<sup>31</sup>

We had some limitations in this study. The most important limitation is that this study was conducted in the critical phase of the COVID-19 outbreak in Iran when we didn't have enough information about COVID-19, so treatment was not conforming to COVID-19. In addition, there might be some problems in the data gathering, group matching, and homogeneity of the patients at the beginning of this outbreak. Considering the above limitations, we didn't design this study as a randomized clinical trial.

## 5 | CONCLUSION

Although the results of this study showed that IFN- $\beta$ 1a can improve the outcomes of severe COVID-19 patients, more adequately-powered randomized controlled trials should be conducted to

determine the initiation time, dose, and duration of treatment of IFN- $\beta$ 1a in severe COVID-19 patients.

## ACKNOWLEDGEMENTS

The authors would like to thank the Clinical Research Development Unit (CRDU) of Loghman Hakim Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran for their help and support in conducting this retrospective observational study. All the authors of this paper are members of the SBMU Taskforce on The COVIFERON. This study received no specific funds from any sector.

## CONFLICT OF INTERESTS

The authors declare that there are no conflict of interests.

## ETHICS STATEMENT

This study was approved by the research ethics committees of the Shahid Beheshti University of Medical Science and the ethical code number is IR. SBMU. RETECH. REC.1399.079.

## AUTHOR CONTRIBUTIONS

Mohammad Fallahzadeh, Ilad Alavi Darazam, and Sajad Besharati prepared the first draft. Mahdi Amir Pourhoseingholi conducted the analysis on data and edited the first draft. Masoud Ghanbari Boroujeni and Ilad Alavi Darazam finalized all drafts, approved the final version of the manuscript, and made the decision to submit the results. Seyed Sina Naghibi Irvani conceived of the study and provided overall guidance. Ilad Alavi Darazam, Shervin Shokouhi, Minoosh Shabani, Mahdi Amirdosara, Mohammadreza Hajjesmaeili, Masoud Mardani, Golshan Mirmomeni, and Masoud Zangi were involved in conducting the trial, recruited patients, and took clinical care of the patients. All other authors gathered data, reviewed and interpreted results, or provided guidance on methodology. All authors critically reviewed and revised the manuscript, and approved the final version of the manuscript.

## DATA AVAILABILITY STATEMENT

The data presented in the manuscript could be available for the journal.

## ORCID

Ilad Alavi Darazam  <http://orcid.org/0000-0002-4440-335X>

## REFERENCES

- Zhu N, Zhang D, Wang W, et al. A novel coronavirus from patients with pneumonia in China, 2019. *N Engl J Med.* 2020;382:727-733.
- Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet.* 2020;395(10223):497-506.
- Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *The Lancet.* 2020;395(10223):507-513.
- Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA.* 2020;323(11):1061-1069.

5. Organization WH. Coronavirus disease 2019 (COVID-19): situation report, 74. 2020.
6. Ebrahimi M, Saki A. Laboratory findings, signs and symptoms, clinical outcomes of Patients with COVID-19 Infection: an updated systematic review and meta-analysis. *medRxiv*. 2020
7. Gottlieb RL, Nirula A, Chen P, et al. Effect of bamlanivimab as monotherapy or in combination with etesevimab on viral load in patients with mild to moderate COVID-19: a randomized clinical trial. *JAMA*. 2021;325(7):632-644.
8. Singh AK, Majumdar S, Singh R, Misra A. Role of corticosteroid in the management of COVID-19: A systemic review and a Clinician's perspective. *Diabetes Metab Syndr*. 2020;14(5):971-978.
9. Dong L, Hu S, Gao J. Discovering drugs to treat coronavirus disease 2019 (COVID-19). *Drug Discov Ther*. 2020;14(1):58-60.
10. World Health Organization Solidarity" clinical trial for COVID-19 treatments. World Health Organization (WHO) Situation reports Geneva: WHO. Accessed April 5, 2020. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/global-research-on-novel-coronavirus-2019-ncov/solidarity-clinical-trial-for-covid-19-treatments>
11. YAVUZ S, Ünal S. Antiviral treatment of COVID-19. *Turk J Med Sci*. 2020;50(SI-1):611-619.
12. Wang M, Cao R, Zhang L, et al. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell Res*. 2020;30(3):269-271.
13. Lim J, Jeon S, Shin H-Y, et al. Case of the index patient who caused tertiary transmission of COVID-19 infection in Korea: the application of lopinavir/ritonavir for the treatment of COVID-19 infected pneumonia monitored by quantitative RT-PCR. *J Korean Med Sci*. 2020;35(6):79.
14. Cascella M, Rajnik M, Cuomo A, Dulebohn SC, Di Napoli R. Features, evaluation, and treatment of coronavirus (COVID-19). *Statpearls [internet]*. 2021
15. Beigel JH, Tomashek KM, Dodd LE, et al. Remdesivir for the Treatment of Covid-19 - Final Report. *N Engl J Med*. 2020;383(19):1813-1826.
16. Hung IF, Lung KC, Tso EY, et al. Triple combination of interferon beta-1b, lopinavir-ritonavir, and ribavirin in the treatment of patients admitted to hospital with COVID-19: an open-label, randomised, phase 2 trial. *Lancet (London, England)*. 2020;395(10238):1695-1704.
17. Li S-f, Gong M-j, Zhao F-r, et al. Type I interferons: distinct biological activities and current applications for viral infection. *Cell Physiol Biochem*. 2018;51(5):2377-96.
18. Stockman LJ, Bellamy R, Garner P. SARS: systematic review of treatment effects. *PLoS Med*. 2006;3(9):343.
19. Rahmzade R, Rahmzadeh R, Hashemian SM, Tabarsi P. Iran's approach to COVID-19: evolving treatment protocols and ongoing clinical trials. *Front Public Health*. 2020;8(523):551889.
20. Weiss P, Murdoch DR. Clinical course and mortality risk of severe COVID-19. *The Lancet*. 2020;395(10229):1014-1015.
21. Yuen C-K, Lam J-Y, Wong W-M, et al. SARS-CoV-2 nsp13, nsp14, nsp15 and orf6 function as potent interferon antagonists. *Emerg Microbes Infect*. 2020;9:1-29.
22. Hadjadj J, Yatim N, Barnabei L, et al. Impaired type I interferon activity and exacerbated inflammatory responses in severe COVID-19 patients. *MedRxiv*. Published online April 23, 2020.
23. Agrawal A. Mechanisms and implications of age-associated impaired innate interferon secretion by dendritic cells: a mini-review. *Gerontology*. 2013;59(5):421-426.
24. Singanayagam A, Loo S-L, Calderazzo M, et al. Antiviral immunity is impaired in COPD patients with frequent exacerbations. *Am J Physiol -Lung Cell Mol Physiol*. 2019;317(6):L893-L903.
25. Spiegel M, Pichlmair A, Mühlberger E, Haller O, Weber F. The antiviral effect of interferon-beta against SARS-coronavirus is not mediated by MxA protein. *J Clin Virol*. 2004;30(3):211-213.
26. Jalkanen J, Hollmén M, Jalkanen S. Interferon beta-1a for COVID-19: critical importance of the administration route. *Crit Care*. 2020;24(1):1-3.
27. Brzoska J, von Eick H, Hündgen M. Interferons in the therapy of severe coronavirus infections: a critical analysis and recollection of a forgotten therapeutic regimen with interferon beta. *Drug Res*. 2020;70(7):291-297.
28. Bellingan G, Maksimow M, Howell DC, et al. The effect of intravenous interferon-beta-1a (FP-1201) on lung CD73 expression and on acute respiratory distress syndrome mortality: an open-label study. *The Lancet Respiratory Medicine*. 2014;2(2):98-107.
29. Group PS. PRISMS-4: long-term efficacy of interferon-beta-1a in relapsing MS. *Neurology*. 2001;56(12):1628-1636.
30. Hung IF, Lung K-C, Tso EY, et al. Triple combination of interferon beta-1b, lopinavir-ritonavir, and ribavirin in the treatment of patients admitted to hospital with COVID-19: an open-label, randomised, phase 2 trial. *The Lancet*. 2020;395(10238):1695-1704.
31. Davoudi-Monfared E, Rahmani H, Khalili H, et al. A randomized clinical trial of the efficacy and safety of interferon  $\beta$ -1a in treatment of severe COVID-19. *Antimicrob Agents Chemother*. 2020;64(9).

**How to cite this article:** Fallahzadeh M, Pourhoseingholi MA, Boroujeni MG, et al. Study of the effects of interferon  $\beta$ -1a on hospitalized patients with COVID-19: SBMU Taskforce on the COVIFERON study. *J Med Virol*. 2022;94:1488-1493. doi:10.1002/jmv.27475