

## cyclo-Tetrakis( $\mu$ -3-acetyl-4-methyl-1H-pyrazole-5-carboxylato- $\kappa^4$ N<sup>2</sup>,O<sup>3</sup>:N<sup>1</sup>,O<sup>5</sup>)tetrakis[aquacopper(II)] tetradeca-hydrate

Sergey Malinkin,<sup>a\*</sup> Irina A. Golenya,<sup>a</sup> Vadim A. Pavlenko,<sup>a</sup> Matti Haukka<sup>b</sup> and Turganbay S. Iskenderov<sup>a</sup>

<sup>a</sup>Kiev National Taras Shevchenko University, Department of Chemistry, Volodymyrska Str. 64, 01601 Kiev, Ukraine, and <sup>b</sup>University of Joensuu, Department of Chemistry, PO Box 111, FI-80101 Joensuu, Finland  
Correspondence e-mail: malinachem@mail.ru


Received 26 July 2011; accepted 1 August 2011

Key indicators: single-crystal X-ray study;  $T = 100$  K; mean  $\sigma(C-C) = 0.003$  Å; disorder in solvent or counterion;  $R$  factor = 0.034;  $wR$  factor = 0.097; data-to-parameter ratio = 24.2.

The title compound,  $[Cu_4(C_7H_6N_2O_3)_4(H_2O)_4] \cdot 14H_2O$ , a tetrานuclear [2 × 2] grid-type complex with  $S4$  symmetry, contains four Cu<sup>II</sup> atoms which are bridged by four pyrazole-carboxylate ligand anions and are additionally bonded to a water molecule. Each Cu<sup>II</sup> atom is coordinated by two O atoms of the carboxylate and acetyl groups, two pyrazole N atoms of doubly deprotonated 3-acetyl-4-methyl-1H-pyrazole-5-carboxylic acid and one O atom of a water molecule. The geometry at each Cu<sup>II</sup> atom is distorted square-pyramidal, with the two N and two O atoms in the equatorial plane and O atoms in the axial positions. O—H···O hydrogen-bonding interactions additionally stabilize the structure. One of the uncoordinated water molecules shows half-occupancy.

### Related literature

For the use of pyrazolate ligands in the preparation of poly-nuclear supramolecular compounds, see: Piguet *et al.* (1997); Krämer *et al.* (2002); Zhang *et al.* (1996); Van der Vlugt *et al.* (2008); Klingele *et al.* (2007); Kovbasyuk *et al.* (2004); Pons *et al.* (2003). For the use of asymmetric ligands in the preparation of heterometallic complexes, see: Moroz *et al.* (2010). For related structures, see: Mokhir *et al.* (2002); Sliva *et al.* (1997); Wörl *et al.* (2005a,b); Świątek-Kozłowska *et al.* (2000). For the preparation of related ligands, see: Sachse *et al.* (2008).



### Experimental

#### Crystal data

$[Cu_4(C_7H_6N_2O_3)_4(H_2O)_4] \cdot 14H_2O$	$Z = 4$
$M_r = 1243.00$	Mo $K\alpha$ radiation
Tetragonal, $I4_1/a$	$\mu = 1.76$ mm <sup>-1</sup>
$a = 13.8502$ (7) Å	$T = 100$ K
$c = 26.280$ (3) Å	$0.35 \times 0.25 \times 0.15$ mm
$V = 5041.1$ (6) Å <sup>3</sup>	

#### Data collection

Bruker SMART APEXII CCD diffractometer	37816 measured reflections
Absorption correction: multi-scan ( <i>SADABS</i> ; Bruker, 2009)	3993 independent reflections
$T_{min} = 0.578$ , $T_{max} = 0.778$	3254 reflections with $I > 2\sigma(I)$
	$R_{int} = 0.037$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$	165 parameters
$wR(F^2) = 0.097$	H-atom parameters constrained
$S = 1.06$	$\Delta\rho_{\text{max}} = 1.20$ e Å <sup>-3</sup>
3993 reflections	$\Delta\rho_{\text{min}} = -0.58$ e Å <sup>-3</sup>

**Table 1**  
Selected geometric parameters (Å, °).

Cu1—N2	1.9495 (16)	Cu1—N1	1.9682 (16)
Cu1—O2	1.9519 (14)	Cu1—O1	2.3938 (15)
Cu1—O4	1.9676 (15)		
N2—Cu1—O2	82.06 (6)	O4—Cu1—N1	91.15 (7)
O2—Cu1—O4	89.18 (6)	N1—Cu1—O1	74.28 (6)
N2—Cu1—N1	97.49 (7)		

**Table 2**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O4—H4O $\cdots$ O5	0.84	1.84	2.680 (3)	173
O4—H4P $\cdots$ O3 <sup>iv</sup>	0.84	2.03	2.868 (2)	177
O5—H5O $\cdots$ O5 <sup>v</sup>	0.88	2.22	2.808 (4)	124
O5—H5P $\cdots$ O7	0.80	1.97	2.766 (3)	171
O6—H6O $\cdots$ O7 <sup>vi</sup>	0.92	1.84	2.752 (2)	177
O6—H6P $\cdots$ O1	0.90	1.99	2.863 (2)	163
O7—H7O $\cdots$ O6 <sup>vii</sup>	0.83	1.92	2.707 (2)	157
O7—H7P $\cdots$ O3	0.83	2.21	3.016 (2)	166
O7—H7P $\cdots$ O2	0.83	2.33	2.951 (2)	132
O8—H8O $\cdots$ O5	0.85	1.81	2.644 (5)	167
O8—H8P $\cdots$ O6	0.81	2.01	2.815 (4)	174

Symmetry codes: (iv)  $-y + \frac{3}{4}, x + \frac{1}{4}, z + \frac{1}{4}$ ; (v)  $-x + 1, -y + \frac{1}{2}, z$ ; (vi)  $x, y + \frac{1}{2}, -z$ ; (vii)  $y - \frac{1}{4}, -x + \frac{3}{4}, z - \frac{1}{4}$ .

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SIR2004* (Burla *et al.*, 2005); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *DIAMOND* (Brandenburg, 2009); software used to prepare material for publication: *SHELXL97*.

Financial support from the State Fund for Fundamental Research of Ukraine (grant No. F40.3/041) and the Swedish Institute (Visby Program) is gratefully acknowledged.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JH2318).

## References

- Brandenburg, K. (2009). *DIAMOND*. Crystal Impact GbR, Bonn, Germany.
- Bruker (2009). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Burla, M. C., Caliandro, R., Camalli, M., Carrozzini, B., Cascarano, G. L., De Caro, L., Giacovazzo, C., Polidori, G. & Spagna, R. (2005). *J. Appl. Cryst.* **38**, 381–388.
- Klingele, J., Prikhodko, A. I., Leibeling, G., Demeshko, S., Dechert, S. & Meyer, F. (2007). *Dalton Trans.* pp. 2003–2008.
- Kovbasyuk, L., Pritzow, H., Krämer, R. & Fritsky, I. O. (2004). *Chem. Commun.* pp. 880–881.
- Krämer, R., Fritsky, I. O., Pritzow, H. & Kovbasyuk, L. A. (2002). *J. Chem. Soc. Dalton Trans.* pp. 1307–1314.
- Mokhir, A. A., Gumienna-Kontecka, E. S., Świątek-Kozłowska, J., Petkova, E. G., Fritsky, I. O., Jerzykiewicz, L., Kapshuk, A. A. & Sliva, T. Yu. (2002). *Inorg. Chim. Acta*, **329**, 113–121.
- Moroz, Y. S., Szyrweil, L., Demeshko, S., Kozłowski, H., Meyer, F. & Fritsky, I. O. (2010). *Inorg. Chem.* **49**, 4750–4752.
- Piguet, C., Bernardinelli, G. & Hopfgartner, G. (1997). *Chem. Rev.* **97**, 2005–2057.
- Pons, J., Sanchez, F. J., Casaby, J., Alvarez-Larena, A., Piniella, J. F. & Ros, J. (2003). *Inorg. Chem. Commun.* **6**, 833–836.
- Sachse, A., Penkova, L., Noel, G., Dechert, S., Varzatskii, O. A., Fritsky, I. O. & Meyer, F. (2008). *Synthesis*, **5**, 800–806.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Sliva, T. Yu., Kowalik-Jankowska, T., Amirkhanov, V. M., Głowiak, T., Onindo, C. O., Fritsky, I. O. & Kozłowski, H. (1997). *J. Inorg. Biochem.* **65**, 287–294.
- Świątek-Kozłowska, J., Fritsky, I. O., Dobosz, A., Karaczyn, A., Dudarenko, N. M., Sliva, T. Yu., Gumienna-Kontecka, E. & Jerzykiewicz, L. (2000). *J. Chem. Soc. Dalton Trans.* pp. 4064–4068.
- Van der Vlugt, J. I., Demeshko, S., Dechert, S. & Meyer, F. (2008). *Inorg. Chem.* **47**, 1576–1585.
- Wörl, S., Fritsky, I. O., Hellwinkel, D., Pritzow, H. & Krämer, R. (2005b). *Eur. J. Inorg. Chem.* pp. 759–765.
- Wörl, S., Pritzow, H., Fritsky, I. O. & Krämer, R. (2005a). *Dalton Trans.* pp. 27–29.
- Zhang, H., Fu, D., Ji, F., Wang, G., Yu, K. & Yao, T. (1996). *J. Chem. Soc. Dalton Trans.* pp. 3799–3805.

## **supplementary materials**

*Acta Cryst.* (2011). E67, m1260-m1261 [doi:10.1107/S1600536811030832]

**cyclo-Tetrakis( $\mu$ -3-acetyl-4-methyl-1H-pyrazole-5-carboxylato- $\kappa^4N^2,O^3:N^1,O^5$ )tetrakis[aquacopper(II)] tetradecahydrate**

**S. Malinkin, I. A. Golenya, V. A. Pavlenko, M. Haukka and T. S. Iskenderov**

**Comment**

Substituted pyrazolate ligands have found widespread use as building blocks for the formation of self-assembled supramolecular coordination complexes with an array of transition metal ions and a variety of different structures, e.g., helical polymers (Piguet *et al.*, 1997; Krämer *et al.*, 2002), so-called [2 × 2] grids (Zhang *et al.*, 1996; Van der Vlugt *et al.*, 2008; Klingele *et al.*, 2007) and other polynuclear structures (Kovbasyuk *et al.*, 2004; Pons *et al.*, 2003). Introduction of different donor substituents in 3- and 5-positions of the pyrazole ring is still rare, and such ligands can be successfully used for the obtaining of oligonuclear heterometallic species (Moroz *et al.*, 2010). Reported here is a new copper(II) complex with [2 × 2] grid-structure based on a novel asymmetric pyrazolate ligand having different substituents (the carboxylic and acetyl groups) in 3- and 5-positions.

In the title compound, (I), the tetranuclear [2 × 2] grid-type complex with S4 symmetry are composed of four Cu<sup>II</sup> ions, four ligands and four metal-bound water molecules (Fig. 1).

Each copper ion is nested in a square-pyramidal environment that is composed of the pyrazolate-N2, deprotonated carboxyl-O2 from a compartment of one ligand molecule and acetyl-O1 atoms, the pyrazolate-N1 from another ligand and one water-O4.

The intermetallic separations pyrazolate-bridged Cu<sup>II</sup> ions is 4.0600 (4) Å which is similar to that seen in the structures reported by Zhang *et al.*, 1996 (4.098 – 4.115 Å), while the distance between diagonal copper atoms is 5.0814 (5) Å, which is more longer to that observed in the structures reported by Klingele *et al.*, 2007 (4.7091 (5) Å) and Van der Vlugt *et al.*, 2008 (4.2308 (6) Å).

The coordinated pyrazolate ligand exhibits C—C, C—N, N—N bond lengths which are normal for bridging pyrazolate rings (Sliva *et al.*, 1997; Świątek-Kozłowska *et al.*, 2000; Mokhir *et al.*, 2002). The C—O bond lengths in the deprotonated carboxylic groups differs significantly (1.239 (2) and 1.292 (2)) which is typical for monodentately coordinated carboxylates (Wörl *et al.*, 2005*a,b*).

A part of the crystal packing of (I) is presented in Fig. 2. In the crystal packing the complex molecules are associated via intermolecular hydrogen bonds that involve the O—H interactions between the coordinated and the solvate water molecules and the non-coordinating carboxylate-O atoms. Thus, the tetranuclear molecules are stacked along the crystallographic *x* and *y* axes, forming the columns. The columns bisect one another at right angles to give a layer-like structure.

**Experimental**

The ligand 5-acetyl-4-methyl-1H-pyrazole-3-carboxylic acid (Sachse *et al.*, 2008) (0.25 g, 1.5 mmol) was added to a solution of Cu(Ac)<sub>2</sub>·H<sub>2</sub>O (0.30 g, 1.5 mmol) in H<sub>2</sub>O–CH<sub>3</sub>OH (50 mL) [80:20 v/v]. The reaction mixture was heated for 30 min with

## supplementary materials

---

constant stirring at 80 °C until completed dissolution of the ligand occurred. The resulting deep blue solution was filtered to remove any undissolved ligand and left at room temperature. Square block dark blue crystals suitable for X-ray diffraction were isolated after standing for several days (yield 0.32 g, 80%). Elemental analysis calc. (%) for C<sub>28</sub>H<sub>40</sub>Cu<sub>4</sub>N<sub>8</sub>O<sub>20</sub>: C 31.64; H 3.79; N 10.54; found: C 31.22; H 3.47; N 10.34.

### Refinement

The O—H and N—H hydrogen atoms were located from the difference Fourier map, and refined with  $U_{\text{iso}} = 1.5 U_{\text{eq}}$ (parent atom). The remaining H atoms were positioned geometrically and were constrained to ride on their parent atoms with C—H = 0.96–0.97 Å, and with  $U_{\text{iso}} = 1.2\text{--}1.5 U_{\text{eq}}$ (parent atom).

### Figures




Fig. 1. The molecular structure of the title compound. H atoms are omitted for clarity.




Fig. 2. A packing diagram for the title compound, showing the columns along the y-axis direction. Copper atoms and water molecules are depicted as the big and the small balls, respectively.

### cyclo-Tetrakis( $\mu$ -3-acetyl-4-methyl-1*H*-pyrazole-5-carboxylato- $\kappa^4N^2,O^3:N^1,O^5$ )tetrakis[aquacopper(II)] tetraadecahydrate

#### Crystal data

[Cu<sub>4</sub>(C<sub>7</sub>H<sub>6</sub>N<sub>2</sub>O<sub>3</sub>)<sub>4</sub>(H<sub>2</sub>O)<sub>4</sub>]·14H<sub>2</sub>O

$D_x = 1.638 \text{ Mg m}^{-3}$

$M_r = 1243.00$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Tetragonal,  $I4_1/a$

Cell parameters from 9905 reflections

Hall symbol: -I 4ad

$\theta = 2.6\text{--}30.3^\circ$

$a = 13.8502 (7) \text{ \AA}$

$\mu = 1.76 \text{ mm}^{-1}$

$c = 26.280 (3) \text{ \AA}$

$T = 100 \text{ K}$

$V = 5041.1 (6) \text{ \AA}^3$

Block, blue

$Z = 4$

$0.35 \times 0.25 \times 0.15 \text{ mm}$

$F(000) = 2560$

## *Data collection*

Bruker SMART APEXII CCD diffractometer	3993 independent reflections
Radiation source: fine-focus sealed tube	3254 reflections with $I > 2\sigma(I)$
flat graphite crystal	$R_{\text{int}} = 0.037$
Detector resolution: 16 pixels mm <sup>-1</sup>	$\theta_{\text{max}} = 30.9^\circ$ , $\theta_{\text{min}} = 1.7^\circ$
$\varphi$ scans and $\omega$ scans	$h = -19 \rightarrow 19$
Absorption correction: multi-scan ( <i>SADABS</i> ; Sheldrick, 2008)	$k = -19 \rightarrow 19$
$T_{\text{min}} = 0.578$ , $T_{\text{max}} = 0.778$	$l = -37 \rightarrow 38$
37816 measured reflections	

## *Refinement*

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.034$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.097$	H-atom parameters constrained
$S = 1.06$	$w = 1/[\sigma^2(F_o^2) + (0.0463P)^2 + 11.3899P]$ where $P = (F_o^2 + 2F_c^2)/3$
3993 reflections	$(\Delta/\sigma)_{\text{max}} < 0.001$
165 parameters	$\Delta\rho_{\text{max}} = 1.20 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\text{min}} = -0.58 \text{ e \AA}^{-3}$

## *Special details*

**Geometry.** All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted R-factor wR and goodness of fit S are based on  $F^2$ , conventional R-factors R are based on F, with F set to zero for negative  $F^2$ . The threshold expression of  $F^2 > 2\text{sigma}(F^2)$  is used only for calculating R-factors(gt) etc. and is not relevant to the choice of reflections for refinement. R-factors based on  $F^2$  are statistically about twice as large as those based on F, and R- factors based on ALL data will be even larger.

## *Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Cu1	0.164669 (16)	0.330841 (17)	0.089037 (8)	0.01628 (8)	
O1	0.19490 (11)	0.47600 (11)	0.13584 (6)	0.0220 (3)	
O2	0.18857 (10)	0.33808 (11)	0.01594 (5)	0.0191 (3)	
O3	0.11824 (11)	0.36118 (11)	-0.05924 (5)	0.0207 (3)	
O4	0.30114 (11)	0.29660 (13)	0.09939 (6)	0.0290 (4)	

## supplementary materials

---

H4O	0.3390	0.2983	0.0745	0.044*	
H4P	0.3283	0.3161	0.1259	0.044*	
O5	0.42451 (14)	0.31765 (18)	0.02160 (7)	0.0496 (6)	
H5O	0.4877	0.3162	0.0229	0.074*	
H5P	0.3994	0.3042	-0.0049	0.074*	
O6	0.34700 (12)	0.60006 (12)	0.10161 (6)	0.0300 (3)	
H6O	0.3363	0.6619	0.0904	0.045*	
H6P	0.2916	0.5694	0.1089	0.045*	
O7	0.32202 (12)	0.28601 (12)	-0.06718 (6)	0.0275 (3)	
H7O	0.3430	0.3107	-0.0939	0.041*	
H7P	0.2651	0.2997	-0.0606	0.041*	
O8	0.4789 (2)	0.4970 (4)	0.04080 (17)	0.0610 (16)	0.50
H8O	0.4561	0.4429	0.0310	0.092*	0.50
H8P	0.4390	0.5278	0.0564	0.092*	0.50
N1	0.13450 (12)	0.30012 (11)	0.16041 (6)	0.0161 (3)	
N2	0.03441 (11)	0.36571 (11)	0.06776 (6)	0.0152 (3)	
C1	0.17729 (14)	0.46349 (14)	0.18133 (8)	0.0191 (4)	
C2	0.18659 (19)	0.54292 (15)	0.21907 (9)	0.0284 (5)	
H2A	0.2408	0.5293	0.2419	0.043*	
H2B	0.1269	0.5478	0.2389	0.043*	
H2C	0.1982	0.6040	0.2012	0.043*	
C3	0.14765 (13)	0.36668 (13)	0.19793 (7)	0.0159 (3)	
C4	-0.07281 (13)	0.38673 (13)	0.00422 (7)	0.0161 (3)	
C5	-0.11564 (15)	0.39273 (16)	-0.04787 (7)	0.0218 (4)	
H5A	-0.0653	0.3807	-0.0733	0.033*	
H5B	-0.1430	0.4572	-0.0531	0.033*	
H5C	-0.1667	0.3442	-0.0513	0.033*	
C6	0.02326 (13)	0.36773 (13)	0.01655 (7)	0.0145 (3)	
C7	0.11406 (13)	0.35477 (13)	-0.01228 (7)	0.0159 (3)	

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Cu1	0.01545 (12)	0.02178 (13)	0.01160 (11)	-0.00004 (8)	-0.00341 (8)	0.00135 (8)
O1	0.0244 (7)	0.0211 (7)	0.0207 (7)	-0.0033 (5)	-0.0084 (5)	0.0037 (5)
O2	0.0159 (6)	0.0270 (7)	0.0145 (6)	-0.0023 (5)	-0.0004 (5)	0.0028 (5)
O3	0.0246 (7)	0.0253 (7)	0.0122 (6)	-0.0018 (6)	0.0017 (5)	0.0031 (5)
O4	0.0222 (7)	0.0459 (10)	0.0190 (7)	0.0079 (7)	-0.0080 (6)	-0.0070 (7)
O5	0.0298 (9)	0.0881 (17)	0.0307 (9)	0.0203 (10)	-0.0046 (8)	-0.0200 (10)
O6	0.0302 (8)	0.0300 (8)	0.0297 (8)	-0.0047 (7)	-0.0066 (7)	0.0033 (7)
O7	0.0244 (7)	0.0294 (8)	0.0288 (8)	-0.0005 (6)	0.0089 (6)	0.0036 (6)
O8	0.0102 (14)	0.102 (4)	0.071 (3)	0.0179 (18)	-0.0138 (16)	-0.068 (3)
N1	0.0212 (8)	0.0149 (7)	0.0121 (7)	0.0011 (6)	-0.0041 (6)	-0.0001 (5)
N2	0.0158 (7)	0.0197 (7)	0.0102 (6)	-0.0028 (5)	-0.0003 (5)	0.0028 (5)
C1	0.0199 (8)	0.0157 (8)	0.0216 (9)	0.0016 (6)	-0.0083 (7)	-0.0002 (7)
C2	0.0416 (13)	0.0164 (9)	0.0270 (10)	-0.0008 (8)	-0.0070 (9)	-0.0035 (8)
C3	0.0188 (8)	0.0150 (8)	0.0138 (8)	0.0031 (6)	-0.0046 (6)	-0.0008 (6)
C4	0.0172 (8)	0.0170 (8)	0.0140 (8)	-0.0046 (6)	-0.0030 (6)	0.0045 (6)

C5	0.0218 (9)	0.0284 (10)	0.0152 (8)	-0.0034 (8)	-0.0070 (7)	0.0038 (7)
C6	0.0164 (8)	0.0158 (8)	0.0112 (7)	-0.0037 (6)	-0.0017 (6)	0.0029 (6)
C7	0.0184 (8)	0.0155 (8)	0.0138 (8)	-0.0037 (6)	0.0009 (6)	0.0025 (6)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

Cu1—N2	1.9495 (16)	O8—H8O	0.8529
Cu1—O2	1.9519 (14)	O8—H8P	0.8080
Cu1—O4	1.9676 (15)	N1—N2 <sup>i</sup>	1.329 (2)
Cu1—N1	1.9682 (16)	N1—C3	1.362 (2)
Cu1—O1	2.3938 (15)	N2—N1 <sup>ii</sup>	1.329 (2)
Cu1—Cu1 <sup>i</sup>	4.0600 (4)	N2—C6	1.355 (2)
Cu1—Cu1 <sup>ii</sup>	4.0600 (4)	C1—C3	1.469 (3)
Cu1—Cu1 <sup>iii</sup>	5.0814 (5)	C1—C2	1.487 (3)
O1—C1	1.232 (3)	C2—H2A	0.9800
O2—C7	1.292 (2)	C2—H2B	0.9800
O3—C7	1.239 (2)	C2—H2C	0.9800
O4—H4O	0.8400	C3—C4 <sup>i</sup>	1.405 (3)
O4—H4P	0.8355	C4—C6	1.394 (2)
O5—H5O	0.8760	C4—C3 <sup>ii</sup>	1.405 (3)
O5—H5P	0.7998	C4—C5	1.494 (3)
O6—H6O	0.9174	C5—H5A	0.9800
O6—H6P	0.8985	C5—H5B	0.9800
O7—H7O	0.8324	C5—H5C	0.9800
O7—H7P	0.8289	C6—C7	1.479 (3)
N2—Cu1—O2	82.06 (6)	H7O—O7—H7P	114.5
N2—Cu1—O4	171.24 (6)	H8O—O8—H8P	111.3
O2—Cu1—O4	89.18 (6)	N2 <sup>i</sup> —N1—C3	108.05 (15)
N2—Cu1—N1	97.49 (7)	N2 <sup>i</sup> —N1—Cu1	130.03 (12)
O2—Cu1—N1	170.07 (6)	C3—N1—Cu1	121.00 (13)
O4—Cu1—N1	91.15 (7)	N1 <sup>ii</sup> —N2—C6	108.91 (15)
N2—Cu1—O1	95.81 (6)	N1 <sup>ii</sup> —N2—Cu1	137.70 (12)
O2—Cu1—O1	115.65 (6)	C6—N2—Cu1	113.30 (12)
O4—Cu1—O1	87.90 (6)	O1—C1—C3	118.16 (17)
N1—Cu1—O1	74.28 (6)	O1—C1—C2	121.73 (18)
N2—Cu1—Cu1 <sup>i</sup>	94.38 (5)	C3—C1—C2	120.11 (18)
O2—Cu1—Cu1 <sup>i</sup>	123.35 (4)	C1—C2—H2A	109.5
O4—Cu1—Cu1 <sup>i</sup>	90.48 (5)	C1—C2—H2B	109.5
N1—Cu1—Cu1 <sup>i</sup>	46.73 (5)	H2A—C2—H2B	109.5
O1—Cu1—Cu1 <sup>i</sup>	120.95 (4)	C1—C2—H2C	109.5
N2—Cu1—Cu1 <sup>ii</sup>	44.57 (5)	H2A—C2—H2C	109.5
O2—Cu1—Cu1 <sup>ii</sup>	125.84 (4)	H2B—C2—H2C	109.5
O4—Cu1—Cu1 <sup>ii</sup>	144.00 (5)	N1—C3—C4 <sup>i</sup>	109.93 (16)
N1—Cu1—Cu1 <sup>ii</sup>	55.97 (5)	N1—C3—C1	116.13 (16)
O1—Cu1—Cu1 <sup>ii</sup>	70.56 (4)	C4 <sup>i</sup> —C3—C1	133.70 (17)

## supplementary materials

---

Cu1 <sup>i</sup> —Cu1—Cu1 <sup>ii</sup>	77.482 (5)	C6—C4—C3 <sup>ii</sup>	103.01 (15)
N2—Cu1—Cu1 <sup>iii</sup>	43.82 (5)	C6—C4—C5	127.02 (18)
O2—Cu1—Cu1 <sup>iii</sup>	100.07 (4)	C3 <sup>ii</sup> —C4—C5	129.97 (17)
O4—Cu1—Cu1 <sup>iii</sup>	139.12 (6)	C4—C5—H5A	109.5
N1—Cu1—Cu1 <sup>iii</sup>	73.39 (5)	C4—C5—H5B	109.5
O1—Cu1—Cu1 <sup>iii</sup>	121.81 (4)	H5A—C5—H5B	109.5
Cu1 <sup>i</sup> —Cu1—Cu1 <sup>iii</sup>	51.259 (3)	C4—C5—H5C	109.5
Cu1 <sup>ii</sup> —Cu1—Cu1 <sup>iii</sup>	51.259 (3)	H5A—C5—H5C	109.5
C1—O1—Cu1	110.22 (12)	H5B—C5—H5C	109.5
C7—O2—Cu1	116.03 (12)	N2—C6—C4	110.08 (16)
Cu1—O4—H4O	119.0	N2—C6—C7	114.16 (15)
Cu1—O4—H4P	118.0	C4—C6—C7	135.66 (17)
H4O—O4—H4P	111.1	O3—C7—O2	123.20 (17)
H5O—O5—H5P	117.6	O3—C7—C6	122.78 (17)
H6O—O6—H6P	111.8	O2—C7—C6	114.02 (15)


Symmetry codes: (i)  $y-1/4, -x+1/4, -z+1/4$ ; (ii)  $-y+1/4, x+1/4, -z+1/4$ ; (iii)  $-x, -y+1/2, z$ .

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D\text{—H}\cdots A$	$D\text{—H}$	$H\cdots A$	$D\cdots A$	$D\text{—H}\cdots A$
O4—H4O $\cdots$ O5	0.84	1.84	2.680 (3)	173.
O4—H4P $\cdots$ O3 <sup>iv</sup>	0.84	2.03	2.868 (2)	177.
O5—H5O $\cdots$ O5 <sup>v</sup>	0.88	2.22	2.808 (4)	124.
O5—H5P $\cdots$ O7	0.80	1.97	2.766 (3)	171.
O6—H6O $\cdots$ O7 <sup>vi</sup>	0.92	1.84	2.752 (2)	177.
O6—H6P $\cdots$ O1	0.90	1.99	2.863 (2)	163.
O7—H7O $\cdots$ O6 <sup>vii</sup>	0.83	1.92	2.707 (2)	157.
O7—H7P $\cdots$ O3	0.83	2.21	3.016 (2)	166.
O7—H7P $\cdots$ O2	0.83	2.33	2.951 (2)	132.
O8—H8O $\cdots$ O5	0.85	1.81	2.644 (5)	167.
O8—H8P $\cdots$ O6	0.81	2.01	2.815 (4)	174.

Symmetry codes: (iv)  $-y+3/4, x+1/4, z+1/4$ ; (v)  $-x+1, -y+1/2, z$ ; (vi)  $x, y+1/2, -z$ ; (vii)  $y-1/4, -x+3/4, z-1/4$ .

Fig. 1



## **supplementary materials**

---

**Fig. 2**

