

Changes in diabetes care indicators: findings from German National Health Interview and Examination Surveys 1997–1999 and 2008–2011

Yong Du,¹ Christin Heidemann,¹ Angelika Schaffrath Rosario,¹ Amanda Buttery,² Rebecca Paprott,¹ Hannelore Neuhauser,¹ Thea Riedel,¹ Andrea Icks,^{3,4,5} Christa Scheidt-Nave¹

To cite: Du Y, Heidemann C, Schaffrath Rosario A, *et al*. Changes in diabetes care indicators: findings from German National Health Interview and Examination Surveys 1997–1999 and 2008–2011. *BMJ Open Diabetes Research and Care* 2015;**3**:e000135. doi:10.1136/bmjdr-2015-000135

► Additional material is available. To view please visit the journal (<http://dx.doi.org/10.1136/bmjdr-2015-000135>).

Received 15 July 2015
Revised 15 September 2015
Accepted 9 October 2015



CrossMark

For numbered affiliations see end of article.

Correspondence to
Dr Christa Scheidt-Nave

ABSTRACT

Objectives: To investigate changes in type 2 diabetes care indicators over time in Germany.

Methods: Adults aged 45–79 years with type 2 diabetes were identified from two national health examination surveys conducted in 1997–1999 (GNHIES98, n=333) and in 2008–2011 (DEGS1, n=526). We examined diabetes care indicators including treatment and preventive targets (glycemic control, blood pressure (BP), total cholesterol (TC), smoking, weight reduction, sports activity), self-management and care process measures (glucose self-monitoring, holding a diabetes passport, annual foot and eye examination; statin use), and the presence of diabetes-specific complications (diabetic nephropathy, retinopathy, neuropathy, diabetic foot, amputations) and comorbid cardiovascular disease (CVD) or chronic kidney disease (CKD). We calculated proportions of persons meeting these care indicators by survey and examined unadjusted and adjusted changes between surveys.

Results: Significant improvement (GNHIES98 vs DEGS1) over time was observed for glycated hemoglobin (HbA1c) <7.0% (53 mmol/mol) (32.4% vs 65.4%), BP <130/80 mm Hg (32.0% vs 47.2%), TC <190 mg/dL (13.5% vs 41.9%), statin use (11.7% vs 35.9%), eye (51.1% vs 78.4%) and foot (48.0% vs 61.4%) examination within the past 12 months, diabetes-specific complications (29.7% vs 21.8%), and CVD (44.5% vs 37.1%). Blood glucose self-monitoring significantly increased (37.4% vs 62.8%), while holding a diabetes passport did not change. Current smoking did not change and obesity rose, although sports activity significantly increased over time. Proportions of adults achieving combination goals of HbA1c, BP, TC, and smoking cessation were low in both surveys in spite of significant improvement.

Conclusions: In Germany, the quality of diabetes care improved over time. There is much room for improvement, in particular regarding preventive goals and diabetes self-management.

INTRODUCTION

Diabetes represents a major threat to public health in many countries of the world. International health organizations have

Key messages

- Quality of care for adults with type 2 diabetes substantially improved in Germany between 1997–1999 and 2008–2011.
- Improvements in quality of care for adults with type 2 diabetes in Germany were most pronounced for process and intermediate outcome measures, such as annual eye examinations and glycemic control.
- Quality of care for adults with type 2 diabetes in Germany still falls short of evidence-based guideline recommendations, in particular with regard to combined therapeutic goal achievement, behavioral risk factor control, and patient self-management.

emphasized the necessity for national plans to improve diabetes prevention and quality of care, but continuous monitoring of quality of care indicators at the population level remains a challenge for many countries.^{1 2} Individualized glycemic control and multifactorial risk reduction are the cornerstones of high-quality diabetes care.^{3–6} Evidence-based guidelines for the management of diabetes recommend lowering glycated hemoglobin (HbA1c) levels to <7.0% (53 mmol/mol) for most people with diabetes, but outline that certain individuals may benefit from more stringent (6.0–6.5%; 42–48 mmol/mol) and less stringent (7.5–8.0%, 58–64 mmol/mol) glycemic control.^{3–6} Optimal individualized diabetes management includes controlling cardiovascular risk factors (eg, hypertension and dyslipidemia), preventive strategies such as annual eye and foot examinations, and lifestyle modifications, in particular smoking cessation, weight reduction and increasing moderate physical activity to at least 150 min/week.^{5 6}

In Germany, about six million adults are estimated to have diabetes.⁷ In an effort to

improve diabetes care, type 2 diabetes disease management programs (DMPs) were initiated nationally in 2003.⁸ These programs incorporated prevention and treatment goals and strategies recommended in international and national guidelines. Tracking quality of care indicators at the population level is essential to help understand successes and failures in prevention and treatment goals for diabetes care and to direct quality improvement initiatives and health policy. To date, no studies have comprehensively investigated the quality of diabetes care over time at the national level. Previous studies of changes over time have been limited to regional population-based surveys and practice-based studies or analyses of data from different statutory sickness funds.^{9–12}

In this study, we examine changes in quality of care indicators for adults in Germany with type 2 diabetes using nationally representative data collected in 1997–1999 and 2008–2011. Diabetes care indicators included therapeutic and preventive goals such as glycemic, blood pressure (BP) and lipid control, diabetes self-management and care process measures, and diabetes comorbidities and complications.

RESEARCH DESIGN AND METHODS

Study design and participants

We analyzed data from two national health interview and examination surveys for adults in Germany conducted in 1997–1999 and 2008–2011 by the Robert Koch Institute, the national public health institute for disease control and prevention in Germany. In both surveys, a two-stage probability cluster sampling procedure was applied to ensure that the survey sample was representative of the resident population in Germany aged 18–79 years. Details of the study design, sampling strategy, and protocol have been previously published.^{13 14} Briefly, in the German National Health Interview and Examination Survey in 1997–1999 (GNHIES98), a sample of 120 German municipalities, representative of municipality sizes and structures in Germany, were selected. In the second stage, age-stratified and sex-stratified random samples of adults aged 18–79 years were drawn from local population registries. The final GNHIES98 sample included 7124 adults (response rate was 61.4%) with complete interview and examination data.¹³ The National Health Interview and Examination Survey for Adults in Germany 2008–2011 (DEGS1) used the same sampling methods as GNHIES98 but included 60 additional municipalities. The DEGS1 sample 18–79 years of age comprised 7987 persons, including 4192 persons who were newly recruited in DEGS1 and 3795 persons who had already participated in GNHIES98. The response rate was 42% for first-time survey participants and 64% for GNHIES98 re-participants.^{14 15} Among DEGS1 study participants 18–79 years, a total of 7115 persons attended one of the study centers and completed both the interview and examination.^{14 15}

The present analysis is confined to participants with known diabetes. We selected those who: (1) answered

‘Yes’ to the question “Have you ever been diagnosed with diabetes by a doctor?” or (2) used antidiabetic medications including insulin and/or oral agents within the past 7 days. Overall, we identified 374 adults with known diabetes from the 7124 participants in GNHIES98 and 591 adults with known diabetes from the 7115 participants with complete interview and examination data in DEGS1. No question on the type of diabetes was included in the surveys.⁷ In order to focus on type 2 diabetes, we confined this analysis to persons with known diabetes 45–79 years of age, excluding 101 (n=38 in GNHIES98; n=63 in DEGS1) persons with diabetes younger than 45 years at the time of the survey. Furthermore, we excluded another five persons with known diabetes (n=3 in GNHIES98; n=2 in DEGS1) who were likely to have type 1 diabetes as they fulfilled a combination of the following criteria: age at diagnosis of diabetes <30 years, insulin prescribed at initial diagnosis and currently using insulin.⁷ The final study population comprised 333 adults in GNHIES98 and 526 adults in DEGS1 aged 45–79 years with known diabetes who are described as having type 2 diabetes (table 1).

Both surveys conform to the principles outlined in the Declaration of Helsinki and the German Federal Data Protection Act. Study protocols were approved by the Federal and State Commissioners for Data Protection and Freedom of Information. DEGS1 was approved by the Charité Universitätsmedizin Berlin ethics committee in September 2008 (number EA2/047/08). All participants provided written informed consent prior to the interview and examination.

Data collection and definition of study characteristics

Data collection methods in DEGS1 were largely comparable to those used in GNHIES98 and have been previously described in detail.¹⁴ Standardized self-administered questionnaires, physician-administered computer-assisted personal interviews (CAPI), and physiological measurements and tests were used to collect information on sociodemographic characteristics (age, sex, region of residence, educational attainment), health-related behaviors (current tobacco use and sports activities in the past 3 months), self-reported physician-diagnosed health conditions, and anthropometric (body weight and height) and biochemical measures.^{13 14}

Individuals with diabetes were asked about their age at first diagnosis, current treatment and treatment at the time of diagnosis, diabetes-specific complications (including diabetic nephropathy, diabetic retinopathy, diabetic neuropathy, diabetic foot and diabetes-related amputation), eye and foot examinations in the past 12 months, self-monitoring of blood glucose, holding a diabetes passport (a national patient-held record of their care), and whether they had ever participated in a diabetes education program. We defined treatment categories as no treatment at all, dietary treatment only, and pharmaceutical treatment (oral antidiabetic agents and/or insulin). Diabetes duration was computed by subtracting the age at diagnosis

from the person's current age and grouped as '<5', '5–14' and '≥15' years.

Participants were asked to bring the original containers of all medication used during the past 7 days to the examination site. This permitted automated recording of unique product identifiers and drug coding according to the WHO "Anatomical Therapeutic Chemical" (ATC) classification system.¹⁶ A medication with an ATC code A10A and A10B was considered as 'insulin' and 'oral agent', respectively.

Body mass index (BMI) was calculated as the ratio of body weight (kg) and body height (m) squared with participants wearing light clothing (GNHIES98) or underwear (DEGS1) and shoes removed.¹⁷ Three standardized BP measurements were taken at 3 minute intervals in upright sitting with a standard mercury sphygmomanometer in GNHIES98 and an oscillometric device (Datascopie Accutorr Plus) in DEGS1. One of three cuff sizes was selected according to the right mid-arm circumference. GNHIES98 measurements were calibrated for comparison with DEGS1 measurements in order to account for device and cuff differences.¹⁸ Random venous blood samples were obtained between 8:00 and 20:00 with participants in the seated position. Full details of specimen handling and laboratory analyses for HbA1c, total cholesterol (TC), and creatinine are provided in the online supplementary appendix.

Educational attainment was classified using the international Comparative Analysis of Social Mobility in Industrial Nations (CASMIN) scale and grouped as 'primary', 'middle', and 'high'.¹⁹ Municipality size was classified as rural town (<5000 inhabitants) and small (5000–<20 000), medium-sized (20 000–<100 000), and large (100 000 inhabitants or more) cities, based on population density according to an established German classification system.¹⁴ Regions of residence were grouped into five commonly described geographical areas: north-west (federal states: Bremen, Hamburg, Niedersachsen, and Schleswig-Holstein); northeast (Berlin, Brandenburg, and Mecklenburg-Vorpommern); central west (Hessen, Nordrhein-Westfalen, Rheinland-Pfalz, and Saarland); central east (Sachsen, Sachsen-Anhalt, and Thüringen); south (Baden-Württemberg and Bayern). Smoking status was categorized as never-smoking, former smoking, and current smoking. The World Health Organization (WHO)-recommended criteria were applied to define normal weight (BMI <25.00 kg/m²), pre-obese (BMI 25.00–29.99 kg/m²), obese class I (BMI ≥30.00–34.99 kg/m²), and obese classes II and III (BMI ≥35).²⁰ Sports activity was assessed by five categories (no sports, <1 hour/week, regularly 1–2 hours/week, regularly 2–4 hours/week, regularly >4 hours/week). This information was aggregated into three categories (no sports, <2 hours/week, ≥2 hours/week). Since the number of people with diabetes engaging in ≥2 hours/week of sports activity in 1998 was small (n=20), we categorized this variable as no sports versus any sports activity for the analysis.

Diabetes care indicators

Indicators were based on clinical guidelines for recommended care^{3–6} including achievement of treatment and preventive goals (control of HbA1c, BP, and TC), patient self-management (glucose self-monitoring, holding a diabetes passport), and care process measures (eye and foot examination within the past 12 months, statin use, angiotensin converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB) use). We also assessed a number of additional preventive targets, including the proportion of persons with type 2 diabetes who were: (1) non-obese (BMI <30 kg/m²); (2) not currently smoking; and (3) engaging in any sports activities.

We considered stringent and less stringent cut-offs for glycemic control ranging from HbA1c <6.5% (48 mmol/mol) to <8.0% (64 mmol/mol) as well as for BP (<130/80 and <140/90 mm Hg) and TC (<190 and <240 mg/dL). Individualized glycemic control takes into consideration the persons' age and presence of complications.⁴ Consistent with previous studies,²¹ we defined the individualized HbA1c target as <8.0% (64 mmol/mol) for adults with diabetes-specific complications or comorbid cardiovascular diseases (CVD), <7.0% (53 mmol/mol) for adults 45–64 years, and <7.5% (58 mmol/mol) for adults 65–79 years without these problems. Furthermore, we defined combined goals of diabetes care as previously described,²¹ with the exception of using total instead of low-density lipoprotein (LDL) cholesterol: goal 1: HbA1c <7.0% (53 mmol/mol), BP <130/80 mm Hg, TC <190 mg/dL, and not currently smoking; goal 2: HbA1c <7.5% (58 mmol/mol), BP <140/90 mm Hg, TC <240 mg/dL, and not currently smoking.

The study population was characterized according to the presence of any self-reported diabetes-specific complication (yes/no) and diabetes-related comorbidities, including self-reported CVD and chronic kidney disease (CKD). Adults reporting a history of physician-diagnosed heart failure, stroke, myocardial infarction, and other coronary heart disease were classified as having CVD. For those without CVD, we calculated the 10-year UK Prospective Diabetes Study (UKPDS) CVD risk in per cent using the original version which considers patients' age, sex, HbA1c, BP, TC, smoking status, and other risk factors.²² CKD was defined according to the International Society of Nephrology²³ based on an estimated glomerular filtration rate (eGFR) <60 mL/min/1.73 m² or the presence of moderate to severe albuminuria. eGFR values were calculated by the CKD-EPI formula²⁴ using measured serum creatinine levels. In German health examination surveys, microalbuminuria was assessed semi-quantitatively using Micral-Test albumin dipstick testing and a cut-off of >50 mg/L was used to define moderate to severe microalbuminuria based on previous evidence on test performance.²⁵

Statistical analysis

SAS V.9.4 survey procedures for complex samples (SAS Institute Inc, Cary, North Carolina, USA) were used for

statistical analyses. Comparisons between the two surveys are based on weighted estimates standardized to the population of 31 December 2010. Survey weights were computed as previously described^{14 15} to account for deviations between the sample and the structure of the general population regarding age, sex, region, nationality, and education.^{7 14 15} In DEGS1, weighting factors additionally included the probability of re-participation in DEGS1 as derived from logistic regression models.¹⁵

Descriptive statistics were used to examine characteristics of adults with type 2 diabetes and diabetes care indicators. Point estimates (means and proportions) and 95% CIs were reported. Rao-Scott corrected Pearson tests for categorical variables and general linear models for continuous variables were used. For each diabetes care indicator, we calculated the absolute changes in proportions (and 95% CI) between surveys, both unadjusted and adjusted for sex, age group, region, community size, educational level, and duration of diabetes. The adjusted changes in proportions were derived from the predictive margins calculated from a logistic regression model.²⁶ First-order interactions between the survey year variable and each covariates were tested separately and interaction terms included in the model if $p < 0.1$. The predictive margins were calculated as the adjusted probabilities predicted by the model, averaged over all participants in the model and assuming that the covariate distribution in GNHIES98 and DEGS1 were identical. The SEs and correlation of the predictive margins were approximated by the SAS LSMEANS statement and used in the calculation of the 95% CI for the adjusted change in proportions. Kernel density estimation was applied to compare weighted HbA1c distributions among persons with diabetes between surveys.

For questions related to medical history, eye and foot examinations, and diabetes complications, an additional 'don't know' answering category was included in DEGS1, whereas in GNHIES98 only 'yes' and 'no' options were available. We therefore treated persons answering 'don't know' in DEGS1 as missing in analyses, but also conducted sensitivity analyses coding these answers as 'no' to ensure consistency of results. Since the results were similar, only results based on coding 'don't know' answers as missing are presented. For each variable, weighted proportion and unweighted n depict the number of participants with complete information. The number of persons with missing values was explicitly stated for each variable. Persons with missing values were excluded from the analyses, with pairwise deletion for descriptive and listwise deletion for multivariable analyses. Statistical significance was defined at $p < 0.05$ based on two-sided tests.

RESULTS

Persons with type 2 diabetes in DEGS1 had higher educational attainment, more frequently engaged in sports activities, were more often ex-smokers, and had higher BMI levels than those in GNHIES98 (table 1). Between

the 1997–1999 and 2008–2011 surveys, mean age at diagnosis declined from 56.3 to 54.6 years ($p = 0.07$) and mean years of diabetes duration increased significantly from 9.2 to 11.3 years ($p = 0.012$). In sex-specific analyses, these differences were statistically significant among women (mean age at diagnosis: 57.3 vs 54.1 years, $p = 0.041$; diabetes duration: 9.8 vs 12.4 years, $p = 0.045$), but not among men (mean age at diagnosis: 55.4 vs 55.0 years, $p = 0.757$; diabetes duration: 8.6 vs 10.3 years, $p = 0.091$).

Figure 1 shows the HbA1c distribution among adults with type 2 diabetes in the two surveys. The HbA1c distribution in DEGS1 has shifted to the left and shows considerably lower variability compared to GNHIES98.

Significantly more persons with type 2 diabetes achieved guideline-recommended stringent and less stringent targets for HbA1c, BP, and TC as well as the individualized HbA1c target and the combined treatment goals in 2008–2011 compared to the 1997–1999 survey period (table 2). The proportion of persons with diabetes reporting any sports activity significantly increased between the two surveys, while the proportion of current non-smokers remained unchanged and the proportion of non-obese persons decreased over time. Improvements in care processes over time were reflected by increasing ACE inhibitor/ARB and statin use, and increasing proportions of persons reporting eye or foot examinations within the past 12 months. Proportions of adults with type 2 diabetes reporting glucose self-monitoring increased over time, but holding a diabetes passport did not significantly differ between the two surveys. A significant decrease between the surveys was found in the proportion of adults reporting diabetes-specific complications or comorbid CVD, but not in the proportion of adults with comorbid CKD (table 2). Among adults without a history of CVD, the 10-year UKPDS risk score declined significantly over time.

DISCUSSION

Community-dwelling adults aged 45–79 years with type 2 diabetes in Germany showed consistent improvements in diabetes care between the survey periods 1997–1999 and 2008–2011. Improvement was seen for treatment targets (HbA1c, BP, serum lipids), statin and ACE inhibitor/ARB use. Self-monitoring of blood glucose levels and uptake of annual eye or foot examinations also improved. Furthermore, the proportion of adults with type 2 diabetes reporting any diabetes-specific complications and comorbid CVD decreased significantly. However, there was no significant decrease in the proportions of adults with diabetes having evidence of CKD. The proportion of people with diabetes and obesity rose over this period and the proportion of those currently smoking was unchanged. Although overall engagement in sports activity significantly improved over time, less than one in five adults in 2008–2011 reported performing more than 2 hours of sports activity per week; this is

Table 1 Characteristics of adults aged 45–79 years with type 2 diabetes in German national health interview and examination surveys 1997–1999 (GNHIES98) and 2008–2011 (DEGS1)

	GNHIES98 (1997–1999) (N=333)			DEGS1 (2008–2011) (N=526)			p Value
	n	Per cent *	95% CI*	n	Per cent*	95% CI*	
<i>Categorical variables (in %)</i>							
Sex							
Men	160	48.7	42.8 to 54.6	298	53.1	47.3 to 58.8	0.221
Women	173	51.3	45.4 to 57.2	228	46.9	41.2 to 52.7	
Age group, years							
45–64	171	43.2	36.8 to 49.8	184	37.6	32.1 to 43.4	0.229
65–79	162	56.8	50.2 to 63.2	342	62.4	56.6 to 67.9	
Municipality size							
Rural town	84	19.3	12.3 to 28.9	111	18.1	11.8 to 26.8	0.628
Small city	75	20.8	13.1 to 31.3	128	26.1	18.6 to 35.2	
Medium-sized city	89	32.4	22.6 to 43.9	150	30.0	22.2 to 39.3	
Large city	85	27.6	18.7 to 38.6	137	25.8	18.4 to 34.8	
Region†							
North-west	27	11.6	6.4 to 20.0	62	15.4	9.7 to 23.5	0.090
Central west	81	30.2	20.9 to 41.4	161	37.9	29.0 to 47.7	
North-east	57	14.5	7.8 to 25.2	73	10.3	6.3 to 16.3	
Central east	99	15.3	9.9 to 22.9	133	14.4	9.4 to 21.4	
South	69	28.5	19.1 to 40.3	97	22.1	15.1 to 31.1	
Educational level‡							
Primary	234	78.6	72.3 to 83.8	280	62.3	56.4 to 67.8	<0.001
Middle	58	14.1	10.4 to 18.7	162	28.2	23.3 to 33.7	
High	25	7.3	4.1 to 12.6	81	9.5	7.0 to 12.8	
Body mass index, kg/m²							
<25	44	12.9	9.1 to 18.0	55	11.7	9.0 to 14.9	0.001
25–29.99	149	48.3	42.6 to 54.0	184	41.1	36.9 to 45.4	
30–34.99	93	25.9	20.4 to 32.2	176	30.0	26.0 to 34.4	
≥35	42	12.9	9.4 to 17.4	107	17.2	14.6 to 20.2	
Smoking status							
Smoker	60	19.5	14.5 to 25.8	85	18.2	14.3 to 22.9	0.033
Ex-smoker	89	27.5	22.2 to 33.6	220	38.0	33.1 to 43.2	
Non-smoker	170	52.9	46.3 to 59.5	217	43.8	38.4 to 49.3	
Sports activity, hours/week							
0 (no sports)	231	76.1	69.0 to 82.0	231	47.7	42.0 to 53.5	<0.001
0–2	66	17.3	12.0 to 24.4	171	34.3	29.4 to 39.5	
≥2	20	6.6	4.0 to 10.7	91	18.0	13.9 to 23.1	
Diabetes treatment pattern							
No treatment at all	42	13.6	9.7 to 18.8	75	17.3	12.7 to 23.1	0.068
Diet only	51	15.6	11.9 to 20.2	54	9.3	6.8 to 12.8	
Insulin only (with or without diet)	33	10.8	7.1 to 16.0	56	11.6	8.5 to 15.7	
Oral agents only (with or without diet)	166	51.9	45.1 to 58.7	262	48.2	42.6 to 53.8	
Both insulin and oral agents (with or without diet)	30	8.0	5.3 to 12.1	67	13.6	10.0 to 18.2	
Diabetes duration, years							
<5	125	37.3	30.4 to 44.7	160	29.7	25.1 to 34.8	0.174
5–14	126	36.5	30.5 to 43.1	210	41.9	36.2 to 47.8	
≥15	77	26.2	20.9 to 32.3	140	28.4	23.5 to 33.8	
Continuous variables (mean)							
Age, years	333	65.5	64.4 to 66.6	526	66.0	65.0 to 67.0	0.549
Age at diagnosis of diabetes, years	328	56.3	54.9 to 57.7	510	54.6	53.1 to 56.0	0.070
Diabetes duration, years	328	9.2	8.1 to 10.3	510	11.3	10.0 to 12.6	0.012

Denominators vary due to the number of persons with missing values. Missing values (GNHIES98, DEGS1): educational level (n=16, n=3), body mass index (n=5, n=4), smoking status (n=14, n=4), sports (n=16, n=33), diabetes treatment pattern (n=11, n=12), age at the initial diagnosis of diabetes and diabetes duration (n=5, n=16).

*Weighted and standardized to the population of 31 December 2010.

†Region: Northwest (federal states: Bremen, Hamburg, Niedersachsen, and Schleswig-Holstein); northeast (Berlin, Brandenburg, and Mecklenburg-Vorpommern); central west (Hessen, Nordrhein-Westfalen, Rheinland-Pfalz, and Saarland); central east (Sachsen, Sachsen-Anhalt, and Thüringen); south (Baden-Württemberg and Bayern).


‡Educational attainment was classified using the international Comparative Analysis of Social Mobility in Industrial Nations (CASMIN) scale.

well below the guideline recommended levels of physical activity (>150 min/week) for adults with diabetes.^{3 6}

There may be several reasons for the observed improvements in some, albeit not all, diabetes care

indicators in Germany. First, there was the introduction of a national type 2 DMP in 2003, between these national health examination surveys 1997–1999 and 2008–2011.⁸ DMP enrollment comprises regular check-up visits and

Figure 1 Density distribution of glycated hemoglobin (HbA1c) among adults aged 45–79 years with type 2 diabetes. German national health interview and examination surveys 1997–1999 (GNHIES98) and 2008–2011 (DEGS1).



aims to promote diabetes education, adherence to treatment goals, and self-management.^{8 27 28} Physicians are contracted under the provision of structural quality requirements and reimbursement is provided with a focus on intermediate outcome measures, such as HbA1c measures. This key national policy change and the introduction of financial incentives for physicians to improve diabetes care are likely to have contributed to the improvements in glycemic control and statin use as observed in the present study. However, we found mixed results regarding diabetes self-management with improvements in glucose self-monitoring but not keeping a diabetes passport. Second, evidence-based national diabetes management and treatment guidelines have been periodically updated between the survey periods and are integral to DMP contracts.⁶ Third, guideline recommendations regarding the diagnostic criteria for type 2 diabetes based on fasting glucose changed in 1998, that is, toward the end of GNHIES98.²⁹ Together, these changes in between survey periods might have contributed to an earlier diagnosis of type 2 diabetes. We found that the age at diagnosis of diabetes was on average 1.7 years earlier in DEGS1 than in GNHIES98. Survey participants detected at an earlier phase of the disease course may have more favorable intermediate outcome measures (HbA1c, BP, cholesterol), and are less likely to have diabetes-specific complications compared with those at a more advanced phase of disease. In sex-specific analyses, a significant difference regarding an earlier age at diagnosis in DEGS1 vs GNHIES98 was confined to women, which may reflect sex differences in care (eg, diagnosis of gestational diabetes) and/or healthcare services utilization.

In Germany, previous studies of changes in diabetes quality of care indicators over time have been confined to regional population-based studies¹¹ or studies among primary care patients.³⁰ In repeated population surveys conducted in southern Germany, recommended eye or

foot examinations among adults with type 2 diabetes significantly increased between 1999–2001 and 2006–2008 (60.5% vs 71.3% and 37.5% vs 55.1% respectively), as did the use of lipid-lowering drugs (18.0% vs 37.9%), and the percentage of adults with diabetes achieving BP targets of <140/90 mm Hg (43.6% vs 70.5%).¹¹ These findings are largely consistent with our results. However, unlike our findings, the proportions of persons with diabetes achieving HbA1c targets of <7.0% (53 mmol/mol) and those with a BMI ≤ 30 kg/m² remained unchanged in the previous study.¹¹ In a study based on data from 110 primary care practices across Germany, the proportion of patients with diabetes achieving HbA1c <6.5% (48 mmol/mol) increased from 31% in 1998 to 36% in 2005, while the percentage of patients with HbA1c $\geq 9.0\%$ (75 mmol/mol) was halved from about 20% to 10% over this time period,³⁰ roughly comparable to our findings from 23.5% in 1997–1999 to 4.6% in 2008–2011 (table 2). Our findings of improvements in diabetes-related complications among our sample are consistent with a variety of studies using various data sources over the past decade in Germany (eg, regional disease registries or hospital discharge data, selected sickness funds) reporting improvements in amputation rates, the incidence of blindness, and cardiovascular risk reduction, particularly among women.^{31–33}

Time trend analyses of diabetes care indicators based on repeated national health surveys have been conducted in the USA. Results from these studies have consistently demonstrated significant improvements in the control of HbA1c, BP, and lipids among adults with diabetes, although absolute changes over the past decade were generally smaller than those observed in the present study.^{21 34–37} Between the 1999–2002 and 2007–2010 NHANES survey waves, the proportion of adults with diabetes achieving HbA1c <7.0% (53 mmol/mol), BP <130/80 mm Hg, and LDL-cholesterol <100 mg/dL increased significantly by 7.9, 11.7, and

Table 2 Absolute change (95% CI) in the prevalence of diabetes care indicators among adults with type 2 diabetes over time, German national health interview and examination surveys 1997–1999 and 2008–2011

	GNHIES98 (1997–1999) (N=333)			DEGS1 (2008–2011) (N=526)			Absolute change (DEGS1-GNHIES98)			
	n	Per cent†	95% CI†	n	Per cent†	95% CI†	Unadjusted		Adjusted*	
							Per cent†	95% CI†	Per cent†	95% CI†
<i>Preventive and therapeutic goals</i>										
<i>HbA1c</i>										
<6.5% (48 mmol/mol)	78	22.9	17.8 to 28.9	256	48.8	42.7 to 55.0	25.9	17.4 to 34.4	27.5	18.8 to 36.3
<7.0% (53 mmol/mol)	105	32.4	26.8 to 38.7	346	65.4	59.6 to 70.7	32.9	24.9 to 41.0	34.8	26.1 to 43.5
<7.5% (58 mmol/mol)	148	45.2	38.8 to 51.7	406	79.1	74.6 to 83.0	33.9	26.1 to 41.8	36.2	26.5 to 45.9
<8.0% (64 mmol/mol)	179	55.3	49.2 to 61.2	451	86.9	82.8 to 90.1	31.6	24.6 to 38.5	35.7	27.0 to 44.4
≥9.0% (75 mmol/mol)	72	23.5	18.0 to 30.1	23	4.6	2.8 to 7.6	-18.9	-25.4 to -12.3	-23.3	-30.3 to -16.3
Individualized HbA1c target	156	49.3	43.2 to 55.5	379	80.7	75.5 to 85.0	31.3	23.2 to 39.5	33.9	25.0 to 42.8
<i>Total cholesterol (mg/dL)</i>										
<190	37	13.5	9.0 to 19.9	224	41.9	36.6 to 47.3	28.3	20.0 to 36.7	25.7	16.2 to 35.1
<240	154	52.0	44.9 to 59.0	442	85.3	80.6 to 89.0	33.3	25.0 to 41.6	32.2	23.9 to 40.6
<i>Blood pressure (mm Hg)</i>										
<130/80	103	32.0	26.2 to 38.4	240	47.2	41.2 to 53.3	15.2	6.4 to 24.0	20.3	11.5 to 29.1
<140/90	181	56.3	49.5 to 62.9	364	69.6	63.8 to 74.8	13.3	4.6 to 21.9	18.2	8.9 to 27.4
Body mass index <30 kg/m ²	193	61.2	55.4 to 66.7	239	45.6	40.0 to 51.4	-15.6	-23.5 to -7.7	-16.6	-26.0 to -7.3
Currently not smoking	259	80.5	74.2 to 85.5	437	81.8	77.1 to 85.7	1.4	-6.0 to 8.7	-1.4	-8.1 to 5.2
Engaging in any sports activity	86	23.9	18.0 to 31.0	262	52.3	46.5 to 58.0	28.4	19.1 to 37.7	28.5	18.0 to 39.0
Combination goal 1	5	1.7	0.5 to 5.1	60	11.4	8.4 to 15.4	9.8	5.8 to 13.7	9.5	6.0 to 13.0
Combination goal 2	31	10.9	7.2 to 16.0	201	39.6	34.0 to 45.4	28.7	21.6 to 35.8	31.1	24.3 to 37.8
<i>Patient self-management</i>										
Self-monitoring of blood glucose	119	37.4	30.8 to 44.6	318	62.8	57.3 to 68.0	25.3	16.8 to 33.9	24.0	15.6 to 32.5
Holding a diabetes passport	172	46.9	39.8 to 54.2	287	54.2	48.4 to 60.0	7.3	-1.2 to 15.9	2.4	-7.1 to 11.8
<i>Care processes</i>										
Last eye examination ≤12 months	170	51.1	44.8 to 57.3	389	78.4	73.6 to 82.6	27.3	19.3 to 35.4	22.2	12.8 to 31.5
Last foot examination ≤12 months	151	48.0	41.2 to 54.9	303	61.4	55.3 to 67.1	13.4	4.0 to 22.8	10.0	0.6 to 19.3
Any ACE inhibitor or ARB	119	34.7	28.6 to 41.3	335	60.7	54.6 to 66.4	26.0	17.2 to 34.7	22.7	12.7 to 32.6
Any lipid-lowering medication	62	18.2	13.8 to 23.7	216	39.6	33.8 to 45.7	21.4	13.7 to 29.0	18.8	10.6 to 27.0
Statin use	40	11.7	8.4 to 16.1	195	35.9	30.3 to 41.9	24.2	17.6 to 30.9	22.5	15.3 to 29.7
<i>Complications/comorbidities/CVD risk</i>										
Any diabetes-specific complication	93	29.7	23.9 to 36.2	107	21.8	17.9 to 26.3	-7.9	-15.6 to -0.2	-9.7	-17.0 to -2.0
Comorbid CVD	138	44.5	38.0 to 51.3	184	37.1	31.3 to 43.3	-7.4	-16.7 to 1.8	-10.3	-21.0 to 0.6
Comorbid chronic kidney disease	120	44.8	37.9 to 51.8	226	40.3	34.8 to 46.0	-4.5	-13.1 to 4.2	-9.1	-20.0 to 1.3
UKPDS risk (in %) for persons without CVD	172	21.0	18.9 to 23.2	303	15.9	14.5 to 17.3	-5.2	-7.7 to -2.7	-6.1	-8.2 to -4.1

Results for absolute changes in bold print denote statistically significant differences based on p values derived from descriptive analyses (unadjusted change) or logistic regression models (adjusted change).

Individualized HbA1c target: <8.0% (64 mmol/mol) for adults with diabetes-specific complications or comorbid CVD, <7.0% (53 mmol/mol) for adults 45–64 years, and <7.5% (58 mmol/mol) for adults 65–79 years without these problems. Combination goal 1: HbA1c <7.0% (53 mmol/mol) and blood pressure <130/80 mm Hg and total cholesterol <190 mg/dL and not currently smoking; Combination goal 2: HbA1c <7.5% (58 mmol/mol) and blood pressure <140/90 mm Hg and total cholesterol <240 mg/dL and not currently smoking.

Denominators vary due to the number of persons with missing values. Missing values (GNHIES98, DEGS1): HbA1c (n=26, n=8), individualized HbA1c target (n=28, n=63), total cholesterol (n=24, n=6), blood pressure (n=1, n=2), combination goal 1 and goal 2 (n=41, n=15), self-monitoring blood glucose (n=5, n=12), holding a diabetes passport (n=5, n=12), eye examination (n=5, n=24), foot examination (n=5, n=34), any lipid-lowering medication and statin use (n=0, n=1), diabetes-specific complication (n=5, n=72), comorbid CVD (n=0, n=25), comorbid chronic kidney disease (n=56, n=7), UKPDS risk (n=23, n=14).

*Adjusted for sex, age group (45–64, 65–79 years), region (northwest, central west, northeast, central east, south), community size (rural town, small, middle-sized, and large cities), educational level (primary, middle, high), duration of diabetes (<5, 5–14, ≥15 years) and significant first-order interactions (p<0.100) as shown in the online supplementary table 1A.

†Weighted and standardized to the population of 31 December 2010.

ARB, angiotensin receptor blocker; CVD, cardiovascular diseases; HbA1c, glycated hemoglobin; UKPDS, UK Prospective Diabetes Study.

20.8 percentage points, respectively,²¹ as compared to 32.9, 15.2, and 28.3 percentage points for HbA1c <7.0% (53 mmol/mol), BP <130/80 mm Hg, and TC <190 mg/dL in the present analysis. Unfortunately, some indicators are not directly comparable between NHANES and the German national health surveys. Unlike NHANES, the present study used TC instead of LDL-cholesterol, as the German surveys did not recruit a random subsample of adults who observed overnight fasting for at least 8 h.^{13 14} Alongside the significantly improved profiles of HbA1c, BP measures, and cholesterol, use of antidiabetic medication, antihypertensive agents, and lipid-lowering drugs significantly increased over time as observed in our study and previously in US population studies.^{34–37} We found that 8.0% of adults with diabetes in GNHIES98 and 13.6% in DEGS1 used a combination of insulin and oral agents, similar to 9.8% and 13.9% of adults with diabetes in the 1999–2004 and 2005–2010 NHANES waves.³⁴ Over the same time period, lipid-lowering drug use increased from 18.2% to 39.6% among adults with diabetes in Germany as shown in this analysis, similar to increases in NHANES participants from 19.5% in 1999–2000 to 42.2% in 2007–2008.³⁸ Consistent with our findings, analyses of data from the US Behavioral Risk Factor Surveillance System (BRFSS) survey waves 2000 and 2008 showed significant improvements with respect to the proportion of adults with diabetes reporting annual foot examinations and glucose self-monitoring.²¹ Unlike our findings, self-reported eye examination among adults with diabetes in BRFSS showed no further improvement over time,²¹ but baseline coverage for this indicator already achieved a much higher level in the USA in the 2000 survey with 75.1% compared to 51.1% in GNHIES98. Data from official health statistics (National Hospital Discharge Survey, US Renal Data System, National Vital Statistics System) have been systematically used to analyze trends in rates of diabetes-related complications in the USA and provide clear evidence for a decline in myocardial infarction, and death from hyperglycemic crisis, stroke, and amputation among people with diabetes between 1990 to 2010.³⁹ This remains a major goal for diabetes surveillance in Germany, where evidence on long-term outcomes is limited so far.⁸

Despite significant improvements in diabetes care observed over the past decade, the current level of care in Germany as in other countries such as the USA,^{21 34–37} Canada,⁴⁰ and Spain⁴¹ falls short of guideline recommendations. Similar to these studies,^{21 40 41} the results of the present analysis based on data from German national health surveys show that about two-thirds of persons with type 2 diabetes reached the target of HbA1c <7.0% (53 mmol/mol), and only 11.4% of patients with diabetes achieved stringent targets in HbA1c, TC, BP, and smoking combined. There was little change in the prevalence of current smokers among adults with diabetes in our study similar to US diabetes population studies.^{21 37} Rising obesity among diabetes populations is a concern

for numerous countries³⁷ and highlights population challenges of guideline-recommended weight reduction. Weight gain among persons with diabetes has been observed in relation to treatment with insulin and certain oral antidiabetic agents.^{42 43} Further insight from longitudinal studies is needed to assess the effect of weight gain on long-term cardiovascular risk and mortality.^{42 43}

The major strength of this analysis is that it provides comprehensive data on changes in diabetes quality of care indicators over time including the time period before (1997–1999) and after (2008–2011) the introduction of DMPs and national evidence-based guideline implementation for type 2 diabetes in Germany. There are several limitations to this study. First, we cannot exclude selection bias, as persons who are severely ill, hospitalized, or institutionalized were not included. Therefore, our results may depict an overly optimistic picture of diabetes quality of care both cross-sectionally and over time. Second, diabetes diagnosis was self-reported and not verified by medical records as occurs in studies using patient registers. However, self-reported physician-diagnosed diabetes provides a valid and internationally established indicator.^{7 34 44} which permits comparisons between countries as well as over time. Third, information on gestational diabetes was collected in DEGS1⁷ but not in GNHIES98. Inclusion of women with gestational diabetes may have reduced the survey-specific rates of persons with diabetes who received pharmacological treatment or preventive eye and foot examinations in this study. Fourth, although data collection methods were kept comparable between surveys as far as possible, changes to analytical methods for physiological measurements remain a challenge to population-based monitoring of BP and biomarkers.^{18 34} Finally, this analysis was limited to intermediate outcome and process indicators of diabetes care available in both surveys. In particular, process of care indicators reflecting self-management and patient education were limited to glucose self-monitoring and holding a diabetes passport in this study.

CONCLUSIONS

In conclusion, many diabetes quality of care indicators have significantly improved over time from 1997–1999 to 2008–2011 in Germany, but a substantial proportion of adults with diabetes do not achieve guideline-recommended targets, particularly around lifestyle interventions such as smoking cessation, weight reduction, and physical activity similar to other international data. Surveillance of national-level diabetes quality indicators is required to evaluate the effects of changes in international guideline updates, national diabetes healthcare policy, and changes to care management. Social inequalities and regional variations in diabetes epidemiology and diabetes quality of care need to be a central focus of national surveillance activities. Most of all, the database regarding subjective quality of care as well as long-

term outcomes and diabetes-related complications needs to be strengthened.

Author affiliations

¹Department of Epidemiology and Health Monitoring, Robert Koch Institute, Berlin, Germany

²King's College London, Faculty of Life Sciences and Medicine, London, UK

³Jean Philippe Assal Group for Health Services Research and Health Economics, German Diabetes Center, Düsseldorf, Germany

⁴Public Health Unit, Faculty of Medicine, Heinrich-Heine University, Düsseldorf, Germany

⁵German Centre for Diabetes Research (DZD), Munich, Germany

Contributors YD helped conceptualize the study, performed the statistical analyses, and drafted the manuscript. CH helped conceptualize the study, reviewed and edited the manuscript, and contributed to the discussion. ASR performed the statistical modeling and contributed to the discussion. AB reviewed and edited the manuscript and contributed to the discussion. RP, HN, and AI reviewed the manuscript and contributed to the discussion. TR was in charge of laboratory measurement quality control and reviewed the manuscript. CSN conceptualized and supervised the study and substantially contributed to the writing of the manuscript. CSN is the guarantor who takes full responsibility for the work as a whole, including the study design, access to data, and the decision to submit and publish the manuscript. All authors read and approved the final version of the manuscript.

Funding National Health Surveys are funded by the Federal Ministry of Health Germany (BMG) as part of the continuous national health monitoring. YD was supported by a research grant from the Kompetenznetz Diabetes mellitus (Competence Network Diabetes mellitus) funded by the Federal Ministry of Education and Research (FKZ 01G1110F).

Competing interests None declared.

Patient consent Obtained.

Ethics approval Charité Universitätsmedizin Berlin ethics committee.

Provenance and peer review Not commissioned; externally peer reviewed.

Data sharing statement No additional data are available specific to type 2 diabetes care among adults in Germany. For the DEGS survey system, additional data are available for research collaborations as described in the DEGS study protocol. <http://www.biomedcentral.com/1471-2458/12/730>

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

1. OECD. Cardiovascular disease and diabetes: policies for better health and quality of care, OECD health policy studies. Paris: OECD Publishing, 2015.
2. World Health Organization (WHO). Global action plan for the prevention and control of noncommunicable diseases 2013–2020. 2013. http://apps.who.int/iris/bitstream/10665/94384/1/9789241506236_eng.pdf?ua=1
3. American Diabetes Association. Standards of medical care in diabetes—2015. *Diabetes Care* 2015;38:S1–93.
4. Inzucchi S, Bergenstal R, Buse J, *et al.* Management of hyperglycemia in type 2 diabetes: a patient-centered approach: position statement of the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care* 2012;35:1364–79.
5. Rydén L, Grant PJ, Anker SD, *et al.* ESC Guidelines on diabetes, pre-diabetes, and cardiovascular diseases developed in collaboration with the EASD: the Task Force on diabetes, pre-diabetes, and cardiovascular diseases of the European Society of Cardiology (ESC) and developed in collaboration with the European Association for the Study of Diabetes (EASD). *Eur Heart J* 2013;34:3035–87.
6. The German Medical Association (Bundesärztekammer, BÄK), the National Association of Statutory Health Insurance Physicians (Kassenärztliche Bundesvereinigung, KBV), and the Association of the Scientific Medical Societies (Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften, AWMF). [German National Disease Management Guideline Type 2 Diabetes], version 4, updated November 2014. <http://www.leitlinien.de/mbd/downloads/nvl/diabetes-mellitus/dm-therapie-1auf1-vers4-lang.pdf>
7. Heidemann C, Du Y, Schubert I, *et al.* [Prevalence and temporal trend of known diabetes mellitus: results of the German Health Interview and Examination Survey for Adults (DEGS1)]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2013;56:668–77.
8. Fuchs S, Henschke C, Blümel M, *et al.* Disease management programs for type 2 diabetes in Germany: a systematic literature review evaluating effectiveness. *Dtsch Arztebl Int* 2014;111:453–63.
9. Rothe U, Müller G, Schwarz PE, *et al.* Evaluation of a diabetes management system based on practice guidelines, integrated care, and continuous quality management in a Federal State of Germany: a population-based approach to health care research. *Diabetes Care* 2008;31:863–8.
10. Berthold HK, Bestehorn KP, Jannowitz C, *et al.* Disease management programs in type 2 diabetes: quality of care. *Am J Manag Care* 2011;17:393–403.
11. Schunk M, Stark R, Reitmeir P, *et al.* [Improvements in type 2 diabetes care? Pooled analysis of survey data in southern Germany (KORA) from 1999–2008]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2011;54:1187–96.
12. Stark R, Schunk M, Meisinger C, *et al.* Medical care of type 2 diabetes in German disease management programmes: a population-based evaluation. *Diabetes Metab Res Rev* 2011;27:383–91.
13. Bellach BM, Knopf H, Thefeld W. Der Bundes-Gesundheitssurvey 1997/98 [The German Health Survey. 1997/98]. *Gesundheitswesen* 1998;60(Suppl 2):S59–68.
14. Scheidt-Nave C, Kamtsiuris P, Göbwald A, *et al.* German health interview and examination survey for adults (DEGS)—design, objectives and implementation of the first data collection wave. *BMC Public Health* 2012;12:730.
15. Kamtsiuris P, Lange M, Hoffmann R, *et al.* [The first wave of the German Health Interview and Examination Survey for Adults (DEGS1): sample design, response, weighting and representativeness]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2013;56:620–30.
16. Knopf H, Grams D. [Medication use of adults in Germany: results of the German Health Interview and Examination Survey for Adults (DEGS1)]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2013;56:868–77.
17. Mensink GB, Schienkiewitz A, Haftenberger M, *et al.* [Overweight and obesity in Germany: results of the German Health Interview and Examination Survey for Adults (DEGS1)]. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2013;56:786–94.
18. Neuhauser HK, Ellert U, Thamm M, *et al.* Calibration of blood pressure data after replacement of the standard mercury sphygmomanometer by an oscillometric device and concurrent change of cuffs. *Blood Press Monit* 2015;20:39–42.
19. Brauns H, Scherer S, Steinmann S. The CASMIN Educational Classification in International Comparative Research. In: Hoffmeyer-Zlotnik JHP, Wolf C, eds. *Advances in cross-national comparison. A European working book for demographic and socio-economic variables*. New York: Kluwer Academic, 2003, 221–44.
20. World Health Organization (WHO). Obesity: preventing and managing the global epidemic. WHO Technical Report Series 894, Geneva, 2000.
21. Ali MK, Bullard KM, Saaddine JB, *et al.* Achievement of goals in U. S. diabetes care, 1999–2010. *N Engl J Med* 2013;368:1613–24.
22. Stevens RJ, Kothari V, Adler AI, *et al.* The UKPDS risk engine: a model for the risk of coronary heart disease in Type II diabetes (UKPDS 56). *Clin Sci (Lond)* 2001;101:671–9.
23. Stevens PE, Levin A. Evaluation and management of chronic kidney disease: synopsis of the kidney disease: improving global outcomes 2012 clinical practice guideline. *Ann Intern Med* 2013;158:825–30.
24. Levey AS, Stevens LA, Schmid CH, *et al.*, CKD-EPI (Chronic Kidney Disease Epidemiology Collaboration). A new equation to estimate glomerular filtration rate. *Ann Intern Med* 2009;150:604–12.
25. Parikh CR, Fischer MJ, Estacio R, *et al.* Rapid microalbuminuria screening in type 2 diabetes mellitus: simplified approach with Micral test strips and specific gravity. *Nephrol Dial Transplant* 2004;19:1881–5.

26. Greenland S. Model-based estimation of relative risks and other epidemiologic measures in studies of common outcomes and in case-control studies. *Am J Epidemiol* 2004;160:301–5.
27. Szecsenyi J, Rosemann T, Joos S, *et al*. German diabetes disease management programs are appropriate for restructuring care according to the chronic care model: an evaluation with the patient assessment of chronic illness care instrument. *Diabetes Care* 2008;31:1150–4.
28. Stock S, Drabik A, Büscher G, *et al*. German diabetes management programs improve quality of care and curb costs. *Health Aff (Millwood)* 2010;29:2197–205.
29. Alberti KG, Zimmet PZ. Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabet Med* 1998;15:539–53.
30. Geller JC, Cassens S, Brosz M, *et al*. Achievement of guideline-defined treatment goals in primary care: the German Coronary Risk Management (CoRiMa) study. *Eur Heart J* 2007;28:3051–8.
31. Genz J, Scheer M, Trautner C, *et al*. Reduced incidence of blindness in relation to diabetes mellitus in southern Germany? *Diabet Med* 2010;27:1138–43.
32. Icks A, Dickhaus T, Hörmann A, *et al*. Differences in trends in estimated incidence of myocardial infarction in non-diabetic and diabetic people: Monitoring Trends and Determinants on Cardiovascular Diseases (MONICA)/Cooperative Health Research in the Region of Augsburg (KORA) registry. *Diabetologia* 2009;52:1836–41.
33. Icks A, Haastert B, Trautner C, *et al*. Incidence of lower-limb amputations in the diabetic compared to the non-diabetic population. Findings from nationwide insurance data, Germany, 2005–2007. *Exp Clin Endocrinol Diabetes* 2009;117:500–4.
34. Selvin E, Parrinello CM, Sacks DB, *et al*. Trends in prevalence and control of diabetes in the United States, 1988–1994 and 1999–2010. *Ann Intern Med* 2014;160:517–25.
35. Casagrande S, Fradkin JE, Saydah SH, *et al*. The prevalence of meeting A1C, blood pressure, and LDL goals among people with diabetes, 1988–2010. *Diabetes Care* 2013;36:2271–9.
36. Wong ND, Patao C, Wong K, *et al*. Trends in control of cardiovascular risk factors among US adults with type 2 diabetes from 1999 to 2010: Comparison by prevalent cardiovascular disease status. *Diab Vasc Dis Res* 2013;10:505–13.
37. Ford ES. Trends in the control of risk factors for cardiovascular disease among adults with diagnosed diabetes: findings from the National Health and Nutrition Examination Survey 1999–2008. *J Diabetes* 2011;3:337–47.
38. Kuznik A, Mardekian J. Trends in utilization of lipid- and blood pressure-lowering agents and goal attainment among the U.S. diabetic population, 1999–2008. *Cardiovasc Diabetol* 2011;10:31.
39. Gregg EW, Li Y, Wang J, *et al*. Changes in diabetes-related complications in the United States, 1990–2010. *N Eng J Med* 2014;370:1514–23.
40. Braga MF, Casanova A, Teoh H, *et al*. Poor achievement of guidelines-recommended targets in type 2 diabetes: findings from a contemporary prospective cohort study. *Int J Clin Pract* 2012;66:457–64.
41. Navarro-Vidal B, Banegas JR, León-Muñoz LM, *et al*. Achievement of cardiometabolic goals among diabetic patients in Spain. A Nationwide Population-Based Study. *PLoS ONE* 2013;8.
42. Russell-Jones D, Khan R. Insulin-associated weight gain in diabetes —causes, effects and coping strategies. *Diabetes Obes Metab* 2007;9:799–812.
43. Ross SA, Dzida G, Vora J, *et al*. Impact of weight gain on outcomes in type 2 diabetes. *Curr Med Res Opin* 2011;27:1431–8.
44. Midthjell K, Holmen J, Bjorndal A, *et al*. Is questionnaire information valid in the study of a chronic disease such as diabetes? The Nord-Trøndelag diabetes study. *J Epidemiol Community Health* 1992;46:537–42.