

Short Paper

Cite this article: Zhang SX, Graf-Vlachy L, Looi KH, Su R, Li J (2020). Social media use as a predictor of handwashing during a pandemic: evidence from COVID-19 in Malaysia.

Epidemiology and Infection **148**, e261, 1–5. <https://doi.org/10.1017/S0950268820002575>

Received: 23 June 2020

Revised: 7 September 2020

Accepted: 21 October 2020

Key words:





COVID-19; epidemic; health information; hygiene; social media use; 2019-nCoV

Author for correspondence:

Stephen X. Zhang,

E-mail: stephen.x.zhang@gmail.com

Social media use as a predictor of handwashing during a pandemic: evidence from COVID-19 in Malaysia

Stephen X. Zhang¹ , Lorenz Graf-Vlachy² , Kim Hoe Looi³ , Rui Su³ and Jizhen Li⁴ 

¹University of Adelaide, Adelaide, Australia; ²ESCP Business School, Berlin, Germany; ³Xiamen University Malaysia, Malaysia and ⁴Tsinghua University, Beijing, China

Abstract

Although handwashing is an effective way to prevent infections, there is scarce evidence on predictors of handwashing during a pandemic. This paper aims to identify behavioural and demographic predictors of handwashing. The study surveyed 674 adults in Malaysia in May 2020 regarding whether the time spent on social media predicted handwashing contingent on gender and number of children. More time spent on social media was positively associated with handwashing for males with three or more children. However, for males without children, social media use was negatively associated with handwashing. The association was not significant for males with one or two children. For females, more time spent on social media was significantly linked to more handwashing only for females with one child. Gender, a traditional predictor of handwashing, was a useful predictor only for those who spent more than three hours per day on social media and had at most one child. Number of children was a novel negative predictor for males who did not use social media and who averaged one hour per day on social media, a positive predictor for males who spent lots of time on social media, but not a predictor for females. In sum, social media use predicts handwashing, and is thus a helpful variable for use in targeted health communication during a pandemic – particularly through social media. Further, more conventional predictors like gender and number of children exhibit contingency effects with social media use.

Handwashing is one of the most basic and effective measures to prevent the spread of disease in general [1] and of upper respiratory infections in particular [2]. Handwashing is notably recommended during pandemics [3]. Unfortunately, compliance remains a constant issue even among physicians and nurses, who are professionally trained to do so [4].

Given that health communication, such as messaging and reminders, has positive effects on handwashing behaviour [5, 6], it is advisable to identify, inform and prompt at-risk populations to regularly wash hands during a pandemic. Moreover, to economise on limited resources while having optimum impact, health organisations, governments and other parties may want to focus their health messaging on less compliant groups.

This study aims to identify predictors of handwashing to help identify such less compliant groups. Specifically, we examine time spent on social media, which is a key channel for health communication during pandemics [7]. Social media has the advantage that less compliant groups, once identified, can be directly targeted with health information via the same platform. However, to our best knowledge, there exist no studies linking social media use to handwashing during a pandemic. This study identifies social media use and several demographic variables as predictors of handwashing that may be useful when deploying targeted health information campaigns on hand hygiene during pandemics.

We conducted a survey of working adults in Malaysia between 2 May and 8 May 2020, six weeks after Malaysia implemented a *cordon sanitaire* preventive measure to contain coronavirus disease-2019 (COVID-19). To collect a sample with broad coverage in Malaysia, a multi-lingual and multi-ethnic country, a questionnaire prepared in English was translated into Malay and Mandarin, the country's major languages. Links to the survey in English, Malay and Mandarin were distributed via WhatsApp, Facebook Messenger and email, via two-stage stratified sampling in terms of geographical area, ethnicity, gender and age. The survey was approved by Tsinghua University (20200322), and all survey participants consented online before enrolling in the survey. The respondents could opt out at any time and were ensured confidentiality and anonymity. No personally identifying information was collected.

The demographic variables of interest in this study comprise of gender, number of children living in the same household, age, educational level and overall health condition [8, 9]. The behavioural variables of interest are social media usage (measured in hours per day)

and frequency of handwashing after touching things outside the home (seven-point Likert scale: 1 = never (0% of the time); 2 = rarely (less than 10% of the time); 3 = occasionally (about 30% of the time); 4 = sometimes (about 50% of the time); 5 = frequently (about 70% of the time); 6 = usually (about 90% of the time); 7 = every time (100% of the time)).

We obtained 674 valid responses from adults across all Malaysian states and federal territories. 51.5% (347) respondents were female and 48.5% (327) male. Almost half of the respondents were living without children in the household (48%, 324), and progressively fewer people were living with an increasing number of children, e.g. 17.1% (115) indicated to be cohabiting with one child and only 0.6% (4) were living with more than five children.

Some 2.8% (19) of respondents indicated they never or rarely washed their hands after touching things outside the household. 45.5% (307) respondents indicated they usually or always washed their hands. 51.6% (348) indicated handwashing frequencies in between. Social media use was widespread in our sample. While 29.7% (200) of respondents reported less than two hours of social media use per day, 29.8% (201) reported between two and just below four hours, and 40.5% (273) reported four or more daily hours. Detailed descriptives for all predictors can be found in [Table 1](#).

We used Stata 16.1 to perform an ordinary least squares (OLS) linear regression on the unweighted data with a significance level of 0.05. All tests were performed two-tailed.

[Table 1](#) shows the regression results for handwashing ($F(10, 663) = 6.81, p = 0.000$). Age ($b = 0.03, 95\% \text{ CI: } 0.02 \text{ to } 0.04, p = 0.000$) and health condition ($b = 0.17, 95\% \text{ CI: } 0.06 \text{ to } 0.28, p = 0.003$) were associated with frequency of handwashing, with older and healthier people washing their hands more frequently. Number of children ($b = -0.22, 95\% \text{ CI: } -0.39 \text{ to } -0.05, p = 0.010$) and time spent on social media per day ($b = -0.09, 95\% \text{ CI: } -0.16 \text{ to } -0.02, p = 0.007$) were significantly negatively related to handwashing.

The relationship between time spent on social media and handwashing, however, was significantly moderated by gender ($b = 0.14, 95\% \text{ CI: } 0.06 \text{ to } 0.23, p = 0.001$) and number of children ($b = 0.06, 95\% \text{ CI: } 0.03 \text{ to } 0.09, p = 0.000$). Furthermore, there was a significant three-way interaction between time spent on social media, gender and number of children ($b = -0.06, 95\% \text{ CI: } -0.11 \text{ to } -0.02, p = 0.006$).

A margin analysis of the interaction results (see [Fig. 1](#)) shows that social media use predicts handwashing, and that this predictor exhibits distinct patterns depending on gender and number of children. First, more time spent on social media was positively associated with the frequency of handwashing for males with three or more children living in the same household (e.g. for those with three children: $b = 0.09, 95\% \text{ CI: } 0.02 \text{ to } 0.17, p = 0.010$). However, for males with no children in the same household, more time spent on social media was negatively associated with frequency of handwashing ($b = -0.09, 95\% \text{ CI: } -0.16 \text{ to } -0.02, p = 0.007$). The association between time spent on social media and handwashing was not significant for males with one or two children. For females in general, the association between social media use and handwashing was significant and positive ($b = 0.05, 95\% \text{ CI: } 0.01 \text{ to } 0.09, p = 0.030$). However, a margin analysis showed that more time spent on social media was significantly linked to more handwashing only for females with one child ($b = 0.05, 95\% \text{ CI: } 0.01 \text{ to } 0.09, p = 0.028$) and only marginally significant ($p < 0.10$) for females without children or with two children. The association between time spent on social

media and handwashing was not significant for women with three or more children.

Second, prior studies frequently identified gender as a predictor of handwashing, finding that women generally wash their hands more frequently than men both during pandemics and in other circumstances [10, 11]. While our results corroborate this direct effect of gender ($b = 0.29, 95\% \text{ CI: } 0.08 \text{ to } 0.50, p = 0.006$), we also found that the gender difference was more nuanced and depended on social media usage and the number of children. Specifically, our study revealed that there was no gender difference in handwashing between females and males who spent three hours or less on social media ($p > 0.10$ across all cases). The gender difference, i.e. the notion that females wash hands more frequently than males do, was significant for those who spent more than three hours per day on social media and had at most one child (e.g. four hours per day on social media and no children: $b = 0.32, 95\% \text{ CI: } 0.05 \text{ to } 0.59, p = 0.020$). The gender difference was not significant for those with two or more children. Hence, the gender difference commonly found in the literature is not universal but instead depends on both social media use and the number of children: Gender is a useful predictor of handwashing largely for people who spend lots of time on social media and have at most one child.

Third, this study unveiled the number of children in the household as an important predictor of handwashing. Number of children negatively predicted handwashing among males who did not use social media ($b = -0.22, 95\% \text{ CI: } -0.39 \text{ to } -0.05, p = 0.010$) and who averaged one hour per day on social media ($b = -0.16, 95\% \text{ CI: } -0.30 \text{ to } -0.02, p = 0.028$). In contrast, number of children positively predicted handwashing among males who spent lots of time on social media ($b = 0.21, 95\% \text{ CI: } 0.09 \text{ to } 0.34, p = 0.001$). For females in general, the number of children did not predict handwashing.

Taken together, these results suggest that research on handwashing should not rely on gender, number of children or social media use alone but must consider all three and their interaction effects simultaneously to yield better predictions. This finding is relevant as social media is an increasingly important information channel but females and males tend to vary in their interests and social circles on social media [12] and people with varying numbers of children may similarly gravitate towards different interest groups. Thus, people are exposed to social information – also regarding pandemics – differently depending on their gender and family situation. This study, as the first to identify social media use as a predictor of handwashing, suggests that it is helpful to take social media usage into account in predictive models of human behaviour during a pandemic. More specifically, we identify social media usage as a marker to screen for people who are vulnerable to COVID-19 infection due to their lack of handwashing. Such people have a higher likelihood to become agents to contract and spread virus. The identification of such people can, in turn, help to ease the pandemic or ‘flatten the curve.’ The identification of vulnerable populations via social media usage is especially promising as it directly enables targeted health communications via social media platforms to those populations.

Our results also provide evidence as to the efficacy of other predictors of handwashing during the COVID-19 pandemic. Extant work found mixed evidence on the association between age and protective behaviours such as handwashing during virus pandemics [10]. Our results suggest that there is indeed a positive relationship between age and handwashing, suggesting more targeted handwashing campaigns should focus on younger

Table 1. Descriptive findings and predictors of handwashing ($n = 674$)

Variables	Frequency (%)	Coefficient (95% CI)
Handwashing: Mean = 5.33, std. dev. = 1.40		
Every time (100% of the time) (7)	190 (28.2%)	-
Usually (about 90% of the time) (6)	117 (17.4%)	
Frequently (about 70% of the time) (5)	192 (28.5%)	
Sometimes (about 50% of the time) (4)	100 (14.8%)	
Occasionally (about 30% of the time) (3)	56 (8.3%)	
Rarely (less than 10% of the time) (2)	16 (2.4%)	
Never (0% of the time) (1)	3 (0.4%)	
Gender (categorical)		
Female	347 (51.5%)	-0.25 (-0.72 to 0.22)
Male	327 (48.5%)	(Reference category)
Age (continuous)		
20–29	103 (15.3%)	0.03*** (0.02 to 0.04)
30–39	198 (29.4%)	
40–49	193 (28.6%)	
50–59	148 (22.0%)	
>59	32 (4.7%)	
Educational level (continuous)		
Primary school completed (1)	0 (0.0%)	-0.04 (-0.23 to 0.14)
Secondary school completed (2)	49 (7.3%)	
College or university completed (3)	410 (60.8%)	
Graduate school completed (4)	215 (31.9%)	
Health condition (continuous)		
Poor (1)	7 (1.0%)	0.17** (0.06 to 0.28)
Fair (2)	74 (11.0%)	
Good (3)	212 (31.5%)	
Very good (4)	252 (37.4%)	
Excellent (5)	129 (19.1%)	
Number of children in household (continuous)		
0	324 (48.0%)	-0.22* (-0.39 to -0.05)
1	115 (17.1%)	
2	101 (15.0%)	
3	76 (11.3%)	
4	41 (6.1%)	
5	13 (1.9%)	
> 5	4 (0.6%)	
Time on social media per day (continuous)		
[0 h; 2 h]	200 (29.7%)	-0.09** (-0.16 to -0.02)
[2 h; 4 h]	201 (29.8%)	
[4 h; 6 h]	119 (17.7%)	
[6 h; 8 h]	70 (10.4%)	
> 8 h	84 (12.5%)	

(Continued)

Table 1. (Continued.)

Variables	Frequency (%)	Coefficient (95% CI)
Interactions		
Time spent on social media per day × gender	-	0.14** (0.06 to 0.23)
Time spent on social media per day × number of children in household	-	0.06*** (0.03 to 0.09)
Gender × number of children in household	-	0.20 (-0.05 to 0.46)
Time spent on social media per day × gender × Number of children in household	-	-0.06** (-0.11 to -0.02)

Note: * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$.




Fig. 1. Predicted handwashing frequency at different values of significant predictors (handwashing frequency: 4 = sometimes (about 50% of the time); 5 = frequently (about 70% of the time); 6 = usually (about 90% of the time); 7 = every time (100% of the time)).

adults in epidemics. Prior work also found that more educated people tend to exhibit greater protective behaviour during pandemics, although some results were inconclusive [10]. In our sample, there was no significant association between education and

handwashing. However, our sample was skewed towards college graduates and cell sizes for other levels of education were small, so we are cautious in interpreting this result. Our nonsignificant findings on education suggest the need for more

research to further examine the predictive effect of education on protective behaviour during epidemics. We did, however, find a significant association between health condition and handwashing. During the COVID-19 pandemic, healthier people indicated greater handwashing frequency – a finding that is new to the literature. Although people with comorbidities suffer more from COVID-19 infections, they behaviorally protect themselves less from the virus due to their less frequent handwashing. This new finding indicates handwashing communication campaigns should specifically target people with suboptimal health conditions.

This study has several limitations that present opportunities for future research. First, the cross-sectional nature of our research precludes claims of causality, which is not a key concern as we are primarily interested in the predictive utility for screening less compliant groups. Second, although our survey was completely anonymous, self-reported handwashing compliance rates might be inflated due to social desirability [11]. Additionally, the anonymity of the survey makes it impossible to test if multiple respondents might be from the same household. Third, we did not explicitly collect data on how frequently respondents left their homes, making it conceivable that some respondents rarely or even never did. Fourth, we strived for broad coverage of the Malaysian population by using two-stage stratified sampling in terms of geographical area, ethnicity, gender and age, but Malaysia is a multi-faceted society and our sample includes a large share of college graduates, so the sample should not be considered strictly representative of the entire population. Fifth, social media usage was very high during the COVID-19 pandemic in Malaysia due to the implementation of a *cordon sanitaire* preventive measure, and future studies may examine our model in less severe pandemics without such a strict lockdown. Sixth, Malaysia is an upper-middle income country, where clean water for handwashing is generally accessible. Studies in countries with heterogeneous access to clean water might yield different results. Finally, this study focused on handwashing as a single hygiene measure. Future research might wish to study additional measures, such as the use of protective equipment or hand sanitiser, either in isolation or in combination with handwashing.

In conclusion, this study identifies several new predictors of handwashing during a pandemic, most notably people's use of social media. Consequently, it helps in identifying less compliant groups and enables health organisations, governments and other stakeholders to deliver more targeted handwashing communications campaigns, especially via social media. As prior research found that reminders [6] and signage [5] about handwashing are effective in increasing handwashing, but that most social media posts during a pandemic contain very little practical advice

[13], targeted social media campaigns based on the results of this study are likely to be effective in increasing handwashing in less compliant groups.

Acknowledgement. We acknowledge the support of Tsinghua University-INDITEX Sustainable Development Fund (Project No. TISD201904).

Data availability statements. Data are available upon request from the authors.

Conflicts of interest. The authors declare that each author has no conflicts of interest with respect to the research, authorship and/or publication of this paper.

References

1. **Jarvis WR** (1994) Handwashing – the Semmelweis lesson forgotten? *The Lancet* **344**, 1311–1312.
2. **World Health Organization Writing Group** (2006) Nonpharmaceutical interventions for pandemic influenza, national and community measures. *Emerging Infectious Diseases* **12**, 88–94.
3. **Rewar S, Mirdha D and Rewar P** (2015) Treatment and prevention of pandemic H1N1 influenza. *Annals of Global Health* **81**, 645–653.
4. **Whitby M, McLaws M-L and Ross MW** (2006) Why healthcare workers don't wash their hands: a behavioral explanation. *Infection Control & Hospital Epidemiology* **27**, 484–492.
5. **Updegraff JA, et al.** (2011) Framing flu prevention – an experimental field test of signs promoting hand hygiene during the 2009-2010 H1N1 pandemic. *Health Psychology* **30**, 295–299.
6. **Morrison LG and Yardley L** (2009) What infection control measures will people carry out to reduce transmission of pandemic influenza? a focus group study. *BMC Public Health* **9**, 258. doi: 10.1186/1471-2458-9-258.
7. **Chan AKM, et al.** (2020) Social media for rapid knowledge dissemination: early experience from the COVID-19 pandemic. *Anaesthesia*. (in press). doi:10.1111/anae.15057 .
8. **DeSalvo KB, et al.** (2006) Assessing measurement properties of two single-item general health measures. *Quality of Life Research* **15**, 191–201.
9. **Roelen CAM, et al.** (2014) Health measures in prediction models for high sickness absence: single-item self-rated health versus multi-item SF-12. *European Journal of Public Health* **25**, 668–672.
10. **Bish A and Michie S** (2010) Demographic and attitudinal determinants of protective behaviours during a pandemic: a review. *British Journal of Health Psychology* **15**, 797–824.
11. **Borchgrevink CP, Cha J and Kim S** (2013) Hand washing practices in a college town environment. *Journal of Environmental Health* **75**, 18–25.
12. **Putzke J, et al.** (2014) Cross-cultural gender differences in the adoption and usage of social media platforms—an exploratory study of Last.FM. *Computer Networks* **75**, 519–530.
13. **Vos SC and Buckner MM** (2016) Social media messages in an emerging health crisis: tweeting bird flu. *Journal of Health Communication* **21**, 301–308.