

Globodera pallida, a non-transgenic invertebrate as a new model for investigating Alzheimer's disease (and other proteinopathies)?

Norah A. Althobaiti*, Farid Mena*, Johnathan J. Dalzell, Brian D. Green*

Biological models of Alzheimer's disease (AD):

Non-human models have contributed tremendously to the understanding of AD and its underlying pathological processes. These models have aided the investigation of the genetic and environmental risk factors. They also have enabled the progression of candidate therapies into human clinical trials. Because of similarities with human brain anatomy and genetics, rodent models have been used extensively to recapitulate some aspects of AD pathology, measure AD-associated behavioral parameters and related nervous system dysfunctions (Eriksen and Janus, 2007). For instance, transgenic mice overexpressing human amyloid precursor protein have furthered the development of the amyloid cascade hypothesis as a central pillar of familial AD. Although considered as advantageous, mammalian models have practical and ethical problems when it comes to high-throughput screening (HTS) of drugs. Therefore, alternative models, including the use of invertebrates with very short lifecycles, represent rapid, simple, and highly cost-effective platforms for HTS or drug target identification (Artal-Sanz et al., 2006). Indeed, the use of such organisms, as opposed to individual cell models, provides advantages for understanding the wider complexity of human pathology. Thereby, prominent invertebrate model organisms, such as *Drosophila melanogaster* (fruit fly), and the nematode *Caenorhabditis elegans* (roundworm), were insightful into the aging process, AD and Parkinson's disease (Newman et al., 2011). The genome of *D. melanogaster* is ~70% homologous to that of humans and shows relatively complex brain and neuronal structures. Its eye is an accessible organ for phenotypic characterization, measuring of behavioral deficits, and determining responses to drug compounds. This fruit fly has unique learning and memory behaviors and can induce the gene expression of proteins considered to be the major hallmarks of AD, such as amyloid-beta (A β) or tau, both leading to AD pathological features and phenotypes. Furthermore, compounds such as GSK3- β inhibitors can rescue axonal transport and locomotion defects in these models. *C. elegans* is another invertebrate model organism employed in AD research. Several attempts have been made to use this free-living nematode (natively occurring in temperate soil environments) to generate transgenic AD models. One transgenic *C. elegans* model expressing A β peptides in muscle has a paralysis transgene-induced phenotype (Dostal and Link, 2010). Other models that neuronally express human A β exhibit phenotypes such as impaired chemotaxis and deficits in associated learning. Reported defects include impaired odor associative learning

behavior, impaired serotonin-stimulated egg-laying, and decreased lifespan evidently resulting from A β toxicity.

Introducing *Globodera pallida* (*G. pallida*) as a novel model for AD (and potentially other proteinopathies):

In this paper, we put forward a non-transgenic invertebrate for newly modeling AD pathology, *G. pallida*, which is one of the best characterized of the potato cyst nematodes (Althobaiti et al., 2021). *G. pallida* is an important model for understanding the biology of potato cyst nematodes, which are responsible for substantial global crop losses (Warnock et al., 2017). The lifecycle of *G. pallida* is described as obligate, involving the cross-fertilization of different hosts, and it has six life stages: egg (or embryo) stage, four juvenile stages (J1 to J4), and a sexually adult stage. Our proposed AD model uses J2s which have a motile, vermiform, round head and fine tail, with bodies ~500 μ m in length, styles of 19–25 μ m, and tails of 37–55 μ m. Nematodes have a structurally simple nervous system but contain a complex array of signaling molecules (e.g., neuropeptides) that perform vital functions such as the maintenance of parasite physiology, feeding, locomotion, and reproduction (Warnock et al., 2017). The nematode 'brain' (Figure 1) consists of a circumpharyngeal nerve ring (CNR) which encircles the oesophagus just behind the pharyngeal bulb. The CNR contains multiple neurons which connect several sensing organs and a nerve trunk that runs longitudinally from the CNR to the head and the

rest of the body. There are two pairs of lengthwise nerve cords on the dorsal and ventral sides, as well as multiple distinct chemosensory neural cell bodies in the ventral nerve cord. Chemosensory organs include amphids, near the head, and phasmids at their posterior. These chemosensory organs are vital to nematode behavior with a large number of neurons because they are sensitive to environmental changes, such as temperature, presence of food, or detection of reproductive mates. It has long been established that the nematode behavior is affected by chemicals exuded from plant roots, and that the sugar mannitol, is a key attractant for *G. pallida*. It is also known that their chemosensory neurons can assimilate exogenous peptides from the external environment (Warnock et al., 2017) – a feature which we have exploited for the development of *G. pallida* as a new model for studying AD pathology. Indeed, our peer-reviewed published research has demonstrated, for the first time, the capacity of *G. pallida* (as a non-genetically modified organism) to naturally assimilate A β peptides (Althobaiti et al., 2021). Crucially, these peptides almost exclusively localized to neurons (including the amphid neurons of the CNR), and this causes measurable impairments of their behavior. We have shown that their normal chemosensing of mannitol, as a food source, can be completely disrupted. The unique feature of such a model system is that it is possible to reliably control the dose, duration, and protocol the organisms' exposure to A β ; additionally, the impact of each specific A β isoform (or combination of isoforms) can be assessed. Chemotaxis is disrupted to the greatest extent by treatment with human A β (1–42) (which is also the most neurotoxic form), and we were able to compare this against the relative harm caused by other A β forms including (1–40), (17–42), (17–40), (1–28) and (1–16). For A β (1–42), we have established a time course for their responsiveness. Pre-incubations for at least 6 hours significantly affect chemosensing, with maximal effect reached by 18–24 hours. Clearly, the absorption and toxicity of A β peptides on the neurons and CNR takes a period to manifest as behavioral changes




Figure 1 | A proposed process for establishing proteinopathy models in *G. pallida* and testing. After hatching and growth to J2 juveniles, the potato cyst nematodes were incubated with a selected peptide or protein (e.g., amyloid beta (1–42)). Assimilation and co-localization to neurons were assessed by immunocytochemistry using an appropriate antibody, and the physiological and behavioral effects were evaluated by a battery of tests. Once the model is characterized, the process can be repeated with libraries of candidate drugs or medicinal plant extracts. AN: Amphidial neuron; BWM: body wall muscle; CNR: circumpharyngeal nerve ring; PHB: pharyngeal bulb; PN: phasmid neurons; SPM: stylet protractor muscles; VNC: ventral nerve cord.

in this nematode. The major contributors to this are the transit times of peptides along neurons (corroborated by immunocytochemical staining), and the time taken for neurotoxicity to occur. In addition to measuring changes in chemosensing, we describe several other quantitative parameters that can be applied in *G. pallida* as an alternative model nematode of AD. Some of these are discussed below and include: the fluorometric measurement of the production of reactive oxygen species (ROS), fluorometric measurement of oxidized/reduced glutathione (GSSG/GSH), quantification of mitochondrial reductase activity, and the visual assessment of their movement and survival (Figure 1). These methods could be combined into a proposed process for establishing proteinopathy models in *G. pallida* and for testing these. Oxidative stress is a factor implicated in the occurrence and progression of AD, and A β is implicated in the production of ROS. In *G. pallida*, ROS significantly rose upon exposure to A β peptide fragments, particularly A β (1–42) and A β (17–42), which are known to be most toxic (Joshi et al., 2005). The methionine at position 35 of A β is thought to be an important amino acid for neurotoxicity and oxidative stress (Navarro-Yepes et al., 2014). The highest levels of ROS production have been elicited with A β (1–16), A β (1–28), A β (1–40), and A β (17–40) peptides. A degree of sensitivity appears to be afforded by measuring ROS in *G. pallida*, since A β (1–16) and A β (1–28) isoforms do not aggregate and are rarely considered to be neurotoxic. Other studies suggest that A β (1–28) may not induce oxidative stress (Butterfield and Boyd-Kimball, 2005), but A β (1–16) has previously been demonstrated to induce ROS formation (Du et al., 2011). In the *G. pallida* system, A β (1–40) and (17–40) induced the lowest amounts of ROS formation, which is in line with the literature. It is postulated that the termination of the peptide sequence at position 40 makes them more soluble, less toxic and less prone to aggregation compared with peptides terminating at position 42. Indeed, it is clear that A β (1–42) and (17–42) cause mitochondrial dysfunction induction, lipid and protein oxidation in neuronal cells culture (Butterfield and Boyd-Kimball, 2005), and A β (1–42) is thought to impact AD pathology to a much greater extent than other smaller fragments of A β (Butterfield and Sultana, 2011). GSH plays an important cellular role in the anti-oxidative activity, but when cells are exposed to increased levels of oxidative stress, it is worth noting that the oxidized glutathione GSSG will accumulate, thereby altering the ratio of GSH to GSSG. Measurement of this ratio in *G. pallida* reinforces the findings of ROS assays. GSSG/GSH ratios were markedly augmented by A β (1–42) in a cumulative manner over 24 hours. This has a disease relevance since brain tissues from AD and mild cognitive impairment subjects have lower levels of antioxidant enzymes, perhaps making it more vulnerable to A β (1–42) induced-toxicity. Relatively high levels of metal ions, oxygen and unsaturated lipids in the brain also make this organ more sensitive and vulnerable to damage from oxidative stress (Pocernich and Butterfield, 2012).

Future potential of *G. pallida* for examining proteinopathies: One might envisage running automated ultra-HTS of thousands of candidate

compounds or medicinal plant extracts, in order to evaluate their potential for countering the neurodegenerative effects of amyloidoses, tauopathies, and synucleinopathies. AD is a key example of both amyloidosis and tauopathy, and thus, studies with tau and phospho-tau are warranted. Other major synucleinopathies include Parkinson's disease (where alpha-synuclein aggregates in the brain stem), Lewy body disease (where Lewy bodies occur in the cortico-limbic and brainstem), and multiple system atrophy (where alpha-synuclein accumulates in basal ganglia, brainstem, and cerebellum). Taken together, studies that examined the assimilation of alpha-synuclein isoforms should be undertaken. Furthermore, *G. pallida* model can be potentially used to examine prion diseases, for instance by examining the role of prion protein in the manifestation of Creutzfeldt-Jakob disease. Another distinctive feature of *G. pallida* is its extreme sensitivity to neuronal RNA interference, which would be a powerful functional genomics tool for studying the underlying mechanisms of proteinopathy toxicity (Dalzell et al., 2010).

Final remarks and perspectives: Although there have been significant advances in understanding neurodegenerative diseases, such as AD, it is worth mentioning that there is still no effective cure. Prevalence of AD has dramatically increased with life expectancy, which has been steadily increasing worldwide. Establishing a novel model system such *G. pallida*, that is cost-effective, practical, non-transgenic (to ensure data reliability), and that provides an HTS platform for screening potential treatments, will hopefully aid in the discovery of effective therapy with minimal adverse effects. Such a disease model provides a means of modulating the dose, duration, sequence, and type of proteinopathy being investigated, and therefore is worthy of further investigation.

Norah A. Althobaiti*, Farid Menaa*, Johnathan J. Dalzell, Brian D. Green*

Biology Department, College of Science and Humanities-Al Quwaiyah, Shaqra University, Al Quwaiyah, Saudi Arabia (Althobaiti NA) Departments of Internal Medicine and Nanomedicine, Fluorotronics-California Innovations Corporation, San Diego, CA, USA (Menaa F) Department of grassland and Plant Science, AgriFood and Biosciences Institute, Northern Ireland, UK (Dalzell JJ) Institute for Global Food Security, School of Biological Sciences, Queen's University Belfast, Belfast, UK (Green BD)

*Correspondence to: Norah A. Althobaiti, PhD, nalthobaiti@su.edu.sa; Farid Menaa, MD, PhD, menateam@gmail.com; Brian D. Green, PhD, b.green@qub.ac.uk.

<https://orcid.org/0000-0001-6893-5521>

(Norah A. Althobaiti)

<https://orcid.org/0000-0002-0258-7322>

(Farid Menaa)

<https://orcid.org/0000-0003-3331-0032>

(Brian D. Green)

Date of submission: November 1, 2021

Date of decision: January 28, 2022

Date of acceptance: February 14, 2022

Date of web publication: April 25, 2022

<https://doi.org/10.4103/1673-5374.341042>

How to cite this article: Althobaiti NA, Menaa F, Dalzell JJ, Green BD (2023) *Globodera pallida*, a non-transgenic invertebrate as a new model for investigating Alzheimer's disease (and other proteinopathies)? *Neural Regen Res* 18(1): 113-114.

Open access statement: This is an open access journal, and articles are distributed under the terms of the Creative Commons AttributionNonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

References

- Althobaiti NA, Menaa F, Albalawi AE, Dalzell JJ, Warnock ND, Mccammick EM, Alsolais A, Alkhaibari AM, Green BD (2021) Assessment and validation of *Globodera pallida* as a novel in vivo model for studying Alzheimer's disease. *Cells* 10:2481.
- Artal-Sanz M, de Jong L, Tavernarakis N (2006) *Caenorhabditis elegans*: a versatile platform for drug discovery. *Biotechnol J* 1:1405-1418.
- Butterfield DA, Boyd-Kimball D (2005) The critical role of methionine 35 in Alzheimer's amyloid β -peptide (1–42)-induced oxidative stress and neurotoxicity. *Biochim Biophys Acta* 1703:149-156.
- Butterfield DA, Sultana R (2011) Methionine-35 of A β (1–42): importance for oxidative stress in Alzheimer disease. *J Amino Acids* 2011:198430.
- Dalzell JJ, McMaster S, Fleming CC, Maule AG (2010) Short interfering RNA-mediated gene silencing in *Globodera pallida* and *Meloidogyne incognita* infective stage juveniles. *Int J Parasitol* 40:91-100.
- Dostal V, Link CD (2010) Assaying β -amyloid toxicity using a transgenic *C. elegans* model. *J Vis Exp* 2010:e2252.
- Du XT, Wang L, Wang YJ, Andreassen M, Zhan DW, Feng Y, Li M, Zhao M, Otzen D, Xue D (2011) A β 1-16 can aggregate and induce the production of reactive oxygen species, nitric oxide, and inflammatory cytokines. *J Alzheimers Dis* 27: 401-413.
- Eriksen JL, Janus CG (2007) Plaques, tangles, and memory loss in mouse models of neurodegeneration. *Behav Genet* 37:79-100.
- Joshi G, Sultana R, Perluigi M, Butterfield DA (2005) In vivo protection of synaptosomes from oxidative stress mediated by Fe²⁺/H₂O₂ or 2, 2-azobis-(2-amidinopropane) dihydrochloride by the glutathione mimetic tricyclodecan-9-yl-xanthogenate. *Free Radic Biol Med* 38:1023-1031.
- Navarro-Yepes J, Zavala-Flores L, Anandhan A, Wang F, Skotak M, Chandra N, Li M, Pappa A, Martinez-Fong D, Del Razo LM (2014) Antioxidant gene therapy against neuronal cell death. *Pharmacol Ther* 142:206-230.
- Newman T, Sinadinos C, Johnston A, Sealey M, Mudher M (2011) Using *Drosophila* models of neurodegenerative diseases for drug discovery. *Expert Opin Drug Discov* 6:129-410.
- Pocernich CB, Butterfield DA (2012) Elevation of glutathione as a therapeutic strategy in Alzheimer disease. *Biochim Biophys Acta* 1822:625-630.
- Warnock ND, Wilson L, Patten C, Fleming CC, Maule AG, Dalzell JJ (2017) Nematode neuropeptides as transgenic nematocides. *PLoS Pathog* 13:e1006237.

C-Editors: Zhao M, Liu WJ, Wang Lu; T-Editor: Jia Y