

Research

Open Access

Analysis of a possible independent origin of triploid *P. formosa* outside of the Río Purificación river system

Susanne Schories¹, Kathrin P Lampert¹, Dunja K Lamatsch^{1,2}, Francisco J García de León³ and Manfred Scharl^{*1}

Address: ¹Physiological Chemistry I, Biocenter, University of Würzburg, Biozentrum, Am Hubland, D-97074 Würzburg, Germany, ²Department of Animal and Plant Sciences, The University of Sheffield, Sheffield S10 2TN, UK and ³Centro de Investigaciones Biológicas del Noroeste, S.C. (CIBNOR, S.C.), Mar Bermejo No. 195, Col. Playa Palo de Santa Rita, La Paz, BCS, 23090, México

Email: Susanne Schories - susanne.schories@biozentrum.uni-wuerzburg.de; Kathrin P Lampert - kathrin.lampert@biozentrum.uni-wuerzburg.de; Dunja K Lamatsch - d.lamatsch@shef.ac.uk; Francisco J García de León - fgarciadl@cibnor.mx; Manfred Scharl^{*} - phch1@biozentrum.uni-wuerzburg.de

* Corresponding author

Published: 15 May 2007

Received: 29 March 2007

Frontiers in Zoology 2007, 4:13 doi:10.1186/1742-9994-4-13

Accepted: 15 May 2007

This article is available from: <http://www.frontiersinzoology.com/content/4/1/13>

© 2007 Schories et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Unisexuality, or all female reproduction, is rare among vertebrates. Studying these exceptional organisms may give useful information with respect to the evolution and maintenance of sexual reproduction. *Poecilia formosa* was the first unisexual vertebrate species to be detected and since then has served as a paradigmatic organism for unisexuality and studies on the evolution of sex. It reproduces through gynogenesis, using sperm of males from related species to trigger parthenogenetic development of the unreduced diploid eggs. Like in other unisexual vertebrates, triploids occur in a certain range of *P. formosa*. It has been suggested that the addition of the host species derived third chromosome set is evolutionary important. Clonal organisms lack sufficient genotypic diversity for adaptive changes to variable environments. Also non-recombining genomes cannot purge deleterious mutations and therefore unisexual organisms should suffer from a genomic decay. Thus, polyploidization leading to triploidy should bring "fresh" genetic material into the asexual lineage. To evaluate the importance of triploidy for maintaining the asexual species, it is important to know whether such an introgression event happens at a reasonable frequency.

Results: In an earlier study it was found that all triploid *P. formosa* in the Río Purificación river system are of monophyletic origin. Here we have analyzed fish from a different river system. Using microsatellite analysis we can show that the triploids from this new location are genetically divergent and most probably of an independent origin.

Conclusion: Our data support the hypothesis that triploidy was not a single chance event in the evolutionary history of *P. formosa* and hence might be a relevant mechanism to increase genotypic divergence and at least partially counteract the genetic degeneration connected to asexuality. It is, however, much rarer than in other asexual vertebrates analyzed so far and thus probably only of moderate evolutionary importance for the maintenance of the asexual breeding complex.

Background

Poecilia formosa is an all female teleost fish species. In fact it was the first asexual vertebrate to be described [1] and has been a paradigmatic organism for studies on the evolution of sexual reproduction and the maintenance of asexuality [2-4]. Its mode of reproduction is gynogenesis, a form of parthenogenesis, in which the embryonic development of unreduced diploid eggs is triggered by sperm from males of closely related species.

Usually the DNA of the sperm does not reach the egg nucleus resulting in clonal offspring. In exceptional cases the exclusion mechanism fails and either small fragments of single chromosomes, designated supernumerary chromosomes or microchromosomes [5], or the full chromosome set of the whole haploid genome of the sperm is included [6-8].

Fish with additional microchromosomes readily can form new clones that stably transmit the microchromosome. This phenomenon has been found in laboratory broods as well as in wild populations and occurs obviously at a reasonable frequency [9]. Interestingly, microchromosome carrying fish have been reported only from the Río Purificación [10].

The situation for the triploids is different: all triploids that occurred in the laboratory so far were sterile, while triploids collected in nature produce all triploid offspring [8,11]. Those triploids have been found in the Río Purificación and its tributaries [12,13], as well as in the hydrologically connected Río Soto la Marina river system. Apart from that only in the river system adjacent to the south, the Río Guayalejo, some stable triploid populations exist [6]. At the periphery of the range of *P. formosa*, triploids appear to be extremely rare. Enigmatic reports exist from the Río Grande (or Río Bravo) basin where during steady large-scale sampling (N = 5000) only four triploids were collected in ten years. Another single triploid was found in the Río Tuxpan system. The rare triploids outside the upper Río Purificación were interpreted to be "sterile first generation triploids" like the laboratory born [13]. Triploids are frequent where *P. formosa* is found together with *P. limantouri*, which serves as sperm donor, but absent or rare where *P. formosa* is sympatric with the alternative sperm donors *P. mexicana* or *P. latipinna*. The hypothesis was put forward that triploids may arise de-novo throughout the whole range, but are only fertile and can survive only in the region, where *P. formosa* occurs together with *P. limantouri* [14], namely in the Ríos Purificación, Soto la Marina and Guayalejo. An alternative hypothesis, however, is that reproductive triploidy is an extremely rare event, which occurred so far only once in an area, where *P. formosa* mates with *P. limantouri*. From there the triploid

fish spread and invaded the neighboring river systems but have not reached the periphery of the range.

A microsatellite based study on fish from the Río Purificación suggested that all triploids that were analyzed belonged to only two clonal lineages and had a monophyletic origin [12]. This stands in contrast to the diploids, which display a high number of different clones [12,15]. It led to the conclusion that the generation of a fertile triploid *P. formosa* is a rare, if not a unique event.

We report here a molecular genetic analysis of a stable population of triploid *P. formosa* outside the Río Purificación system. The allelic and genotypic variability of the triploid lineages and the question how recently they might have emerged was studied. Phylogenetic analysis revealed that the triploid clones from the new locality most probably are of independent origin of the Río Purificación clones. Thus fertile triploids may have been generated at least twice, but this event is much less frequent contrary to other asexual vertebrates.

Results

In a genetic survey of natural populations of *P. formosa* we analyzed 114 and 100 animals collected in 2002 and 2005, respectively, at a site in the Río Guayalejo (G) river system. This river system is the next south but is hydrogeographically separated from the Río Purificación and Río Soto la Marina system (P). Flow cytometric measurements of DNA content revealed that 14 individuals of the 2002 (12%) and 16 individuals of the 2005 sample (16%) were triploids.

The eleven microsatellite loci used to genotype the triploid clones detected very different levels of variability (Table 1). Combining all loci the allelic diversity was higher in triploids from Río Purificación (P) than in triploids from Río Guayalejo (G) (Table 1). Triploids from both sites shared 35% of all alleles found in triploid individuals.

In a previous study on the triploids from the Río Purificación system the triploid genotypes were separated in two lineages based on their overall genotypic differences in the three microsatellite loci used [12]. The lineages were designated Y and Z. Following this nomenclature the new lineage from G on the basis of the same three microsatellite loci was named X. When additional microsatellite loci were analyzed in this study they revealed that each of the three lineages includes several clonal genotypes. The clonal variation between individual genotypes was mainly due to the variation at the two most variable loci mCA20 and MSD23. The overall genotypic variation was lower in G (4 different genotypes) than in P (20 different genotypes) (Table 2). The recapture rate of clones was also

Table 1: Variability of microsatellite loci in different sites (G – Río Guayalejo; P – Río Purificación) and years.

Locus	site – year				total	% shared alleles
	G – 2002	G – 2005	P – 2002	P – 2005		
N	11	15	26	13	65	
mCA20	2	2	16	12	17	11.76
MSD23	6	6	9	9	14	42.86
SatI	3	3	5	5	8	12.5
KonDI5	3	3	5	5	6	33.33
PR39	3	3	4	4	5	40
mCA16	2	2	3	3	3	66.67
MATG31	3	3	3	3	4	50
MATG38	3	3	4	4	4	75
MATG44	2	2	3	3	3	66.67
MATG61	3	3	3	3	4	50
MATG78	2	2	2	2	3	33.33

Given are numbers of alleles for each locus and in total for all triploids found. Only individuals that could be genotyped at all 11 loci were taken into account. The percentages of shared alleles between sites for each locus are given.

higher in G: three of the four clones found in 2002 could be found again in 2005, while in P of the 15 clones found in 2002 only three were found again in 2005. In and between the years none of the triploid clones found in one site could be found in the other site (Table 2). The lower genotypic variability in G was confirmed by a frequency analysis according to Menken et al. [16]: the proportion of distinguishable clones in G (0.28 (mean of both years)) was lower than in P (0.60 (mean of both years)). The clonal diversity value (mean of both years) was also lower in G (0.57) than in P (0.88) (Table 3).

In order to find out whether the triploids from the Guayalejo and from the Purificación system have a monophyletic origin, a phylogenetic tree using the Neighbour Joining method (NJ) method based on d_{mu}^2 [17] was performed (Fig. 1). This revealed a clear differentiation between the X lineage found at G and the Y and Z lineages found at P.

The separation of the triploids in G and P could also be shown in a principal component analysis: one significant axis ($P = 0.039$) was found that clearly separated the X genotype from the Y and Z genotypes and explained 59% of the variability found in the data (Fig. 2). A second axis separating the Z from the Y genotypes was found to be not significant ($P = 0.858$). A search for a closely related diploid genotype (meaning a diploid sharing all alleles with a triploid) for each triploid lineage revealed that one and the same diploid genotype was closest to all triploid lineages (X, Y, and Z). Such a diploid genotype was found at P (P11). This was confirmed by the neighbor joining tree (Fig. 1).

Mitochondrial DNA did not show any variability in the cytochrome b or the D-loop sequences analyzed. The same haplotype was found in the diploids as well as in the triploids from all sampling sites.

Discussion

In this study we have analyzed a population of diploid and triploid *P. formosa* outside the Río Purificación system. From the high number of triploids found in both years (12% in 2002 and 16% in 2005) we conclude that in the Río Guayalejo system triploids actually are a stable part of the population. They were as common as in the formerly known Río Purificación field sites [12]. This is also in the range reported in the literature for the occurrence of triploids identified earlier at these and other sites [6,14,18].

Triploid lineages in this study were analyzed using 11 microsatellites as well as mtDNA. Mitochondrial DNA was found to be not variable enough to convey any information about the formation of the novel triploid lineage found in G. For all three investigated segments of the mitochondrial genome a single haplotype was present in diploid as well as triploid individuals from P and G. This may be due to the assumed single monophyletic origin of all diploid *P. formosa* [4,19,20].

Three microsatellites had been used in a former study to analyze clonal diversity of triploids at several sites of P and had uncovered only two different lineages: Y and Z [12]. Interestingly, two of the additional microsatellites used in this study proved to be very variable and showed that both triploid lineages show a certain degree of clonal variability. The high number of triploid genotypes detected in the P site is concordant with the findings from multilocus DNA fingerprints, which predicted a high genetic variability for triploids [12].

Triploids from the G site form a clearly distinct lineage and are less variable than triploids from the P site. The origin and the low diversity of the triploids in this site can be explained either by the hypothesis that an ancient population immigrated from G, diverged and then went

Table 2: Genotypes of all triploids found at the two sites

genotypes														
#	site	year	mCA20	MSD 23	SatI	KonDI5	PR39	mCA16	mATG31	mATG38	mATG44	mATG61	mATG78	
X1	1	G	2002	115-115- 196	205 -217-245	102-124-152	255-280- 282	141- 153 -166	<u>203-203-221</u>	70 -73-79	134-170-185	<u>115-115-140</u>	133- 142 -145	192 -198
X2	7	G	2002/2005	115-115- 196	205 -229-241	102-124-152	255-280- 282	141- 153 -166	<u>203-203-221</u>	70 -73-79	134-170-185	<u>115-115-140</u>	133- 142 -145	192 -198
X3	15	G	2002/2005	115-115- 196	205 -229-245	102-124-152	255-280- 282	141- 153 -166	<u>203-203-221</u>	70 -73-79	134-170-185	<u>115-115-140</u>	133- 142 -145	192 -198
X4	3	G	2002/2005	115-115- 196	205 -233-245	102-124-152	255-280- 282	141- 153 -166	<u>203-203-221</u>	70 -73-79	134-170-185	<u>115-115-140</u>	133- 142 -145	192 -198
Y1	1	P	2002	115-115-208	201-217-241	116-122-144	255-258-270	141-147-166	191-203-221	73-76-79	134-170-185	115-118-140	133-139-145	198-201
Y2	2	P	2002	115-115-252	205-217-245	116-122-144	255-258-270	141-147-166	191-203-221	73-76-79	134-170-185	115-118-140	133-139-145	198-201
Y3	2	P	2002	115-115-258	205-217-253	116-122-144	255-258-270	141-147-166	191-203-221	73-76-79	134-170-185	115-118-140	133-139-145	198-201
Y4	4	P	2002/2005	115-115-236	209-213-245	116-122-144	255-258-270	141-147-166	191-203-221	73-76-79	134-170-185	115-118-140	133-139-145	198-201
Y5	1	P	2002	115-115-240	209-213-245	116-122-144	255-258-270	141-147-166	191-203-221	73-76-79	134-170-185	115-118-140	133-139-145	198-201
Y6	6	P	2002/2005	115-115-242	209-213-245	116-122-144	255-258-270	141-147-166	191-203-221	73-76-79	134-170-185	115-118-140	133-139-145	198-201
Y7	2	P	2002	115-115-244	209-213-245	116-122-144	255-258-270	141-147-166	191-203-221	73-76-79	134-170-185	115-118-140	133-139-145	198-201
Y8	1	P	2002	115-115-246	209-213-245	116-122-144	255-258-270	141-147-166	191-203-221	73-76-79	134-170-185	115-118-140	133-139-145	198-201
Y9	1	P	2002	115-115-248	209-213-245	116-122-144	255-258-270	141-147-166	191-203-221	73-76-79	134-170-185	115-118-140	133-139-145	198-201
Y10	2	P	2002	115-115-250	209-213-245	116-122-144	255-258-270	141-147-166	191-203-221	73-76-79	134-170-185	115-118-140	133-139-145	198-201
Y11	2	P	2002	115-115-254	209-213-245	116-122-144	255-258-270	141-147-166	191-203-221	73-76-79	134-170-185	115-118-140	133-139-145	198-201
Y12	3	P	2002	115-115-258	209-213-245	116-122-144	255-258-270	141-147-166	191-203-221	73-76-79	134-170-185	115-118-140	133-139-145	198-201
Y13	1	P	2002	115-115-262	209-213-245	116-122-144	255-258-270	141-147-166	191-203-221	73-76-79	134-170-185	115-118-140	133-139-145	198-201
Y14	3	P	2002/2005	115-115-264	209-213-245	116-122-144	255-258-270	141-147-166	191-203-221	73-76-79	134-170-185	115-118-140	133-139-145	198-201
Y15	3	P	2002	115-115-268	209-213-245	116-122-144	255-258-270	141-147-166	191-203-221	73-76-79	134-170-185	115-118-140	133-139-145	198-201
Y16	1	P	2005	115-115-242	201-209-245	116-122-144	255-258-270	141-147-166	191-203-221	73-76-79	134-170-185	115-118-140	133-139-145	198-201
Y17	1	P	2005	115-115-264	201-213-245	116-122-144	255-258-270	141-147-166	191-203-221	73-76-79	134-170-185	115-118-140	133-139-145	198-201
Y18	1	P	2005	115-115-246	209-213-251	116-122-144	255-258-270	141-147-166	191-203-221	73-76-79	134-170-185	115-118-140	133-139-145	198-201
Y19	1	P	2005	115-115-236	211-217-245	116-122-144	255-258-270	141-147-166	191-203-221	73-76-79	134-170-185	115-118-140	133-139-145	198-201
Z1	1	P	2005	115-115- <i>188</i>	209-233-245	<i>114-120-144</i>	255-263-280	147- <i>149</i> -166	191-203-221	73-76-79	134-170- <i>182</i>	115-115-140	133-139-145	198-201

Given are the genotype name, the number of individuals found to have this exact genotype, the site the genotypes were found (P – Río Purificación, G – Río Guayalejo), the year and the allele sizes (bp) for all 11 microsatellite loci. Alleles that are present in the triploid X lineage from G but not in the triploid Y and Z lineages in P are marked in bold typeface. Genotypes that differ between P and G even though all alleles are present in both populations are underlined. Alleles differing between the P lineages Y and Z are marked in italic typeface.

Table 3: Menken analyses of clonal diversity of the triploid lineages in the two sampling sites.

	P 2002	P 2005	G 2002	G 2005
#ind	26	13	11	15
#clones	15	8	4	3
ENC	12.52	6.26	2.20	2.42
PDC	0.58	0.62	0.36	0.20
Unique	7.00	5	1	0
CD	0.92	0.84	0.55	0.59
CE	0.78	0.78	0.55	0.81

For both years and sites the number of triploid individuals sampled (#ind), the number of different clones found within those individuals (#clones), the effective number of clones (ENC), the proportion distinguishable clones, the number of unique clones (unique), the clonal diversity (CD) and the clonal evenness (CE) are given.

through a bottleneck or that the triploids are of recent origin at the P site. Such an independent triploidisation at P seems to be the most likely explanation because our results show a clear separation of G (X) and P (Y+Z) genotypes in the NJ tree as well as in the principal component analyses. Even though triploids from G share several alleles with the triploids from P those alleles are very common in the diploid population and might therefore not have been inherited from a common ancestor.

Interestingly, the closest related diploid genotype to all three triploid lineages was found at P. This result was supported by the NJ tree. While diploid P and G genotypes were found to be mixed within separated branches the three diploid genotypes (P4, P7, P11) closest to the triploids clearly came from P. We hesitate to assume, solely based on this information, that the X genotype was derived from P because the frequency of diploid genotypes varies greatly between years and the diploid genotype P11 or the related true ancestral diploid genotype leading to the triploid lineage X might not have been collected so far at the P site or may no longer be around.

An explanation for the fact that the triploid lineages at the P and the G site are closest related to the same diploid genotype might be that successful triploidisation may only be possible in a certain diploid genotypic background, probably even a single clonal lineage. It would also fit the observation that the triploids, which occur quite frequently under laboratory conditions are always sterile [11] probably due to non-matching genotypes. In the hybridogenetic unisexual fish *Poeciliopsis* convincing evidence has been produced that triploid clones are generated de-novo quite frequently, but that the third genome was added always to the same allodiploid clone of a certain biotype [21].

A bottleneck due to breakdown of an old P derived triploid population (that invaded to G) and a recent recovery




Figure 1
Neighbour joining tree of individual genetic distances between diploid and triploid genotypes in P and G.

Shown are the triploid genotypes from P (YYY and ZZZ) and G (XXX) and several closely related diploid genotypes from both sites (P1–11; G1–8). Branches with a bootstrap support of less than 60% were collapsed. Bootstrapping support was derived from 1000 iterations.




Figure 2
Principal component analyses based on individual genotypes of the triploids found in the different sites.
 The X axis is significant ($P = 0.039$) and explains 59% of the variability found in the data. The Y axis is not significant ($P = 0.858$). Levels of significance were derived from 10000 iterations. (Solid line – genotypes from P; dotted line – genotypes from G).

from a genotypically uniform small founder population is less likely because migration of a triploid genotype from P to G appears to be impossible due to geographic barriers: the G river system is the next south to the P river system but it is hydrogeographically completely separated by a high mountain range. In this context it is interesting to note that this border between the Río Purificación/Río Soto la Marina system and the Guayalejo/Panuco river basin efficiently prevents transgression of fish species. It is the northern limit of *P. latipunctata* and *Xiphophorus variatus* and the southern limit of *X. xiphidium* [22].

We can of course not exclude that the triploid fish were artificially introduced into the Guayalejo system by human activities a long time ago, diverged here, and then from a genetic bottleneck re-established themselves from a highly divergent clone. River mouth hopping obviously happened in the spreading of diploid *P. formosa* [23] but is also unlikely as a source of triploids at G as triploids are not recorded despite extensive sampling from the lower Río Soto la Marina and the estuaries of both rivers. In addition, triploids apparently survive and are perpetuated only in the clear waters of upstream habitats [14].

Although the triploid lineage at G appears to represent an independent polyploidization event, this is only the second to be described in many samples analyzed to date. This is surprisingly rare compared to other asexual breeding complexes.

Multiple independent origins of gynogenetic triploid lineages have been found in the *Poeciliopsis* complex [24]

and for the hybrid spined loaches of the genus *Cobitis* [25]. Here, multiple hybridization events of a diploid asexual fish with the host species established reproductive triploidy [25], leading to several independent triploid clonal lineages even at one and the same spot [24]. Also in the mixed ploidy unisexual breeding complexes of the Iberian minnow and the North-American genus *Phoxinus*, allotriploids are formed from diploid ancestors quite frequently [26,27].

Triploidy in general is very often connected to parthenogenesis. Parthenogenetic triploid systems are widespread and many examples are found amongst ants, lizards, salamanders, grasshoppers, fish, annelids, plathelminthes, nematodes, and plants. (For reviews see [28-32]). In those cases where the origin of the polyploid state is known to be due to "genome addition", it is a rather frequent event [28].

In summary, the triploids from the Río Purificación are according to our present knowledge all monophyletic derivatives of the same single introgression event [12]. On the basis of our analysis we favor the hypothesis that the triploids from the Guayalejo system arose from a second independent hybridization event of *P. formosa* with its sexual host species *P. mexicana*. It is also possible that the third genome is derived from *P. latipunctata*, which is a third host species for *P. formosa* and occurs in the Río Guayalejo and its tributaries [33]. However, this species has not been detected so far at the locality where the triploids were collected. It will be interesting to search at further sites in the future, whether other independent populations of triploid *P. formosa* exist.

Conclusion

The microsatellite analysis identified the triploid fish from the Río Guayalejo as being genetically divergent from the triploids of the Río Purificación. The most likely explanation is that these fish arose from independent paternal introgression events, through which an entire haploid genome of the gonochoristic host species was added to the non-recombining diploid genome of *P. formosa*. The fact that this introgression event was not a singular one, but might be much more frequent, supports the hypothesis that triploidy might be a relevant mechanism through which the new genetic variation is introduced. This can supplement the generally low genotypic diversity for adaptive changes to variable environments or parasites that clonal organisms suffer from [34,35]. In addition, accumulation of deleterious mutations as a result of the lack of recombination in an asexual organism could be at least partly compensated. It is, however, intriguing that reproductive triploidy occurs obviously extremely rare in *P. formosa*, compared to the other mode of paternal introgression, namely microchromosomes. Thus triploidy

appears to be of less evolutionary importance for the long term survival of this asexual species.

Methods

Sampling

The collection site is a ford approximately 1 km southeast of the village of Emiliano Zapata at the Río Guayalejo (23° 16.624'N 98° 56.315'W). Individuals were captured using seine nets. One (5 mm²) piece of the fast regenerating dorsal fin was cut from each individual and stored in ethanol (70%) for transportation to the laboratory. Immediately after taking the biopsy, all animals were released. Samples were taken in February 2002 and in March 2005.

Sample processing

All fins were cut in half to use one part to determine ploidy levels via flow cytometry and the second part to extract DNA for PCR analyses.

For flow cytometric measurements the fins were stained with the fluorescent dye DAPI, and ploidy levels were determined using chicken blood as a reference, essentially as described [36].

For microsatellite analysis DNA was extracted using 100 mM NaCl/0.5% Sarcosyl buffer and chelex (20%), according to [37]. The following primers were used: Sat1, KonD15, PR39 [12] and mCA16, mCA20, MSD23, mATG31, mATG38, mATG44, mATG61, mATG78 (originally from Walter et al 2004 [38] as reported in Lampert et al. 2006 [39]). From the primer pairs, one primer was Cy5 labelled (MWG, Ebersberg, Germany) at the 5' end, to allow detection by an ALF Express sequencer (Amersham Biosciences, Freiburg, Germany) (see below). PCR reactions: 300 sec of denaturing, 30 sec at 94°C, 30 sec at annealing temperature (52°C for KonD15/58°C for Sat1/55°C for all other primers) and 30 sec at 72°C for 35 cycles followed by a 300 sec elongation step. PCR products were analyzed on an ALF Express sequencer using an external marker (50–500 bp in 50 bp steps) (Amersham Biosciences, Freiburg). Internal standards were used to facilitate fragment alignment. ALF Express created files (.alx) were converted into alf files using the appropriate program ALX2ALF. .alf files were analyzed via the FM (Fragment Manager) program (Amersham Biosciences, Freiburg).

To confirm the morphological identification of the fish as either *P. formosa* or *P. mexicana* a diagnosis by PCR with primers for the *src* gene was performed [20]. Primers ScrE and ScrF give one product of approximately 650 bp from *P. mexicana* DNA and two products of 650 bp and 700 bp from *P. formosa*. Homo- or heterozygosity of individuals

and hence species identity can easily be determined on 2% agarose gels.

Extracted DNA was also used to investigate variability of mitochondrial DNA (mtDNA) in the triploids found in both sites. We sequenced two parts of the cytochrome b gene and parts of the D-loop sequence in triploids and in two diploids from the two sites (P and G). The diploid genotypes used for the comparison of mtDNA genes were picked due to their very different microsatellite allelic pattern. For cytochrome b we used the primers L14724/H15149 and L15513/H16100 and for the D-loop primer pairs L15995/H16498 were used (all from Meyer et al. 2004 [40]). Amplification was done using the PCR conditions recommended by Meyer et al. We directly sequenced the PCR products using the appropriate H and L PCR primers. Sequencing protocol using the DTSC method followed the Beckman-Coulter instructions. Sequences were analyzed using a Beckman-Coulter capillary sequencer and the Ceq8000 software provided by the company. Sequence alignments were done using the BioEdit program [41].

Data analyses

Triploid genotypes found at the Río Guayalejo site (G) were compared to the triploid genotypes from the Río Purificación site (P) from the same years for levels of allelic and genotypic variability. Only individuals that could be genotyped at all 11 microsatellite loci were included in the statistical analyses. Genetic diversity among triploids was described following Menken et al. (1995) [16]: the proportion of distinguishable clones ($pdc = \text{number of clones divided by sample size}$; a value of 1.00 means every individual has a different genotype), the effective number of clones ($enc = 1/\sum p_i^2$, p_i – frequency of the i th clone in a population), clonal diversity ($cd = 1 - \sum p_i^2$) and clonal evenness ($ce = (1/\sum p_i^2)/Nc - Nc$ number of clones (Clonal evenness (ce) gives a relative and therefore comparable measure for the distribution of clones. A ce value of one means all clones are equally distributed.) were calculated.

We used the program POPULATIONS [42] to calculate individual distances based on Goldstein's d_{mu}^2 [17] between genotypes. To enable the inclusion of diploid genotypes and to avoid an "oversampling" of the Y branch only three microsatellite loci (Sat1, KonD15, PR39) were included in the calculations. Individual distances between genotypes were used to create a neighbor-joining (NJ) tree. Bootstrapping values for the NJ tree were obtained from 1000 iterations. In addition, we performed a principal component analysis based on all 11 microsatellite loci using the program PCAGen [43] of individual genotypes. 10.000 iterations were used to obtain p-values for the axes found. As the program PCAGen only accepts diploid gen-

otypes we re-coded the triploid genotypes that were based on 11 microsatellite loci as 16 diploid loci treating the alleles as independent units (e.g. 010203 040506 (two triploid loci) => 0102 0304 0506 (three diploid loci)). Results of the analysis did not change if loci were re-coded in a different order (e.g. starting from the last locus or random).

Competing interests

The author(s) declare that they have no competing interests.

Authors' contributions

SS carried out part of the molecular genetic and cytological analyses. KPL participated in the field data collection, carried out part of the molecular genetic and cytological analyses, performed the statistical analyses and drafted the manuscript. DKL participated in the field data collection and coordination of the study. FJGL participated in the field data collection and coordination of the study. MS conceived the study, designed and participated in its coordination, organized and participated in the field study, and drafted the manuscript. All authors read and approved the final manuscript.

Acknowledgements

We are grateful to the Mexican government for issuing permit no. 210696-213-03 to collect *P. formosa* and to Maximilian Schartl, Matthias Stoeck, Ute Hornung and Alexander Froschauer for help in the field. The technical assistance of Petra Fischer and Melanie Hidding is gratefully acknowledged. We thank Georg Schneider, Hugo Schwind and Petra Weber for breeding of the fish in the laboratory, and Monika Niklaus-Ruiz for help in preparing the manuscript. Financial support for this study was granted by the DFG (SFB 567 Mechanismen der interspezifischen Interaktion von Organismen) and Fonds der Chemischen Industrie. We thank Michael Schmid for the possibility to use the flow cytometer.

References

- Hubbs CL, Hubbs LC: **Apparent parthenogenesis in nature, in a form of fish of hybrid origin.** *Science* 1932, **76**:628-630.
- Schlupp I: **The evolutionary ecology of gynogenesis.** *Annual Review of Ecology Evolution and Systematics* 2005, **36**:399-417.
- Schlupp I, Nanda I, Dobler M, Lamatsch DK, Epplen JT, Parzefall J, Schmid M, Schartl M: **Dispensable and indispensable genes in an ameiotic fish, the Amazon molly *Poecilia formosa*.** *Cytogenet Cell Genet* 1998, **80**(1-4):193-198.
- Turner BJ: **The evolutionary genetics of a unisexual fish, *Poecilia formosa*.** In *Mechanisms of Speciation* Edited by: Barigozzi C. New York, Liss; 1982:265-305.
- Schartl M, Nanda I, Schlupp I, Wilde B, Epplen JT, Schmid M, Parzefall J: **Incorporation of Subgenomic Amounts of DNA as Compensation for Mutational Load in a Gynogenetic Fish.** *Nature* 1995, **373**(6509):68-71.
- Balsano JS, Abramoff P, Darnell RM: **Electrophoretic Evidence of Triploidy Associated with Populations of Gynogenetic Teleost *Poecilia-Formosa*.** *Copeia* 1972:292-297.
- Rasch EM, Darnell RM, Kallman KD, Abramoff P: **Cytophotometric evidence for triploidy in hybrids of the gynogenetic fish, *Poecilia formosa*.** *J Exp Zool* 1965, **160**:155-170.
- Schultz RJ, Kallman KD: **Triploid Hybrids between the All-female Teleost *Poecilia formosa* and *Poecilia sphenops*.** *Nature* 1968, **219**:280-282.
- Nanda I, Schlupp I, Lamatsch D, Lampert K, Schmid M, Schartl M: **Stable Inheritance of host species-derived microchromosomes in the gynogenetic fish, *Poecilia formosa*.** 2007:submitted.
- Lamatsch DK, Nanda I, Schlupp I, Epplen JT, Schmid M, Schartl M: **Distribution and stability of supernumerary microchromosomes in natural populations of the Amazon molly, *Poecilia formosa*.** *Cytogenet Genome Res* 2004, **106**(2-4):189-194.
- Nanda I, Schartl M, Feichtinger W, Schlupp I, Parzefall J, Schmid M: **Chromosomal evidence for laboratory synthesis of a triploid hybrid between the gynogenetic teleost *Poecilia formosa* and its host species.** *J Fish Biol* 1995, **47**:619-623.
- Lampert KP, Lamatsch DK, Epplen JT, Schartl M: **Evidence for a monophyletic origin of triploid clones of the Amazon molly, *Poecilia formosa*.** *Evolution Int J Org Evolution* 2005, **59**(4):881-889.
- Rasch EM, Balsano JS: **Trihybrids related to the unisexual molly fish, *Poecilia formosa*.** In *Evolution and Ecology of Unisexual Vertebrates* Edited by: Dawley RM, Bogart JB. Albany, NY, New York State Museum; 1989.
- Balsano JS, Rasch EM, Monaco PJ: **The Evolutionary Ecology of *Poecilia formosa* and Its Triploid Associate.** In *Ecology and Evolution of Livebearing Fishes (Poeciliidae)* Edited by: Meffe GK, Snelson FFJ. Englewood Cliffs, Prentice Hall; 1989:277-297.
- Turner BJ, Elder JF, Laughlin TF, Davis WP: **Genetic-Variation in Clonal Vertebrates Detected by Simple-Sequence DNA Fingerprinting.** *Proceedings of the National Academy of Sciences of the United States of America* 1990, **87**(15):5653-5657.
- Menken SB, Smit E, Den Nijs HJCM: **Genetical Population Structure in Plants: Gene Flow between Diploid Sexual and Triploid Asexual Dandelions (*Taraxacum* Section *Ruderalia*).** In *Evolution Volume 49 (6)*. The Society for the Study of Evolution; 1995:1108-1118.
- Goldstein DB, Riuz Linares A, Cacalli-Sforza LL, Feldman MW: **An Evaluation of Genetic Distances for Use With Microsatellite Loci.** *Genetics* 1995, **139**:463-471.
- Turner BJ, Balsano JS, Monaco PJ, Rasch EM: **Clonal Diversity and Evolutionary Dynamics in a Diploid-Triploid Breeding Complex of Unisexual Fishes (*Poecilia*).** *Evolution* 1983, **37**(4):798-809.
- Avise JC, Trexler JC, Travis J, Nelson WS: ***Poecilia mexicana* is the recent female parent of the unisexual fish *P. formosa*.** *Evolution* 1991, **46**(6):1530-1533.
- Schartl M, Wilde B, Schlupp I, Parzefall J: **Evolutionary origin of a parthenoform, the Amazon molly *Poecilia formosa*, on the basis of a molecular genealogy.** *Evolution* 1995, **49**(5):827-835.
- Quattro JM, Avise JC, Vrijenhoek RC: **Mode of Origin and Sources of Genotypic Diversity in Triploid Gynogenetic Fish Clones (*Poeciliopsis*, *Poeciliidae*).** *Genetics* 1992, **130**(3):621-628.
- Kallman KD, Kazianis S: **The Genus *Xiphophorus* in Mexico and Central America.** *Zebrafish* 2006, **3**(3):271-285.
- Schlupp I, Parzefall J, Schartl M: **Biogeography of the Amazon molly, *Poecilia formosa*.** *J Biogeogr J Biogeogr* 2002, **29**(1):1-6.
- Mateos M, Vrijenhoek RC: **Independent origins of allotriploidy in the fish genus *Poeciliopsis*.** *Journal of Heredity* 2005, **96**(1):32-39.
- Janko K, Kotlik P, Rab P: **Evolutionary history of asexual hybrid loaches (*Cobitis*: *Teleostei*) inferred from phylogenetic analysis of mitochondrial DNA variation.** *Journal of Evolutionary Biology* 2003, **16**(6):1280-1287.
- Alves MJ, Coelho MM, Collares-Pereira MJ: **Evolution in action through hybridisation and polyploidy in an Iberian freshwater fish: a genetic review.** *Genetica* 2001, **111**(1-3):375-385.
- Goddard KA, Dawley RM, Dowling T: **Origin and genetic relationships of diploid, triploid, and diploid-triploid mosaic biotypes in the *Phoxinus eos-neogaeus* unisexual complex.** In *Evolution and ecology of unisexual vertebrates Volume Bull. 466*. Edited by: Dawley RM, Bogart JP. Albany: New York State Museum; 1989:268-280.
- Beukeboom LW, Vrijenhoek RC: **Evolutionary genetics and ecology of sperm-dependent parthenogenesis.** *Journal of Evolutionary Biology* 1998, **11**(6):755-782.
- Dawley RM: **An introduction to unisexual vertebrates.** In *Evolution and ecology of unisexual vertebrates* Edited by: Dawley RM, Bogart JB. Albany, NY, New York State Museum; 1989:1-18.
- Kearney M: **Hybridization, glaciation and geographical parthenogenesis.** *Trends in Ecology & Evolution* 2005, **20**(9):495-502.

31. Lundmark M: **Polyploidization, hybridization and geographical parthenogenesis.** *Trends in Ecology & Evolution* 2006, **21(1)**:9-9.
32. Stock M, Lamatsch DK, Steinlein C, Epplen JT, Grosse WR, Hock R, Klapperstuck T, Lampert KP, Scheer U, Schmid M, Schartl M: **A bisexually reproducing all-triploid vertebrate.** *Nat Genet* 2002, **30(3)**:325-328.
33. Niemeitz A, Kreutzfeldt R, Schartl M, Parzefall J, Schlupp I: **Male mating behaviour of a molly, *Poecilia latipunctata*: a third host for the asexual Amazon molly, *Poecilia formosa*.** *Acta Ethol* 2002, **5**:45-49.
34. Maynard Smith J: **The Evolution of Sex.** Cambridge University Press; 1978.
35. Van Valen L: **A new evolutionary law.** *Evolutionary Theory* 1973, **1**:1-30.
36. Lamatsch DK, Steinlein C, Schmid M, Schartl M: **Noninvasive determination of genome size and ploidy level in fishes by flow cytometry: detection of triploid *Poecilia formosa*.** *Cytometry* 2000, **39(2)**:91-95.
37. Altschmied J, Hornung U, Schlupp I, Gadau J, Kolb R, Schartl M: **Isolation of DNA suitable for PCR for field and laboratory work.** *Biotechniques* 1997, **23(2)**:228-229.
38. Walter RB, Rains JD, Russell JE, Guerra TM, Daniels C, Johnston DA, Kumar J, Wheeler A, Kelnar K, Khanolkar VA, Williams EL, Hornecker JL, Hollek L, Mamerow MM, Pedroza A, Kazianis S: **A microsatellite genetic linkage map for *xiphophorus*.** *Genetics* 2004, **168(1)**:363-372.
39. Lampert KP, Lamatsch DK, Schories S, Hopf A, De Leon FJG, Schartl M: **Microsatellites for the gynogenetic Amazon molly, *Poecilia formosa*: useful tools for detection of mutation rate, ploidy determination and overall genetic diversity.** *Journal of Genetics* 2006, **85(1)**:67-71.
40. Meyer MK, Schneider K, Radda AC, Wilde B, Schartl M: **A new species of *Poecilia*, subgenus *Mollienesia*, from upper Río Cahabón system, Guatemala, with remarks on the Nomenclature of *Mollienesia petenensis* Günther, 1866 (Teleostei: Cyprinodontiformes: Poeciliidae).** *Zool Abh Dresden* 2004, **54**:145-154.
41. Hall TA: **BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT.** *Nucl Acids Symp Ser* 1999, **41**:95-98.
42. Langella O: **Populations 1.2.28.** 1999 [http://bioinformatics.org/project/?group_id=84].
43. Goudet J: **PCA-Gen Version 1.2, Institute of Ecology, Biology Building, UNIL, Lausanne, Switzerland.** 1999 [<http://www2.unil.ch/popgen/softwares/pcagen.htm>].

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

