

Case report

Primary squamous cell carcinoma of the breast: a case report and review of literature

Youssef Seddik^{1,&}, Sami Aziz Brahmi¹, Said Afqir¹

¹Medical Oncology Department, University Hospital Mohammed VI, Oujda, Morocco

[&]Corresponding author: Youssef Seddik, Medical Oncology Department, University Hospital Mohammed VI, Oujda, Morocco

Key words: Primary squamous cell carcinoma, breast cancer, aggressive tumor

Received: 24/01/2015 - Accepted: 06/02/2015 - Published: 18/02/2015

Abstract

Primary squamous cell carcinoma is a well known malignancy of the skin and other organs composed of squamous cells, which are normally not found inside the breast. Therefore, a primary squamous cell carcinoma of the breast is an exceedingly uncommon phenomenon and the management of this type of disease is still unclear. We report the case of a 43-year-old Moroccan woman, without significant medical history, presented an infected mass of 9 cm in the left breast associated with ipsilateral axillary lymphadenopathy. The mass's surgical biopsy revealed a triple negative primary squamous cell carcinoma of the breast. She underwent a neoadjuvant chemotherapy using 5 Fluoro-Uracil and platinum. After three courses, she presented a contralateral breast progression and apparition of metastasis at D10. She received one course of a palliative chemotherapy based on weekly paclitaxel stopped because of her performance status deterioration. She died 7 months after her admission.

Pan African Medical Journal. 2015; 20:152 doi:10.11604/pamj.2015.20.152.6188

This article is available online at: <http://www.panafrican-med-journal.com/content/article/20/152/full/>

© Youssef Seddik et al. The Pan African Medical Journal - ISSN 1937-8688. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Primary squamous cell carcinoma (PSCC) is a well known malignancy of the skin and other organs composed of squamous cells, which are normally not found inside the breast. Therefore, a primary squamous cell carcinoma of the breast is an exceedingly uncommon; it represents less than 0.1% of all breast carcinomas [1]. The diagnosis is established when squamous cell carcinoma is the only malignancy found in the breast specimen, metastases from another primary are excluded, and the tumor does not originate from the skin of the breast [2]. Because of the rarity of this neoplasm, data on its diagnosis, treatment and prognosis is limited to isolated case reports and case series. We report a case of PSCC of the breast and review of the literature.

Patient and observation

We report the case of a 43-year-old Moroccan woman, without history of skin or breast cancer, nor any skin, oro-pharynx, or anal lesions. Her family history was not significant for breast cancer. She was admitted to our center eight months ago for a mass in the left breast which had been growing rapidly. A physical examination revealed a left breast infected mass, measuring 9x8x9 cm in size, with some cystic areas, associated with ipsilateral axillary lymphadenopathy (**Figure 1**). Mammography demonstrated an irregular opacity with skin infiltration, approximately 9 cm in its greatest dimension, which was classified as BI-RADS 5. The right breast was negative on palpation and mammography. The mass's surgical biopsy revealed a primary squamous cell carcinoma of the breast (**Figure 2**). Immunohistochemical evaluation was negative for estrogen receptor (ER), progesterone receptor (PgR) and HER2. Metastatic disease was ruled out via Total body CT scan, and bone scintigraphy. The serum level of tumor marker CA 15-3 was 290 U/ml (≤ 30). She underwent three courses of neoadjuvant therapy: the first course using cisplatin (at a dose of 75 mg/m² as a 1-hour intravenous infusion on day 1) and an infusion of 5-fluoro-Uracil as bolus (at a dose of 400 mg/m² per day for 3 days) every 3 weeks. She presented a grade 3 vomiting and febrile neutropenia which was treated by vancomycin with good evolution. Because of the side effects, we substitute, in the two other courses, cisplatin by Carboplatin (at an area under the curve of 5 mg per milliliter per minute, as a 1-hour intravenous infusion on day 1) and we substitute 5-fluorouracil bolus by continuous infusion (at a dose of 2400 mg/m² for 2 days) every 3 weeks. The evaluation investigations based on CT scan showed a contralateral breast progression and apparition of bone metastasis at D10 (**Figure 3**). She received a palliative chemotherapy based on weekly paclitaxel at a dose of 80 mg/m², stopped after one course because of her general performance status deterioration. She died 7 months after her admission.

Discussion

Primary squamous cell carcinoma (PSCC) of the breast is very rare. It represents less than 0.1 % of all breast carcinomas [1]. Diagnosis is retained when no other neoplastic components such as ductal or mesenchymal elements are present in the lesion, the tumor origin is independent of the overlying skin and nipple and absence of an associated primary Squamous cell carcinoma in a second site (oral cavity bronchus, esophagus, renal pelvis, bladder, ovary, and cervix) [2]. The histogenesis of this malignancy is controversial [3]. It has been suggested that it may be a very extreme form of squamous

cell metaplasia, developing into an adenocarcinoma. Squamous cell metaplasia is also seen in cysts, chronic inflammations, abscesses and adenofibromas. Therefore, PSCC is divided into pure squamous cell carcinoma and that mixed with adenocarcinoma. PSCC of the breast has been diagnosed in adult women of ages ranging from 29 years to 90 years [3]. No case was described in the male. The PSCC tend to be relatively large (> 4 cm) at diagnosis and cystic in 50% of the cases [1]. Some authors have noted only rare involvement of the lymph nodes [1-4]. There are no typical findings on the mammogram, ultrasound may show a complicated cyst or an inflammatory process [5]. Our patient was 43 year old, without significant medical history, she had an infected mass of 9cm associated with ipsilateral axillary lymphadenopathy, without typical findings on the mammogram. A surgical biopsy procedure is usually required to establish the diagnosis of squamous cell carcinoma which is generally characterized by histologically identified keratinization and intercellular bridges [6]. The breast Squamous cell carcinoma is generally a high-grade and triple negative tumor (ER, PR and HER2-negative) [1,3,7-9]. The immunohistology of our case is consistent with those findings. Ruohong Shui and colleagues [10] reported three cases of primary squamous cell carcinoma of the breast with an unusual "basal-HER2" phenotype. This tumor overexpresses usually the Epidermal Growth Factor Receptor EGFR [11]. There are no treatment recommendations because of the rarity of this distinct type of breast cancer. Most patients undergo a mastectomy with lymph nodes dissection if possible. Conservation surgery is not usually feasible because of the locally advanced of the disease [12]. Adjuvant Chemotherapy is not usually been used. Rostock et al. suggests that PSCC of the breast is not sensitive to chemotherapeutic agents commonly used for ductal carcinoma such as methotrexate, cyclophosphamide, 5-fluoro-Uracil and anthracycline [13]. Hennessy et al [12] also reported no benefit to neoadjuvant therapy using anthracycline/taxane-based regimens. In contrast, a remarkable response was reported in a patient who received neoadjuvant therapy using cisplatin and 5-fluoro-Uracil. Our patient had progression after chemotherapy regimen based on 5-fluoro-Uracil and platinum. A good response on metastatic disease has been reported in one patient who received cisplatin and 5-FU, but this has never been investigated in other report [14]. Radiation therapy has been shown little benefit, despite that squamous cell carcinoma are generally radio sensitive [3]. The PSCC of the breast is usually a hormone receptor and HER2 -negative tumor [10] this means that hormone based therapy and HER-2 targeted therapy may not be effective in this tumor. The high frequency of EGFR positivity is interesting [11] the use of anti-EGFR therapy should be explored. The prognosis of this malignancy remains pejorative with a five years overall survival of 51% +/-13% [15] the short period from diagnosis and the death of our patient (7 months) demonstrates that this tumor is very aggressive.

Conclusion

Primary squamous cell carcinoma of the breast is a rare and aggressive disease often treatment-refractory. Clinical trials including large series of this tumor are needed to increase our knowledge and to improve patient's outcome.

Competing interests

The authors declare no competing interests.

Authors' contributions

Youssef Seddik contributed to the treatment of the patient and wrote the article. **Sami Aziz Brahmi** and **Said Afqir** contributed not only to the treatment of the patient but also to the analysis and interpretation of the case. All authors read and approved the final manuscript.

Acknowledgements

Authors acknowledge Dr. Benmira Khalil who provides us the figure of histological examination.

Figures

Figure 1: a left breast infected mass, measuring 9x8x9 cm in size, with some cystic areas

Figure 2: squamous cell carcinoma of the breast

Figure 3: CT scan showed a contralateral breast progression and apparition bone metastasis at D10

References

1. Gupta G, Malani AK, Weigand RT, Rangenini G. Pure primary squamous cell carcinoma of the breast: a rare presentation and clinicopathologic comparison with usual ductal carcinoma of the breast. *Pathol Res Pract*. 2006 Feb; 23 (6):465-9. **PubMed | Google Scholar**
2. Macia M, Ces JA, Becerra E, Novo A. Pure squamous cell carcinoma of the breast: report of a case diagnosed by aspiration cytology. *Acta Cytol*. 1989 Mar-Apr; 33(2):201-4. **PubMed | Google Scholar**
3. Behranwala KA, Nasiri N, Abdullah N et al. Squamous cell carcinoma of the breast: clinico-pathologic implications and outcome. *Eur J Surg Oncol*. 2003 May; 29(4):386-9. **PubMed | Google Scholar**
4. Peison B, Benish B, Chung-Loyh. Pure Squamous Cell Carcinoma of the breast. *N J Med*. 1991 Apr; 88(4):273-5. **PubMed | Google Scholar**
5. Damin AP, Nascimento FC, Andreola JB, Cerutti TH, Roehe A, Damin DC. Primary epidermoid carcinoma of the breast presenting as a breast abscess and sepsis. *Sao Paulo Med J*. 2011 Dec; 129(6):424-7. **PubMed | Google Scholar**
6. Yamaguchi R, Horii R, Maeda I, et al. Clinicopathologic study of 53 metaplastic breast carcinomas: their elements and prognostic implications. *Hum Pathol*. 2010 May;41(5):679-8. **PubMed | Google Scholar**
7. Siegelmann-Danieli N, Murphy TJ, Meschter SC et al. Primary pure squamous cell carcinoma of the breast. *Clin Breast Cancer*. 2005 Aug;6(3):270-2. **PubMed | Google Scholar**
8. Cardoso F, Leal C, Meira A, Azevedo R, Mauricio MJ, Leal Da Silva JM, Lopes C, Pinto Ferreira E. Squamous cell carcinoma of the breast. *Breast*. 2000 Dec; 9(6):315-9. **PubMed | Google Scholar**
9. Wrightson WR, Edwards MJ, McMasters KM. Primary squamous cell carcinoma of the breast presenting as a breast abscess. *Am Surg*. 1999 Dec; 65(12):1153-5. **PubMed | Google Scholar**
10. Ruohong Shui, Anqi Li, Fei Yang, Xiaoyan Zhou, Baohua Yu, Xiaoli Xu, Wentao Yang. Primary squamous cell carcinoma of the breast with unusual basal-HER2 phenotype. *Int J Clin Exp Pathol*. 2014 Jul; 7(8):5203-5209. **PubMed | Google Scholar**
11. Grenier J, Soria JC, Mathieu MC, Andre F, Abdelmoula S, Velasco V, Morat L, Besse B, Du-nant A, Spielmann M and Delalogue S. Differential immunohistochemical and biological profile of squamous cell carcinoma of the breast. *Anticancer Res*. 2007 Jan-Feb; 27(1B):547-55. **PubMed | Google Scholar**
12. Hennessy BT, Krishnamurthy S, Giordano S, Buchholz TA, Kau SW, Zhigang D. Squamous cell carcinoma of the breast. *J Clin Oncol*. 2005 Nov 1; 23(31):7827-35. **PubMed | Google Scholar**
13. Rostock RA, Bauer TW, Eggleston JC. Primary squamous carcinoma of the breast: A review. *Breast*. 1984; 10:27-31. **PubMed | Google Scholar**
14. Hiramatsu K, Kato K, Hirata A, Matsuba H, Hara T, Ito T, et al. A resected case of squamous cell carcinoma of the breast successfully treated by FU plus cisplatin (CDDP) adjuvant therapy against recurrent metastases. *Gan To Kagaku Ryoho*. 2007 Mar;34(3):443-6. **PubMed | Google Scholar**
15. Nayak A, Wu Y, Gilcrease MZ. Primary Squamous Cell Carcinoma of the Breast: Predictors of Locoregional Recurrence and Overall Survival. *Am J Surg Pathol*. 2013 Jun;37(6):867-73. **PubMed | Google Scholar**



Figure 1: a left breast infected mass, measuring 9x8x9 cm in size, with some cystic areas




Figure 2: squamous cell carcinoma of the breast




Figure 3: CT scan showed a contralateral breast progression and apparition bone metastatsis at D10