



LETTER TO THE EDITOR

Open Access

# Stress ulcer prophylaxis in critically ill poisoned patients

Omid Mehrpour<sup>1,2\*</sup>, Bita Dadpour<sup>2</sup> and Nasim Zamani<sup>3</sup>

## Letter to the editor

Management of critically ill patients in the intensive care unit (ICU) is an important issue in medicine [1,2]. These patients may have clinically important bleeding, which is associated with an increased mortality rate. Mechanical ventilation may also increase the risk of bleeding in the upper gastrointestinal system. Sucralfate, histamine-2-receptor antagonists, and proton pumps inhibitors (PPIs) are widely used for stress ulcer prophylaxis in critically ill adults [3]. Available data shows that PPIs are safe and efficacious for increasing intragastric pH in critically ill patients [4]. Since the superiority of the PPIs over histamine-2-receptor antagonists or sucralfate is not yet proved, they should only be used as an alternative to these agents.

Poisoned patients may need to be admitted in ICU because of their critical condition. Poisoning with fatal agents such as aluminum phosphide, paraquat, organophosphate, or massive pharmaceutical drug overdoses usually needs ICU care and sometimes mechanical ventilation [5,6]. A question is therefore arisen: which of these agents is better to be used for ulcer prophylaxis in poisoned patients?

Affection of the cytochrome P450 (CYP) by toxins or pharmaceutical drugs is an important point in answering this question. Generally, most of the toxicants have a liver metabolism [7]. The number of the known P450 enzyme inducers or inhibitors exceeds 1000, but P450 3A4 is the most frequent CYP in the human liver and is known to metabolize the majority of drugs whose biotransformation is known [8]. So, inhibition or induction of CYP may increase or decrease the final metabolites of the toxicants and may have a value in the treatment of poisoned patients in the ICU.

Histamine-2-receptor antagonists are known as CYP inhibitors [9] and PPIs are inducers of CYP [8]. PPIs can therefore be recommended for the intoxications in which a CYP inhibitor is involved (such as antidepressant intoxication) or the medications which are toxic in origin but induce non-toxic metabolites. Histamine-2 receptor antagonists may be more useful in the poisonings with CYP inducers (such as carbamazepine or Phenobarbital toxicities) or in cases whose metabolites are more toxic than the parent medication (such as acetaminophen, tramadol, etc.). This may affect the patients' prognosis as well as their hospital stay. We, therefore, would like to encourage toxicologists to provide studies to evaluate such potential benefits of each type of these ulcer protectors in each poisoning.

## Competing interests

The authors declare that they have no competing interests.

## Authors' contributions

BD and NZ did bibliography and drafted the article. OM gave the idea and completed/edited/revised the article. All authors read and approved the final manuscript.

## Author details

<sup>1</sup>Medical Toxicology and Drug Abuse Research Center (MTDRC), Birjand, University of Medical Science, Pasdaran Avenue, Birjand, Iran. <sup>2</sup>Addiction Research Centre, Mashhad University of Medical Toxicology, Mashhad, Iran.

<sup>3</sup>Department of Clinical Toxicology, Loghman Hakim Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Received: 23 November 2012 Accepted: 24 November 2012

Published: 27 November 2012

## References

1. Honarmand H, Abdollahi M, Ahmadi A, Javadi MR, Khoshaiand MR, Tabeeifar H, Mousavi S, Mahmoudi L, Radfar M, Najafi A, Mojtabahedzadeh M: Randomized trial of the effect of intravenous paracetamol on inflammatory biomarkers and outcome in febrile critically ill adults. *DARU* 2012, 20:12.
2. Mirrahimi B, Hamishehkar H, Ahmadi A, Mirjalili MR, Aghamohammadi M, Najafi A, Abdollahi M, Mojtabahedzadeh M: The efficacy of magnesium sulfate loading on microalbuminuria following SIRS: One step forward in dosing. *DARU* 2012, 20:7.
3. Steinberg KP: Stress-related mucosal disease in the critically ill patient: risk factors and strategies to prevent stress-related bleeding in the intensive care unit. *Crit Care Med* 2002, 30(6 Suppl):S362-S364.
4. Jung R, MacLaren R: Proton-pump inhibitors for stress ulcer prophylaxis in critically ill patients. *Ann Pharmacother* 2002, 36(12):1929-1937.

\* Correspondence: omid.mehrpor@yahoo.com.au

<sup>1</sup>Medical Toxicology and Drug Abuse Research Center (MTDRC), Birjand, University of Medical Science, Pasdaran Avenue, Birjand, Iran

<sup>2</sup>Addiction Research Centre, Mashhad University of Medical Toxicology, Mashhad, Iran

Full list of author information is available at the end of the article

5. Taghaddosinejad F, Sheikhzadi A, Yaghmaei A, Mehrpour O, Schwake L: Epidemiology and treatment of severe poisoning in the intensive care unit: lessons from a One-year prospective observational study. *J Clinic Toxicol* 2012, S1:007. doi:10.4172/2161-0495.S1-007.
6. Mehrpour O, Jafarzadeh M, Abdollahi M: A systematic review of aluminium phosphide poisoning. *Arh Hig Rada Toksikol* 2012, 63:61–73.
7. McGraw J, Waller D: Cytochrome P450 variations in different ethnic populations. *Expert Opin Drug Metab Toxicol* 2012, 8(3):371–382.
8. Masubuchi N, Li AP, Okazaki O: An evaluation of the cytochrome P450 induction potential of pantoprazole in primary human hepatocytes. *Chem Biol Interact* 1998, 114:1–13.
9. Petroianu GA, Hasan MY, Nurulain SM, Arafat K, Sha Ullah M, Naseer O: Protective agents in acute high-dose organophosphate exposure: comparison of ranitidine with pralidoxime in rats. *J Appl Toxicol* 2005, 25:68–73.

doi:10.1186/2008-2231-20-80

Cite this article as: Mehrpour et al.: Stress ulcer prophylaxis in critically ill poisoned patients. *DARU Journal of Pharmaceutical Sciences* 2012 20:80.

Submit your next manuscript to BioMed Central  
and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

