

REVIEW

The chondrodystrophic dog: A clinically relevant intermediate-sized animal model for the study of intervertebral disc-associated spinal pain

Kelly Thompson¹ | Sarah Moore¹ | Shirley Tang² | Matthew Wiet² | Devina Purmessur^{2,3} 

¹Department of Veterinary Clinical Sciences, The Ohio State University, Columbus, Ohio

²Department of Biomedical Engineering, College of Engineering, The Ohio State University, Columbus, Ohio

³Department of Orthopedics, College of Medicine, The Ohio State University, Columbus, Ohio

Correspondence

Devina Purmessur, Department of Biomedical Engineering, The Ohio State University, Columbus Ohio, Davis Heart and Lung Research Institute, 473 W 12th Avenue, Columbus, OH 43210
Email: devina.purmessur@osumc.edu

Funding information

National Institutes of Health, Grant/Award Number: 1R01AR064157

Low back pain (LBP) is the leading cause of disability worldwide, with an estimated 80% of the American population suffering from a painful back condition at some point during their lives. The most common cause of LBP is intervertebral disc (IVD) degeneration (IVDD), a condition that can be difficult to treat, either surgically or medically, with current available therapies. Thus, understanding the pathological mechanisms of IVDD and developing novel treatments are critical for improving outcome and quality of life in people living with LBP. While experimental animal models provide valuable mechanistic insight, each model has limitations that complicate translation to the clinical setting. This review focuses on the chondrodystrophic canine clinical model of IVDD as a promising model to assess IVD-associated spinal pain and translational therapeutic strategies for LBP. The canine IVD, while smaller in size than human, goat, ovine, and bovine IVDs, is larger than most other small animal IVDD models and undergoes maturational changes similar to those of the human IVD. Furthermore, both dogs and humans develop painful IVDD as a spontaneous process, resulting in similar characteristic pathologies and clinical signs. Future exploration of the canine model as a model of IVD-associated spinal pain and biological treatments using the canine clinical model will further demonstrate its translational capabilities with the added ethical benefit of treating an existing veterinary patient population with IVDD.

KEYWORDS

degeneration, pain, preclinical models

1 | INTRODUCTION

The Global Burden of Disease study of 2015 identified lower back pain (LBP) as the leading cause of disability worldwide.¹ Indeed, it is estimated that 80% of Americans will suffer from a painful back condition at some point during their lives.¹ The most common cause is degeneration of the intervertebral disc (IVD), with a prevalence of 39%-42% in LBP patients.² IVD degeneration (IVDD) can occur as a natural part of the aging process, or as a consequence to cell-mediated responses to structural failure.^{3,4} Medical management of LBP is primarily focused on the treatment of symptomatic pain, which may alleviate clinical signs but does not restore IVD function. Surgical

management strategies can address structural problems with the disc, but have failure rates as high as 40% due to incomplete decompression, spinal instability, alteration of the vertebral column's biomechanics, epidural scar tissue formation or iatrogenic nerve injury.^{5,6} Thus, there is a critical need to investigate the pathophysiology of painful IVDD and resultant LBP in order to develop safe, successful and less invasive clinical treatments.

Experimental animal models are vital for understanding the mechanisms of painful IVDD and for the early stages of developing therapeutic strategies; however, these models have some limitations that may impact their predictive value when translating an intervention from the laboratory to the clinical setting. Of recent interest from

the translation perspective is the canine “chondrodystrophic” (CD) model of IVDD, which is a spontaneous clinical model of IVDD occurring in pet dogs. IVDD affects upwards of 20% of certain breeds of dogs such as the miniature dachshund.⁷ In North America alone, between 20,000 and 30,000 cases of spontaneously occurring IVDD in pet dogs are managed by veterinary spinal specialists each year.⁸ As a result, the canine IVDD population is highly amenable to large scale veterinary clinical studies which can be conducted with adherence to CONSORT guidelines, National Institutes of Health (NIH) standards for scientific transparency and rigor, and can closely recapitulate a human clinical trial condition.^{9,10} Of significant note, unlike most other animal models, dogs with spontaneous IVDD commonly present to the veterinary clinic with IVDD-associated pain.¹¹ Degeneration of the canine IVD also occurs spontaneously, making pathologic processes associated with IVDD in this species highly relevant to the human condition. Additionally, the size of a typical dog with spontaneous IVDD addresses many “scaling up” issues encountered in the direct translation of therapies from rodents to humans. Lastly, the use of pet dogs with spontaneous IVDD can contribute to reduction, refinement, and replacement of experimental animal IVD models (3R principles) by providing an ethically responsible source of nucleus pulposus (NP) material collected during surgical dissection and cells, tissues or whole discs collected during autopsy with consent from owners of IVDD-affected dogs. For all of these reasons, there has been a recent increase in the interest of using spontaneously occurring IVDD in pet dogs as a unique model through which to conduct properly designed veterinary preclinical studies for IVDD treatment prior to entering the human clinical setting.

This narrative review focuses on models of painful IVDD with an emphasis on the utility of the CD canine clinical model of IVDD as a potential translational research tool for studies investigating IVD-associated spinal pain. It compares the major structural, cell, and tissue level characteristics across species in both health and disease together with pain-associated behavior. Lastly, it reviews current treatment strategies that have utilized cells, tissue, and whole IVDs from CD dogs and outlines areas of opportunity for further exploration of the clinical canine model of painful IVDD to maximize its value as a translational model.

2 | THE HEALTHY IVD

The general structure of the IVD is well-conserved across vertebrate species, and serves to counteract the compressive forces of the body that are placed upon the vertebral column while also acting as joints to promote motion and flexibility.¹² Under healthy conditions, the IVD is composed of the annulus fibrosus (AF), which consists of concentric lamellae of collagen type I, elastin, and fibroblast-like cells, and encloses the proteoglycan-rich NP.¹³ The NP and AF are formed from the embryonic notochord and surrounding mesenchyme, respectively, during development of the IVD.^{13,14} The dense collagen lamellar structure of the AF provides tensile strength and stability to the vertebral column by contributing to overall rigidity.¹⁵ The NP is a hydrated mixture of cells, aggrecan and collagen type II.¹⁶ Aggrecan helps to imbibe water, creating a gelatinous NP core that distributes load across the disc and absorbs compressive forces.¹⁷ The

transitional zone (TZ) or inner AF region separates the AF from the NP in the mature IVD.¹⁸ The IVD is avascular, relying on the permeability of the adjacent cartilaginous endplates (CEPs) to receive nutrition.^{19–22} The CEP is a layer of hyaline cartilage covering the vertebral bodies that isolates the disc from the rest of the vertebral column.²² In addition to being avascular, the healthy IVD is aneural, and can be viewed as a site of immune-privilege.¹¹

3 | DEGENERATION OF THE INTERVERTEBRAL DISC

3.1 | Growth and maturation

Figure 1 demonstrates maturation of the IVD, from the young IVD (Figure 1A) through the early (Figure 1B) and late stages of IVDD (Figure 1C-E). Many species such as rodents, pigs, rabbits and non-CD (NCD) dogs have IVDs that retain notochordal cells throughout adulthood.^{11,23} However, during growth or adolescence, the notochordal cells within the NP of the human and CD canine IVD, as well as intermediate and large sized animals such as sheep, cattle, and goat are replaced with chondrocyte-like cells (CLCs)^{24,25} as seen in the gelatinous and less translucent NP in Figure 1B. CD dogs typically lose their notochordal cells as they reach skeletal maturity, usually within the first year of life, while humans lose theirs by approximately 10 years of age.²⁶ Since notochordal cells protect the NP from degradation and apoptosis, their loss is associated with an imbalance in matrix turnover and catabolism of the IVD.²⁷ In contrast, rodents, rabbits and most NCD dogs retain a population of notochordal cells within the IVD and thus maintain a healthy balance of matrix turnover, resulting in preservation of the structural integrity and function of the IVD.²⁷ While there is an association between a loss of notochordal cells and disc degeneration, this has not been unequivocally established and there are likely additional factors (ie, genetics, lifestyle, comorbidities) that are involved.

3.2 | Mild changes

Proteoglycan content of the NP is maximized during young adulthood and begins a slow and steady decline shortly thereafter due to increased fragmentation and simultaneous increase in collagen content.^{20,27} The demarcation between the NP and AF becomes less apparent as the collagen fibrils of the AF encroach upon the NP (Figure 1C). During this transition, the overall cellularity of the IVD declines and the disc becomes more fibrous with increased collagen deposition.²⁸

3.3 | Moderate and late-stage changes

The avascular nature of the IVD causes it to lack the innate ability to appropriately heal and repair.^{11,21} Changes in the NP as a result of maturation affect the biomechanics of the IVD and ultimately lead to “wear and tear,” whereby continued strenuous activities enhance matrix degradation over synthesis.²⁹ This inability to counteract compressive forces places additional strain on the AF, leads to AF degeneration including cleft and crack formation, and reduces its contributions to the overall strength of the vertebral column (Figure 1D).¹⁷ Aging also leads to calcification of the CEP, which reduces porosity within the CEP




FIGURE 1 Intervertebral disc (IVD) maturation from young to early and late stage IVDD where the first column shows illustrative representations throughout the stages (A-E), middle column shows Pfirrmann grading via IVD magnetic resonance imaging (MRI) and last column showing Thompson grading of canine IVD. AF, annulus fibrosus; CEP, cartilaginous end plate; NP, nucleus pulposus. Pfirrmann grade and Thompson grade images adapted with permission from Bergknut et al. American Journal of Veterinary Research, 2011;72:899

structure itself and decreases disc access to nearby blood vessels and nutrition, enhancing a decline in cellularity. Lactic acid accumulation alters the cellular microenvironment within the disc, slowing matrix production but not matrix enzyme activity.³⁰

Upregulation of inflammatory and catabolic responses in degeneration can lead to increased expression of vascular endothelial growth factor (VEGF) and nerve growth factor (NGF), resulting in the invasion of nerves and blood vessels (Figure 1E).³¹ These changes in disc environment lead to changes in structure function as well as contribute to pain mechanisms.^{11,32}

4 | EXPERIMENTAL ANIMAL MODELS OF IVDD

Experimental models have provided valuable insight into the mechanisms underlying disc degeneration as well as the ability to assess therapeutic interventions. Popular experimental animal models of

IVDD include the mouse, rabbit, rat, pig, sheep, goat, and cattle.^{11,23,33} While these models have contributed to the current understanding of IVDD pathophysiology, each presents a unique set of limitations that hinder the translation of experimental results to the human condition. Unlike the human IVD where notochordal cells are replaced during adolescence, mice, rabbits, and pigs retain their supply of notochordal cells through adulthood, while cows and sheep lose theirs rapidly after birth.^{34–36} These fundamental developmental differences mean that IVDD does not typically occur spontaneously and must be artificially induced through injury.¹¹ Common artificial IVD injury models include annular stab, puncture, chemical or mechanical induction, altered nutrient supply, or genetic modification.^{23,37} The physiological changes following artificial induction may differ from the spontaneous process occurring in people.¹¹ An exception to this is the spontaneous occurrence of IVDD in the sand rat; however, its small IVD size relative to the human IVD presents both logistical challenges and translational difficulties.³³

TABLE 1 Experimental animal models of LBP in comparison with human IVDs

Animal models of LBP	Experimental models									
	CD Canine ^{11,30,38,39}	NCD Canine ^{11,30,39,40}	Rodents ^{23,27,34,41-43}	Rabbits ^{27,37,41}	Cattle ^{37,41}	Pig ^{35,41}	Sand rat ^{33,44,45}	Goat ^{37,46}	Sheep ^{36,47}	
Characteristics of the human IVD in healthy and diseased										
Healthy IVD										
Hydrated NP	X	X	X	X	X	X	X	X	X	X
Fibrous AF	X	X	X	X	X	X	X	X	X	X
Large notochordal cells	X	X	X	X	X	X	X	X	X	X
Avascular/aneural	X	X	X	X	X	X	X	X	X	X
Degenerate IVD										
Occurs spontaneously	X	X	X	X	X	X	X	X	X	X
Matrix degradation	X	X	X	X	X	X	X	X	X	X
Decreased number of cells	X	X	X	X	X	X	X	X	X	X
Dimension reduction (disc height)	X	X	X	X	X	X	X	X	X	X
Nerve ingrowth	X	X	X	X	X	X	X	X	X	X
Loss of nutrition	X	X	X	X	X	X	X	X	X	X
Pain	X	X	X	X	X	X	X	X	X	X
Increased axial loading	X	X	X	X	X	X	X	X	X	X
Inflammation	X	X	X	X	X	X	X	X	X	X
Immune infiltration	-	-	X	X	X	X	X	X	X	X
Intermediate-large size	X	X	X	X	X	X	X	X	X	X
Clinical patient model	X	X	X	X	X	X	X	X	X	X

Thus, in seeking out an animal model of IVDD relevant to studying the human condition, there are a number of aspects that must be considered. While the use of bipedal animals may better mimic the biomechanics of the human IVD, research with nonhuman primates is ethically challenging and often cost prohibitive; as a result, the use of quadrupeds is much more practical. However, cows, pigs, goat, and sheep IVDs have limited motion capabilities and flexibility compared to that of the human IVD.³⁶ Animal and IVD size is also important for the translation of pathology results and calculations for therapeutic administration. Intermediate or large sized animals are often preferable, as they minimize anatomical differences as well as errors associated with scaling up of doses for use in clinical studies. Lastly, translational IVDD models should ideally offer the ability to assess and measure pain resulting from degenerative processes. While rodents are useful for pain assessment, they do not meet the previously stated criteria, and pain cannot be adequately measured in the rabbit, cow or pig models.

Table 1 summarizes the criteria necessary for modeling human IVDD with a comparison of animal models discussed in this review. As seen, the CD canine model most closely matches the human condition for reasons to be discussed in the following section. Characteristics that are consistent across species of all sizes in the healthy IVD include the avascular and aneural nature, hydrated NP, and fibrous AF. For the degenerate IVD these include matrix degradation, decreases in cellularity, alterations in disc height, inflammation, loss of nutrition, and increased axial loading. Differences between species center largely around the presence or absence of notochordal cells through adulthood, whether disease occurs spontaneously or must be induced, and whether validated objective assessments of pain exist. Large gaps include the assessment of neurovascular ingrowth as a mechanism of pain and limited number of spontaneous intermediate to large-sized animal models to study pain in the degenerate IVD.

5 | THE CANINE CLINICAL MODEL OF IVDD

5.1 | Hansen type I and type II

IVDD-associated pain and neurologic dysfunction are a common clinical problem in pet dogs.⁸ Spontaneous canine IVDD occurs in what has been classically believed to be two clinically distinct forms, described by Hansen as type I and type II IVDD.⁴⁸ Hansen type I IVDD is described as dehydration, degeneration and dystrophic calcification of the NP of the IVD, and is most commonly observed in CD dog breeds such as the dachshund, beagle, shih tzu, lhasa apso, and Pekingese. Within these breeds, the supply of notochordal cells within the NP matrix is replaced with CLCs, often as early as 2 months of age.^{48,49} Typically, CD dogs display clinical signs of IVDD between 3 and 7 years of age, when dehydration of the NP places additional stress on the dorsal AF leading to rupture and extrusion of NP matrix into the vertebral canal. The IVDs of the cervical or thoracolumbar spine are the most commonly affected sites.^{48,49} Dachshunds are the breed most commonly affected with type I IVDD, with a breed prevalence as high as 20%.⁷

IVDD-like pathologies in NCD dogs are associated more with age-related changes, as notochordal cells are typically retained within the NP throughout adulthood. However, certain large breed dogs such as the German shepherd and Labrador retriever may develop what has been termed Hansen type II IVDD later in life.^{40,50,51} This tends to occur in regions of the vertebral column susceptible to “wear and tear” such as the caudal lumbar and lumbosacral regions.¹¹ Specifically, the oblique angle created by the joint of the seventh lumbosacral vertebrae and first sacral vertebrae is subject to a larger amount of workload compared to the other areas of the spine, which makes it especially vulnerable.⁵⁰ Hansen type II IVDD has been classically described as fibroid degeneration and dorsal thickening of the AF; however, recent work suggests that the term chondroid metaplasia may be more appropriate to describe these pathologic changes.^{38,48,52} Specifically, IVDD involves replacement of notochordal cells of the NP by CLCs in both CD and NCD species. Increased collagen content of the NP causes AF fibers to split, allowing plasma and fluid from the NP to accumulate between the AF fibers. Over time, the AF thickens due to a buildup of pressure in the area and bulges dorsally into the vertebral canal, causing spinal cord or nerve root compression.^{40,50,51} Figure 2 depicts the discussed differences in the two types of Hansen herniation in canine IVDs.

IVDD by either mechanism can result in significant neurologic manifestations, including neck or back pain, radiculopathy, and in severe cases, paralysis.⁵³ It should be noted that while severe neurologic abnormalities can result, particularly from Hansen type I IVDD in approximately 10%-15% of cases, a substantial portion of dogs with IVDD present with spinal pain as the only clinical sign of IVDD.⁵⁴ While both canine spontaneous models of IVDD have potential value, Hansen type I IVDD in CD dogs may have the most direct relevance to the human condition. The pathophysiology is most similar to IVDD that occurs in people as changes are not just age-related but can be described as a “cell-mediated response to structural failure.”⁴ For this reason, the remainder of this review will focus on Hansen type I IVDD (referred to as IVDD from here on).

6 | CLINICAL RELEVANCE TO IVDD AND IVD-ASSOCIATED SPINAL PAIN

The canine model of IVDD complements experimental models by offering confirmation of promising laboratory findings in a

spontaneous model of disease. Beyond its value as a confirmatory tool, the model offers additional distinct advantages. This spontaneous model of IVDD has a high degree of biologic relevance because its onset, related to the upregulation of degradation pathways within the IVD, is similar to the pathology of human IVDD.^{11,38,39} The depletion of notochordal cells in CD dog IVDs during growth, replacement with CLCs, and subsequent increase in fibrous content provide for a similar NP composition to the mature human IVD. As a result, both the canine and human IVDs undergo similar pathophysiological changes during degeneration.^{11,38,39}

A small number of studies have begun characterizing disc degeneration in the canine model, including comparisons between the CD and NCD breeds as shown in Table 2. While these studies characterize the degenerate process in NCD and CD species they do not highlight the similarities and differences in painful pathologies between the two. Both dogs and people diagnosed with IVDD present with clinical signs or symptoms of pain. In people, LBP is often debilitating, and can occur with or without concurrent neurological deficits. Either scenario results in a substantial impact on quality of life. CD pet dogs with IVDD are frequently presented to veterinary specialists with clinical signs of spinal pain (with or without neurologic deficits) that disrupts daily activities. This similarity offers an important advantage with respect to clinical relevance of this model.^{31,48}

Because pet dogs with spontaneous IVDD represent a clinical population, they offer the ability to pursue longitudinal studies of biologic changes associated with aging and degeneration of the IVD, as well as interventional preclinical studies examining biological therapies in a genetically and environmentally diverse population that closely mirrors a human clinical trial setting.³⁰ Dogs with IVDD are typically diagnosed and managed in a similar fashion to people.⁸ Cross sectional imaging such as magnetic resonance imaging (MRI) is often performed for diagnosis, and management consists of symptomatic treatment of pain using nonsteroidal anti-inflammatory drugs or steroids, opioids, or other drugs targeted at pain and inflammation. Spinal surgery is also often performed for canine “patients” with neurologic deficits or refractory pain.⁸

The high prevalence of IVDD in certain breeds such as the dachshund provides a platform to expediently conduct preclinical studies that can be designed to adhere to CONSORT standards for human clinical trial design and reporting. For example, our institution alone manages approximately 200 cases per year of canine IVDD, most of which would be available for enrollment in treatment studies. Several




FIGURE 2 Herniation of the intervertebral disc (IVD): Hansen type I (left) and type II (right) IVDD with nucleus pulposus (NP) protrusion through and AF (annulus fibrosus) rupture in type I and protrusion of AF into the vertebral canal in type II. TZ, transitional zone

TABLE 2 Characterization models for the canine IVD

Research questions	Experimental conditions				Measurements	NCD	CD	Outcomes
	Cells	Tissue	Organ	In vivo Groups				
Effects of hypoxia on NC organization ⁵⁵	X			Monolayer and 3D; hypoxia (3.5% O ₂) and normoxia (21% O ₂)	Histology; matrix production	X		Under hypoxia NCs organize themselves and produce matrix similar to in vivo; not in normoxia
Investigated Wnt/B-catenin signalling ⁵⁶	X	X		Healthy and early degeneration	Histology; B-catenin expression; qRT-PCR for T, KRT8, axin2, cyclin D and c-myc	X	X	Dual role of B-catenin in NC-rich progenitor cells and also in early disease
Gene expression profiling of early intervertebral disc degeneration ⁵⁷	X	X		NCs; mixed NC + CLC; CLCs	Histology; microarray; qRT-PCR for T, KRT8 and Wnt target genes; B-catenin and caveolin-1 expression	X	X	Early degeneration involves down-regulation of Wnt signaling and caveolin-1 expression—Essential to physiology and preservation of NCs
Osmolarity and clustering regulate NC phenotype ⁵⁸	X			(DMEM)/F12 (300 mOsm/L; a-MEM (300 mOsm/L); a-MEM (400 mOsm/L)	NC morphology and matrix (histology); qRT-PCR for T, KRT8 and 18 and matrix genes; DNA/GAG	X		Culturing NCs in native clusters and high osmolarity media retain NC phenotype
Proteomic and biomechanical characterization ⁵⁹		X		NCD and CD	iTRAQ proteomics of secretome; western blot; histology/IHC; matrix and biomechanics	X	X	Differences in ECM proteins between species - decorin, biglycan, fibronectin, fibromodulin and HAPLN1; CD less stiff than NCD
Characterization of inflammatory profile in the healthy and degenerate canine IVD ³⁹		X		Healthy and degenerate	Levels of PGE2, cytokines, chemokines, and matrix components; histology and COX-2 expression	X	X	PGE2 and CCL2 levels in degenerated IVDs significantly higher than healthy IVDs; COX-2 increased with grade degeneration
Discectomy model of cervical disc degeneration ⁶⁰			X	Discectomy versus adjacent control IVD	Histology; MRI; radiographs		X	Discectomy induced degenerative changes; loss disc height, modic changes and sclerosis
Whole genome screening for skeletal dysplasia and disc degeneration ⁶¹	Blood	X		Skeletal dysplasia within 1 breed; IVD degeneration across multiple breeds	GWAS; genotyping; qRT-PCR; semi-qRT-PCR	X	X	FGF4 retrogene on CFA12 responsible for chondrodystrophy and IVD degeneration
Inflammatory profile of herniated canine IVDs ⁶²		X		Herniated (H), affected nonherniated (NH) disc, and adjacent nonaffected (NA) disc; control discs	qRT-PCR and protein expression of inflammatory cytokines; neurological assessment?	?	?	Gene—IL-6 and TNF α up-regulation and IL-1b down-regulation with herniation; protein expression varied for IL-6 and associated with positive outcomes; infiltration of monocytes and macrophages

recent large-scale multicenter placebo controlled randomized veterinary studies have been performed using the canine spontaneous model of IVDD.^{63,64} While these studies focused on neuroprotective strategies aimed at treating the 10%-15% of dogs with IVD that can develop severe neurologic complications rather than treatments targeted specifically at disc degeneration, they still serve as important proof of concept regarding feasibility of large-scale studies using this disease model.⁵⁴

Studies in the dog model of IVDD can be performed using clinically relevant outcome measures shared across species including quality of life assessments, measures of neuropathic pain, and

locomotor outcomes. Techniques such as owner-derived quality of life questionnaires, quantitative sensory testing (QST), various locomotor scoring systems and kinematic gait assessments have all been validated for use in dogs with IVDD.^{65,66} Because these outcome measures mirror those of human trials, positive results may be more predictive of translational success. The utility of canine translation models for the study of chronic pain has recently gained attention in the literature; however, most canine pain studies to date have focused on osteoarthritis while the potentially valuable scenario of chronic disc-associated pain in dogs with IVD has been under-explored.⁶⁷⁻⁶⁹

TABLE 3 Therapeutic models for the canine IVD

Research model	Experimental conditions							Outcomes
	Cells	Tissue	Organ	In vivo	NCD	CD	Pain	
Therapeutic models								
Cell therapies								
Autologous disc chondrocyte transplantation ⁸⁶	X			X	?	?		Cells were viable and proliferative after transplantation; produced matrix & maintained disc height
Wharton's jelly cell transplantation ⁸⁷	X			X		X		Cells were viable after 24 weeks post transplantation into degeneration induced NP of beagles. WJC treated beagles had smaller disc reduction, well preserved structure and ACAN/COL2/SOX-9 gene upregulation compared to nontreated groups.
Adipose stem cells ⁸⁸	X			X	?	?	X	Autologous adipose tissue derived stem cells promoted disc regeneration; produced matrix and maintained disc height.
BMP2 and MSCs ⁸⁹	X				X	X		BMP2 showed regenerative effects on chondrocyte-like-cells with more healthy NP matrix vs TGF-β1. Adding MSCs to BMP2 treated cells did not show additional regenerative effects on either CD or NCD.
Chondrocyte transplantation ⁹⁰	X			X	?	?	X	Cells were viable and proliferative after transplantation, produced ECM and maintained disc height. All 3 pain scores showed significant reduction of low back pain.
Transplant activate NP cells ⁹¹	X			X		X		Cryopreserved vs activated NP cells showed no difference in treatment of in-vivo canine models. Disc height was better maintained compared to nontreated groups, cells maintained viability and slowed disc degeneration.
Notochordal conditioned media								
Canine NCCM on bovine NP cells ⁹²	X				X	X		Difference in proteoglycan production was seen with different NCCM concentrations. However, there was no dose dependency of NCCM for cell proliferation. NC cells maintained phenotype in masses in suspension and in monolayer.
Canine NCCM contains CTGF and increases proteoglycan ⁹³	X				X	X		NC cells contain growth factor CTGF (upregulates aggrecan, versican, HAS-2 gene). Found no difference in CTGF gene expression in NCD vs CD canine NC cells. Study suggests CTGF as anabolic factor and dependent on population of NC cells in disc.
NCCM protects NP cells from degradation and apoptosis ⁹⁴	X				X			NC cell secreted factors prevent NP apoptosis via inhibition of caspase-9 and -3/9. Degradation prevented via upregulation of anabolic and matrix protection genes.
Molecular therapy; NCCM characterization ⁹⁵	X			X	X			Found TGF-β and CTGF to be major hubs in protein interaction networks. Treatment with TGF-β1 and CTGF in vitro promoted ECM synthesis, increased cell proliferation and decreased cell death. Injection of TGF-β1 and CTGF in rat tail injury restored NP.
Canine NC conditioned media effects on arthritic chondrocytes ⁹⁶	X				X			NCCM restored cartilage matrix production of end-stage human OA chondrocytes and suppressed production of inflammatory mediators. NCCM was age and disease dependent based on human donors >55y.o
Canine NCCM ⁹⁷	X				X	X		NCCM increased NP cell proliferation, GAG production, and increased NP phenotypic gene expression. BMS cells showed increased GAG production in NCCM but no gene level effects and did not increase GAG content in NP cells compared to NCCM alone.
Canine NCCM ⁹⁸	X				X	X		NC cells did not maintain phenotype in culture of alginate beads. NC and NP cell coculture ECM content and anabolic gene expression showed no difference. MSCs and NC coculture showed increased GAG content and Brachyury T expression.
Bioactive ligands								
BMP7 ⁹⁹	X			X	X			hBMP7 transfected NP cells injected into cryopreserved IVDs and implanted in dogs. Treated dogs maintained structural integrity of disc, ECM and biomechanical properties.
IL-10 and TGFβ ¹⁰⁰	X					X		Treatment suppressed IL-1β and TNF-α and inflammatory responses.
BMP7 ¹⁰¹	X			X		X	X	rhBMP-7 treatment in vitro increased matrix production and gene expression of ACAN and COL2A1. However, no regenerative effects were observed for in vivo treatments at IVD. Extra-discal bone formation observed.
Caveolin and repair ¹⁰²	X				X	X		Caveolin-1-null mice had collagen rich ECM and fewer NCs with high apoptosis activity compared to wild-type mice. Found high caveolin-1 expression and cell dead in degenerate canine IVDs. Yet, caveolin-1 silencing decreased GAG content but rescued by caveolin-1

(Continues)

TABLE 3 (Continued)

Research model	Experimental conditions							Outcomes
	Cells	Tissue	Organ	In vivo	NCD	CD	Pain	
Link-N ¹⁰³	X				X	X		scaffolding domain (CSD). CSD treatment also increased TGF- β /pSmad2 signaling. Human and canine link-N showed species-specific effects on chondrocyte like cells but both induced negligible GAG deposition in canine CLCs.
Hydrogels								
Disc replacement cervical spine ¹⁰⁴	X	X		X		X		Disc height retention and physiological hydration, matrix production and integration into host tissue after 16 weeks. Still lacks mechanical properties compared to native tissue.
Poly(ϵ -caprolactone-co-lactide)-b-poly(ethylene glycol)-bpoly(ϵ -caprolactone-co-lactide) hydrogel + celecoxib ¹⁰⁵	X	X		X		X	X	No adverse reaction to hydrogel injection. 9/10 dogs showed back pain reduction, 3/10 dogs had recurring pain after 3 months.
Polyester amide microspheres ¹⁰⁶	X			X		X		No degenerative changes occurred post injection of PEAM compared to nontreatment groups, good cyto-compatibility in vitro.
Poly-N-isopropylacrylamide MgFe-layered double hydroxide hydrogel and celecoxib ¹⁰⁷	X	X		X		X	X	Good biocompatibility and safe application of hydrogel. However, controlled release of CXB had only limited inhibition of PGE2 and resulted in mild IVD degeneration.

QST describes a collection of techniques used commonly in human clinical studies of LBP to quantify pain, assess sensory abnormalities, and document treatment effects. The type of stimulus evaluated using QST varies, with mechanical, thermal, and vibrational stimuli reported for use in a variety of animal disease models and in people⁷⁰⁻⁷². Several recent studies have reported the use of both mechanical QST (with an electronic von Frey anesthesiometer, von Frey filaments, or other devices) and thermal sensory testing to document sensory abnormalities in dogs with IVDD.^{73,74} As mentioned above, these studies have also focused on the small percentage of dogs with IVDD presenting with severe neurologic deficits; however, they demonstrate feasibility of QST using the canine IVDD model and suggest that translation of these

protocols to dogs presenting only with back pain is likely possible. There are also several clinical metrology instruments (CMI) that have been validated for use as owner-derived pain assessments in veterinary studies. The Canine Brief Pain Inventory (CBPI) and the Helsinki Chronic Pain Index were both developed specifically to evaluate chronic pain in dogs with osteoarthritis but have direct relevance to canine spinal pain.^{75,76} Activity monitors similar to the Fitbit are available for dogs and have been validated for monitoring step counts, active minutes, and intensity of activity.⁷⁷⁻⁷⁹ Various canine activity monitors have been used as surrogate markers of decreased mobility associated with chronic pain in dogs and lend themselves to outcomes assessment in clinical studies of canine IVDD.⁸⁰

Healthy




FIGURE 3 Comparison of the healthy and degenerate human (left) and canine intervertebral disc (IVD) (right) on the gross and molecular level with neurovascular ingrowth, decreased in chondrocyte-like cells and broken aggrecan and collagen II in the degenerate nucleus pulposus (NP)




FIGURE 4 Cells within the degenerate canine intervertebral disc (IVD) where nucleus pulposus (NP) tissue was surgically removed from herniated canine IVDs (A-D); schematic of herniated IVD (a); safranin O fast staining of cell clusters in diseased disc (B) and hematoxylin and eosin staining of red blood cells in granulation tissue suggestive of angiogenesis (C); giemsa staining of mast cells (dark blue/purple, $\times 40$) (D)

7 | CHALLENGES ASSOCIATED WITH THE CANINE MODEL

While the canine clinical model of IVDD may hold significant promise in enhancing translational efficiency, a few differences in canine and human vertebral column structure must be considered. Minor differences include number of vertebrae and overall size: the typical human vertebral column has 12 thoracic and 5 lumbar IVDs, while the canine has 13 thoracic and 7 lumbar IVD. Additionally, human CEPs are thicker than those of the dog due to expanded layers of chondrocytes.^{81,82} This results in an increased number of CEP irregularities compared to those observed dogs.¹¹ The canine vertebral growth plates close at skeletal maturity (~8 months) whereas in people, secondary ossification centers (ring apophysis) develop during teenage years then close at skeletal maturity.⁸³ The canine vertebral column contains growth plates throughout, which are responsible for the majority of vertebral growth. This species difference is relevant for histologic and imaging-based grading of IVDD and CEP changes, but is unlikely to affect translational relevance.¹⁸ Differences in spinal biomechanics between canine and human, namely the quadrupedal nature of the dog placing the spine at a horizontal nature vs the vertical human spine due to bipedalism, must also be considered. Recent investigations however, have demonstrated that the axial loading effects are similar across the vertebral column in both species, suggesting that biomechanical differences may be less relevant than might be expected.^{11,29,84} Indeed, ligaments and muscle play a key role in the stabilization of the spine both in bipeds and quadrupeds.⁸⁵

8 | EXPERIMENTAL STUDIES USING THE CANINE MODEL—BIOLOGICAL STRATEGIES AND TREATMENTS

Current available treatment options for both dogs and people experiencing IVDD are either largely conservative and aim to manage pain and discomfort, or highly invasive surgeries which do little or nothing to stop or prevent further degeneration and do not attempt disc restoration. Physiotherapy is often recommended to help both dogs and people build core strength and retain normal everyday

movements through relief of compression on the IVDD affected areas.⁸ Given the potential benefits as a translational model, several recent studies have utilized the canine model to investigate regenerative therapies for IVDD via cell therapies, notochordal conditioned media (NCCM), ligands (growth factors and gene therapy) and various hydrogels as summarized in Table 3.

Cell based therapies are a common means of targeting matrix regeneration of the IVD, including the use of autologous IVD cells, CLCs, bone marrow- and adipose-derived mesenchymal stem cells (MSCs), and differentiated cells. The studies in Table 3 have demonstrated the positive regenerative potential of CLCs via increases in cell viability, matrix production and disc height integrity; however, CLC treatment is limited in both human and canine patients based on the stage of IVDD and is most beneficial at earlier stages.^{86,90,91} Recently, the safety and feasibility of autologous bone marrow-derived MSCs to treat disc degeneration was evaluated in the canine clinical model of IVDD, which demonstrated the safety of intradiscal injection of MSC.¹⁰⁸ Small animal experiments have demonstrated degeneration reduction in the progression of IVDD post treatment with MSCs, and additional studies in the beagle model have proven useful in inhibiting IVDD, with the potential for promoting continued avascularity.¹⁰⁹ Similarly, adipose derived stem cells enhance disc regeneration⁸⁸ along with Wharton's jelly cells (WJCs) which are capable of differentiating into NP cells in coculture.⁸⁷

Significantly, there are a number of studies using NCCM on canine IVD cells, which have shown to decrease apoptosis of NP cells, increase cell proliferation, slow degradation of ECM and promote ECM synthesis.^{94,95,97,98,110} Additionally, several growth factors from notochordal cells have been identified, including TGF- $\beta 1$ and CTGF that aid in matrix production.^{89,92,95} Growth factor treatments such as GDFs/BMPs including caveolin also promote matrix biosynthesis resulting in regeneration of healthy disc tissue.^{89,102,111} Furthermore, with regards to gene therapy, exploration of the immune and inflammatory responses and pathways that follow can function as targets to slow the degenerative process of the IVD.^{100,103}

Within the field of tissue engineering, several cell seeded hydrogels have been proposed to treat IVDD, which include replacement of the entire or part of the disc in vivo or mainly for use as drug control release systems. Some examples are biocompatible hydrogels

such as PCLA-PEG-PCLA (poly(ϵ -caprolactone-co-lactide)-b-poly(ethylene glycol)-b-poly(ϵ -caprolactone-co-lactide)), agarose, and polyester amides.^{104–106} Additionally, hydrogels have also been used to encapsulate drugs for delivery into the IVD.^{105–107}

These various therapies, as summarized, demonstrate the potential of the CD dog model as a clinically relevant model to evaluate regenerative IVDD therapies which can later translate to human conditions. However, unexplored is the potential of using organ culture as a tool to investigate and screen biological therapies for IVDD.

9 | OPPORTUNITIES FOR FUTURE EXPLORATION

When critically evaluating canine spontaneous IVDD for suitability as an animal model of IVDD, what remains largely unknown is the contribution of immune and inflammatory responses to the disease process in dogs. Degeneration of the human IVD has been shown to trigger responses that elicit an infiltration of mast cells (MCs) and macrophages (M ϕ s) into the disc, particularly in injured areas and regions of granulation.¹¹² MCs are involved in the body's first responses to injury, whereby they release granules of bioactive ligands to stimulate healing and repair.^{112,113} Likewise, when propagated by injury, M ϕ s release cytokines such as TNF- α , IL-1 β , and prostaglandins.³⁹ These cytokines upregulate catabolic processes that breakdown IVD matrix and limit regenerative processes. The specific roles of MCs and M ϕ s in the pathogenesis of IVDD in people and dogs, however, remain to be elucidated. Coupled with these responses are pain predictors such as substance P and NGF, which can facilitate nerve ingrowth and pain experienced by canine and human patients with IVDD.^{31,114} Further studies are required to better understand whether the inflammatory responses observed in human IVDD are modeled by the canine clinical disease.

10 | ARE CHONDRODYSSTROPHIC DOGS A GOOD TRANSLATIONAL MODEL FOR HUMAN IVDD?

This review summarizes and evaluates the criteria necessary for qualifying animal models of IVDD for use in translational application to the human condition, as is visible in Table 1. As summarized in Figure 3, the biochemical and cellular composition of the IVD is similar across the CD canine and human species. Aside from thinner CEPs and smaller overall size of the canine IVD, the structures of the healthy discs are very similar. Using histological stains as a comparison, cell clusters and angiogenesis along with the infiltration of MCs are similar pathologies that also occur in the human degenerate IVD (Figure 4).¹¹⁵ While further studies are required to investigate the presence of neurovascular ingrowth, immune infiltration and mechanisms underlying IVD-associated pain in the canine degenerate IVD as occurs in the human, other characteristics of human IVDD are commonly present in the CD canine model. Additionally, the similarities in this clinical population as well as diagnostic and treatment

methods demonstrate the potential suitability of CD canine IVDD as a model for human IVDD.

11 | CONCLUSION

A number of experimental animal models exist that attempt to recapitulate IVDD in people. Available models are limited by notochordal cell populations, small size, and artificial induction of the disease that do not mimic spontaneous mechanisms of degeneration. This review highlights aspects of the spontaneous canine clinical model of IVDD that make it attractive as a preclinical model for translational studies. Future work should focus on defining the inflammatory and symptomatic profiles associated with painful IVDD to allow for the better understanding of how these relate to those observed in human IVDD.

ACKNOWLEDGEMENTS

The authors would like to thank Timothy Vojt for his assistance with drafting the figures for this review. The authors would like to acknowledge funding through NIH grant 1R01AR064157.

Author contributions

K.A.T., S.A.M., S.T., M.G.W., and D.P. contributed to drafting the article or revising it critically for important intellectual content and final approval of the version to be submitted.

Conflict of interest

The authors have no conflicts to declare.

ORCID

Devina Purmessur  <http://orcid.org/0000-0002-7445-1678>

REFERENCES

1. Vos T, Barber RM, Bell B, et al. Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease study 2013. *Lancet*. 2015;386:743–800.
2. Schwarzer A, Aprill C, Derby R, Fortin J, Kine G. The prevalence and clinical features of internal disc disruption in patients with chronic low back pain. *Spine (Phila Pa 1976)*. 1995;20(17):1878–1883.
3. Malik K, Cohen S, Walega D, Benzon H. Diagnostic criteria and treatment of discogenic pain: a systematic review of recent clinical literature. *Spine J*. 2013;13:1675–1689.
4. Adams MA, Roughley PJ. What is intervertebral disc degeneration, and what causes it? *Spine (Phila Pa 1976)*. 2006;31:2151–2161
5. North RB, Campbell JN, James CS, et al. Failed back surgery syndrome: 5-year follow-up in 102 patients undergoing repeated operation. *Neurosurgery*. 1991;28:685–691.
6. Fritsch EW, Heisel J, Rupp S. The failed back surgery syndrome: reasons, intraoperative findings, and long-term results: a report of 182 operative treatments. *Spine*. 1996;21:626–633.
7. Lappalainen AK, Vaittinen E, Junnila J, Laitinen-Vapaavuori O. Intervertebral disc calcifications intervertebral disc disease in dachshunds radiographically screened for intervertebral disc calcifications. *Acta Vet Scand*. 2014;56:89.

8. Moore SA, Early PJ, Hettlich BF. Practice patterns in the management of acute intervertebral disc herniation in dogs. *J Small Anim Pract.* 2016;57:409-415.
9. Altman DG, Schulz KF, Moher D, et al. The revised CONSORT statement for reporting randomized trials: explanation and elaboration. *Ann Intern Med.* 2001;134:663-694.
10. Filipovich AH, Weisdorf D, Pavletic S, et al. National Institutes of Health consensus development project on criteria for clinical trials in chronic graft-versus-host disease: I. Diagnosis and staging working group report. *Biol Blood Marrow Transplant.* 2005;11:945-956.
11. Bergknut N, Rutges JPHJ, Kranenburg HC. The dog as an animal model for intervertebral disc degeneration? *Spine (Phila Pa 1976).* 2011;37:351-358.
12. Boszczyk BM, Boszczyk AA, Putz R. Comparative and functional anatomy of the mammalian lumbar spine. *Anat Rec.* 2001;264:157-168.
13. Ghosh P. *The Biology of the Intervertebral Disc.* CRC Press Boca Raton, FL 1988.
14. Smith LJ, Nerurkar NL, Choi K-S, Harfe BD, Elliott DM. Degeneration and regeneration of the intervertebral disc: lessons from development. *Dis Model Mech.* 2010;4(1):31-41.
15. Melrose J, Ghosh P, Taylor TK. A comparative analysis of the differential spatial and temporal distributions of the large (aggrecan, versican) and small (decorin, biglycan, fibromodulin) proteoglycans of the intervertebral disc. *J Anat.* 2001;198:3-15.
16. Trout JJ, Buckwalter JA, Moore KC, Landas SK. Ultrastructure of the human intervertebral disc. I. Changes in notochordal cells with age. *Tissue Cell.* 1982;14:359-369.
17. Meij BP, Bergknut N. Degenerative lumbosacral stenosis in dogs. *Vet Clin Small Anim.* 2010;40:983-1009.
18. Bergknut N, Meij BP, Hagman R, et al. Intervertebral disc disease in dogs—part 1: a new histological grading scheme for classification of intervertebral disc degeneration in dogs. *Vet J.* 2013;195:156-163.
19. Trout JJ, Buckwalter J a, Moore KC. Ultrastructure of the human intervertebral disc. II. Cells of the nucleus pulposus. *Anat Rec.* 1982;204:307-314.
20. Buckwalter JA. Aging and degeneration of the human intervertebral disc. *Spine (Phila Pa 1976).* 1995;20:1307-1314.
21. Bibby SRS, Jones DA, Ripley RM, Urban JPG. Metabolism of the intervertebral disc: effects of low levels of oxygen, glucose, and pH on rates of energy metabolism of bovine nucleus pulposus cells. *Spine (Phila Pa 1976).* 2005;30:487-496.
22. Taylor JR. Growth of human intervertebral discs and vertebral bodies. *J Anat.* 1975;120:49-68.
23. Iatridis JC, Mente PL, Stokes IAF, Aronsson DD, Alini M. Compression-induced changes in intervertebral disc properties in a rat tail model. *Spine (Phila Pa 1976).* 1999;24:996-1002.
24. Hunter CJ, Matyas JR, Duncan NA. The Notochordal cell in the nucleus pulposus: a review in the context of tissue engineering. *Tissue Eng.* 2003;9:667-677.
25. Daly C, Ghosh P, Jenkin G, Oehme D, Goldschlager T. A review of animal models of intervertebral disc degeneration: pathophysiology, regeneration, and translation to the clinic. *Biomed Res Int.* 2016;2016:1-14.
26. Oegema TR. The role of disc cell heterogeneity in determining disc biochemistry: a speculation. *Biochem Soc Trans.* 2002;30:839-844.
27. Antoniou J, Steffen T, Nelson F. The human lumbar intervertebral disc: evidence for changes in the biosynthesis and denaturation of the extracellular matrix with growth, maturation, ageing, and degeneration. *J Clin.* 1996;98(4):996-1003.
28. Ariga K, Miyamoto S, Nakase T, et al. The relationship between apoptosis of endplate chondrocytes and aging and degeneration of the intervertebral disc. *Spine (Phila Pa 1976).* 2001;26:2414-2420.
29. O'Connell GD, Vresilovic EJ, Elliott DM. Human intervertebral disc internal strain in compression: the effect of disc region, loading position, and degeneration. *J Orthop Res.* 2011;29:547-555.
30. Bach FC, Willems N, Penning LC, Ito K, Meij BP, Tryfonidou MA. Potential regenerative treatment strategies for intervertebral disc degeneration in dogs. *BMC Vet Res.* 2014;10:3.
31. Purmessur D, Freemont AJ, Hoyland JA. Research article expression and regulation of neurotrophins in the nondegenerate and degenerate human intervertebral disc. *Arthritis Res Ther.* 2008;10:1-9.
32. Amonoo-Kuofi HS. Morphometric changes in the heights and anteroposterior diameters of the lumbar intervertebral discs with age. *J Anat.* 1991;175:159-168.
33. Gruber H, Johnson T, Norton H, Jr EH. The sand rat model for disc degeneration: radiologic characterization of age-related changes: cross-sectional and prospective analyses. *Spine (Phila Pa 1976).* 2002;27:230-234.
34. Bowles RD, Gebhard HH, Härtl R, Bonassar LJ. Tissue-engineered intervertebral discs produce new matrix, maintain disc height, and restore biomechanical function to the rodent spine. *Proc Natl Acad Sci U S A.* 2011;108:13106-13111.
35. Omlor G, Nerlich A, Wilke H, Pfeiffer M, Lorenz H. A new porcine in vivo animal model of disc degeneration: response of anulus fibrosus cells, chondrocyte-like nucleus pulposus cells, and notochordal nucleus. *Spine (Phila Pa 1976).* 2009;34:2730-2739.
36. Wilke H, Kettler A, Claes L. Are sheep spines a valid biomechanical model for human spines? *Spine (Phila Pa 1976).* 1997;22:2365-2374.
37. Alini M, Eisenstein S, Ito K, Little C, Kettler A. Are animal models useful for studying human disc disorders/degeneration? *Eur Spine.* 2008;17:2-19.
38. Bergknut N, Smolders LA, Grinwis GCM, et al. Intervertebral disc degeneration in the dog. Part 1: anatomy and physiology of the intervertebral disc and characteristics of intervertebral disc degeneration. *Vet J.* 2015;195:282-291. <https://doi.org/10.1016/j.tvjl.2012.10.024>.
39. Willems N, Tellegen AR, Bergknut N, et al. Inflammatory profiles in canine intervertebral disc degeneration. *BMC Vet Res.* 2016;12(1):10. <https://doi.org/10.1186/s12917-016-0635-6>.
40. Cherrone KL, Dewey CW, Acvs D, Coates JR, Bergman RL. A retrospective comparison of cervical intervertebral disk disease in non-chondrodystrophic large dogs versus small dogs. *J Am Anim Hosp Assoc.* 2004;40:316-320.
41. Molinos M, Almeida CR, Caldeira J, Cunha C, Goncalves RM, Barbosa MA. Inflammation in intervertebral disc degeneration and regeneration. *J R Soc Interface.* 2015;12:20141191.
42. Aoki Y, Ohtori S, Ino H, et al. Disc inflammation potentially promotes axonal regeneration of dorsal root ganglion neurons innervating lumbar intervertebral disc in rats. *Spine (Phila Pa 1976).* 2004;29:2621-2626.
43. Aoki Y, Ohtori S, Nakagawa K, Nakajima A. Neural mechanisms of Discogenic back pain: how does nerve growth factor play a key role? *Korean J Spine.* 2011;8:83-87.
44. Gruber HE, Ingram JA, Hanley EN. Immunolocalization of thrombospondin in the human and sand rat intervertebral disc. *Spine (Phila Pa 1976).* 2006;31:2556-2561.
45. Gruber HE, Ingram JA, Hoelscher G, Zinchenko N, Norton HJ, Hanley EN. Brain-derived neurotrophic factor and its receptor in the human and the sand rat intervertebral disc. *Arthritis Res Ther.* 2008;10:R82.
46. Detiger SEL, Helder MN, Smit TH, Hoogendoorn RJW. Adverse effects of stromal vascular fraction during regenerative treatment of the intervertebral disc: observations in a goat model. *Eur Spine J.* 2015;24:1992-2000.
47. Tong W, Lu Z, Qin L, et al. Cell therapy for the degenerating intervertebral disc. *Transl Res.* 2017;181:49-58.
48. Hansen H. A pathologic-anatomical study on disc degeneration in dog: with special reference to the so-called enchondrosis intervertebralis. *Acta Orthop Scand.* 1952;23:1-130.
49. Brisson BA. Intervertebral disc disease in dogs. *Vet Clin North Am Small Anim Pract.* 2010;40:829-858.
50. Seiler GS, Hani H, Busato AR, Lang J. Facet joint geometry and intervertebral disk degeneration in the L5-S1 region of the vertebral column in German shepherd dogs. *Am J Vet Res.* 2002;63:86-90.
51. Macias C, Mckee WM, May C, Innes JF. Thoracolumbar disc disease in large dogs: a study of 99 cases. *J Small Anim Pract.* 2002;43:439-446.
52. Hansen T, Smolders L, Tryfonidou M. The myth of fibroid degeneration in the canine intervertebral disc: a histopathological comparison

- of intervertebral disc degeneration in chondrodystrophic and. *Veterinary*. 2017;54:945-952.
53. Kranenburg HC, Grinwis GC, Bergknot N, et al. Intervertebral disc disease in dogs—part 2: comparison of clinical, magnetic resonance imaging, and histological findings in 74 surgically treated dogs. *Vet J*. 2013;195:164-171.
 54. Granger N, Carwardine D. Acute spinal cord injury. Tetraplegia and paraplegia in small animals. *Vet Clin North Am Small Anim Pract*. 2014;44:1131-1156.
 55. Erwin WM, Heras FL, Islam D, Fehlings MG, Inman RD. The regenerative capacity of the notochordal cell: tissue constructs generated in vitro under hypoxic conditions. *J Neurosurg Spine*. 2009;10:513-521.
 56. Smolders LA, Meij BP, Riemers FM, et al. Canonical Wnt signaling in the notochordal cell is upregulated in early intervertebral disk degeneration. *J Orthop Res*. 2012;30:950-957.
 57. Smolders LA, Meij BP, Onis D, et al. Gene expression profiling of early intervertebral disc degeneration reveals a down-regulation of canonical Wnt signaling and caveolin-1 expression: implications for development of regenerative strategies. *Arthritis Res Ther*. 2013;15:R23.
 58. Spillekom S, Smolders LA, Grinwis GCM, et al. Increased Osmolarity and cell clustering preserve canine Notochordal cell phenotype in culture. *Tissue Eng Part C Methods*. 2014;20:652-662.
 59. Erwin WM, DeSouza L, Funabashi M, et al. The biological basis of degenerative disc disease: proteomic and biomechanical analysis of the canine intervertebral disc. *Arthritis Res Ther*. 2015;17:240.
 60. Grunert P, Moriguchi Y, Grossbard BP, Ricart Arbona RJ, Bonassar LJ, Härtl R. Degenerative changes of the canine cervical spine after discectomy procedures, an in vivo study. *BMC Vet Res*. 2017;13:1-10.
 61. Brown EA, Dickinson PJ, Mansour T, et al. *FGF4* retrogene on CFA12 is responsible for chondrodystrophy and intervertebral disc disease in dogs. *Proc Natl Acad Sci U S A*. 2017;114:11476-11481. <https://doi.org/10.1073/pnas.1709082114>.
 62. Monchaux M, Forterre S, Spreng D, Karol A, Forterre F, Wuertz-Kozak K. Inflammatory processes associated with canine intervertebral disc herniation. *Front Immunol*. 2017;8:1681.
 63. Levine J, Cohen N, Heller M, Fajt V, Levine G. Efficacy of a metalloproteinase inhibitor in spinal cord injured dogs. *PLoS One*. 2014;9:e96408.
 64. Olby NJ, Muguet-Chanoit AC, Lim JH, et al. A placebo-controlled, prospective, randomized clinical trial of polyethylene glycol and methylprednisolone sodium succinate in dogs with intervertebral disk herniation. *J Vet Intern Med*. 2016;30:206-214.
 65. Budke CM, Levine JM, Kerwin SC, Levine GJ, Hettlich BF, Slater MR. Evaluation of a questionnaire for obtaining owner-perceived, weighted quality-of-life assessments for dogs with spinal cord injuries. *J Am Vet Med Assoc*. 2008;233:925-930.
 66. Moore S, Granger N, Olby N. Targeting translational successes through CANSORT-SCI: using pet dogs to identify effective treatments for spinal cord injury. *J Neurotrauma*. 2017;34:2007-2018.
 67. Cimino Brown D. What can we learn from osteoarthritis pain in companion animals? *Clin Exp Rheumatol*. 2017;35:S53-S58.
 68. Lascelles BDX, Brown DC, Maixner W, Mogil JS. Spontaneous painful disease in companion animals can facilitate the development of chronic pain therapies for humans. *Osteoarthritis Cartilage*. 2017;26:1-9. <https://doi.org/10.1016/j.joca.2017.11.011>.
 69. Klinck MP, Mogil JS, Moreau M, et al. Translational pain assessment: could natural animal models be the missing link? *Pain*. 2017;158:1633-1646.
 70. Imoto K, Takebayashi T, Kanaya K, Kawaguchi S, Katahira G, Yamashita T. Quantitative analysis of sensory functions after lumbar discectomy using current perception threshold testing. *Eur Spine J*. 2007;16:971-975.
 71. Rolke R, Magerl W, Campbell KA, et al. Quantitative sensory testing: a comprehensive protocol for clinical trials. *Eur J Pain*. 2006;10:77-88.
 72. Samuelsson L, Lundin A. Thermal quantitative sensory testing in lumbar disc herniation. *Eur Spine J*. 2002;11:71-75.
 73. Song RB, Basso DM, da Costa RC, Fisher LC, Mo X, Moore SA. Von Frey anesthesiometry to assess sensory impairment after acute spinal cord injury caused by thoracolumbar intervertebral disc extrusion in dogs. *Vet J*. 2016;209:144-149.
 74. Gorney AM, Blau SR, Dohse CS, et al. Mechanical and thermal sensory testing in normal chondrodystrophoid dogs and dogs with spinal cord injury caused by thoracolumbar intervertebral disc herniations. *J Vet Intern Med*. 2016;30:627-635.
 75. Brown DC, Boston RC, Coyne JC, Farrar JT. Ability of the canine brief pain inventory to detect response to treatment in dogs with osteoarthritis. *J Am Vet Med Assoc*. 2008;233:1278-1283.
 76. Hielm-Björkman AK, Rita H, Tulamo RM. Psychometric testing of the Helsinki chronic pain index by completion of a questionnaire in Finnish by owners of dogs with chronic signs of pain caused by osteoarthritis. *Am J Vet Res*. 2009;70:727-734.
 77. Yashari JM, Duncan CG, Duerr FM. Evaluation of a novel canine activity monitor for at-home physical activity analysis. *BMC Vet Res*. 2015;11:146.
 78. Clark K, Caraguel C, Leahey L, Béraud R. Evaluation of a novel accelerometer for kinetic gait analysis in dogs. *Can J Vet Res*. 2014;78:226-232.
 79. Martin KW, Olsen AM, Duncan CG, Duerr FM. The method of attachment influences accelerometer-based activity data in dogs. *BMC Vet Res*. 2017;13:48.
 80. Scott RM, Evans R, Conzemius MG. Efficacy of an oral nutraceutical for the treatment of canine osteoarthritis. A double-blind, randomized, placebo-controlled prospective clinical trial. *Vet Comp Orthop Traumatol*. 2017;30:318-323.
 81. Fletcher TF, Kitchell RL. Anatomical studies on the spinal cord segments of the dog. *Am J Vet Res*. 1966;27:1759-1767.
 82. Netter FH. Atlas of human anatomy. *Exp Neurol*. 2006;97:152-160.
 83. Humzah MD, Soames RW. Human intervertebral disc: structure and function. *Anat Rec*. 1988;220:337-356.
 84. Iatridis JC, Nicoll SB, Michalek AJ, Walter BA, Gupta MS. Role of biomechanics in intervertebral disc degeneration and regenerative therapies: what needs repairing in the disc and what are promising biomaterials for its repair? *Spine J*. 2013;13:243-262.
 85. Smit TH. The use of a quadruped as an in vivo model for the study of the spine - biomechanical considerations. *Eur Spine J*. 2002;11:137-144.
 86. Ganey T, Libera J, Moos V, et al. Disc chondrocyte transplantation in a canine model: a treatment for degenerated or damaged intervertebral disc. *Spine (Phila Pa 1976)*. 2003;28:2609-2620.
 87. Zhang Y, Tao H, Gu T, et al. The effects of human Wharton's jelly cell transplantation on the intervertebral disc in a canine disc degeneration model. *Stem Cell Res Ther*. 2015;6:1-15.
 88. Ganey T, Hutton WC, Moseley T, Hedrick M, Meisel H. Intervertebral disc repair using adipose tissue-derived stem and regenerative cells experiments in a canine model. *Spine (Phila Pa 1976)*. 2009;34:2297-2304.
 89. Bach FC, Miranda-Bedate A, van Heel FWM, et al. Bone morphogenetic protein-2, but not mesenchymal stromal cells, exert regenerative effects on canine and human nucleus pulposus cells. *Tissue Eng Part A*. 2017;23:233-242.
 90. Meisel HJ, Siodla V, Ganey T, Minkus Y, Hutton WC, Alasevic OJ. Clinical experience in cell-based therapeutics: disc chondrocyte transplantation: a treatment for degenerated or damaged intervertebral disc. *Biomol Eng*. 2007;24:5-21.
 91. Nukaga T, Sakai D, Tanaka M, Hiyama A, Nakai T, Mochida J. Transplantation of activated nucleus pulposus cells after cryopreservation: efficacy study in a canine disc degeneration model. *Eur Cell Mater*. 2016;31:95-106.
 92. Erwin WM, Inman RD. Notochord cells regulate intervertebral disc chondrocyte proteoglycan production and cell proliferation. *Spine (Phila Pa 1976)*. 2006;31:1094-1099.
 93. Erwin WM, Ashman K, O'Donnel P, Inman RD. Nucleus pulposus notochord cells secrete connective tissue growth factor and up-regulate proteoglycan expression by intervertebral disc chondrocytes. *Arthritis Rheum*. 2006;54:3859-3867.
 94. Erwin WM, Islam D, Inman RD, Fehlings MG, Tsui FWL. Notochordal cells protect nucleus pulposus cells from degradation and

- apoptosis: implications for the mechanisms of intervertebral disc degeneration. *Arthritis Res Ther*. 2011;13:R215.
95. Matta A, Karim MZ, Iseman DE, Erwin WM. Molecular therapy for degenerative disc disease: clues from secretome analysis of the notochordal cell-rich nucleus pulposus. *Sci Rep*. 2017;7:45623.
 96. Bach FC, Miranda-Bedate A, van Heel FWM, et al. Bone morphogenetic protein-2, but not mesenchymal stromal cells, exert regenerative effects on canine and human nucleus pulposus cells. *Tissue Eng*. 2016;23:233-242. <https://doi.org/10.1089/ten.TEA.2016.0251>.
 97. de Vries SAH, Potier E, van Doeselaar M, Meij BP, Tryfonidou MA, Ito K. Conditioned medium derived from notochordal cell-rich nucleus pulposus tissue stimulates matrix production by canine nucleus pulposus cells and bone marrow-derived stromal cells. *Tissue Eng Part A*. 2015;21:1077-1084.
 98. Arkesteijn IIT, Smolders LA, Spillekom S, et al. Effect of coculturing canine notochordal, nucleus pulposus and mesenchymal stromal cells for intervertebral disc regeneration. *Arthritis Res Ther*. 2015;17:60.
 99. Gu T, Shi Z, Wang C, et al. Human bone morphogenetic protein 7 transfected nucleus pulposus cells delay the degeneration of intervertebral disc in dogs. *J Orthop Res*. 2017;35:1311-1322.
 100. Li W, Liu T, Wu L, et al. Blocking the function of inflammatory cytokines and mediators by using IL-10 and TGF- β : a potential biological immunotherapy for intervertebral disc degeneration in a beagle model. *Int J Mol Sci*. 2014;15:17270-17283.
 101. Willems N, Bach FC, Plomp SGM, et al. Intradiscal application of rhBMP-7 does not induce regeneration in a canine model of spontaneous intervertebral disc degeneration. *Arthritis Res Ther*. 2015;17:137.
 102. Bach FC, Zhang Y, Miranda-Bedate A, et al. Increased caveolin-1 in intervertebral disc degeneration facilitates repair. *Arthritis Res Ther*. 2016;18:59.
 103. Bach FC, Laagland LT, Grant MP, et al. Link-N: the missing link towards intervertebral disc repair is species-specific. *PLoS One*. 2017;12:1-21.
 104. Moriguchi Y, Mojica-Santiago J, Grunert P, et al. Total disc replacement using tissue-engineered intervertebral discs in the canine cervical spine. *PLoS One*. 2017;12:1-18.
 105. Tellegen AR, Willems N, Beukers M, et al. Intradiscal application of a PCLA-PEG-PCLA hydrogel loaded with celecoxib for the treatment of back pain in canines: what's in it for humans? *J Tissue Eng Regen Med*. 2017;12:642-652. <https://doi.org/10.1002/term.2483>.
 106. Willems N, Mihov G, Grinwis GCM, et al. Safety of intradiscal injection and biocompatibility of polyester amide microspheres in a canine model predisposed to intervertebral disc degeneration. *J Biomed Mater Res: Part B Appl Biomater*. 2017;105:707-714.
 107. Willems N, Yang HY, Langelaan MLP, et al. Biocompatibility and intradiscal application of a thermoreversible celecoxib-loaded poly-N-isopropylacrylamide MgFe-layered double hydroxide hydrogel in a canine model. *Arthritis Res Ther*. 2015;17:214.
 108. Steffen F, Smolders L, Roentgen A. Bone marrow-derived mesenchymal stem cells as autologous therapy in dogs with naturally occurring intervertebral disc disease: feasibility, safety and preliminary results. *Tissue Eng Part C Methods*. 2017;23(11):643-651.
 109. Sakai D, Mochida J, Iwashina T, et al. Regenerative effects of transplanting mesenchymal stem cells embedded in atelocollagen to the degenerated intervertebral disc. *Biomaterials*. 2006;27:335-345.
 110. Muller C, Gaines B, Gruen M, et al. Evaluation of clinical metrology instrument in dogs with osteoarthritis. *J Vet Intern Med*. 2016;30:836-846.
 111. Purmessur D, Schek RM, Abbott RD, Ballif BA, Godburn KE, Iatridis JC. Notochordal conditioned media from tissue increases proteoglycan accumulation and promotes a healthy nucleus pulposus phenotype in human mesenchymal stem cells. *Arthritis Res Ther*. 2011;13:R81.
 112. Peng B, Hao J, Hou S, et al. Possible pathogenesis of painful intervertebral disc degeneration. *Spine (Phila Pa 1976)*. 2006;31:560-566.
 113. Habtemariam A, Virri J, Grönblad M, Seitsalo S, Karaharju E. The role of mast cells in disc herniation inflammation. *Spine (Phila Pa 1976)*. 1999;24:1516-1520.
 114. Le Maitre CL, Pockert A, Buttle DJ, Freemont AJ, Hoyland JA. Matrix synthesis and degradation in human intervertebral disc degeneration. *Biochem Soc Trans*. 2007;35:652-655.
 115. Wiet MG, Piscioneri A, Khan SN, Ballinger MN, Hoyland JA, Purmessur D. Mast cell-intervertebral disc cell interactions regulate inflammation, catabolism and angiogenesis in discogenic back pain. *Sci Rep*. 2017;7:1-14.

How to cite this article: Thompson K, Moore S, Tang S, Wiet M, Purmessur D. The chondrodystrophic dog: A clinically relevant intermediate-sized animal model for the study of intervertebral disc-associated spinal pain. *JOR Spine*. 2018;1:e1011. <https://doi.org/10.1002/jsp2.1011>