

CORRECTION

# Correction: Selective Chemokine Receptor Usage by Central Nervous System Myeloid Cells in CCR2-Red Fluorescent Protein Knock-In Mice

Noah Saederup, Astrid E. Cardona, Kelsey Croft, Makiko Mizutani, Anne C. Coteleur, Chia-Lin Tsou, Richard M. Ransohoff, Israel F. Charo


The Supporting Information file S1 Text is incorrectly duplicated as [S1 Table](#). Please see the correct [S1 Table](#) here.

## Supporting information

**S1 Table. Primers used for tail-DNA genotyping.** \*Primers A-B amplify the wild type allele, whereas primers B-C identify the knockout allele.  
(PDF)

## Reference

1. Saederup N, Cardona AE, Croft K, Mizutani M, Coteleur AC, Tsou C-L, et al. (2010) Selective Chemokine Receptor Usage by Central Nervous System Myeloid Cells in CCR2-Red Fluorescent Protein Knock-In Mice. *PLoS ONE* 5(10): e13693. doi:[10.1371/journal.pone.0013693](https://doi.org/10.1371/journal.pone.0013693) PMID: [21060874](https://pubmed.ncbi.nlm.nih.gov/21060874/)



## OPEN ACCESS

**Citation:** Saederup N, Cardona AE, Croft K, Mizutani M, Coteleur AC, Tsou C-L, et al. (2017) Correction: Selective Chemokine Receptor Usage by Central Nervous System Myeloid Cells in CCR2-Red Fluorescent Protein Knock-In Mice. *PLoS ONE* 12(4): e0176931. <https://doi.org/10.1371/journal.pone.0176931>

**Published:** April 27, 2017

**Copyright:** © 2017 Saederup et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.