

4-[(4-Chlorophenyl)(hydroxy)methylidene]isochromane-1,3-dione

Akoun Abou,^{a*} Abdoulaye Djandé,^b Bintou Sessouma,^b
Adama Saba^b and Rita Kakou-Yao^a

^aLaboratoire de Cristallographie et Physique Moléculaire, UFR SSMT, Université de Cocody 22 BP 582 Abidjan 22, Côte d'Ivoire, and ^bLaboratoire de Chimie Bio-organique et Phytochimie, Université de Ouagadougou 03 BP 7021 Ouagadougou 03, Burkina Faso
Correspondence e-mail: abou_akoun@yahoo.fr


Received 10 November 2011; accepted 14 November 2011

Key indicators: single-crystal X-ray study; $T = 298\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.053; wR factor = 0.163; data-to-parameter ratio = 17.0.

In the title compound, $\text{C}_{16}\text{H}_9\text{ClO}_4$, the six-membered heterocyclic ring adopts a screw-boat conformation. The benzene rings are oriented to each other at a dihedral angle of $59.26(9)^\circ$. The molecular structure exhibits a ring motif, *viz.* $S(6)$, owing to an intramolecular $\text{O}-\text{H}\cdots\text{O}$ hydrogen bond. The presence of $\text{C}-\text{H}\cdots\text{O}$ contacts generates an infinite chain along [001]. Also present are $\pi-\pi$ stacking interactions between neighbouring isochromanedione benzene rings [centroid–centroid distance = $3.746(1)\text{ \AA}$], and $\text{C}-\text{O}\cdots\pi$ interactions [$\text{O}\cdots\text{centroid} = 3.934(2)\text{ \AA}$].

Related literature

For the biological activity of isochromanones, see: Bianchi *et al.* (2004); Buntin *et al.* (2008). For $\pi-\pi$ stacking interactions, see: Janiak (2000). For hydrogen-bond motifs, see: Bernstein *et al.* (1995). For puckering parameters, see: Cremer & Pople (1975).



Experimental

Crystal data

$\text{C}_{16}\text{H}_9\text{ClO}_4$
 $M_r = 300.68$
Monoclinic, $P2_1/c$

$a = 15.4973(4)\text{ \AA}$
 $b = 5.9631(1)\text{ \AA}$
 $c = 14.4526(3)\text{ \AA}$

$\beta = 102.661(1)^\circ$
 $V = 1303.12(5)\text{ \AA}^3$
 $Z = 4$
Mo $K\alpha$ radiation

$\mu = 0.31\text{ mm}^{-1}$
 $T = 298\text{ K}$
 $0.40 \times 0.30 \times 0.20\text{ mm}$

Data collection

Nonius KappaCCD diffractometer
12213 measured reflections
3248 independent reflections
2693 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.037$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.053$
 $wR(F^2) = 0.163$
 $S = 1.08$
3248 reflections
191 parameters
H-atom parameters constrained
 $\Delta\rho_{\text{max}} = 0.25\text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.31\text{ e \AA}^{-3}$

Table 1
Hydrogen-bond geometry (\AA , $^\circ$).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O14—H14 \cdots O12	0.82	1.76	2.492 (2)	148
C3—H3 \cdots O11 ⁱ	0.93	2.56	3.288 (2)	136

Symmetry code: (i) $x, -y - \frac{1}{2}, z + \frac{1}{2}$.

Data collection: *COLLECT* (Nonius, 2001); cell refinement: *DENZO/SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO/SCALEPACK*; program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1994); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*, *publCIF* (Westrip, 2010) and *WinGX* (Farrugia, 1999).

We thank the Laboratoire de Physique des Interactions Ioniques et Spectropôle, Université de Provence, and Université Paul Cézanne, Faculté des Sciences et Techniques de Saint Jérôme, Marseille, France, for the use of their facilities.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK5016).

References

- Altomare, A., Cascarano, G., Giacovazzo, G., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.
- Bernstein, J., Davis, R. E., Shimoni, L. & Chang, N.-L. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 1555–1573.
- Bianchi, D. A., Blanco, N. E., Carrillo, N. & Kaufnam, T. S. (2004). *J. Agric. Food Chem.* **52**, 1923–1927.
- Buntin, K., Rachid, S., Scharfe, M., Blöcker, H., Weissman, K. J. & Müller, R. (2008). *Angew. Chem. Int. Ed. Engl.* **47**, 4595–4599.
- Cremer, D. & Pople, J. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Janiak, J. (2000). *J. Chem. Soc. Dalton Trans.* pp. 3885–3896.
- Nonius (2001). *COLLECT*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
- Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

supplementary materials

Acta Cryst. (2011). E67, o3349 [doi:10.1107/S160053681104829X]

4-[(4-Chlorophenyl)(hydroxy)methylidene]isochromane-1,3-dione

A. Abou, A. Djandé, B. Sessouma, A. Saba and R. Kakou-Yao

Comment

The title molecule is related to isochromanone derivatives that are generally known as regulators of plant growth (Bianchi *et al.*, 2004). Depending on their chemical structure and concentration they can act either as inhibitors or stimulators of these processes. Some substituted isochromanones isolated from myxobacteria strains were introduced as anti-fungal agents (Buntin *et al.*, 2008).

The structure of the title compound (I) (Fig. 1) consists of two planar benzene rings with the maximum deviations from the best planes of 0.035 (2) Å for atom C1 (benzene ring C1—C6) and ± 0.007 (2) Å for atoms C15 and C16 (benzene ring C15—C20). An *S*(6) ring motif (Bernstein *et al.*, 1995), arises from an intramolecular O—H \cdots O hydrogen bond to generate a planar pseudo six-membered ring (maximum deviation from planarity being 0.059 (2) Å for atom C13) to result in a tricyclic ring system (Fig. 1). The dihedral angles between two benzene rings is 59.26 (9) and that between the pseudo six-membered ring and benzene ring C1—C6 is 13.65 (9)°. The heterocyclic ring C1/C6/C7/O8/C9/C10 adopts a screw-boat conformation as judged from the puckering parameters (Cremer & Pople, 1975): $Q = 0.0952$ (19) Å, $\theta = 67.5$ (11)° and $\varphi = 228.4$ (12)°. Furthermore, intermolecular C—H \cdots O contacts (Table 1) link molecules into infinite chains through along [001] (Fig. 2).

The supramolecular aggregation is completed by the presence of C—O \cdots π interactions ($O12\cdots Cg3[x, 1/2 - y, -1/2 + z] = 3.934$ (2) Å, $C9\cdots O12 = 83.48$ (12)°, where $Cg3$ is the centroid of the benzene ring C15—C20, and $\pi\cdots\pi$ stacking between two parallel isochromanedione-benzene C1—C6 rings; in the latter, the centroid \cdots centroid distance, ($Cg2\cdots Cg2$) ($-x, -y, -z$) of 3.746 (1) Å, is less than 3.8 Å, the maximum regarded as relevant for $\pi\cdots\pi$ interactions (Janiak, 2000) (Fig.3).

Experimental

To a solution of 4-chlorobenzoyl chloride (4.10^{-2} mol) in dry tetrahydrofuran (150 ml), was added dried triethylamine (0.12 mol) and homophthalic anhydride (4.10^{-2} mol) in small portions over 30 min. The mixture was then refluxed for 3 h and poured in 300 ml of chloroform. The solution was acidified with dilute hydrochloric acid until the pH was 2–3. The organic layer was extracted, washed with water, dried over $MgSO_4$ and the solvent removed. The crude product was recrystallized from a chloroform-hexane (1/1, v/v) mixture. Yellow crystals were obtained in a good yield: 90%; *M.pt.* 432–433 K.

Refinement

H atoms were placed in calculated positions (O—H = 0.82 Å and C—H = 0.93 Å) and refined using a riding model approximation with $U_{iso}(\text{H})$ constrained to 1.2 (aromatic) or 1.5 (O—H) times U_{eq} of the respective parent atom.

supplementary materials

Figures




Fig. 1. The molecular structure of (I) showing the atomic labeling scheme with displacement ellipsoids drawn at the 50% probability level. H atoms are shown as spheres of arbitrary radius. Dashed lines indicate an hydrogen bond.




Fig. 2. Crystal packing, viewed down the b axis, showing parallel chains along the c direction. Dashed lines indicate C—H···O contacts. H atoms not involved in hydrogen bonds have been omitted for clarity.




Fig. 3. A view of the crystal packing, showing C—O···π and π···π stacking interactions (dashed lines). The yellow dots are centroids of rings. H atoms have been omitted for clarity.

4-[(4-Chlorophenyl)(hydroxy)methylidene]isochromane-1,3-dione

Crystal data

$C_{16}H_9ClO_4$	$F(000) = 616$
$M_r = 300.68$	$D_x = 1.533 \text{ Mg m}^{-3}$
Monoclinic, $P2_1/c$	Melting point = 432–433 K
Hall symbol: -P 2ybc	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 15.4973 (4) \text{ \AA}$	Cell parameters from 12213 reflections
$b = 5.9631 (1) \text{ \AA}$	$\theta = 1.4\text{--}29.0^\circ$
$c = 14.4526 (3) \text{ \AA}$	$\mu = 0.31 \text{ mm}^{-1}$
$\beta = 102.661 (1)^\circ$	$T = 298 \text{ K}$
$V = 1303.12 (5) \text{ \AA}^3$	Prism, yellow
$Z = 4$	$0.40 \times 0.30 \times 0.20 \text{ mm}$

Data collection

Nonius KappaCCD diffractometer	2693 reflections with $I > 2\sigma(I)$
Radiation source: fine-focus sealed tube	$R_{\text{int}} = 0.037$

graphite	$\theta_{\max} = 29.0^\circ, \theta_{\min} = 1.4^\circ$
φ and ω scans	$h = -20 \rightarrow 20$
12213 measured reflections	$k = -7 \rightarrow 7$
3248 independent reflections	$l = -19 \rightarrow 19$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.053$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.163$	H-atom parameters constrained
$S = 1.08$	$w = 1/[\sigma^2(F_o^2) + (0.085P)^2 + 0.4082P]$ where $P = (F_o^2 + 2F_c^2)/3$
3248 reflections	$(\Delta/\sigma)_{\max} < 0.001$
191 parameters	$\Delta\rho_{\max} = 0.25 \text{ e \AA}^{-3}$
0 restraints	$\Delta\rho_{\min} = -0.31 \text{ e \AA}^{-3}$
36 constraints	

Special details

Geometry. All s.u.'s (except the s.u. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell s.u.'s are taken into account individually in the estimation of s.u.'s in distances, angles and torsion angles; correlations between s.u.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell s.u.'s is used for estimating s.u.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Cl21	0.46136 (4)	0.22385 (12)	0.41605 (4)	0.0768 (3)
O8	0.15001 (11)	0.2734 (3)	-0.20791 (9)	0.0600 (4)
C1	0.16846 (10)	0.1431 (3)	-0.01749 (10)	0.0355 (3)
C6	0.11986 (11)	0.0094 (3)	-0.09040 (10)	0.0412 (4)
O12	0.22812 (11)	0.5783 (3)	-0.16990 (11)	0.0688 (5)
C2	0.16289 (11)	0.0881 (3)	0.07551 (10)	0.0391 (4)
H2	0.1898	0.1804	0.1254	0.047*
C10	0.21737 (11)	0.3377 (3)	-0.04175 (11)	0.0401 (4)
O14	0.30865 (10)	0.6583 (2)	-0.00417 (11)	0.0631 (4)
H14	0.2881	0.6813	-0.0606	0.095*
C7	0.11070 (14)	0.0730 (3)	-0.18995 (12)	0.0525 (5)
C13	0.27868 (11)	0.4635 (3)	0.02079 (13)	0.0437 (4)
C19	0.40760 (12)	0.1419 (3)	0.22901 (14)	0.0479 (4)

supplementary materials

H19	0.4368	0.0053	0.2416	0.057*
O11	0.07046 (13)	-0.0255 (3)	-0.25801 (10)	0.0769 (5)
C5	0.07423 (13)	-0.1807 (3)	-0.07083 (13)	0.0507 (4)
H5	0.0429	-0.2681	-0.1203	0.061*
C4	0.07567 (13)	-0.2382 (3)	0.02135 (15)	0.0521 (5)
H4	0.0482	-0.3690	0.0349	0.062*
C16	0.32267 (13)	0.5515 (3)	0.19262 (14)	0.0504 (4)
H16	0.2949	0.6900	0.1802	0.060*
C15	0.32154 (11)	0.4012 (3)	0.11887 (12)	0.0413 (4)
C20	0.36492 (12)	0.1967 (3)	0.13808 (13)	0.0452 (4)
H20	0.3651	0.0961	0.0889	0.054*
C17	0.36473 (13)	0.4967 (3)	0.28416 (14)	0.0547 (5)
H17	0.3648	0.5966	0.3336	0.066*
C3	0.11838 (12)	-0.1000 (3)	0.09398 (12)	0.0454 (4)
H3	0.1169	-0.1350	0.1563	0.054*
C18	0.40658 (12)	0.2924 (3)	0.30156 (13)	0.0484 (4)
C9	0.19987 (13)	0.4065 (3)	-0.13974 (12)	0.0499 (4)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Cl21	0.0704 (4)	0.1024 (5)	0.0503 (3)	0.0049 (3)	-0.0029 (3)	0.0100 (3)
O8	0.0778 (10)	0.0730 (10)	0.0291 (6)	0.0145 (7)	0.0118 (6)	0.0079 (6)
C1	0.0389 (7)	0.0382 (8)	0.0293 (7)	0.0059 (6)	0.0070 (6)	-0.0002 (5)
C6	0.0459 (8)	0.0452 (9)	0.0303 (7)	0.0107 (7)	0.0032 (6)	-0.0060 (6)
O12	0.0821 (11)	0.0719 (10)	0.0570 (8)	0.0075 (8)	0.0251 (7)	0.0298 (7)
C2	0.0427 (8)	0.0448 (9)	0.0290 (7)	-0.0061 (6)	0.0059 (6)	-0.0030 (6)
C10	0.0452 (8)	0.0422 (8)	0.0347 (7)	0.0065 (6)	0.0124 (6)	0.0057 (6)
O14	0.0663 (9)	0.0489 (8)	0.0720 (10)	-0.0103 (7)	0.0107 (7)	0.0188 (7)
C7	0.0646 (11)	0.0585 (11)	0.0313 (8)	0.0215 (9)	0.0034 (7)	-0.0057 (7)
C13	0.0444 (8)	0.0379 (8)	0.0508 (9)	0.0024 (6)	0.0147 (7)	0.0074 (7)
C19	0.0424 (8)	0.0427 (9)	0.0569 (10)	0.0037 (7)	0.0075 (7)	0.0054 (7)
O11	0.1078 (13)	0.0786 (11)	0.0337 (7)	0.0195 (9)	-0.0076 (7)	-0.0149 (7)
C5	0.0531 (10)	0.0472 (10)	0.0456 (9)	0.0004 (8)	-0.0030 (7)	-0.0133 (7)
C4	0.0534 (10)	0.0446 (10)	0.0550 (11)	-0.0112 (8)	0.0052 (8)	-0.0023 (8)
C16	0.0512 (10)	0.0372 (9)	0.0597 (11)	0.0044 (7)	0.0056 (8)	-0.0054 (7)
C15	0.0384 (8)	0.0365 (8)	0.0483 (9)	-0.0032 (6)	0.0081 (7)	0.0014 (6)
C20	0.0472 (9)	0.0387 (9)	0.0493 (9)	0.0025 (7)	0.0097 (7)	-0.0033 (7)
C17	0.0549 (11)	0.0544 (11)	0.0523 (10)	0.0022 (8)	0.0068 (8)	-0.0133 (8)
C3	0.0476 (9)	0.0502 (10)	0.0376 (8)	-0.0064 (7)	0.0074 (7)	0.0035 (7)
C18	0.0384 (8)	0.0576 (11)	0.0473 (9)	-0.0039 (7)	0.0054 (7)	0.0028 (8)
C9	0.0552 (10)	0.0583 (11)	0.0392 (8)	0.0153 (8)	0.0171 (7)	0.0120 (7)

Geometric parameters (\AA , $^\circ$)

Cl21—C18	1.7349 (19)	C13—C15	1.475 (2)
O8—C9	1.365 (3)	C19—C20	1.375 (3)
O8—C7	1.391 (3)	C19—C18	1.383 (3)
C1—C6	1.402 (2)	C19—H19	0.9300

C1—C2	1.405 (2)	C5—C4	1.371 (3)
C1—C10	1.469 (2)	C5—H5	0.9300
C6—C5	1.397 (3)	C4—C3	1.384 (3)
C6—C7	1.465 (2)	C4—H4	0.9300
O12—C9	1.231 (2)	C16—C17	1.380 (3)
C2—C3	1.373 (2)	C16—C15	1.390 (2)
C2—H2	0.9300	C16—H16	0.9300
C10—C13	1.381 (2)	C15—C20	1.390 (2)
C10—C9	1.442 (2)	C20—H20	0.9300
O14—C13	1.330 (2)	C17—C18	1.377 (3)
O14—H14	0.8200	C17—H17	0.9300
C7—O11	1.197 (2)	C3—H3	0.9300
C9—O8—C7	124.50 (14)	C6—C5—H5	120.1
C6—C1—C2	116.77 (15)	C5—C4—C3	119.43 (17)
C6—C1—C10	119.35 (14)	C5—C4—H4	120.3
C2—C1—C10	123.78 (14)	C3—C4—H4	120.3
C5—C6—C1	121.45 (15)	C17—C16—C15	120.40 (17)
C5—C6—C7	117.68 (16)	C17—C16—H16	119.8
C1—C6—C7	120.75 (17)	C15—C16—H16	119.8
C3—C2—C1	121.06 (15)	C16—C15—C20	119.22 (16)
C3—C2—H2	119.5	C16—C15—C13	120.09 (16)
C1—C2—H2	119.5	C20—C15—C13	120.61 (16)
C13—C10—C9	116.26 (16)	C19—C20—C15	120.63 (17)
C13—C10—C1	126.15 (14)	C19—C20—H20	119.7
C9—C10—C1	117.58 (15)	C15—C20—H20	119.7
C13—O14—H14	109.5	C18—C17—C16	119.31 (18)
O11—C7—O8	116.02 (18)	C18—C17—H17	120.3
O11—C7—C6	127.0 (2)	C16—C17—H17	120.3
O8—C7—C6	116.98 (16)	C2—C3—C4	121.07 (16)
O14—C13—C10	121.90 (16)	C2—C3—H3	119.5
O14—C13—C15	111.74 (16)	C4—C3—H3	119.5
C10—C13—C15	126.31 (15)	C17—C18—C19	121.25 (18)
C20—C19—C18	119.17 (17)	C17—C18—Cl21	119.68 (16)
C20—C19—H19	120.4	C19—C18—Cl21	119.05 (15)
C18—C19—H19	120.4	O12—C9—O8	114.66 (16)
C4—C5—C6	119.89 (16)	O12—C9—C10	125.43 (19)
C4—C5—H5	120.1	O8—C9—C10	119.89 (17)
C2—C1—C6—C5	-5.3 (2)	C17—C16—C15—C20	-1.4 (3)
C10—C1—C6—C5	178.26 (15)	C17—C16—C15—C13	-178.23 (17)
C2—C1—C6—C7	170.79 (15)	O14—C13—C15—C16	52.6 (2)
C10—C1—C6—C7	-5.7 (2)	C10—C13—C15—C16	-129.9 (2)
C6—C1—C2—C3	5.7 (2)	O14—C13—C15—C20	-124.18 (18)
C10—C1—C2—C3	-177.98 (15)	C10—C13—C15—C20	53.3 (2)
C6—C1—C10—C13	-169.15 (16)	C18—C19—C20—C15	0.3 (3)
C2—C1—C10—C13	14.6 (3)	C16—C15—C20—C19	0.8 (3)
C6—C1—C10—C9	11.3 (2)	C13—C15—C20—C19	177.65 (16)
C2—C1—C10—C9	-164.88 (16)	C15—C16—C17—C18	0.8 (3)
C9—O8—C7—O11	-178.66 (18)	C1—C2—C3—C4	-1.7 (3)

supplementary materials


C9—O8—C7—C6	3.6 (3)	C5—C4—C3—C2	-3.1 (3)
C5—C6—C7—O11	-3.0 (3)	C16—C17—C18—C19	0.3 (3)
C1—C6—C7—O11	-179.20 (19)	C16—C17—C18—Cl21	178.84 (15)
C5—C6—C7—O8	174.43 (15)	C20—C19—C18—C17	-0.9 (3)
C1—C6—C7—O8	-1.8 (2)	C20—C19—C18—Cl21	-179.41 (14)
C9—C10—C13—O14	11.1 (3)	C7—O8—C9—O12	-179.24 (16)
C1—C10—C13—O14	-168.40 (16)	C7—O8—C9—C10	2.3 (3)
C9—C10—C13—C15	-166.14 (16)	C13—C10—C9—O12	-7.6 (3)
C1—C10—C13—C15	14.3 (3)	C1—C10—C9—O12	172.00 (17)
C1—C6—C5—C4	0.7 (3)	C13—C10—C9—O8	170.70 (16)
C7—C6—C5—C4	-175.45 (18)	C1—C10—C9—O8	-9.7 (2)
C6—C5—C4—C3	3.6 (3)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O14—H14 \cdots O12	0.82	1.76	2.492 (2)	148
C3—H3 \cdots O11 ⁱ	0.93	2.56	3.288 (2)	136

Symmetry codes: (i) $x, -y-1/2, z+1/2$.

Fig. 1



supplementary materials

Fig. 2




Fig. 3

