

Editorial



Myocardial Efficiency: A Reliable Load-independent Parameter of Cardiac Performance?

In-Chang Hwang , MD^{1,2}

¹Department of Cardiology, Cardiovascular Center, Seoul National University Bundang Hospital, Seongnam, Korea

²Department of Internal Medicine, Seoul National University College of Medicine, Seoul, Korea

OPEN ACCESS

► See the article “An Analysis of Myocardial Efficiency in Patients with Severe Asymptomatic Mitral Regurgitation” in volume 28 on page 267.

Received: Jul 16, 2020

Accepted: Jul 29, 2020

Address for Correspondence:

In-Chang Hwang, MD

Department of Internal Medicine, Seoul National University College of Medicine, Cardiovascular Center, Seoul National University Bundang Hospital, 82 Gumi-ro 173beon-gil, Bundang-gu, Seongnam 13620, Korea.

E-mail: inchang.hwang@gmail.com

Copyright © 2020 Korean Society of Echocardiography

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

In-Chang Hwang 

<https://orcid.org/0000-0003-4966-3924>

Conflict of Interest

The author has no financial conflicts of interest.

As an mechanical engine, the efficiency of heart can be summarized as the ratio of external work (EW) and the oxygen uptake of myocardium in a given time, which was initially suggested by Bing et al in 1949.^{1,2)} The definition of myocardial efficiency (MEf), in usual clinical and research settings, include only left ventricle (LV), while the work and metabolism of right ventricle and atria are neglected. Also, the MEf accounts the EW of LV, but not the potential energy of LV which indicates the work required to expand the LV to its actual minimal volume (LV end-systolic volume). Under these assumptions, the MEf is calculated using the product of EW of LV and heart rate, divided by the product of myocardial oxygen consumption (mVO_2) and LV mass (LVM) (**Figure 1**). The EW, or stroke work (SW) indicates the area contained in a pressure-volume loop of LV (**Figure 1**), and can be approximately estimated as a product of stroke volume (SV) and end-systolic pressure, or a product of SV and mean arterial pressure (MAP).

The mVO_2 can be measured with invasive cardiac catheterization, or with the use of radioactive isotope; inhaled ^{15}O -labeled oxygen gas, or generally, injected ^{11}C -acetate. After intravenous injection, the ^{11}C -acetate is transformed to ^{11}C -acetyl-CoA in myocardium and enters the tricarboxylic acid cycle where it is metabolized to ^{11}C - CO_2 .¹⁾ The myocardial time-activity curve shows a biexponential clearance (rapid phase [k_1] and slow phase [k_2]), and the rate of ^{11}C - CO_2 clearance is monoexponential under steady hemodynamic conditions and is linearly related to oxygen uptake, which indicates the myocardial oxidative metabolism (**Figure 1**). Thus, the clearance rate of ^{11}C -activity (k_2) is used for the calculation of mVO_2 , which is in the denominator of the MEf equation. As a simplified method, Beanlands et al.³⁾ introduced an alternative measure of efficiency index, the work-metabolic index (WMI), using the monoexponential time constant (k_{mono}) of the linear part of time-activity curve (**Figure 1**). The measurement of SV, as well as LVM, is performed by non-invasive tests including echocardiography, magnetic resonance, computed tomography, or positron emission tomography (PET).⁴⁻⁶⁾

Previous studies showed that the MEf is decreased in patients with hypertension, heart failure (HF), aortic stenosis (AS), and mitral regurgitation (MR).⁶⁻¹³⁾ Furthermore, therapeutic intervention to these volume and/or pressure-overloaded conditions alleviate the MEf




Figure 1. Non-invasive measurement of myocardial efficiency. (A) In the non-invasive calculation of MEF, the EW which indicates the area contained in the PV loop is approximated as a product of SV and MAP. (B) The time constants (k_1 , k_2 , and k_{mono}) are obtained from the myocardial time-activity curve of ^{11}C -acetate PET. The WMI is calculated using the simplified k_{mono} of decay for ^{11}C -acetate, and the mVO_2 (which is in the denominator of MEF equation) is calculated from the washout rate of ^{11}C -activity (k_2).

MEf: myocardial efficiency; EW: external work; PV: pressure-volume; SV: stroke volume; MAP: mean arterial pressure; k_1 : rapid phase; k_2 : slow phase; k_{mono} : monoexponential time constant; PET: positron emission tomography; WMI: work-metabolic index; mVO_2 : myocardial oxygen consumption; LV: left ventricular; PE: potential energy; ESPVR: end-systolic pressure-volume relationship; EDPVR: end-diastolic pressure-volume relationship; TCA: tricarboxylic acid; SBP: systolic blood pressure; SVi: stroke volume index; HR: heart rate; LVM: left ventricular mass.

through decreased mVO_2 , reduced LVM, and increased EW. For example, the afterload reduction by aortic valve replacement improved MEF through LV reverse remodeling and reduced mVO_2 , in conjunction with an increase in EW, in patients with severe AS.⁹⁾ The improvement of WMI was also observed in patients with severe MR at 1-year after MV replacement.¹¹⁾ These studies support the potential usefulness and feasibility of MEF in various cardiac diseases.

In this paper of *Journal of Cardiovascular Imaging*, the authors compared the MEF in patients with severe MR and healthy volunteers, using parameters obtained from echocardiography.¹⁴⁾ Although the LV-ejection fraction (EF) and forward SV were preserved in patients with MR, the mVO_2 was significantly increased, and thus, the MEF was markedly reduced. Adding to the main finding on the reduced MEF in patients with severe MR, the authors further reported that the patients with severe MR who were on both beta-blockers and renin-angiotensin system (RAS) blockers tended to have a higher MEF, compared to the patients without these medications. This finding corresponds to the established benefits of these cardioprotective medications: beta-blockers reduce mVO_2 , and RAS blockers reduce afterload and augment forward SV. Also, it is in concordance with previous studies that reported the improved MEF by the use of beta-blockers in patients with HF, which was accompanied with improved LV-EF and increased SW, and reduced oxygen consumption.¹⁵⁾¹⁶⁾

Theoretically, the concept of MEF is convincing as it considers the actual EW corrected to the consumption of oxygen, and also includes the loading condition (product of SV and MAP). Adding to this, the MEF has further strengths over traditional parameters: MEF reveals the metabolic status of myocardium, which may enable a more sensitive detection of benefits of medical therapy in a diseased myocardium. Indeed, traditional parameters are largely based on the motion of myocardium, which should be the final stage output from the impaired myocardial energetics and disturbed loading conditions. Furthermore, the increasing number of facilities equipped with cyclotron is a factor that causes the utilization of MEF to increase in the future.

Despite these potential strengths of MEF, there are several issues to be further investigated for its wider clinical application. First, the reference range of MEF significantly varies according to the methods of measurement. The standardization and validation between modalities (echocardiography, magnetic resonance, computed tomography, and PET) are required. Second, the equation of MEF calculation is based on several assumptions and approximations. This is not limited to the innate assumptions of the imaging modalities, but also to the assumptions for the SW of the heart. Third, supporting evidence for the assessment of MEF and its changes according to therapeutic interventions is still limited to a few disease groups.

At present, it can be suggested that the MEF has a potential as a useful non-invasive, load-independent tool for assessment of LV performance, and for the monitoring of treatment response. Future studies investigating the role of MEF in various disease population are warranted. Also, it is important to understand the characteristics of MEF and related parameters, how to measure them, and the remaining limitations.

REFERENCES

1. Sørensen J, Harms HJ, Aalen JM, Baron T, Smiseth OA, Flachskampf FA. Myocardial efficiency: a fundamental physiological concept on the verge of clinical impact. *JACC Cardiovasc Imaging* 2020;13:1564-76. [PUBMED](#) | [CROSSREF](#)
2. Bing RJ, Hammond MM, Handelsman JC, et al. The measurement of coronary blood flow, oxygen consumption, and efficiency of the left ventricle in man. *Am Heart J* 1949;38:1-24. [PUBMED](#) | [CROSSREF](#)
3. Beanlands RS, Bach DS, Raylman R, et al. Acute effects of dobutamine on myocardial oxygen consumption and cardiac efficiency measured using carbon-11 acetate kinetics in patients with dilated cardiomyopathy. *J Am Coll Cardiol* 1993;22:1389-98. [PUBMED](#) | [CROSSREF](#)
4. Harms HJ, Tolbod LP, Hansson NH, et al. Automatic extraction of forward stroke volume using dynamic PET/CT: a dual-tracer and dual-scanner validation in patients with heart valve disease. *EJNMMI Phys* 2015;2:25. [PUBMED](#) | [CROSSREF](#)
5. Harms HJ, Stubkjær Hansson NH, Tolbod LP, et al. Automatic extraction of myocardial mass and volume using parametric images from dynamic nongated PET. *J Nucl Med* 2016;57:1382-7. [PUBMED](#) | [CROSSREF](#)
6. Harms HJ, Hansson NH, Kero T, et al. Automatic calculation of myocardial external efficiency using a single ¹¹C-acetate PET scan. *J Nucl Cardiol* 2018;25:1937-44. [PUBMED](#) | [CROSSREF](#)
7. Beanlands RS, Armstrong WF, Hicks RJ, et al. The effects of afterload reduction on myocardial carbon 11-labeled acetate kinetics and noninvasively estimated mechanical efficiency in patients with dilated cardiomyopathy. *J Nucl Cardiol* 1994;1:3-16. [PUBMED](#) | [CROSSREF](#)

8. Kim IS, Izawa H, Sobue T, et al. Prognostic value of mechanical efficiency in ambulatory patients with idiopathic dilated cardiomyopathy in sinus rhythm. *J Am Coll Cardiol* 2002;39:1264-8.
[PUBMED](#) | [CROSSREF](#)
9. Güçlü A, Knaapen P, Harms HJ, et al. Myocardial efficiency is an important determinant of functional improvement after aortic valve replacement in aortic valve stenosis patients: a combined PET and CMR study. *Eur Heart J Cardiovasc Imaging* 2015;16:882-9.
[PUBMED](#) | [CROSSREF](#)
10. Hansson NH, Sörensen J, Harms HJ, et al. Metoprolol reduces hemodynamic and metabolic overload in asymptomatic aortic valve stenosis patients: a randomized trial. *Circ Cardiovasc Imaging* 2017;10:e006557.
[PUBMED](#) | [CROSSREF](#)
11. Chow BJ, Abunassar JG, Ascah K, et al. Effects of mitral valve surgery on myocardial energetics in patients with severe mitral regurgitation. *Circ Cardiovasc Imaging* 2010;3:308-13.
[PUBMED](#) | [CROSSREF](#)
12. Sugiki T, Naya M, Manabe O, et al. Effects of surgical ventricular reconstruction and mitral complex reconstruction on cardiac oxidative metabolism and efficiency in nonischemic and ischemic dilated cardiomyopathy. *JACC Cardiovasc Imaging* 2011;4:762-70.
[PUBMED](#) | [CROSSREF](#)
13. Laine H, Katoh C, Luotolahti M, et al. Myocardial oxygen consumption is unchanged but efficiency is reduced in patients with essential hypertension and left ventricular hypertrophy. *Circulation* 1999;100:2425-30.
[PUBMED](#) | [CROSSREF](#)
14. Güvenç RÇ, Aruğaslan E, Güvenç TS, Karadeniz FÖ, Kaşıkçıoğlu H, Çam N. An analysis of myocardial efficiency in patients with severe asymptomatic mitral regurgitation. *J Cardiovasc Imaging* 2020;28:267-78.
[CROSSREF](#)
15. Eichhorn EJ, Heesch CM, Barnett JH, et al. Effect of metoprolol on myocardial function and energetics in patients with nonischemic dilated cardiomyopathy: a randomized, double-blind, placebo-controlled study. *J Am Coll Cardiol* 1994;24:1310-20.
[PUBMED](#) | [CROSSREF](#)
16. Beanlands RS, Nahmias C, Gordon E, et al. The effects of β_1 -blockade on oxidative metabolism and the metabolic cost of ventricular work in patients with left ventricular dysfunction: a double-blind, placebo-controlled, positron-emission tomography study. *Circulation* 2000;102:2070-5.
[PUBMED](#) | [CROSSREF](#)