


RESEARCH

Open Access



First identification and molecular subtyping of *Blastocystis* sp. in zoo animals in southwestern China

Lei Deng^{1†}, Jingxin Yao^{1†}, Shanyu Chen^{1†}, Tingmei He², Yijun Chai¹, Ziyao Zhou¹, Xiaogang Shi², Haifeng Liu¹, Zhijun Zhong¹, Hualin Fu¹ and Guangneng Peng^{1*} 

Abstract

Background: *Blastocystis* sp. is an anaerobic protozoan that parasitizes many animal hosts and the human gastrointestinal tract, and its pathogenicity is controversial. Captive wildlife may be potential reservoirs for human infection with *Blastocystis* sp. The present study was performed to investigate the prevalence and subtype distribution of *Blastocystis* sp. in zoo animals in Sichuan Province, southwestern China.

Methods: A total of 420 fresh fecal samples were collected from 54 captive wildlife species in four zoos in southwestern China between June 2017 and September 2019. The prevalence and subtype (ST) genetic characteristics of *Blastocystis* sp. were determined by PCR amplification of the barcode region of the *SSU* rRNA gene and phylogenetic analysis.

Results: Overall, 15.7% (66/420) of the animal samples and 20.7% (14/54) of the species tested were shown to be infected with *Blastocystis* sp. The highest prevalence of *Blastocystis* sp. was found in Panzhihua Zoo (24.3%), which was significantly higher than that in Chengdu Zoo (6.9%), and Xichang Zoo (2.9%) ($P < 0.05$). There are also significant differences in the prevalence of *Blastocystis* sp. among different species ($P < 0.05$), and the highest of *Blastocystis* sp. prevalence was observed in white-cheeked gibbon, black great squirrel, and red giant flying squirrel (100%). Subtype analysis of *Blastocystis* sp. revealed nine subtypes, including six zoonotic STs (ST1-5, and ST8) and three animal-specific STs (ST10, ST14, and ST17), with ST17 as the predominant subtype (26/66) in *Blastocystis* sp.-positive isolates.

Conclusions: To our knowledge, this is the first report on the prevalence and subtype distribution of *Blastocystis* sp. among captive wildlife in zoos in southwestern China. This study highlights that these animals may serve as reservoirs for human *Blastocystis* sp. infections.

Keywords: *Blastocystis* sp., Captive wildlife, Subtypes, Phylogenetic analysis, China

Background

Blastocystis sp., belonging to the phylum stramenopiles, is a common anaerobic eukaryotic protist that inhabits the gastrointestinal tract of a wide range of hosts, including humans. It is estimated that this parasite has colonized 1 to 2 billion people worldwide based on epidemiological surveys [1]. *Blastocystis* sp. is mainly transmitted through the fecal-oral route, food, and water [2–4]. There is supporting evidence that some human infections

*Correspondence: pgn.sicau@163.com

†Lei Deng, Jingxin Yao, Shanyu Chen equal contributors

¹The Key Laboratory of Animal Disease and Human Health of Sichuan Province, College of Veterinary Medicine, Sichuan Agricultural University, Chengdu 611130, Sichuan, People's Republic of China
Full list of author information is available at the end of the article



© The Author(s) 2021. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

may be caused by zoonotic transmission of *Blastocystis* sp. [5, 6].

Based on polymorphisms of small subunit (*SSU*) gene of *Blastocystis* sp., 22 subtypes (STs) consisting of ST1 to ST17, ST21, and ST23 to ST26 have been identified in humans and domestic and wild animals worldwide [7]. ST1-9 and -12 are able to infect humans, with ST1-4 being the most common, accounting for more than 90% of human *Blastocystis* sp. infections [8]. Interestingly, the prevalence of different STs among countries and among regions within the same country seems to vary greatly [9]. Remarkable genetic diversity was observed among various STs [10], and different STs exhibit diverse biological features, such as drug resistance, pathogenicity, and effects on microbiota [11–13].

Although *Blastocystis* sp. has been reported > 100 years, the clinical significance of this common parasite remains controversial [14]. Accumulating evidence shows *Blastocystis* sp. long-term colonization in asymptomatic carriers, accompanied with a healthy gut microbiota [15, 16], suggesting that it should be regarded as a member of the normal intestinal microbiota. *Blastocystis* sp. has been found in patients with irritable bowel syndrome (IBS) and inflammatory bowel disease (IBD), [17, 18], but not the presence of the protist that causes them. *In vitro* experiments using cell lines have also determined the potential pathogenicity of some specific STs of *Blastocystis* sp., such as disrupting the epithelial barrier [19] by increasing the epithelial permeability [20, 21]. Moreover, experimental infections with *Blastocystis* sp. in mouse models have shown that it can decrease the abundance of beneficial bacteria *Bifidobacterium* and *Lactobacillus* [12].

Blastocystis sp. has been reported in a substantial number of animal hosts, including livestock, companion animals, and captive wildlife, with greatly varying prevalence [22–24]. In recent years, several important intestinal zoonotic pathogens (e.g., *Cryptosporidium*, *Giardia*, and Microsporidia) have been reported in captive wildlife in China [25, 26], highlighting wildlife may be potential reservoirs for human to contract these infectious agents. However, less information is currently available regarding the prevalence and subtype distribution of *Blastocystis* sp. in zoo animals in China [27]. The purpose of the present study was thus to determine the genetic characteristics and subtype distribution of *Blastocystis* sp. in various zoo animals in southwestern China to better assess its zoonotic potential.

Methods

Sample collection

A total of 420 fresh fecal samples were collected from wildlife in Chengdu Zoo ($n=144$), Ya'an Zoo ($n=204$),

Xichang Zoo ($n=35$), and Panzhihua Zoo ($n=37$) between June 2017 and September 2019 in Sichuan Province, southwestern China (Fig. 1). The collected samples include a large variety of mammalian groups and several avian species. The animals were either housed individually or in monospecific groups of 5–10 individuals sharing the same enclosures. For those animals housed individually, only one sample was collected per animal. In group housing, between two and five samples were collected, each of which was considered as individual sample. All the fresh fecal samples were collected by zookeepers before the cleaning of animal cages in the morning and were strictly controlled to minimize potential contamination between animal species. Feces samples from some avian species were collected carefully directly on the ground or in their nests. All fecal samples were collected in sterilized plastic containers using disposable sterile gloves and preserved at 4 °C until DNA extraction.

DNA extraction

Fecal specimens were sieved and washed three times with distilled water by centrifugation at 3000g for 5 min. Genomic DNA was extracted using a QIAamp DNA

Stool Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's instructions. DNA quality was verified by NanoDrop (Thermo Fisher Scientific, Carlsbad, CA, USA) measurements. DNA was eluted in 50 µl of nuclease-free water and stored at – 20 °C until use.

PCR amplification

All extracted DNA was screened for the presence of *Blastocystis* sp. by a polymerase chain reaction (PCR) amplification of the barcode region of the *SSU* rRNA gene, using primers RD5 (5'-ATCTGGTTGATCCTGCCAGT-3') and BhRDr (5'-GAGCTTTTAACTGCAACAACG-3') [28]. The PCR mixture (25 µl) contained 12.5 µl Taq PCR Master Mix (Sangon Biotech Co., Ltd., Shanghai, China), 1 µl each primer (0.4 µM), 2 µl genomic DNA, 1.5 mM MgCl₂, and nuclease-free water up to desired volume. The PCR was started at 94 °C for 4 min followed by 30 cycles of 95 °C for 15 s, 60 °C for 15 s, and 72 °C for 30 s, with an extension at 72 °C for 5 min. Positive and negative controls were included in all the PCR tests. PCR products were subjected to 1.5% agarose gel (AddGene, Watertown, MA, USA) electrophoresis and visualized by staining with SYBR Safe DNA Gel Stain (Thermo Fisher Scientific).

Sequencing and phylogenetic analysis

PCR products with the expected size (~600 bp) were excised from the agarose gel and purified using a QIAquick Gel Extraction Kit (Qiagen) according to the manufacturer's instructions. Purified products were

directly sequenced on an ABI PRISM™ 3730 DNA Analyzer (Applied Biosystems, USA), using a BigDye Terminator v3.1 Cycle Sequencing kit (Applied Biosystems, Foster, CA, USA).

Nucleotide sequences obtained in the present study were subjected to BLAST searches (<http://www.ncbi.nlm.nih.gov/blast/>), and the reference sequences were downloaded from the GenBank database. *Blastocystis* sp. subtypes were identified by BLAST searches (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>), and the alleles were identified at the *Blastocystis* sp. database (<http://pubmlst.org/blastocystis>). The sequences obtained in this study and reference sequences were aligned using the MUSCLE algorithm of MEGA-X [29]. The ambiguous positions were removed from the alignment, and then the alignment was trimmed using MEGA 6 (<http://www.megasoftware.net/>) before phylogenetic analysis. ST11 is not available in the barcode region, and ST21 and ST23-26 need further data for determination, so we did not include these subtypes in the phylogenetic analysis [7]. The phylogenetic tree was constructed with the neighbor-joining (NJ) method using the Mega 6 software. Evolutionary distances were calculated using the Kimura two-parameter model. Bootstrap analysis (with 1000 replicates) was performed to define the robustness of the findings. *Proteromonas lacertae* was used as outgroup for the phylogenetic analysis.

Statistical analysis

The difference in *Blastocystis* sp. prevalence between different zoos and the order of animals was analyzed with the chi-square (χ^2) test, using SPSS 20.0 (IBM, Chicago, IL, USA). The difference was considered statistically significant when $P < 0.05$. Ninety-five percent confidence intervals (95% CIs) and odds ratios (ORs) were also calculated to explore the strengths of association between *Blastocystis* sp. positivity and each factor.

Results

Prevalence of *Blastocystis* sp. in captive wild animals

It is worth noting that the prevalences herein are the minimum since we used barcoding primers rather than qPCR. In the present study, 66 of 420 (15.7%) fecal samples collected from four animal zoos in China were determined to be *Blastocystis* sp. positive by PCR amplification of the barcode region of the *SSU* rRNA gene. Specifically, 10 of 144 (6.9%) animals sampled from Chengdu Zoo, 46 of 204 (22.5%) samples from Ya'an Zoo, 1 of 35 (2.9%) samples from Xichang Zoo, and 9 of 37 (24.3%) samples collected from Panzhihua Zoo were *Blastocystis* sp. positive (Table 1). The difference in *Blastocystis* sp. prevalence was significant in four zoos ($P < 0.05$). The prevalence of *Blastocystis* sp. in nonhuman primates (NHPs) was 30.5%, which is higher than that in Rodentia (18.2%), Artiodactyla (12.8%), birds (8.8%), and Carnivora (5.9%).




Fig. 1 Geographical distribution of the sampled cities (filled circle) in Sichuan Province, Southwestern China

Similarly, the difference of *Blastocystis* sp. prevalence among different order animals was significant ($P < 0.05$).

In general, of the 54 species tested in this study, 14 (20.7%) were positive for *Blastocystis* sp. (Table 2). Specifically, of the 29 species tested at the Chengdu Zoo, 6 (20.7%) were positive for *Blastocystis* sp. At the Ya'an Zoo, the prevalence of the parasite was 33.3% (5/15) among the species screened; 7.7% (1/13) and 33.3% (3/9) species at Xichang Zoo and Panzhihua Zoo in the present study were shown to be infected with *Blastocystis* sp. respectively.

Interestingly, the prevalence of *Blastocystis* sp. varies greatly among different species (Table 3). The highest *Blastocystis* sp. prevalence was observed in white-cheeked gibbon (100%, 4/4). By comparison, sika deer and green peafowl showed lower *Blastocystis* sp. prevalence, accounting for 9.1% and 8.3% respectively.

Subtype distributions of *Blastocystis* sp. in captive wild animals

Among the 66 *Blastocystis* sp.-positive samples, 9 subtypes were identified, including 6 zoonotic STs (ST1-5, and ST8) and 3 animal-specific STs (ST10, ST14, and ST17). ST17 (allele 160) (39.4%, 26/66) was the dominant subtype in zoo animals examined in the present study (Table 4), followed by ST1 (allele 1) (22.7%, 15/66), ST5 (allele 118) (10.6%, 7/66), ST2 (allele 15) (9.1%, 6/66), and ST8 (allele 21) (6.1%, 4/66). ST3 (allele 34), ST4 (allele 42), ST10 (allele 152), and ST14 (allele 157) were only found in two fecal samples respectively (Table 4). Notably, ST1 has the widest host range in zoo animals, detected in rhesus macaque, ring-tailed lemur, leopard, and sika deer (Table 4). Meanwhile, ST17 was identified in four species of animals, including squirrel monkey, Asiatic black bear, and chinchilla (Table 4).

Genetic characteristics of *Blastocystis* sp. subtypes

The identity analysis of the *SSU* rRNA gene revealed that 15 sequences of ST1 isolates identified in NHPs, leopard, and sika deer were identical to those from Philippine long-tailed macaque in the Philippines (KY929113). Similarly, six ST2 sequences from NHPs (ring-tailed lemur, and white-cheeked gibbon) showed 100% identity with GenBank sequences MN585877 (from human in Brazil) and MF581567 (from human in Angola). ST3 and ST4 sequences had the largest identity (99.62% and 98.97%) related to that from human in China (MT645668), and human in Czech Republic (MT042813), with two and six nucleotide substitutions respectively. Meanwhile, one leopard and six ostrich-derived ST5 sequences had 98.95% identity with that from an ostrich in China (MK930351), with six single-nucleotide polymorphisms (SNPs). In terms of four pigeon-derived ST8 isolates, the

sequences had 99.65% identity with that from human in Brazil (MN585867), with two nucleotide substitutions being observed. The sequences of ST10 and ST14 from alpaca were identical to the GenBank sequences MN316600 and MN316667, both from goat in Iran respectively. For 26 ST17 isolates, two representative sequences were obtained from NHPs, Asiatic black bear, and chinchillas. The sequence (MN227379) of ST17 isolates showed 100% identity to the sequence of ST17 isolated from a gundi in Libya (KC148208). The sequence (MN227380) exhibited 99.15% identity to the sequence of ST17 isolated from a chinchilla in China (MN124518), with five SNPs.

Phylogenetic analysis of *Blastocystis* sp.

A total of 10 representative sequences were obtained from 66 *Blastocystis* sp. isolates in the present study. The sequences obtained in this study showed high identity with the reference sequences of *Blastocystis* sp. in GenBank. Newly acquired sequences belong to ST1, ST2, ST3, ST4, ST5, ST8, ST10, ST14, and ST17. ST1 and ST2 along with sequences originating from humans and *Macaca* clustered together. ST3 and ST8 grouped together with sequences mainly from humans. ST4 clustered together with sequences from rats and humans. ST14 along with sequences isolated from goat and cattle grouped together. ST5 along with sequences isolated from ostrich, cattle, and human clustered together. ST10 formed a clade with sequences from dog, cattle, and goat. Similarly, ST17 formed a separate branch with sequences from gundi and chinchilla (Fig. 2).

Discussion

Blastocystis sp. is a common intestinal protozoan parasite with uncertain pathogenicity. It is believed that zoonotic STs often spread between animals and humans, as some STs of animal origin are a large potential reservoir for human infections [30–32]. *Blastocystis* sp. has been identified in many animal hosts, such as NHPs, pigs, cattle, sheep, goats, dogs, birds, and various captive wildlife [23, 24, 33]. The prevalence of *Blastocystis* sp. in zoo animals examined in this study was 15.7% (66/420), which was lower than that in captive wild animals in Qinling Mountains, China (40.2%, 200/497) [34], in zoo animals in the UK (35.5%, 82/231) [35], in zoo animals in Japan (39.0%, 46/118) [36], in various captive animals in France (32.2%, 99/307) [37], and in zoo animals in Western Australia (42%, 32/76) [5], while the *Blastocystis* sp. prevalence in the present study was higher than that in zoo animals in three cities in China (6.0%, 27/450) [27]. However, it is difficult to explain the discrepancy of *Blastocystis* sp. prevalence between different countries or within the same country because many factors contribute to the

Table 1 Factors associated with the prevalence of *Blastocystis* sp. in zoo animals in China

Factors	No. of examined	No. of positive	Prevalence (%) (95% CI)	OR (95% CI)	P value
Zoo					
Ya'an Zoo	204	46	22.5% (16.8–28.3)	0.906 (0.399–2.056)	0.813
Chengdu Zoo	144	10	6.9% (2.8–11.1)	0.232 (0.086–0.624)	0.004
Xichang Zoo	35	1	2.9% (0–8.4)	0.092 (0.011–0.767)	0.027
Panzhuhua Zoo	37	9	24.3% (10.5–38.1)	Reference	
Host					
Primates	128	39	30.5% (22.5–38.4)	4.541 (2.201–9.370)	0.000
Carnivora	85	5	5.9% (0.8–10.9)	0.648 (0.217–1.936)	0.437
Rodentia	33	6	18.2% (5.0–31.3)	2.303 (0.782–6.779)	0.130
Artiodactyla	39	5	12.8% (2.3–23.3)	1.524 (0.495–4.691)	0.463
Perissodactyla	4	0	0		
Diprotodontia	3	0	0		
Erinaceidae	3	0	0		
Birds	125	11	8.8% (3.8–13.8)	Reference	
Total	420	66	15.7% (12.2–19.2)		

effect the prevalence, such as the size of examined samples, animal species, or management methods.

The prevalence of *Blastocystis* sp. among the four zoos was significantly different in the present study. The prevalence of *Blastocystis* sp. in Panzhuhua Zoo was the highest (24.3%), which was significantly higher than that in Chengdu Zoo (6.9%) and Xichang Zoo (2.9%) ($P < 0.05$). The difference in the prevalence of *Blastocystis* sp. may be related to the zoo's sanitary conditions and management methods. The sanitary conditions are relatively poor, and there is no good deworming and immunization program in the Panzhuhua Zoo. In addition, the host and number of samples collected in each zoo are different, which may also be one of the reasons for the difference in *Blastocystis* sp. prevalences.

In the present study, 9 *Blastocystis* STs, including ST1-5, ST8, ST10, ST14, and ST17, were identified in 66 *Blastocystis* sp.-positive samples from captive wildlife. Of them, four subtypes (ST1-3, and ST17) were identified in NHPs (rhesus macaque, ring-tailed lemur, squirrel monkey, and white-cheeked gibbon). Generally, ST1-3 has been frequently determined in NHPs, while ST17 was identified to a lesser extent [34, 37]. Interestingly, a more recent study showed zoo animals and staff were infected with ST1-3 and showed high consistency with known sequences from NHPs and humans, highlighting that these STs have zoonotic potential [38]. In China, ST1, ST2, ST3, ST4, ST5, ST9, ST10, and ST13 were identified in NHPs [27, 31, 39, 40], with ST1 and ST2 being the most common. In this study, *Blastocystis* sp. was first discovered in squirrel monkey, and subtype analysis showed all isolates from squirrel monkey were ST17.

To date, several zoonotic STs have been identified in Carnivora, such as ST1-5, ST7-8, and ST10 [32, 33, 38]. In the current study, sequences obtained from Carnivora belonged to ST1, ST5, and ST17. These isolates were infecting the leopard (ST1 and ST5) and Asiatic black bear (ST17). As previously reported, the animals belonging to the order of Carnivora such as South American coati, Arctic fox, and dogs were determined to be infected with ST1 [38, 41]. Hussein *et al.* reported that ST1 inoculated into Wistar rats could cause moderate and severe degrees of pathological changes, suggesting the potential pathogenicity of this subtype [42]. ST5 was the most predominant subtype in pigs [43], but it was also identified in various animals, such as NHPs, cattle, sheep, rodents, and birds [23]. However, ST5 was rarely found in carnivores, and only a few wild carnivores, such as Northern tiger cat, and meerkat were determined to have ST5 infection [27, 38]. In addition, ST5 infection has also been reported in stray and domestic dogs in India [44] and the Philippines [45]. ST17 has been identified in gundi in Libya [46] and in cattle in the USA [47]. While there is no study on ST17 infections in Carnivora, this is the first report of ST17 infection in Asiatic black bear in China, demonstrating a wider host range of this subtype.

Previous studies reported many animals in the order of Artiodactyla harboring the *Blastocystis* sp., such as pigs, cattle, sheep, goats, camels, and deer [23, 33]. The majority of STs (ST1-7, ST10, ST13-15, and ST17) have been identified in Artiodactyla to date [46, 48]. Among them, ST10 was the most common subtype in cattle in the US [49], Denmark [50], and China [51, 52]. In the present study, ST1 was found in sika deer. Several studies

Table 2 Animal samples collected from various hosts from four different zoos in Sichuan Province, southwestern China

Host	Scientific name	No. of examined	YA	CD	XC	PZH	No. of <i>Blastocystis</i> -positive
Primates							
De Brazza's monkey	<i>Cercopithecus neglectus</i>	1	1				0
Rhesus macaque	<i>Macaca mulatta</i>	17	15	2			10
Ring-tailed lemur	<i>Lemur catta</i>	13	13				6
Squirrel monkey	<i>Saimiri sciureus</i>	93	93				19
White-cheeked gibbon	<i>Hylobates leucogenys</i>	4	4				4
Carnivora							
African lion	<i>Panthera leo</i>	14	14				0
Bengal tiger	<i>Panthera tigris</i>	2	1			1	0
Leopard	<i>Panthera pardus</i>	3	3				2
Lynx	<i>Lynx lynx</i>	1	1				0
Asiatic black bear	<i>Ursus thibetanus</i>	12				12	3
Brown bear	<i>Ursus arctos</i>	1				1	0
Asiatic wild dog	<i>Cuon alpinus</i>	2	2				0
Arctic fox	<i>Vulpes lagopus</i>	3		3			0
Wolf	<i>Canis lupus Linnaeus</i>	2		2			0
Raccoon dog	<i>Nyctereutes procyonoides</i>	2		2			0
Eurasian badger	<i>Meles meles</i>	2		2			0
Eurasian river otter	<i>Lutra lutra</i>	2		2			0
Ferret	<i>Mustela putorius furo</i>	2		2			0
Red panda	<i>Ailurus fulgens</i>	17	17				0
Raccoon	<i>Procyon lotor</i>	18	13	5			0
Coati	<i>Nasua olivacea</i>	1		1			0
Civet	<i>Paguma sp.</i>	1		1			0
Rodentia							
Chipmunk	<i>Tamias sp.</i>	5		5			0
Prairie dogs	<i>Cynomys sp.</i>	3		3			0
Polatouche	<i>Pteromys volans</i>	2		2			0
Black great squirrel	<i>Ratufa bicolor</i>	1		1			1
Red giant flying squirrel	<i>Petaurista petaurista</i>	1		1			1
Capybara	<i>Hydrochoerus hydrochaeris</i>	4	4				0
Guinea pig	<i>Cavia porcellus</i>	2		2			0
Chinchilla	<i>Chinchilla lanigera</i>	6		6			4
Beaver	<i>Castor fiber</i>	4		4			0
Nepal porcupine	<i>Hystrix brachyura subcristata</i>	3		3			0
Hamster	<i>Cricetulus sp.</i>	2		2			0
Artiodactyla							
Sika deer	<i>Cervus nippon</i>	11		3	3	5	1
Red muntjac	<i>Muntiacus muntjak</i>	7		7			0
Fallow deer	<i>Dama dama</i>	3		3			0
Roe deer	<i>Capreolus pygargus</i>	1		1			0
Alpaca	<i>Vicugna pacos</i>	11	11				4
Two-humped camel	<i>Camelus bactrianus</i>	4			2	2	0
Yellow cattle	<i>Bos taurus domestica</i>	2			2		0
Perissodactyla							
Horse	<i>Equus caballus</i>	3			1	2	0
Common zebra	<i>Equus burchellii</i>	1				1	0

Table 2 (continued)

Host	Scientific name	No. of examined	YA	CD	XC	PZH	No. of <i>Blastocystis</i> -positive
Diprotodontia							
Parma wallaby	<i>Macropus parma</i>	3		3			0
Erinaceidae							
Hedgehog	<i>Erinaceus amurensis</i>	3		3			0
Galliformes							
Green peafowl	<i>Pavo muticus</i>	12			5	7	1
Chicken	<i>Gallus gallus</i>	51		41	10		0
Ring-necked Pheasant	<i>Phasianus colchicus</i>	1			1		0
Guinea fowl	Numididae	3			3		0
Struthioniformes							
Common ostrich	<i>Struthio camelus</i>	19	12		1	6	6
Psittaciformes							
Green-winged macaw	<i>Ara chloroptera</i>	1			1		0
Columbiformes							
Common pigeon	<i>Columba livia</i>	34		31	3		4
Anseriformes							
Goose	<i>Anser cygnoides domesticus</i>	2			2		0
Strigiformes							
Eurasian eagle owl	<i>Bubo bubo</i>	1			1		0
Ratitae							
Cassowary	Casuarius sp.	1		1			0
Total		420	204	144	35	37	66

YA = Ya'an Zoo; CD = Chengdu Zoo; XC = Xichang Zoo; PZH = Panzhihua Zoo

Table 3 Prevalence of *Blastocystis* sp. among different species

Species	No. of examined	No. of positive	Prevalence (%)	YA	CD	XC	PZH
Squirrel monkey	93	19	20.4	ST17 (19)			
Rhesus macaque	17	10	58.8	ST1 (10)			
White-cheeked gibbon	4	4	100	ST2 (3); ST3 (1)			
Ring-tailed lemur	13	6	46.2	ST1 (3); ST2 (3)			
Chinchilla	6	4	66.7		ST17 (4)		
Red giant flying squirrel	1	1	100		ST4 (1)		
Black great squirrel	1	1	100		ST4 (1)		
Alpaca	11	4	36.4	ST10 (2); ST14 (2)			
Sika deer	11	1	9.1				ST1 (1)
Asiatic black bear	12	3	25				ST17 (3)
Leopard	3	2	66.7	ST1 (1); ST5 (1)			
Common pigeon	34	4	11.8		ST8 (4)		
Green peafowl	12	1	8.3			ST3 (1)	
Common ostrich	19	6	31.6	ST5 (1)			ST5 (5)
Total	237	66	27.8	ST17 (19); ST1 (14); ST2 (6); ST5 (2); ST10 (2); ST14 (2); ST3 (1)	ST8 (4); ST17 (4); ST4 (2)	ST3 (1)	ST5 (5); ST17 (3); ST1 (1)

YA = Ya'an Zoo; CD = Chengdu Zoo; XC represents Xichang Zoo; PZH = Panzhihua Zoo

Table 4 Subtype distributions from different animal species

Host	<i>Blastocystis</i> sp. STs										Sequences
	1	2	3	4	5	8	10	14	17		
Primates											
Rhesus macaque	10										10
Ring-tailed lemur	3	3									6
Squirrel monkey									19		19
White-cheeked gibbon		3	1								4
Carnivora											
Leopard	1				1						2
Asiatic black bear									3		3
Rodentia											
Chinchilla									4		4
Red giant flying squirrel				1							1
Black great squirrel				1							1
Artiodactyla											
Sika deer	1										1
Alpaca								2	2		4
Birds											
Common ostrich					6						6
Common pigeon						4					4
Green peafowl			1								1
Total	15	6	2	2	7	4	2	2	26		66

have reported on *Blastocystis* sp. infections in deer, with different subtype distributions. For example, sika deer, fallow deer, and white-lipped deer were reported to be infected with ST10 in China [34]. ST4 and ST10 were also reported in red deer, and muntjac deer were found with ST14 infection in the UK [53]. Strikingly, a rare ST (ST13) was also determined in a mouse deer in the UK [46] and in Java mouse-deer in France [37]. The distribution of STs in alpaca in the present study was consistent with a previous study in the Qinglin Mountains in China, in which all isolates identified belonged to ST10 and ST14 [34]. Similarly, ST10 infection in alpaca was also reported in French zoos [37]. Overall, these data suggest that deer and alpaca may serve as natural hosts of *Blastocystis* sp.


In this study, ST4 and ST17 were identified in rodents, corroborating previous data on pet rodents in Sichuan Province [54]. ST4 was originally isolated from a healthy Wistar rat in Singapore [55], and rodents were proposed to be a main reservoir of ST4. Recent studies confirmed that other STs, such as ST1-3, ST5, ST7, ST8, ST10, and ST17, can also be found in rodents [46, 53]. ST4 infection has been reported for the first time in red giant flying squirrel and black great squirrel, expanding its host range. The observation of ST17 in shrew-faced squirrels in the United Arab Emirates suggests rodents may be the

natural host of this subtype [56]. In the present study, ST17 was also observed in chinchilla in China for the first time to our knowledge, indicating a novel host for this subtype.

Regarding the non-mammalian groups, birds have already been considered potential reservoirs of *Blastocystis* sp. transmission to humans [57]. It is believed that birds usually harbor ST6 and ST7, which are considered ‘avian STs’ because of their predominance in this host group. Nevertheless, of the 11 avian isolates characterized in the present study, none were identified as belonging to the ‘avian’ ST6 or ST7. Six of them belonged to ST5, four to ST8, and one to ST3; these results are similar to those of a previous study in birds in French zoos where ST5 was the dominant subtype [37]. It should be noted that our previous study reported that ST8 was the predominant subtype in captive birds in Sichuan Province [58], suggesting that this subtype may circulate among birds in the investigated areas. However, the transmission characteristics of these zoonotic subtypes warrant further examination in future studies.

Conclusions

The present study described the prevalence, subtype distribution and genetic characterizations of *Blastocystis* sp. for the first time in zoo animals in southwestern China.



The data demonstrated that *Blastocystis* sp. could be maintained and transmitted between wildlife. Moreover, the zoonotic subtypes were identified in captive wildlife suggesting that these animals may serve as natural reservoirs for human *Blastocystis* sp. infections. The present results could provide fundamental information for the evaluation of potential zoonotic transmission between wildlife and humans.

Abbreviations

SSU rRNA: Small subunit ribosomal RNA; STs: Subtypes; IBS: Irritable bowel syndrome; IBD: Inflammatory bowel disease; PCR: Polymerase chain reaction; ORs: Odds ratios; NHPs: Nonhuman primates; SNPs: Single nucleotide polymorphisms.

Acknowledgements

We thank Wuyou Wang for the comments on the manuscript and Tong Jie Xin for critical editing. Lei Deng was the recipient of a scholarship from the Chinese Scholarship Council (CSC).

Authors' contributions

This study was conceived and designed by GP. Experiments were performed by LD, JY, and SC. Fecal samples were collected by TH, XS, ZZ (Zhijun Zhong), and YC. Data were analyzed by ZZ (Ziyao Zhou), HF, and HL. All authors have read and approved the submitted version of this manuscript.

Funding

This work was funded by the Chengdu Giant Panda Breeding Research Foundation (CPF2017-12) and Special Fund for Forestry Reform and Development of Wolong Special Administrative Region of Sichuan Province (510000-02-064387).

Availability of data and materials

The nucleotide sequences generated in the present study have been deposited in GenBank (<https://www.ncbi.nlm.nih.gov/>) under accession numbers MN227371, MN227378, MN227374, MN227376, MN227362, MN227363, MW078483, MW078491, MN227379, and MN227380. The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

This study protocol was reviewed and approved by the Research Ethics Committee and the Animal Ethical Committee of Sichuan Agricultural University. Permission was obtained from zoo managers prior to collection of fecal specimens.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹ The Key Laboratory of Animal Disease and Human Health of Sichuan Province, College of Veterinary Medicine, Sichuan Agricultural University, Chengdu 611130, Sichuan, People's Republic of China. ² Sichuan Wolong National Natural Reserve Administration, Aba 623006, Sichuan, People's Republic of China.

Received: 13 October 2020 Accepted: 1 December 2020

Published online: 06 January 2021

References

- Scanlan PD, Stensvold CR. *Blastocystis*: getting to grips with our guileful guest. *Trends Parasitol.* 2013;29:523–9.
- Tan KSW. New insights on classification, identification, and clinical relevance of *Blastocystis* spp. *Clin Microbiol Rev.* 2008;21:639.
- Li L, Tan TC, Karmacharya BM, Govind SK. *Blastocystis* sp.: waterborne zoonotic organism, a possibility? *Parasit Vectors.* 2012;5:130–130.
- Leelayoova S, Siripattanapipong S, Thathaisong U, Naaglor T, Taamasri P, Piyaraj P, et al. Drinking water: a possible source of *Blastocystis* spp. subtype 1 infection in schoolchildren of a rural community in central Thailand. *Am J Trop Med Hyg.* 2008;79:401–6.
- Parkar U, Traub RJ, Vitali S, Elliot A, Levecke B, Robertson I, et al. Molecular characterization of *Blastocystis* isolates from zoo animals and their animal-keepers. *Vet Parasitol.* 2010;169:8–17.

6. Wang W, Owen H, Traub RJ, Cuttall L, Inpankaew T, Bielefeldt-Hohmann H. Molecular epidemiology of *Blastocystis* in pigs and their in-contact humans in Southeast Queensland, Australia, and Cambodia. *Vet Parasitol*. 2014;203:264–9.
7. Stensvold CR, Clark CG. Pre-empting Pandora's Box: *Blastocystis* subtypes revisited. *Trends Parasitol*. 2020;36:229–32.
8. Stensvold CR, Tan KSW, Clark CG. *Blastocystis*. *Trends Parasitol*. 2020;36:315–6.
9. Clark CG, van der Giezen M, Alfellani MA, Stensvold CR. Recent developments in *Blastocystis* research. *Adv Parasitol*. 2013;82:1–32.
10. Gentekaki E, Curtis BA, Stairs CW, Klimeš V, Eliáš M, Salas-Leiva DE, et al. Extreme genome diversity in the hyper-prevalent parasitic eukaryote *Blastocystis*. *PLoS Biol*. 2017;15:e2003769.
11. Ajjampur SS, Tan KS. Pathogenic mechanisms in *Blastocystis* spp.—interpreting results from in vitro and in vivo studies. *Parasitol Int*. 2016;65:772–9.
12. Yason JA, Liang YR, Png CW, Zhang Y, Tan KSW. Interactions between a pathogenic *Blastocystis* subtype and gut microbiota: *in vitro* and *in vivo* studies. *Microbiome*. 2019;7:30.
13. Rajamanikam A, Hooi HS, Kudva M, Samudi C, Kumar S. Resistance towards metronidazole in *Blastocystis* sp.: a pathogenic consequence. *PLoS ONE*. 2019;14:e0212542.
14. Andersen LO, Stensvold CR. *Blastocystis* in health and disease: Are we moving from a clinical to a public health perspective? *J Clin Microbiol*. 2016;54:524–8.
15. Scanlan PD, Stensvold CR, Rajilić-Stojanović M, Heilig HG, De Vos WM, O'Toole PW, et al. The microbial eukaryote *Blastocystis* is a prevalent and diverse member of the healthy human gut microbiota. *FEMS Microbiol Ecol*. 2014;90:326–30.
16. Audebert C, Even G, Cian A, Group BI, Loywick A, Merlin S, et al. Colonization with the enteric protozoa *Blastocystis* is associated with increased diversity of human gut bacterial microbiota. *Sci Rep*. 2016; 6: 25255.
17. Shirvani G, Fasihi-Harandi M, Raiesi O, Bazargan N, Zahedi MJ, Sharifi I, et al. Prevalence and molecular subtyping of *Blastocystis* from patients with irritable bowel syndrome, inflammatory bowel disease and chronic urticaria in Iran. *Acta Parasitol*. 2020;65:90–6.
18. Peña S, Carrasco G, Rojas P, Castillo D, Ozaki LS, Mercado R. Determination of subtypes of *Blastocystis* sp. in Chilean patients with and without inflammatory bowel syndrome, A preliminary report. *Parasite Epidemiol Control*. 2020;8:e00125.
19. Wu Z, Mirza H, Tan KS. Intra-subtype variation in enteroadhesion accounts for differences in epithelial barrier disruption and is associated with metronidazole resistance in *Blastocystis* subtype-7. *PLoS Negl Trop Dis*. 2014;8:e2885.
20. Mirza H, Wu Z, Teo JD, Tan KS. Statin pleiotropy prevents rho kinase-mediated intestinal epithelial barrier compromise induced by *Blastocystis* cysteine proteases. *Cell Microbiol*. 2012;14:1474–84.
21. Puthia MK, Sio SW, Lu J, Tan KS. *Blastocystis ratti* induces contact-independent apoptosis, F-actin rearrangement, and barrier function disruption in IEC-6 cells. *Infect Immun*. 2006;74:4114–23.
22. Rauff-Adedotun AA, Mohd Zain SN, Farah Haziqah MT. Current status of *Blastocystis* sp. in animals from Southeast Asia: a review. *Parasitol Res*. 2020; 1–12.
23. Jiménez PA, Jaimes JE, Ramírez JD. A summary of *Blastocystis* subtypes in North and South America. *Parasit Vectors*. 2019;12:376.
24. Mohammadpour I, Bozorg-Ghalati F, Gazzonis AL, Manfredi MT, Motazedian MH, Mohammadpour N. First molecular subtyping and phylogeny of *Blastocystis* sp. isolated from domestic and synanthropic animals (dogs, cats and brown rats) in southern Iran. *Parasit Vectors*. 2020;13:365.
25. Li W, Deng L, Yu X, Zhong Z, Wang Q, Liu X, et al. Multilocus genotypes and broad host-range of *Enterocytozoon bieneusi* in captive wildlife at zoological gardens in China. *Parasit Vectors*. 2016;9:395.
26. Li J, Qi M, Chang Y, Wang R, Li T, Dong H, et al. Molecular characterization of *Cryptosporidium* spp., *Giardia duodenalis*, and *Enterocytozoon bieneusi* in captive wildlife at Zhengzhou Zoo. *China J Eukaryot Microbiol*. 2015;62:833–9.
27. Li XD, Zou Y, Pan J, Liang QL, Zeng Z, Meng YM, et al. Prevalence and subtypes of *Blastocystis* sp. infection in zoo animals in three cities in China. *Parasitol Res*. 2020;119:465–71.
28. Scicluna SM, Tawari B, Clark CG. DNA Barcoding of *Blastocystis*. *Protist*. 2006;157:77–85.
29. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol*. 2018;35:1547–9.
30. Sharifi Y, Abbasi F, Shahabi S, Zareei A, Mikaeili F, Sarkari B. Comparative genotyping of *Blastocystis* infecting cattle and human in the south of Iran. *Comp Immunol Microbiol Infect Dis*. 2020;72:101529.
31. Zhu W, Wei Z, Li Q, Lin Y, Yang H, Li W. Prevalence and subtype diversity of *Blastocystis* in human and nonhuman primates in North China. *Parasitol Res*. 2020;119:2719–25.
32. Li J, Karim MR, Li D, Rahman Sumon SMM, Siddiki S, Rume FI, et al. Molecular characterization of *Blastocystis* sp. in captive wildlife in Bangladesh National Zoo: non-human primates with high prevalence and zoonotic significance. *Int J Parasitol Parasites Wildl*. 2019;10:314–20.
33. Deng L, Chai Y, Zhou Z, Liu H, Zhong Z, Hu Y, et al. Epidemiology of *Blastocystis* sp. infection in China: a systematic review. *Parasite*. 2019;26:41.
34. Zhao GH, Hu XF, Liu TL, Hu RS, Yu ZQ, Yang WB, et al. Molecular characterization of *Blastocystis* sp. in captive wild animals in Qinling Mountains. *Parasitol Res*. 2017;116:2327–33.
35. Betts EL, Gentekaki E, Tsaousis AD. Exploring micro-eukaryotic diversity in the gut: Co-occurrence of *Blastocystis* subtypes and other protists in zoo animals. *Front Microbiol*. 2020;11:288.
36. Abe N, Nagoshi M, Takami K, Sawano Y, Yoshikawa H. A survey of *Blastocystis* sp. in livestock, pets, and zoo animals in Japan. *Vet Parasitol*. 2002;106:203–12.
37. Cian A, El Safadi D, Osman M, Moriniere R, Gantois N, Benamrouz-Vanneste S, et al. Molecular epidemiology of *Blastocystis* sp. in various animal groups from two French zoos and evaluation of potential zoonotic risk. *PLoS ONE*. 2017;12:e0169659.
38. Oliveira-Arbex AP, David ÉB, Tenório MDS, Cicchi PJP, Patti M, Coradi ST, et al. Diversity of *Blastocystis* subtypes in wild mammals from a zoo and two conservation units in southeastern Brazil. *Infect Genet Evol*. 2020;78:104053.
39. Ma L, Qiao H, Wang H, Li S, Zhai P, Huang J, et al. Molecular prevalence and subtypes of *Blastocystis* sp. in primates in northern China. *Transbound Emerg Dis*. 2020;26:46.
40. Li TC, Li Z, Zhang YL, Chen WJ, Dong XL, Yang JF, et al. Assessment of the subtypes and the zoonotic risk of *Blastocystis* sp. of experimental macaques in Yunnan province, southwestern China. *Parasitol Res*. 2020;119:741–8.
41. Wang J, Gong B, Liu X, Zhao W, Bu T, Zhang W, et al. Distribution and genetic diversity of *Blastocystis* subtypes in various mammal and bird species in northeastern China. *Parasit Vectors*. 2018;11:522.
42. Hussein EM, Hussein AM, Eida MM, Atwa MM. Pathophysiological variability of different genotypes of human *Blastocystis hominis* Egyptian isolates in experimentally infected rats. *Parasitol Res*. 2008;102:853–60.
43. Song JK, Hu RS, Fan XC, Wang SS, Zhang HJ, Zhao GH. Molecular characterization of *Blastocystis* from pigs in Shaanxi province of China. *Acta Trop*. 2017;173:130–5.
44. Wang W, Cuttall L, Bielefeldt-Ohmann H, Inpankaew T, Owen H, Traub RJ. Diversity of *Blastocystis* subtypes in dogs in different geographical settings. *Parasit Vectors*. 2013;6:215.
45. Belleza ML, Cadacio JL, Borja MP, Solon JA, Padilla MA, Tongol-Rivera PN, et al. Epidemiologic study of *Blastocystis* infection in an urban community in the Philippines. *J Environ Publ Heal*. 2015;2015:894297.
46. Alfellani MA, Taner-Mulla D, Jacob AS, Imeede CA, Yoshikawa H, Stensvold CR, et al. Genetic diversity of *Blastocystis* in livestock and zoo animals. *Protist*. 2013;164:497–509.
47. Maloney JG, Lombard JE, Urie NJ, Shivley CB, Santin M. Zoonotic and genetically diverse subtypes of *Blastocystis* in US pre-weaned dairy heifer calves. *Parasitol Res*. 2019;118:575–82.
48. Wawrzyniak I, Poirier P, Viscogliosi E, Dionigia M, Texier C, Delbac F, et al. *Blastocystis*, an unrecognized parasite: an overview of pathogenesis and diagnosis. *Therap Adv Infect Dis*. 2013;1:167.
49. Ronald F, Monica S, Dumitru M. Detection of concurrent infection of dairy cattle with *Blastocystis*, *Cryptosporidium*, *Giardia*, and *Enterocytozoon* by molecular and microscopic methods. *Parasitol Res*. 2012;111:1349–55.
50. Stensvold CR, Alfellani MA, Nørskov-Lauritsen S, Prip K, Victory EL, Maddox C, et al. Subtype distribution of *Blastocystis* isolates from synanthropic and zoo animals and identification of a new subtype. *Int J Parasitol*. 2009;39:473–9.
51. Zhu W, Tao W, Gong B, Yang H, Li Y, Song M, et al. First report of *Blastocystis* infections in cattle in China. *Vet Parasitol*. 2017;246:38–42.
52. Wang J, Gong B, Yang F, Zhang W, Zheng Y, Liu A. Subtype distribution and genetic characterizations of *Blastocystis* in pigs, cattle, sheep and

- goats in northeastern China's Heilongjiang Province. *Infect Genet Evol.* 2018;57:171–6.
53. Betts EL, Gentekaki E, Thomasz A, Breakell V, Carpenter AI, Tsaousis AD. Genetic diversity of *Blastocystis* in non-primate animals. *Parasitol.* 2018;145:1228–34.
54. Chai Y, Deng L, Liu H, Yao J, Zhong Z, Fu H, et al. First subtyping of *Blastocystis* sp. from pet rodents in southwestern China. *Int J Parasitol Parasites Wildl.* 2020;11:143–8.
55. Chen XQ, Singh M, Ho LC, Moe KT, Tan SW, Yap EH. A survey of *Blastocystis* sp. in rodents. *Lab Anim Sci.* 1997;47:91–4.
56. AbuOdeh R, Ezzedine S, Madkour M, Stensvold CR, Samie A, Nasrallah G, et al. Molecular subtyping of *Blastocystis* from diverse animals in the United Arab Emirates. *Protist.* 2019;170:125679.
57. Noël C, Dufernez F, Gerbod D, Edgcomb VP, Delgado-Viscogliosi P, Ho LC, et al. Molecular phylogenies of *Blastocystis* isolates from different hosts: implications for genetic diversity, identification of species, and zoonosis. *J Clin Microbiol.* 2005;43:348–55.
58. Deng L, Yao JX, Liu HF, Zhou ZY, Chai YJ, Wang WY, et al. First report of *Blastocystis* in giant pandas, red pandas, and various bird species in Sichuan province, southwestern China. *Int J Parasitol Parasites Wildl.* 2019;9:298–304.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

