

Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Hirji, Zogg, Nguyen Commentary

society at large, and the health care team. 4 Modeling studies from throughout the pandemic have further emphasized that it will take time for hospitals to clear their backlog, depending on their respective operative capacity in the post-COVID period<sup>5</sup> and the potential consequences that waiting could have on patients requiring more urgent operative care. We believe that a stepwise/tiered reopening approach that focuses on minimizing infection risk is prudent and will simultaneously ensure that patients in need of care are not endlessly caught in the crosshairs of surgeons' and hospitals' struggle with how best to answer the difficult question of when is the right time to operate or wait.

#### References

- 1. Nguyen TC. Impact of COVID-19 on adult cardiac surgery: an analysis of the STS ACSD. Presented at: The 57th Annual Meeting of the Society of Thoracic Surgeons, January 29-31, 2021 [virtual].
- 2. Cardiothoracic Interdisciplinary Research Network, COVIDSurg Collaborative. Early outcomes and complications following cardiac surgery in patients testing positive for coronavirus disease 2019: an international cohort study. J Thorac Cardiovasc Surg. 2021;162:e355-72.
- 3. COVIDSurg Collaborative, GlobalSurg Collaborative. Timing of surgery following SARS-CoV-2 infection: an international prospective cohort study. Anaesthesia, 2021:76:748-58.
- 4. Haft JW, Atluri P, Ailawadi G, Engelman DT, Grant MC, Hassan A, et al. Adult cardiac surgery during the COVID-19 pandemic: a tiered patient triage guidance statement. J Thorac Cardiovasc Surg. 2020;160:452-5.
- 5. Salenger R, Etchill EW, Ad N, Matthew T, Alejo D, Whitman G, et al. The surge after the surge: cardiac surgery post-COVID-19. Ann Thorac Surg. 2020;110:

## See Article page e355.


# Commentary: Cardiac surgery in **COVID** patients: Figuring it out as we go

Sarah A. Clark, MD, and Nicholas R. Teman, MD

Since the outbreak of the novel SARS-CoV-2 (COVID-19) viral pandemic in late 2019, the entire world has had to make decisions, ranging from how to treat the profound respiratory complications of COVID-19 pneumonia to how to maintain a globalized economy with significant travel and trade restrictions, based on very limited information. Despite our limited knowledge of this disease, many of those decisions have had an enormous impact on our dayto-day lives, forcing everyone to grapple with uncertainty and changing norms. Although our knowledge of this virus and how to manage it has grown exponentially over the last year, this brief communication by the Cardio-Thoracic



Check for updates



Sarah A. Clark, MD, and Nicholas R. Teman, MD

## CENTRAL MESSAGE

COVID infection in patients undergoing cardiac surgery is associated with significant morbidity and mortality, but more information is needed to guide clinical decision making.

Interdisciplinary Research Network and COVIDSurg Collaboration serves as a reminder of how little we know about COVID-19 and its sequelae.<sup>1</sup>

In this multicenter international observational cohort study, unadjusted outcomes of patients undergoing cardiac surgery with known COVID-19 infection within the perioperative period (7 days preoperatively up to 30 days postoperatively) were examined. Patients who contracted COVID-19 in the perioperative period exhibited higher than normal rates of respiratory complications, including prolonged mechanical ventilation and pneumonia. The authors also report

From the Division of Cardiac Surgery, University of Virginia Health System, Charlot-

Disclosures: The authors reported no conflicts of interest.

The Journal policy requires editors and reviewers to disclose conflicts of interest and to decline handling or reviewing manuscripts for which they may have a conflict of interest. The editors and reviewers of this article have no conflicts of interest.

Received for publication April 19, 2021; revisions received April 19, 2021; accepted for publication April 20, 2021; available ahead of print April 27, 2021.

Address for reprints: Nicholas R. Teman, MD, Division of Cardiac Surgery, University of Virginia Health System, PO Box 800679, 1215 Lee St, Charlottesville, VA 22908 (E-mail: NRT4C@virginia.edu).

J Thorac Cardiovasc Surg 2021;162:e374-5

<sup>0022-5223/\$36.00</sup> 

Copyright © 2021 by The American Association for Thoracic Surgery https://doi.org/10.1016/j.jtcvs.2021.04.058

Clark and Teman Commentary

a 20.8% mortality among patients who were diagnosed with COVID-19 within 30 days of surgery. These findings underscore the incredible clinical impact that COVID-19 infection can have, even among patients undergoing "minor" procedures. These data should give all cardiac surgeons pause, and highlight the importance of vigilance to prevent perioperative COVID-19 infection.

This study has some limitations, which are a reflection of its accelerated nature and a desire for timely results in the face of an oft-changing pandemic. The series includes only patients operated on before July 2020, relatively early in the pandemic, when our methods of diagnosing, treating, and preventing this disease were very much in their infancy. Furthermore, the study lacked a comparison group and risk stratification of the study population. Unfortunately, with this study design, it is unclear whether the patients who contracted COVID-19 were at higher risk for any pulmonary complication. Without a comparison group or risk stratification of the patients, it is also difficult to accurately interpret the high mortality rate, given that 64.7% of these operations were performed emergently, 70.0% of the patients were men, and 62.3% of the patients were over age 60, all of which are established risk factors for mortality.<sup>3-5</sup> The authors did, however, attempt to mitigate some of these confounders by including a subgroup analysis in which they compared characteristics between survivors and nonsurvivors.

Despite its limitations in data and design, this study is a step toward a better understanding of COVID-19 and represents the start of this growing body of literature. The authors attempted to draw logical conclusions from their limited data and use that to make recommendations for improved patient care during this pandemic. We can use studies such as these as stepping stones and keep figuring things out as we go, just as everyone has had to do navigate this post-COVID world.

### References

- The Cardiothoracic Interdisciplinary Research Network and COVIDSurg Collaborative. Early outcomes and complications following cardiac surgery in COVID-19 positive patients: an international cohort study. *J Thorac Cardiovasc Surg.* 2021;162:e355-72.
- Allou N, Bronchard R, Guglielminotti J, Dilly MP, Provenchere S, Lucet JC, et al. Risk factors for postoperative pneumonia after cardiac surgery and development of a preoperative risk score. Crit Care Med. 2014;42:1150-6.
- Hagl C, Khaladj N, Peterss S, Martens A, Kutschka I, Goerler H, et al. Acute treatment of ST-segment elevation myocardial infarction: is there a role for the cardiac surgeon? *Ann Thorac Surg.* 2009;88:1786-92.
- Abramov D, Tamariz MG, Sever JY, Christakis GT, Bhatnagar G, Heenan AL, et al. The influence of gender on the outcome of coronary artery bypass surgery. *Ann Thorac Surg.* 2000;70:800-5.
- Loop FD, Lytle BW, Cosgrove DM, Goormastic M, Taylor PC, Golding LA, et al. Coronary artery bypass graft surgery in the elderly: indications and outcome. Cleve Clin J Med. 1988:55:23-34.