

# The prevalence and distribution of hypodontia in a sample of Qatari patients

Hayder Abdalla Hashim and Sozan Al-Said

## ABSTRACT

**Objective:** The aims of this study were to determine the prevalence of hypodontia in the permanent dentition in a sample of Qatari patients attending a dental center and to compare the results with the reported findings of other populations.

**Materials and Methods:** Orthodontic files including orthopantomographs of 1000 patients (655 females and 345 males, 11–36-year-old) were examined and inspected for evidence of hypodontia.

**Results:** The prevalence of hypodontia in the present Qatari sample was 7.8%; 6.9% was in males and 8.2% in females. Hypodontia was found more frequently in the maxilla than in the mandible. The distribution of missing teeth was noticed in the left side more than the right side. The most frequently missing teeth were the maxillary lateral incisors followed by the mandibular second premolars, maxillary second premolars, and mandibular left lateral incisor. The majority of patients with hypodontia had one or two teeth missing, but rarely more than four teeth were missing in the same patient. Bilateral missing teeth in the current study was commonly seen in the maxillary lateral incisor (14.1%) followed by mandibular second premolar (12.8%) and maxillary second premolar (6.4%).

**Conclusions:** The prevalence of hypodontia in a sample of Qatari individuals was within the range reported in the literature for other populations. The incidence of hypodontia in the anterior segment requires multidisciplinary team approach (orthodontic and prosthodontic) to restore the esthetic and function and improve patient self-esteem.

**Key words:** Hypodontia, prevalence, Qatari

## INTRODUCTION AND REVIEW OF THE LITERATURE

Hypodontia is generally defined as the developmental absence of one or more teeth, excluding the third molars, either in primary or permanent dentition. Researchers have used a variety of terminology to describe the condition, such as a reduction in teeth number, teeth aplasia, congenitally

missing teeth, the absence of teeth, agenesis of teeth, and lack of teeth.<sup>[1-11]</sup> The missing teeth are those which have failed to erupt clinically in the oral cavity and had no sign of appearance in radiographs. The cause is usually disturbance during the early stages of tooth development.<sup>[3,12]</sup> Hypodontia is one of the most common human dental developmental anomalies.<sup>[12-15]</sup>

Many methods of classification have been reported in the literature.<sup>[12-14,16-23]</sup> Some researchers have found the congenital absence of teeth to occur either as an isolated family form or as an inherited form. The inherited form could be autosomal-dominant, autosomal-recessive or an X-linked

Department of Orthodontics, Dental Centre,  
Rumailah Hospital, Doha, Qatar

**Address for correspondence:** Prof. Hayder Abdalla Hashim,  
Dental Centre, Rumailah Hospital, P. O. Box 3050, Doha, Qatar.  
E-mail: hahashim78@yahoo.com

Access this article online	
Quick Response Code:	Website: <a href="http://www.jorthodsci.org">www.jorthodsci.org</a>
	DOI: 10.4103/2278-0203.176651

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

**For reprints contact:** [reprints@medknow.com](mailto:reprints@medknow.com)

**How to cite this article:** Hashim HA, Al-Said S. The prevalence and distribution of hypodontia in a sample of Qatari patients. *J Orthodont Sci* 2016;5:1-6.

trait.<sup>[20]</sup> Others have defined the congenital absence of teeth according to the number of missing teeth.<sup>[17,21,23-25]</sup>

Hypodontia refers to the condition where there is an absence of fewer than six teeth. However, the term oligodontia is usually used to describe a larger number of missing teeth (six or more). Anodontia is the complete absence of teeth.

Dhanrajani<sup>[5]</sup> classified hypodontia according to the severity of the condition following the method of previous researchers.<sup>[13,26]</sup> He used "mild to moderate hypodontia" to denote agenesis of two to five teeth, and referred to the absence of six or more teeth, excluding the third molars, as "severe hypodontia." He defined the term oligodontia as the absence of multiple teeth, usually associated with systemic disorders.<sup>[5]</sup>

Many other researchers have used similar methods of classifying the congenital absence of teeth.<sup>[19,27]</sup> In general, they identified three categories of hypodontia, excluding third molars, as follows: Mild with one or two missing teeth, moderate with 3–5 missing teeth and severe with six or more missing teeth.

Hypodontia is also classified as either isolated hypodontia or syndromic hypodontia. Isolated hypodontia refers to those cases without syndrome.<sup>[28,29]</sup> Thus, hypodontia can occur either as part of a syndrome or as a nonsyndromic, familial form. In the latter form, it occurs as an isolated trait, affects variable numbers of teeth and appears either sporadically or as an inherited condition within a family pedigree.<sup>[23,29]</sup>

The prevalence of agenesis of permanent teeth has been reported in different races and countries.<sup>[30-37]</sup> After third molars, agenesis of lower second premolars and upper lateral incisors is the most common type of hypodontia.<sup>[35]</sup> The reported prevalence of hypodontia in orthodontic patients was different between studies, ranging from 2.7%<sup>[35]</sup> to 11.3%.<sup>[36]</sup> According to Jorgenson,<sup>[3]</sup> the mandibular second premolar is the tooth most frequently absent after the third molar, followed by the maxillary lateral incisor and maxillary second premolar, for Europeans.

The data for hypodontia, excluding the third molars, in both genders combined varies from 0.3% in the Israeli population<sup>[38]</sup> to 11.3% in the Irish<sup>[39]</sup> and 11.3% in Slovenian populations.<sup>[36]</sup> Al-Ajwadi<sup>[40]</sup> reported that the hypodontia in Iraqi is mostly seen in upper lateral incisors and lower second premolar followed by lower central incisors, upper central incisors, and finally the lower lateral incisors.

Recently, Hassan *et al.* reported that the prevalence of hypodontia of in Sudanese sample was 5.1%. Hypodontia was found considerably more frequently in the mandible than in the maxilla. The distribution of missing teeth was noticed in the left side more than in the right side. The most frequently missing teeth were the maxillary lateral incisors, followed by the mandibular second premolars, maxillary second premolars, and mandibular left lateral incisor. The majority of patients

had two or three teeth missing, but rarely more than five teeth missing.<sup>[41]</sup>

More recent, Abdel Jawad *et al.* conducted a study of hypodontia among Qatari patients. They reported that the prevalence of hypodontia was 6.2% (females 8% and males 4.2%;  $P < 0.05$ ). The majority of patients had one or two missing teeth. There were no significant differences between right and left sides for any particular tooth. The most frequently missing teeth were maxillary lateral incisors (36.2%), followed by mandibular second premolar (32.6%) and maxillary second premolar (20.2%). Hypodontia was more commonly found unilaterally than bilaterally (63.2% and 44.3%, respectively).<sup>[42]</sup>

### Aims of the Study

Literature search revealed only one study was done to investigate the prevalence of hypodontia in the permanent dentition in Qatari population.<sup>[43]</sup> Therefore, the aims of this retrospective cross-sectional study were to document the prevalence, distribution of hypodontia in permanent dentition in a sample of Qatari patients selected from a dental center with different age range and to compare the results with the reported findings of other populations.

## MATERIALS AND METHODS

### Sample

The sample consisted of 1000 Qatari patients, of which 345 males and 655 females, who attended the Dental Center at Rumaila Hospital, Doha, Qatar. The patient file (panoramic radiograph, specific periapical radiographs, and anamnestic data) was considered the only source of information used to diagnose hypodontia in this study.

### Criteria of Selection

- Qatari national
- Age range from 11 to 36 year.

### Exclusion Criteria

- Patients with any syndrome and/or cleft lip/palate
- Poor radiographic image quality.

### Methods

The second author evaluated all radiographs on a standard radiographic illuminated viewer. The radiographic findings were checked with patients' records to assure that the missing tooth had not been extracted. All information including age, gender, clinical findings were documented in the file, number, and type of missing tooth, the site of the agenesis (maxilla or mandible, right or left side), being unilateral or bilateral was determined and recorded in a special form. A tooth was identified as congenitally missing when there was no evidence that it had been extracted and no mineralization of the tooth crown could be recognized on the orthopantomogram.<sup>[21]</sup> If an accurate diagnosis of hypodontia could not be made, the file was excluded.

**Statistical Analysis**

To compare and evaluate the difference between male and female patients, maxilla and mandible, and right and left sides in both jaws Student's *t*-test was performed. The level of significance was set at 0.05.

**RESULTS**

A total of 1000 patients' records that fulfilled the criteria of selection were included. Patients age ranged from 10 to 26 years of age, 655 were females with an average age of 14.5-year-old, and 345 were males with an average age of 16.4-year-old.

Out of 1000 patient's records reviewed, 78 exhibited congenital absence of one or more teeth (24 male and 54 female). The prevalence of hypodontia in the studied Qatari population sample was 7.8% in which 6.9% was in male and 8.2% in female [Figure 1].

Figure 2 shows the distribution of missing one tooth as it was found in 11 (3.2%) males and 24 females (3.6%) patients, two missing teeth were noticed in 10 (2.9%) males and 18 (2.7%) females, three missing teeth were observed in 1 (0.3%) male and 6 (0.9%) in females and in patient with four teeth


congenitally missing teeth were exhibited in 2 (0.6%) males and in 6 (0.9%) females [Figure 2]. This indicates that females were affected more than males, but no significant difference was observed ( $P > 0.05$ ).

Figures 3 and 4 demonstrate the distribution of hypodontia in upper and lower jaw. The upper jaw revealed a higher percentage of hypodontia than the lower jaw. On the other hand, the left side in both jaws showed a higher percentage of missing teeth compared to the right side, but no significant difference was noted ( $P > 0.05$ ).


Figure 5 exhibits the distribution of hypodontia in the right and the left side versus bilateral in upper and lower jaw. The bilateral hypodontia revealed a higher percentage for maxillary lateral incisor, maxillary second premolar, and mandibular second premolar; whereas, it was equal for maxillary first premolar and mandibular central incisor. On the other hand, the unilateral hypodontia of mandibular lateral incisor showed a higher percentage compared to the bilateral.

**DISCUSSION**


This is a descriptive cross-sectional study to determine the prevalence of hypodontia in a sample of Qatari patients.




**Figure 1:** Number of patients with hypodontia and their distribution according to gender ( $n = 78$ )




**Figure 2:** Congenitally missing teeth distribution among males and females patients with hypodontia



**Figure 3:** Distribution of congenitally missing teeth in the upper jaw



**Figure 4:** Distribution of congenitally missing teeth in the lower jaw



**Figure 5:** Distribution of hypodontia in the right, left side and bilateral

The etiology of hypodontia is usually disturbance during the early stages of tooth development.<sup>[3,12]</sup> Crown calcification starts at the age of 3 years and usually completes at the age of 6 years.<sup>[44-47]</sup> However, there is variation in the development of some teeth (e.g., premolars).<sup>[48]</sup> Therefore, it is very difficult to decide whether the tooth is missing or not before the age of 9 years, especially among males.<sup>[49]</sup> This was confirmed in an investigation conducted by Wisth *et al.*<sup>[50]</sup> They observed that the prevalence of missing teeth was higher at the age of 7 years (7.1%) compared when the same sample was re-examined at the age of 9 years (6.6%). This is the reason why in the present study, patients above 10 years of age were selected.

The prevalence of hypodontia in the primary dentition was found to be very low. The range has generally been between 0.1% and 0.9% of the population.<sup>[2]</sup> However, in the permanent dentition a wide range of prevalence values (4.6–12.6%) was reported. The results of this study revealed that hypodontia prevalence in the permanent dentition is 7.8% in Qatari sample, which falls within the range reported in the literature. Further, this finding is consistent with other reported prevalence values among Danish (7.8%) and Iceland (7.9%) populations.<sup>[51,52]</sup> However, slightly smaller value was observed among the Turkish population (7.5%).<sup>[53]</sup> Whereas, in Sudanese and Qatari samples, the prevalence was less (5.1% and 6.2%, respectively).<sup>[41,42]</sup> On the other hand, higher prevalence values were reported in two German studies (12.6%)<sup>[54]</sup> and (11.3%)<sup>[36]</sup> and very low prevalence value was noticed in French (1.9%)<sup>[55]</sup> and Malaysian studies (2.8%).<sup>[56]</sup>

Gender dimorphism was investigated and the result indicated that the hypodontia was higher in females than males. This finding is in agreement with several previous studies conducted in different populations.<sup>[52,57-59]</sup> However, Yildray *et al.*<sup>[54]</sup> stated “although others reported a higher incidence in females than in males, even though we determined significant differences for some teeth.” Further, the result of the present investigation revealed that females were dominant in cases with one tooth, two teeth, three teeth, and more than four missing teeth.

The comparison of the congenitally missing teeth between the right and left sides of the current study showed that the left side in the upper jaw was more affected than the right side. However, this was not observed in the lower jaw. This finding was in disagreement with the finding of Fekonja who reported that the missing teeth were more commonly absent on the right side than on the left side.<sup>[36]</sup>

Many studies have demonstrated that there is no consistent finding as to which jaw has more missing teeth.<sup>[31,35,40,51,52,61]</sup> This study supports the finding that more teeth were missing in the maxilla than in the mandible. However, other investigators reported more absence in the mandible.<sup>[7,42,60,62]</sup>

Most investigators observed the predominance of bilateral congenitally missing teeth to the extent as being as twice as unilateral missing.<sup>[31,63]</sup> However, this was not in line with the finding observed in this study. Bilateral missing teeth in the current study was commonly seen in the maxillary lateral incisor (14.1%) followed by mandibular second premolar (12.8%) and maxillary second premolar (6.4%). This finding was consistent with study carried out in Qatari orthodontic and pediatric patients.<sup>[43]</sup>

Further, the maxillary lateral incisor was the most frequently missing tooth, and the mandibular second premolar was in the second rank followed by the maxillary second premolar and mandibular lateral incisor. This coincides with the result conducted in Turkish population sample,<sup>[54]</sup> and in Qatari orthodontic patients.<sup>[43]</sup> On the other hand, others have reported the difference in the sequence of most frequently affected teeth. They reported that the most frequent missing tooth was the mandibular second premolar, followed by the maxillary second premolar, the maxillary lateral incisor, and the mandibular central incisor.<sup>[50,52,53]</sup>

Furthermore, the result of this study revealed that the agenesis of maxillary and mandibular canines and second molars were very rare. Furthermore, the same result was consistent with other reported previous investigations.<sup>[35,55,62,63]</sup>

Hence, many studies were carried out to evaluate the prevalence of hypodontia among different populations.<sup>[30,32]</sup> A wide range of prevalence values (4.6–12.6%) was reported. This wide range could be attributed to geographic, gender, racial and genetic differences and also to the large differences in the sample sizes and the criteria of selection. All these difference might play an important role in these various reported results of hypodontia<sup>[31,32,36,54]</sup> and making the comparison of this study result very limited with other previous studies. The above-mentioned factors may justify the difference of the prevalence of hypodontia in Qatari sample (7.8%) in the present study and the previous study among Qatari orthodontic and pediatric patients collected from a different place (6.2%).<sup>[42]</sup>

Most patients seek orthodontic treatment to restore their facial esthetic more than the function. However, esthetic problems, periodontal damage, malocclusion, and alveolar bone deficiency are some of the complications following hypodontia. Some studies stated that also anterior hypodontia has a significant effect on skeletal relationships.<sup>[44]</sup> However, each of these problems can be an indication for orthodontic treatment. For example, agenesis of maxillary lateral incisor impairs dental esthetics and function from a very young age.<sup>[37]</sup>

Missing permanent teeth in the anterior or posterior region represents a clinical problem and requires a multidisciplinary approach to cover the needs of the patients. The end result of this approach is restoring the esthetic and function and improving the self-esteem of the patient. Therefore, by early detection of missing teeth, proper diagnosis and treatment plan can be performed with a multidisciplinary team approach.

## CONCLUSIONS

- The prevalence of hypodontia in this study (7.8%) was within the range reported in the literature.
- The maxillary lateral incisor was the most frequently missing tooth, and the mandibular second premolar was in the second rank followed by the maxillary second premolar and mandibular lateral incisor.
- Hypodontia was found more frequently in the maxilla than in the mandible, and the distribution of missing teeth was noticed in the left side more than in the right side.
- Most common bilaterally missing tooth was maxillary lateral incisor (9.7%), followed by mandibular second premolar (6.9%) and maxillary second premolar (3.5%).

## Acknowledgment

We would like to thank Hamad Medical Corporation/ Hamad Medical Research Center for sponsoring this research (Sub-Research Proposal Number 15041/150) and Dr. Hashim AL Hussain for his help and support during the preparation of this manuscript.

## Financial Support and Sponsorship

Nil.

## Conflicts of Interest

There are no conflicts of interest.

## REFERENCES

- Hunstadbraten K. Hypodontia in the permanent dentition. *ASDC J Dent Child* 1973;40:115-7.
- Brook AH. Dental anomalies of number, form and size: their prevalence in British schoolchildren. *J Int Assoc Dent Child* 1974;5:37-53.
- Jorgenson RJ. Clinician's view of hypodontia. *J Am Dent Assoc* 1980;101:283-6.
- Zhu JF, Marcusamer M, King DL, Henry RJ. Supernumerary and congenitally absent teeth: A literature review. *J Clin Pediatr Dent* 1996;20:87-95.
- Dhanrajani PJ. Hypodontia: Etiology, clinical features, and management. *Quintessence Int* 2002;33:294-302.
- McKeown HF, Robinson DL, Elcock C, al-Sharood M, Brook AH. Tooth dimensions in hypodontia patients, their unaffected relatives and a control group measured by a new image analysis system. *Eur J Orthod* 2002;24:131-41.
- Kirzioglu Z, Köselcer Sentut T, Ozay Ertürk MS, Karayilmaz H. Clinical features of hypodontia and associated dental anomalies: A retrospective study. *Oral Dis* 2005;11:399-404.
- Wu CC, Wong RW, Hagg U. A review of hypodontia: The possible etiologies and orthodontic, surgical and restorative treatment options conventional and futuristic. *Hong Kong Dent J* 2007;4:113-21.
- Swinnen S, Bailleul-Forestier I, Arte S, Nieminen P, Devriendt K, Carels C. Investigating the etiology of multiple tooth agenesis in three sisters with severe oligodontia. *Orthod Craniofac Res* 2008;11:24-31.
- Brook AH, Elcock C, Aggarwal M, Lath DL, Russell JM, Patel PI, *et al.* Tooth dimensions in hypodontia with a known PAX9 mutation. *Arch Oral Biol* 2009;54 Suppl 1:S57-62.
- Brook AH, Griffin RC, Smith RN, Townsend GC, Kaur G, Davis GR, *et al.* Tooth size patterns in patients with hypodontia and supernumerary teeth. *Arch Oral Biol* 2009;54 Suppl 1:S63-70.
- Pemberton TJ, Das P, Patel PI. Hypodontia: Genetics and future perspectives. *Braz J Oral Sci* 2005;4:695-706.
- Hobkirk JA, Brook AH. The management of patients with severe hypodontia. *J Oral Rehabil* 1980;7:289-98.
- Vastardis H. The genetics of human tooth agenesis: New discoveries for understanding dental anomalies. *Am J Orthod Dentofacial Orthop* 2000;117:650-6.
- De Coster PJ, Marks LA, Martens LC, Huyseune A. Dental agenesis: Genetic and clinical perspectives. *J Oral Pathol Med* 2009;38:1-17.
- Schalk-van der Weide Y, Steen WH, Bosman F. Distribution of missing teeth and tooth morphology in patients with oligodontia. *ASDC J Dent Child* 1992;59:133-40.
- Goodman JR, Jones SP, Hobkirk JA, King PA. Hypodontia: 1. Clinical features and the management of mild to moderate hypodontia. *Dent Update* 1994;21:381-4.
- Schalk-van der Weide Y, Bosman F. Tooth size in relatives of individuals with oligodontia. *Arch Oral Biol* 1996;41:469-72.
- Brook AH, Elcock C, al-Sharood MH, McKeown HF, Khalaf K, Smith RN. Further studies of a model for the etiology of anomalies of tooth number and size in humans. *Connect Tissue Res* 2002;43:289-95.
- Mostowska A, Kobiela A, Biedziak B, Trzeciak WH. Novel mutation in the paired box sequence of PAX9 gene in a sporadic form of oligodontia. *Eur J Oral Sci* 2003;111:272-6.
- Nunn JH, Carter NE, Gillgrass TJ, Hobson RS, Jepson NJ, Meechan JG, *et al.* The interdisciplinary management of hypodontia: Background and role of paediatric dentistry. *Br Dent J* 2003;194:245-51.
- Polder BJ, Van't Hof MA, Van der Linden FP, Kuijpers-Jagtman AM. A meta-analysis of the prevalence of dental agenesis of permanent teeth. *Community Dent Oral Epidemiol* 2004;32:217-26.
- Cobourne MT. Familial human hypodontia – Is it all in the genes? *Br Dent J* 2007;203:203-8.
- Burzynski NJ, Escobar VH. Classification and genetics of numeric anomalies of dentition. *Birth Defects Orig Artic Ser* 1983;19:95-106.
- Schalk van der Weide Y, Prah-Andersen B, Bosman F. Tooth formation in patients with oligodontia. *Angle Orthod* 1993;63:31-7.
- Brook AH. A unifying aetiological explanation for anomalies of human tooth number and size. *Arch Oral Biol* 1984;29:373-8.
- Hobkirk JA, Gill DS, Jones SP, Hemmings KW, Bassi GS, O'Donnell AL, *et al.* Hypodontia: A Team Approach to Management. 1<sup>st</sup> ed. Chichester: Wiley-Blackwell; 2011.
- Arte S, Nieminen P, Apajalahti S, Haavikko K, Thesleff I, Pirinen S. Characteristics of incisor-premolar hypodontia in families. *J Dent Res* 2001;80:1445-50.
- Tan SP, van Wijk AJ, Prah-Andersen B. Severe hypodontia: Identifying patterns of human tooth agenesis. *Eur J Orthod* 2011;33:150-4.
- Gupta SK, Saxena P, Jain S, Jain D. Prevalence and distribution of selected developmental dental anomalies in an Indian population. *J Oral Sci* 2011;53:231-8.
- Endo T, Ozoe R, Kubota M, Akiyama M, Shimooka S. A survey of

- hypodontia in Japanese orthodontic patients. *Am J Orthod Dentofacial Orthop* 2006;129:29-35.
32. Chung CJ, Han JH, Kim KH. The pattern and prevalence of hypodontia in Koreans. *Oral Dis* 2008;14:620-5.
  33. González-Allo A, Campoy MD, Moreira J, Ustrell J, Pinho T. Tooth agenesis in a Portuguese population. *Int Orthod* 2012;10:198-210.
  34. Medina AC. Radiographic study of prevalence and distribution of hypodontia in a pediatric orthodontic population in Venezuela. *Pediatr Dent* 2012;34:113-6.
  35. Silva Meza R. Radiographic assessment of congenitally missing teeth in orthodontic patients. *Int J Paediatr Dent* 2003;13:112-6.
  36. Fekonja A. Hypodontia in orthodontically treated children. *Eur J Orthod* 2005;27:457-60.
  37. Kavadia S, Papadiochou S, Papadiochos I, Zafiriadis L. Agenesis of maxillary lateral incisors: A global overview of the clinical problem. *Orthodontics (Chic)* 2011;12:296-317.
  38. Rosenzweig KA, Garbarski D. Numerical aberrations in the permanent teeth of grade school children in Jerusalem. *Am J Phys Anthropol* 1965;23:277-83.
  39. O'Dowling IB, McNamara TG. Congenital absence of permanent teeth among Irish school-children. *J Ir Dent Assoc* 1990;36:136-8.
  40. Al-Ajwadi SA. An orthopantomographic study of hypodontia in permanent teeth of Iraqi patients. *Med Dent J* 2009;6:139-44.
  41. Hassan DA, Abuaffan AH, Hashim HA. Prevalence of hypodontia in a sample of Sudanese orthodontic patients. *J Orthod Sci* 2014;3:63-7.
  42. Abed Al Jawad FH, Al Yafei H, Al Sheeb M, Al Emadi B, Al Hashimi N. Hypodontia prevalence and distribution pattern in a group of Qatari orthodontic and pediatric patients: A retrospective study. *Eur J Dent* 2015;9:267-71.
  43. Bergendal B. Oligodontia ectodermal dysplasia – On signs, symptoms, genetics, and outcomes of dental treatment. *Swed Dent J Suppl* 2010;205:13-78, 7-8.
  44. Mostowska A, Biedziak B, Jagodzinski PP. Novel MSX1 mutation in a family with autosomal-dominant hypodontia of second premolars and third molars. *Arch Oral Biol* 2012;57:790-5.
  45. Zhu J, Yang X, Zhang C, Ge L, Zheng S. A novel nonsense mutation in PAX9 is associated with sporadic hypodontia. *Mutagenesis* 2012;27:313-7.
  46. Acharya PN, Jones SP, Moles D, Gill D, Hunt NP. A cephalometric study to investigate the skeletal relationships in patients with increasing severity of hypodontia. *Angle Orthod* 2010;80:511-8.
  47. Nordgarden H, Jensen JL, Storhaug K. Reported prevalence of congenitally missing teeth in two Norwegian counties. *Community Dent Health* 2002;19:258-61.
  48. Aasheim B, Ogaard B. Hypodontia in 9-year-old Norwegians related to need of orthodontic treatment. *Scand J Dent Res* 1993;101:257-60.
  49. Ng'ang'a RN, Ng'ang'a PM. Hypodontia of permanent teeth in a Kenyan population. *East Afr Med J* 2001;78:200-3.
  50. Wisth PJ, Thunold K, Bøe OE. Frequency of hypodontia in relation to tooth size and dental arch width. *Acta Odontol Scand* 1974;32:201-6.
  51. Rølling S. Hypodontia of permanent teeth in Danish schoolchildren. *Scand J Dent Res* 1980;88:365-9.
  52. Magnússon TE. An epidemiologic study of dental space anomalies in Icelandic schoolchildren. *Community Dent Oral Epidemiol* 1977;5:292-300.
  53. Sisman Y, Uysal T, Gelgor IE. Hypodontia. Does the prevalence and distribution pattern differ in orthodontic patients? *Eur J Dent* 2007;1:167-73.
  54. Behr M, Proff P, Leitzmann M, Pretzel M, Handel G, Schmalz G, *et al.* Survey of congenitally missing teeth in orthodontic patients in Eastern Bavaria. *Eur J Orthod* 2011;33:32-6.
  55. Nik-Hussein NN. Hypodontia in the permanent dentition: A study of its prevalence in Malaysian children. *Aust Orthod J* 1989;11:93-5.
  56. Davis PJ. Hypodontia and hyperdontia of permanent teeth in Hong Kong schoolchildren. *Community Dent Oral Epidemiol* 1987;15:218-20.
  57. Altug-Atac AT, Erdem D. Prevalence and distribution of dental anomalies in orthodontic patients. *Am J Orthod Dentofacial Orthop* 2007;131:510-4.
  58. Bäckman B, Wahlin YB. Variations in number and morphology of permanent teeth in 7-year-old Swedish children. *Int J Paediatr Dent* 2001;11:11-7.
  59. Dolder E. Deficient dentition. Statistical survey. *Dent Rec* 1937;57:142-3.
  60. Silverman NE, Ackerman JL. Oligodontia: A study of its prevalence and variation in 4032 children. *ASDC J Dent Child* 1979;46:470-7.
  61. Amini F, Rakhshan V, Babaei P. Prevalence and pattern of hypodontia in the permanent dentition of 3374 Iranian orthodontic patients. *Dent Res J (Isfahan)* 2012;9:245-50.
  62. Bhardwaj VK, Veerasha KL, Sharma KR. Prevalence of malocclusion and orthodontic treatment needs among 16 and 17 year-old school-going children in Shimla city, Himachal Pradesh. *Indian J Dent Res* 2011;22:556-60.
  63. Ajami BA, Shabzendedar M, Mehrjerdian M. Prevalence of hypodontia in nine- to fourteen-year-old children who attended the Mashhad School of Dentistry. *Indian J Dent Res* 2010;21:549-51.

## Author Help: Online submission of the manuscripts

Articles can be submitted online from <http://www.journalonweb.com>. For online submission, the articles should be prepared in two files (first page file and article file). Images should be submitted separately.

### 1) First Page File:

Prepare the title page, covering letter, acknowledgement etc. using a word processor program. All information related to your identity should be included here. Use text/rtf/doc/pdf files. Do not zip the files.

### 2) Article File:

The main text of the article, beginning with the Abstract to References (including tables) should be in this file. Do not include any information (such as acknowledgement, your names in page headers etc.) in this file. Use text/rtf/doc/pdf files. Do not zip the files. Limit the file size to 1 MB. Do not incorporate images in the file. If file size is large, graphs can be submitted separately as images, without their being incorporated in the article file. This will reduce the size of the file.

### 3) Images:

Submit good quality color images. Each image should be less than 4096 kb (4 MB) in size. The size of the image can be reduced by decreasing the actual height and width of the images (keep up to about 6 inches and up to about 1800 x 1200 pixels). JPEG is the most suitable file format. The image quality should be good enough to judge the scientific value of the image. For the purpose of printing, always retain a good quality, high resolution image. This high resolution image should be sent to the editorial office at the time of sending a revised article.

### 4) Legends:

Legends for the figures/images should be included at the end of the article file.